**Analysis of a rare variant of mitotic gene *TAO3* reveals its meiotic**

**interactors**

Saumya Gupta[*], Aparna Radhakrishnan[*], Rachana Nitin[*], Pandu Raharja-Liu[†], Gen

Lin[‡], Lars M. Steinmetz[‡,§,**], Julien Gagneur[†], Himanshu Sinha[*,1]

[*]Department of Biological Sciences, Tata Institute of Fundamental Research, Mumbai

400005, India

[†]Gene Center, Ludwig-Maximilians-Universität, 81377 Munich, Germany

[‡]European Molecular Biology Laboratory, Genome Biology Unit, 69117 Heidelberg,

Germany

[§]Department of Genetics, Stanford University School of Medicine, Stanford,

California 94305

[**]Stanford Genome Technology Center, Stanford University, Palo Alto, California

94304

[1]Corresponding author: Department of Biological Sciences, Tata Institute of

Fundamental Research, Homi Bhabha Road, Colaba, Mumbai 400005, India. Email:

hsinha@tifr.res.in

Running title:

Analyzing functional role of a rare variant

Keywords:

Rare variant, *TAO3*, meiosis, transcriptome profiling, allelic variant

## ABSTRACT

Genome-wide association studies have successfully identified thousands of common variants associated with complex traits and diseases. If only common variants are considered, a significant proportion of heritability in common diseases remains unexplained. One of the sources considered to explain this missing heritability are the rare genetic variants. Studying the consequences of rare genetic alterations offers additional opportunity for predicting molecular mediators underlying pathway deregulation. Here, we characterized the functional role of a rare variant having a significant contribution in sporulation efficiency variation. This causal variant is present in the coding sequence of *TAO3*, encoding a putative scaffolding protein conserved from yeast to humans and component of RAM network in yeast involved in mitosis. We observed that the role of *TAO3* allele on meiosis is independent of *ACE2*, a transcription factor regulated by RAM network during mitosis. By expressing *TAO3* causal allele conditionally during sporulation and quantitatively measuring cell cycle progression, we determined its role within the first 6 hours in meiosis, which coincides with cells entering into meiotic cell division. Time-resolved genome-wide gene expression analysis identified genes showing early and increasing expression trend during sporulation to be target genes of *UME6*, a crucial meiotic regulator. Genes regulating *UME6*, and other target genes expressed early, specifically, in the presence of causal *TAO3*, were chosen as candidate genes. Allele-specific functional validations identified *ERT1* (regulator of switch from fermentation to respiration) and *PIP2*, (regulator involved in beta-oxidation of fatty acids) as the mediating genes associated with *TAO3* causal allele and responsible for sporulation efficiency variation. Our study uncovered interesting link between *TAO3* and regulators of metabolic cues that modulate the switch between multiple developmental phenotypes, *viz.* mitosis to meiosis. Although a small proportion in a population might contain rare variants, identification of novel regulators of sporulation from our study highlighted the significance of studying rare genetic variants to obtain novel insights into the phenotype and disease biology.

61                                    **INTRODUCTION**

The genetic architecture of complex traits is not completely understood. The

'common disease, common variant' rationale of genome-wide association studies

(GWAS) is being challenged owing to the limited proportion of disease heritability

explained by common variants, giving rise to the 'missing heritability' problem in

quantitative genetics (Manolio *et al.* 2009; Zuk *et al.* 2014). Not considering the effect

of rare variants has been suggested as one of the potential contributor of this 'hidden'

instead of 'missing' heritability (Saint Pierre and Génin 2014). This view has been

substantiated by the identification of multiple rare variants that confer considerable

risk in diseases such as autism, schizophrenia and epilepsy (Stankiewicz and Lupski

2010). Since the initial genomic technologies were more adept at identifying common

variants, sequencing methods and rare variant association studies are now being

increasingly used to identify rare variants (Cirulli and Goldstein 2010; Zuk *et al.*

2014). Once identified, characterizing the functional role of rare variants associated

with complex diseases has the potential for revealing new biology.

Common or rare, if genetic variants are in the core regulatory pathways underlying

the trait, understanding the biological mechanism of phenotypic modulation is more

direct. However, many genetic variants are identified in the ancillary biological

processes. These processes are not part of the core regulatory pathways underlying the

trait, but could impinge on the phenotype by being associated with the core pathways.

For instance, gene identified by GWAS for face shape variation is a gap junction

protein, which indirectly affects the mechanical load taken up by face bones (Liu *et*

*al.* 2012). Ancillary pathways, hence, provide a good hub for accumulation of genetic

variation. Their location in the regulatory network confers an advantage in regulating

the phenotype while the core pathway genes could be too important to be varied, and

hence probably under a negative selection pressure. This could be one of the reasons

why crucial face forming gene of *Shh* has not been identified for facial shape

development (Hallgrimsson *et al.* 2014). Therefore, characterization of variants, not

part of the core biological processes associated with the phenotype is challenging.

Here, we characterized the functional role of a rare variant in the novel mitotic gene,

*TAO3*, in yeast sporulation efficiency variation. We studied genome-wide gene

expression dynamics in presence of both the alleles (common and causative rare

alleles) of *TAO3 (G4477C)* during sporulation, and identified the core meiotic

pathways and the ancillary pathways associated to sporulation showing differential

expression. Our approach showed regulatory pathways involved in nutrient

metabolism impinging on the core sporulation pathway, contributing to trait variation.

**MATERIALS AND METHODS**

*Yeast strains and media*

The yeast strains were grown in standard conditions at 30˚C in YPD (1% Yeast

extract, 2% Bacto peptone, 2% dextrose). Whole-genome resequencing of the strain

YAD331 (Deutschbauer and Davis 2005) with S288c strain as the reference strain,

identified two additional polymorphisms (Supporting Figure S1, Supporting Table

S1). Three consecutive backcrosses were performed between the haploid obtained

from this strain and the haploid reference strain, to remove the secondary

polymorphisms. The sole genetic difference between the reference S288c strain and

the backcrossed allele replacement strain was at *TAO3(G4477C)* position, which was

confirmed by performing PCR-based sequencing 650bp up and downstream around

the two secondary polymorphisms and the *TAO3* polymorphism region. This

backcrossed strain was diplodized to make it homozygous at *TAO3(4477C)* position

and was termed as "T strain" in this study. The diploid parental strain S288c was

termed as "S strain" in the study. All gene deletions in the study were made in the

haploids of T and S strains, except the ones made in SK1 strain (Supporting Table

S2). Deletions were performed and verified as described previously (Goldstein and

McCusker 1999; Gietz and Woods 2002). All the experiments in this study were

performed using the diplodized parent strains and their diploid derivatives. For

replacing the endogenous *TAO3* promoter (-150 to -1bp upstream start site) in T

strain, with a tetracycline-responsive promoter, a *tetO7*-based promoter substitution

cassette containing *kanMX4* was amplified from the plasmid pCM225 (Bellí *et al.*

1998b). Diploid T strain with this *tetO7*-based cassette is termed $P_{Tet}$-*TAO3(4477C)*

strain. The primers for sequencing, deletions and their confirmations are listed in

Supporting Table S3.

*Phenotyping*

Sporulation efficiency estimation at 48h, progression through meiotic landmark

events Meiosis I (MI) and Meiosis II (MII) and its quantitation was done as described

previously (Gupta *et al.* 2015). For quantitation of meiotic landmarks in T strain,

parametric curves assuming delayed and 1st order kinetics were fitted to the DAPI-

stained meiotic progression time course data and fitting uncertainties were estimated

by bootstrapping (Supporting File S1). Cell cycle progression data for S288c and SK1

strains was taken from Gupta *et al.* (2015) (Figure 1D-E). Conditional expression of

*TAO3(4477C)* was performed by constructing $P_{Tet}$-*TAO3* strain (details in Supporting

File S1), which was responsive to tetracycline-analogue doxycycline (Bellí *et al.*

1998a; b). Addition of 2μg/ml doxycycline was utilized to decrease the activity of the

gene, unless any other concentration was stated. For each strain, minimum three

biological replicates were used and the experiment was carried minimum two times.

Approximately 300 cells were counted per replicate. Fold difference was calculated as

the ratio of mean sporulation efficiencies of the two strains A and B when the

sporulation efficiency of A is greater than of B.

*Statistical test for calculating sporulation efficiency*

For comparing sporulation efficiency, two statistical tests were used: the pair test and

the interaction test. The pair test tests the null hypothesis that two given strains (S and

T) have the same sporulation efficiency.

The number $y_{i,k}$ of sporulated cells (4-nuclei count) among the total number of cells

$n_{i,k}$ of strain $i$ in replicate experiment $k$ was modeled with a quasi-binomial

generalized linear model using the *logit* link function and subject to a common log-

odd ratio $\beta_i$ between replicates, *i.e.*:

$$\log\left(\frac{\mu_{i,k}}{n_{i,k} - \mu_{i,k}}\right) = \beta_i \text{ for all } k,$$

where, $\mu_{i,k} = E(y_{i,k})$.

The pair test tests the null hypothesis of equality of log odd-ratios for two strains $i$ and

$j$, i.e. $H_0 : \beta_i = \beta_j$.

In the case of S and T strains, the interaction test tests the null hypothesis that the

effect of mutation A is independent of the effect of mutation B, taking the T strain as

reference background. This test, thus, compares four strains: mutation A only,

mutation B only, both A and B and neither A nor B (T strain). Here, the strain S was

considered as a T strain mutated for *TAO3(4477)*. For every interaction test, we

considered the dataset of the four strains of interest and fitted a quasi-binomial

generalized linear model using the *logit* link function and subject to:

168
$$\log\left(\frac{\mu_{i,k}}{n_{i,k} - \mu_{i,k}}\right) = \beta_0 + \beta_A A_i + \beta_B B_i + \beta_{A,B} A_i B_i \text{ for all } k,$$

where, $A_i$ and $B_i$ are indicator variables of the mutations A and B in strain $i$

respectively. The interaction test tested the null hypothesis that the odd ratio of

sporulation in the double mutant equals the product of the odd ratios of each mutation,

i.e. $H_0 : \beta_{A,B} = 0$.

Both the pair test and the interaction test were implemented in the statistical language

R with the function *glm()* assuming a constant variance function fitted by maximizing

the quasi-likelihood and using the t-test on tested parameters (Gupta *et al.* 2015).

***Whole genome gene-expression profiling***

Sporulating yeast cell collection at 0h, 30m, 45m, 1h10m, 1h40m, 2h30m, 3h50m,

5h40m and 8h30m (logarithmic time-series), RNA isolation and cDNA preparation

were performed as described (Xu *et al.* 2009). Samples were hybridized to *S.*

*cerevisiae* yeast tiling array (Affymetrix, Cat# 520055). Arrays at each time point for

both the strains were normalized together using *vsn* normalization method (Huber *et*

*al.* 2002).

***Whole genome gene-expression analysis***

Within each strain, the $\log_2$ expression values obtained were smoothed using *locfit* at

optimized bandwidth parameter $h = 1.2$ (Supporting Figure S2), base transformed for

each transcript by subtracting the expression value at each time point from the

baseline value at time point t = 0h ($t_0$) (Supporting Table S4. This $\log_2$ fold change

value with respect to $t_0$ is described as "expression" throughout the manuscript. For

identifying the genes showing temporal differential expression between T and S

strains (Supporting Table S5, method implemented in EDGE software was used,

which calculated statistically significant changes in expression between T and S strain

over time (Storey *et al.* 2005). The differentially expressed genes were clustered

according to their temporal expression patterns using time abstraction clustering

algorithm implemented in the TimeClust software (Magni *et al.* 2008) (see Supporting

File S1). Four major clusters were identified in each strain: Cluster I (Early trend),

Cluster II (Increasing trend), Cluster III (Late trend), Cluster IV (Repressing trend)

(Supporting Table S6). The transcription factors regulating a cluster of genes were

extracted using the YEASTRACT database (Teixeira *et al.* 2014). Only those

transcription factors were considered as candidate genes whose target genes were

significantly enriched in the corresponding cluster ($P \leq 0.05$, odds ratio $\geq 1.5$).

YEASTRACT database was also used to obtain the regulation matrix of yeast, for

identifying target genes of regulators in this study such as *UME6*. Target genes for

*ACE2* were obtained from Nelson *et al.* (2003). Significantly enriched Gene Ontology

terms by biological process (Bonferroni corrected $P < 0.05$, Table 1) were obtained

from SGD Yeastmine (Balakrishnan *et al.* 2012).

***Data availability***

The array data for T strain reported has been deposited in ArrayExpress

(http://www.ebi.ac.uk/arrayexpress/) with accession number E-MTAB-3889. The

whole genome sequence data for T strain has been deposited in the European

Nucleotide Archive (http://www.ebi.ac.uk/ena/) with the accession number

PRJEB8698. The rest of the data is available as Supporting Information. Array data

and whole genome sequence data for S strain were downloaded from Gupta *et al.*

(2015).

                              **RESULTS**

*Role of TAO3 causative allele in sporulation efficiency variation*

The causative allele in *TAO3(4477C)*, identified in high sporulating SK1 strain

(Deutschbauer and Davis 2005), was not present in other high sporulating natural and

laboratory yeast strains in the SGRP collection (Figure 1A). This indicated that

*TAO3(4477C)* was a rare variant for high sporulation efficiency. Substitution of this

causative single nucleotide polymorphism *(G4477C)* in 7,131bp long *TAO3*, in the

low sporulating S288c strain, was sufficient to increase its sporulation efficiency by

three-fold ($P = 1.8 \times 10^{-10}$, pair test in Methods, Figure 1B) compared to the S288c

strain (*TAO3(4477G)*), reconfirming a significant role of *TAO3(4477C)* SNP in

sporulation efficiency variation (Deutschbauer and Davis 2005). The increased

sporulation efficiency in T strain saturated within 48h and did not increase beyond

three-folds even when kept in sporulation medium for a week (Figure 1C).

Quantitative analysis of cell cycle progression in T strain predicted that the difference

between T and S strain occurred already during their entry into meiosis, since both,

the time to initiate meiosis and rate of transition from $G_1/G_0$ into MI, were

significantly different between the two strains (Figures 1D-E, Supporting Figure S3).

Studying the progression of meiotic phases also showed that the T strain initiated

meiosis within 12h (Figures 1D-E). To further resolve this 12h temporal phase when

*TAO3(4477C)* could have an effect on the phenotype, this allele was conditionally

expressed during sporulation (see Methods). The sporulation efficiency of the

tetracycline-responsive T strain ($P_{Tet}$-*TAO3(4477C)* strain) in the presence of 2μg/ml

doxycycline throughout the sporulation period was equivalent to the S strain (Figure

1F). Shorter periods of inactivation of *TAO3* activity (in presence of doxycycline)

showed that a 3h inhibition indeed reduced sporulation efficiency but this was

significantly different ($P = 0.02$) from 48h inhibition, although, a 6h inhibition was

equivalent to 48h inhibition. This showed that *TAO3(4477C)* allele affected

sporulation efficiency within the first 6h in sporulation.

*Role of TAO3 in meiosis is distinct from its role during mitosis*

A part of the RAM (Regulation of Ace2p activity and cellular Morphogenesis)

signaling network, Tao3 is required for activation and localization of an NDR protein

kinase, Cbk1, another essential component of RAM network (Du and Novick 2002;

Hergovich *et al.* 2006). This signaling network (consisting of Cbk1, Hym1, Kic1,

Mob2, Tao3) contributes to various important aspects of mitotic cellular growth,
particularly in cell separation by regulating transcription factor Ace2, in polarized
growth (Nelson *et al.* 2003) and in cellular progression, through a Ace2-independent
pathway (Bogomolnaya *et al.* 2006). Ace2 is a transcription factor that peaks early in
mitosis and is involved in $G_1$/S transition (Spellman *et al.* 1998). In S288c-
background strains, the RAM network genes are essential and hence the affect of their
deletion in sporulation efficiency has not been previously studied (Deutschbauer *et al.*
2002). Here, we tested if this upstream mitotic function of RAM network was also
involved in meiotic regulation.

To determine the unique molecular effects of the causative *TAO3* allele during
sporulation, we compared genome-wide expression between T and S strains (see
Methods). Genome-wide gene expression was measured for both these strains from 0h
to 8h30m in sporulation medium (see Methods). Over one thousand gene transcripts
(1,122 including non-coding SUTs, Supporting Table S5) were identified to be
statistically significant in gene expression as a function of time between the two
strains (FDR cut-off 10%). However, in this set, none of the RAM network genes
showed significant differential expression in the presence of *TAO3(4477C)*. A few of
*ACE2*-regulated genes showed differential expression in the presence of
*TAO3(4477C)* (11 genes labeled green in Figure 2A), however, deletion of *ace2Δ* had
no affect on the sporulation efficiency of T strain (Figure 2B). Furthermore, even in
the high sporulating parent SK1, *ace2Δ*, which showed a lesser clumping phenotype
during growth in rich medium, had no effect on sporulation efficiency (Figure 2B).
Since previously Ace2-independent effect of RAM network on cellular polarization
have been observed (Nelson *et al.* 2003), therefore, RAM network could still be
involved in meiosis independently of Ace2. Since by reducing *TAO3(4477C)* activity
only during the growth phase in rich medium (in glucose) had no effect the
sporulation efficiency (Figure 1F), hence we concluded that the causative allele of
*TAO3* might affect sporulation efficiency by interacting with a set of genes unique
from the gene set involved during mitosis.

While studying the expression profiles of RAM network genes, we also observed a
higher expression of *TAO3* in the presence of the causative polymorphism in the T
strain, compared to the S strain ($P = 0.004$, Figure 2C). By altering *TAO3* expression,

using different concentrations of doxycycline, we found that higher than native

expression of *TAO3* increased sporulation efficiency (no doxycycline added, $P < 1 \times$

$10^{-4}$) and concordantly, a decrease in expression (increased doxycycline

concentration) reduced sporulation efficiency (Figure 2D). These results suggested an

interesting possibility of how the coding polymorphism in the transcriptional activator

*TAO3* might be involved in sporulation efficiency variation by affecting its gene

expression.

***Global gene expression variation during sporulation in presence of causative TAO3***

***allele***

While majority of regulation of sporulation occurs during both early and middle

stages of meiosis (Neiman 2011), even late genes are known to affect sporulation

efficiency in yeast (Deutschbauer *et al.* 2002). Between T and S strains, all of these

classes of sporulation genes, *i.e.*, early (*IME2*, *HOP1*, *DMC1*), middle and late

(*NDT80*, *SPR3*, *SPS1*) showed differential expression (Figure 3A, Supporting Table

S5). Most of the known sporulation genes (*IME2, DMC1, NDT80, SMK1, SWM1,*

*RAD6, REC8*), including various regulators of meiosis (*IME1, IME2, NDT80*) were

enriched ($P = 5.5 \times 10^{-12}$) in the cluster showing an increasing expression (Cluster II)

during sporulation in T strain (Figure 3B). Approximately 50% of genes in this cluster

(including *IME1*, *IME2*, *DMC1*, *ECM11*, *NDT80*) showed a similar trend in S strain

also (Supporting Figure S4, Supporting Table S6). Our cell cycle progression results

showed that lesser number of cells in S strain entered meiosis compared to T strain

(Figure 1C). Higher expression of meiotic and meiosis-associated genes in T strain

also reflected more cells entering meiosis. These results indicated that the early

effects of *TAO3* allele on sporulation were important for regulation of sporulation.

The early expressing Cluster I genes of T strain belonged to biological processes

regulating entry in sporulation, such as carbohydrate metabolic process, ion transport,

mitochondrion organization and cellular respiration (Table 1). Sparse overlap (7%,

Supporting Figure S4) was observed in genes of Cluster I between T and S strains,

with many meiosis-associated pathway genes (carbohydrate metabolic process and

mitochondrial organization) showing repression in S (Table 1, Supporting Figure S5).

We next studied the genes regulating sporulation genes in Cluster II and of those

getting expressed earlier than these sporulation genes, *viz.* early genes in Cluster I

(Supporting Tables S7, S8).

***Identifying candidate genes mediating the allele specific effects of TAO3 during***

***sporulation using the temporal gene expression data***

Regulators of genes upregulated either early or increasingly uniquely in T strain as

time progresses in sporulation were enriched in nutrient metabolism and chromatin

modification - the biological processes that are important for initiation of meiosis

(Neiman 2011). *UME6*, a transcriptional repressor whose degradation is required for

meiotic progression (Kassir *et al.* 2003) and is known to cause sporulation defects in

both S288c and SK1 strains, was also identified. This core sporulation gene is also

associated with various other important biological processes initiating meiosis

(Lardenois *et al.* 2015) and was identified as a regulator for most of the biological

processes occurring early in the presence of *TAO3(4477C)* (Table 2). From the list of

regulators (Supporting Tables S7, S8), we identified those regulators that were

upstream to *UME6* also (Figure 4A, Supporting Table S9). These sporulation-

associated regulators were selected as candidate causal mediating genes. They

included *ERT1*, regulator of carbon source utilization (Turcotte *et al.* 2009) involved

in the switch from fermentation to respiration in glucose-limiting conditions (Gasmi

*et al.* 2014); *OAF1,* regulator of lipid metabolism forming a protein complex with

*PIP2* (Karpichev and Small 1998); and *DAL81,* regulator of nitrogen degradation

pathway (Marzluf 1997). Interestingly, similar to *UME6*, *OAF1* target genes were

repressed (Cluster IV) in S strain (Supporting Table S10). Earlier work in S and SK1

strains has shown up regulation of *ERT1*, *PIP2* and *DAL81* in SK1 strain during

sporulation (Primig *et al.* 2000), though their deletion in S strain did not effect

sporulation efficiency (Deutschbauer *et al.* 2002).

A few other sporulation-associated regulators (Supporting Tables S7, S8) that were

not upstream *UME6* were also considered as candidate genes to be further

investigated. These were: *GAT3*, a transcription factor involved in spore wall

assembly (Lin *et al.* 2013); *RSC2*, required for the expression of mid-late sporulation

genes (Bungard *et al.* 2004); *GAT1* and *PHO4*, regulators of nitrogen and phosphorus

metabolism, respectively. These sporulation genes are not known to have an effect on

sporulation efficiency in S288c strain (Deutschbauer *et al.* 2002). Another putative

causal mediating gene whose target genes were enriched in the Cluster II uniquely in

*TAO3(4477C)* allele was *XBP1*, a transcriptional repressor induced in stress and

starvation, involved in chromatin modification and $G_1$ to S cell cycle progression. In SK1 strain, *XBP1* was highly induced during meiosis and its deletion reduced sporulation efficiency (Mai and Breeden 2000), whereas, *xbp1Δ* in S strain showed no effect on its sporulation efficiency (Deutschbauer *et al.* 2002). Interestingly, *XBP1* regulates *VAC8*, a vacuolar membrane protein required for efficient sporulation (Neiman 2005; Tang *et al.* 2006) that is also known to interact with *TAO3* by yeast-two hybrid studies (Tang *et al.* 2006).

In order to functionally validate a few of the candidate genes identified from the above analysis, we used a genetic model described previously (Gupta *et al.* 2015). According to this model, if a gene has no effect on sporulation efficiency, its deletion would not affect on the phenotype in T strain. If a gene had an independent role in sporulation, reduction in sporulation efficiency by its deletion would be independent of the *TAO3* background, and we would observe an additive effect irrespective of the background. Any significant deviation from this expectation would imply dependence on the genotype, such as the case of epistasis where deleting the gene in S background would not lead to decreased sporulation efficiency. While no difference was observed for *GAT1* (Supporting Figure S6), single deletions of *ERT1*, *PIP2* and *GAT3* reduced the mean sporulation efficiency in the T strain significantly, by about 1.5-fold ($P = 2.1 \times 10^{-12}$, $P = 6.1 \times 10^{-13}$, $P = 9.6 \times 10^{-10}$ respectively, pair test in Methods, Figure 4B). Significant interaction terms (see Methods) were obtained between the genetic background (S and T) and *ert1Δ* and *pip2Δ* ($P = 2.3 \times 10^{-4}$, $P = 0.04$), but not for *gat3Δ*. This showed that the effect of *ert1Δ* and *pip2Δ* on sporulation efficiency was specific to *TAO3(4477C)*. These analyses concluded that the meiotic role of allelic variant of *TAO3* was complex and was effecting sporulation efficiency by interacting with regulatory pathways involved in carbon metabolism (*ERT1*) and membrane component proteins involved in beta-oxidation (*OAF1-PIP2*).

## DISCUSSION

Strong effects on phenotypic variation have been observed as a consequence of rare coding variants (Cohen *et al.* 2004; 2005). *TAO3(4477C)*, a rare variant specific to the high sporulating laboratory yeast strain of SK1 isolated from soil, is not present in the other yeast strains in the SGRP collection (Liti *et al.* 2009). Here, we showed the applicability of temporal genome-wide transcriptome profiling approach (Gupta *et al.*

2015) to understand more about regulation of yeast sporulation efficiency variation

for the rare causal genetic variant *TAO3*.

Tao3, conserved from yeast to humans (Hergovich *et al.* 2006) has been functionally

annotated for mitotic cell division (Du and Novick 2002; Nelson *et al.* 2003), until it

got mapped for sporulation efficiency variation (Deutschbauer and Davis 2005). The

SK1 allele of *TAO3* has also been identified to affect growth variation in glycerol and

high temperature (Wilkening *et al.* 2014). Tao3 localizes to polarized bud sites during

mitosis (Nelson *et al.* 2003). Here, we observed genetic interaction of *TAO3(4477C)*

with membrane-associated *ERT1*. Thus, it would be interesting to ascertain the

localization of *TAO3* variants during sporulation. Similar to other scaffolding proteins

like *Fry* (*Drosophila*) and *SAX-2* (*C. elegans*), Tao3 has multiple conserved

Armadillo-like repeats (Hergovich *et al.* 2006) and the causal sporulation variant is

present in one of these domains. Since overexpression of *TAO3(4477G)* is known to

negatively affect the number of cells transitioning from $G_1$ to S phase during mitosis

(Bogomolnaya *et al.* 2006), a higher expression of the causative *TAO3* allele

specifically in meiosis raised the interesting possibility of this expression level

variation to be involved in $G_1$ to S phase transition in meiosis.

In complex diseases, combinations of molecular perturbations, which may vary in

different patients, may be dysregulating similar components of a cellular system.

Using a reductionist view of disease biology, many drug candidates have been

successfully identified by studying the core pathways associated to a disease (Schadt

*et al.* 2009). However, variation in the ancillary molecular players impinging on the

core pathways and residing in the genetic bottleneck could also contribute variation in

the phenotype (Lorenz and Cohen 2014). Genetic variation in mitosis-associated

genes of *MKT1* and *TAO3* are examples of the non-canonical pathways associated

with sporulation efficiency variation. Moreover, the novel metabolic and

mitochondrial regulators identified for *MKT1* (Gupta *et al.* 2015) and in this work for

*TAO3* as the mediating pathways causing variation in sporulation efficiency, are not

part of the core sporulation regulatory framework. Their modulation by genetic

variants provides important implications for understanding the rewiring of cellular

networks underlying phenotypic variation. In this study, *ERT1*, regulator of carbon

catabolite activation of transcription involved in the shift between fermentation to

respiratory growth (Gasmi *et al.* 2014) and *OAF1-PIP2,* regulators of lipid
metabolism necessary for the expression of sporulation specific genes *SPS18* and
*SPS19* (Gurvitz *et al.* 2009), were identified as mediating pathways contributing to
efficiency in meiosis. The molecular pathways in which these regulators were
involved in, emphasized the role of pleiotropic metabolic pathways that act as a
switch between multiple developmental phenotypes (Granek *et al.* 2011), to also have
important functional roles in variation in these phenotypes. Since, Tao3 is a
scaffolding protein involved in mitosis, and our findings link this protein in meiosis,
this puts forth an interesting hypothesis for its dual role of in recruiting different
molecular players in the two developmental phenotypes. Identification of these
ancillary pathways contributing to variation in meiosis suggested that a systems view
is necessary to understand the underlying complex biology of phenotypes including
various diseases. This systems-based study provides support to the emerging view of
disease as a complex network of interconnected pathways (Schadt *et al.* 2009), which
could be used to refine the clinical translation of GWAS results to understand disease
mechanism.
440
441

444   **Table 1.** Comparison of functional GO categories of differentially expressed genes in

445   T strain clusters with S strain. See Supporting Table S6 for full list of genes in each

446   cluster.

| Cluster | Functional GO category | Genes |
|---|---|---|
| Early in T strain (Cluster I) | Carbohydrate metabolic process | *DOG1, YPI1* |
| | Ion transport | *AVT4, DAL5* |
| | Mitochondrion organization | *PPE1, UPS3* |
| | Cellular respiration | *COX5B* |
| Early in T strain (Cluster I) repressed in S strain (Cluster IV) | Carbohydrate metabolic process | *ALG6, DEP1, DOG1, TPS3, YPI1* |
| | Mitochondrial organization | *ATG33, COX20, PPE1, UPS3* |

450
451 **Table 2.** Functional GO classification of regulators of differentially expressed genes
452 showing early and increasing expression only in T strain. See Supporting Tables S7,
453 S8 for full list of genes.
454

| Functional GO category | Regulators | *P* value |
|---|---|---|
| Carbon metabolism | *ERT1, OAF1, PIP2, MIG1, MIG2* | $1.9 \times 10^{-6}$ |
| Nitrogen catabolite regulation | *DAL81, DAL82, GAT1, UME6* | $1.7 \times 10^{-5}$ |
| Chromatin modification | *ISW1, PHO2, PHO4, UME6, OAF1, XBP1, SIF2, RSC2* | $1.4 \times 10^{-5}$ |

455
456

457
**ACKNOWLEDGEMENTS**

We thank Manu Tekkedil for help with whole genome sequencing sample

preparation. This research was supported by Tata Institute of Fundamental Research

intramural funds and Department of Biotechnology grant

BT/PR14842/BRB/10/881/2010 (H.S.); Bavarian Research Center for Molecular

Biosystems and Bundesministerium für Bildung und Forschung through the

Juniorverbund in der Systemmedizin "mitOmics" grant FKZ 01ZX1405A (J.G.); the

National Institutes of Health, Deutsche Forschungsgemeinschaft and a European

Research Council Advanced Investigator Grant (L.M.S.). The funders had no role in

study design, data collection and analysis, decision to publish, or preparation of the

manuscript.

**FIGURE LEGENDS**

**Figure 1. Role of *TAO3* in sporulation efficiency**

(A) Comparison of genomic sequence of *TAO3* (4,441-4,500) across the SGRP

collection (Liti *et al.* 2009). The 4,477th position of *TAO3* consists of the sporulation

causative variant, where identical nucleotides are indicated by the same color. Identity

indicates the percentage match between the nucleotides in the shown region of the

gene. The strains are ordered according to their mean sporulation efficiency (Tomar *et*

*al.* 2013): high (60-100%), intermediate (10-60%), low (0-10%) and ND (not

determined).

(B) Bar plots represents the mean sporulation efficiency, after 48h, of the SK1, T and

S strains. The sporulation efficiency data is indicated as circles.

(C) Line graphs represent mean sporulation efficiency of S, T and SK1 strains

measured till saturation, *i.e.*, till sporulation efficiency did not vary for 3 consecutive

days.

(D) Percentage of 1-, 2- and 4-nuclei states of the T strain (y-axis) versus time in

sporulation medium (x-axis). 1-nucleus stage is indicated as red circles ($G_0/G_1$ phase),

2-nuclei state as yellow circles (completion of MI phase) and blue circles is 4-nuclei

stage (completion of MII phase).

(E) Bootstrap distribution of the time to initiate meiosis and rate of transition from

$G_1/G_0$ into MI, estimated from time courses in (D). See Methods for details.

(F) Conditional expression of *TAO3(4477C)* during sporulation in $P_{Tet}$-*TAO3(4477C)*

strain (depicted as $P_{Tet}$). Y-axis is mean sporulation efficiency in 48h. No doxycycline

in growth (YPD) or spo (YPA + sporulation) medium is depicted as "-" condition on

x-axis and addition of doxycycline is depicted as "+" in that condition. "+3h"

condition in Spo implies doxycycline was throughout in the growth medium and in

the sporulation medium till 3h, after which cells were sporulated in the absence of

doxycycline. "+6h" condition implies doxycycline was throughout in the growth

medium and in the sporulation medium till 6h, after which cells were sporulated in the

absence of doxycycline. One-way ANOVA with Bonferroni's multiple comparison

tests was performed to test significance.

**Figure 2. Role of *TAO3* in meiosis is distinct from its role during mitosis**

(A) Heatmap showing gene expression of RAM network genes and Ace2-regulated genes in T and S strains. Gene names in green show differential expression (data in Supporting Tables S4, S5).

(B) Bar plots represent mean sporulation efficiency, after 48h, of the SK1 and T wild type (wt) and *ace2Δ* deletion strains. Pair and interaction tests (described in Methods) were performed to test significance.

(C) Expression profile (log$_2$ fold change t$_0$) of *TAO3* is given in the y-axis for T (purple) and S strains (red) and the x-axis denotes the time in sporulation medium. (data in Supporting Tables S4, S5)

(D) Bar plots represent mean sporulation efficiency of P$_{Tet}$-*TAO3(4477C)* strain in absence and presence of different concentrations of doxycycline (x-axis), compared to S strain. One-way ANOVA with Bonferroni's multiple comparison tests was performed to test significance.

**Figure 3. Global gene expression variation in presence of causative *TAO3* allele**

(A) Temporal heat map of meiotic genes in T and S strains. The gene names shown in green are differentially expressed in the presence of *TAO3(4477C)*.

(B) The expression profile (log$_2$ fold change t$_0$) for the meiotic landmark genes is given in the y-axis and the x-axis denotes the time in sporulation medium. Red line represents the expression profile of the respective gene in S strain and blue line is the same in T strain.

(C) Heat map of the T and S strains showing differentially expressed gene across time within each cluster. Each row represents a single gene and columns are time points of each strain (for gene list in each cluster see Supporting Table S6). The order of genes in the two strains is based on the clustering of the T strain. Functional GO categories of genes in each cluster are shown on left. The boxplots shown on right represent the average expression profile of each cluster in the T and S strain. The number of genes in each cluster in a strain is indicated in brackets.

**Figure 4. Identifying candidate genes mediating the allele specific effects of *TAO3* during sporulation using the temporal gene expression data**

(A) Regulatory network of candidate genes predicted to mediate the effects of *TAO3(4477C)* in sporulation. The candidate mediating genes are shown as bigger

nodes (large circles), with their target genes (small circles) connected to them as

straight lines. The box contains the protein network interactions of the candidate

genes with core sporulation gene *UME6*, obtained from YEASTRACT (see Methods).

Colors inside the nodes were calculated as an average of the first six time points in

sporulation (early phase). For complete list of interacting genes and their expression

values, see Supporting Tables S9 and S4, respectively.

(B) Bar plots represent mean sporulation efficiency, after 48h, of the T and S wild

type (wt) and *ert1Δ*, *pip2Δ* and *gat3Δ* strains. Pair and interaction tests (see Methods)

were performed to test significance.

547
<div align="center">**SUPPORTING INFORMATION**</div>

**Supporting Files**

**File S1.** Detailed methods

**Supporting Tables**

**Table S1.** Whole genome resequencing results for the *TAO3* allele replacement strain

**Table S2.** Strains names

**Table S3.** Primer names

**Table S4.** Smoothed expression data, base transformed with respect to $t_0$ for T and S
strains

**Table S5.** Differentially expressed genes between T and S strains, with their *P* and *Q*
values calculated using EDGE

**Table S6.** Genes in each cluster using TimeClust

**Table S7.** Transcription factors regulating unique early (Cluster I) genes of the T
strain

**Table S8.** Transcription factors regulating unique increasing (Cluster II) genes of the
T strain

**Table S9.** Differentially expressed target genes of regulators of candidate genes
mediating the affect of *TAO3*

**Table S10.** Transcription factors regulating unique repressing (Cluster IV) genes of
the S strain

**Supporting Figures**

**Figure S1.** Whole genome resequencing of *TAO3* allele replacement strain (YAD331,
(Deutschbauer and Davis 2005) in comparison to S288c reference strain

**Figure S2.** Smoothing of normalized temporal data using *locfit*

**Figure S3.** Mathematical modeling to identify stage of meiosis affected by *TAO3*
causal allele

**Figure S4.** Comparison of genes showing early (Cluster I) and increasing trend
(Cluster II) between T and S strains

**Figure S5.** Genes showing early expression in T strain show expression at later time
points or repressed in S strain

**Figure S6.** Sporulation efficiency of *gat1Δ* in T strain

**REFERENCES**

Balakrishnan R., Park J., Karra K., Hitz B. C., Binkley G., Hong E. L., Sullivan J.,
Micklem G., Cherry J. M., 2012  YeastMine—an integrated data warehouse for
Saccharomyces cerevisiae data as a multipurpose tool-kit. Database 2012:
bar062–bar062.
Bellí G., Garí E., Aldea M., Herrero E., 1998a  Functional analysis of yeast essential
genes using a promoter-substitution cassette and the tetracycline-regulatable dual
expression system. Yeast 14: 1127–1138.
Bellí G., Garí E., Piedrafita L., Aldea M., Herrero E., 1998b  An activator/repressor
dual system allows tight tetracycline-regulated gene expression in budding yeast.
Nucleic Acids Research 26: 942–947.
Bogomolnaya L. M., Pathak R., Guo J., Polymenis M., 2006  Roles of the RAM
signaling network in cell cycle progression in Saccharomyces cerevisiae. Curr
Genet 49: 384–392.
Bungard D., Reed M., Winter E., 2004  RSC1 and RSC2 are required for expression
of mid-late sporulation-specific genes in Saccharomyces cerevisiae. Eukaryotic
Cell 3: 910–918.
Cirulli E. T., Goldstein D. B., 2010  Uncovering the roles of rare variants in common
disease through whole-genome sequencing. Nat Rev Genet 11: 415–425.
Cohen J. C., Kiss R. S., Pertsemlidis A., Marcel Y. L., 2004  Multiple rare alleles
contribute to low plasma levels of HDL cholesterol. Science 305: 869–872.
Cohen J., Pertsemlidis A., Kotowski I. K., Graham R., Garcia C. K., Hobbs H. H.,
2005  Low LDL cholesterol in individuals of African descent resulting from
frequent nonsense mutations in PCSK9. Nat. Genet. 37: 161–165.
Deutschbauer A. M., Davis R. W., 2005  Quantitative trait loci mapped to single-
nucleotide resolution in yeast. Nat. Genet. 37: 1333–1340.
Deutschbauer A. M., Williams R. M., Chu A. M., Davis R. W., 2002  Parallel
phenotypic analysis of sporulation and postgermination growth in Saccharomyces
cerevisiae. Proceedings of the National Academy of Sciences 99: 15530–15535.
Du L.-L., Novick P., 2002  Pag1p, a novel protein associated with protein kinase
Cbk1p, is required for cell morphogenesis and proliferation in Saccharomyces
cerevisiae. Mol. Biol. Cell 13: 503–514.
Gasmi N., Jacques P.-E., Klimova N., Guo X., Ricciardi A., Robert F., Turcotte B.,
2014  The switch from fermentation to respiration in Saccharomyces cerevisiae
is regulated by the Ert1 transcriptional activator/repressor. Genetics 198: 547–
560.
Gietz R., Woods R. A., 2002  Transformation of yeast by lithium acetate/single-
stranded carrier DNA/polyethylene glycol method. Meth. Enzymol. 350: 87–96.
Goldstein A. L., McCusker J. H., 1999  Three new dominant drug resistance cassettes
for gene disruption in Saccharomyces cerevisiae. Yeast 15: 1541–1553.
Granek J. A., Kayıkçı Ö., Magwene P. M., 2011  Pleiotropic signaling pathways
orchestrate yeast development. Current Opinion in Microbiology 14: 1–6.
Gupta S., Radhakrishnan A., Raharja-Liu P., Lin G., Steinmetz L. M., Gagneur J.,
Sinha H., 2015  Temporal expression profiling identifies pathways mediating
effect of causal variant on phenotype. PLoS Genet 11: e1005195.
Gurvitz A., Suomi F., Rottensteiner H., Hiltunen J. K., Dawes I. W., 2009  Avoiding
unscheduled transcription in shared promoters: Saccharomyces cerevisiae Sum1p
represses the divergent gene pair SPS18-SPS19 through a midsporulation element

(MSE). FEMS Yeast Res 9: 821–831.

Hallgrimsson B., Mio W., Marcucio R. S., Spritz R., 2014   Let's Face It—Complex
Traits Are Just Not That Simple. PLoS Genet 10: e1004724.

Hergovich A., Stegert M. R., Schmitz D., 2006   NDR kinases regulate essential cell
processes from yeast to humans. Nature Reviews Molecular Cell Biology 7: 253–
264.

Huber W., Heydebreck von A., Sültmann H., Poustka A., Vingron M., 2002
Variance stabilization applied to microarray data calibration and to the
quantification of differential expression. Bioinformatics 18 Suppl 1: S96–104.

Karpichev I. V., Small G. M., 1998   Global Regulatory Functions of Oaf1p and
Pip2p (Oaf2p), Transcription Factors That Regulate Genes Encoding Peroxisomal
Proteins in Saccharomyces cerevisiae. Molecular and Cellular Biology 18: 6560–
6570.

Kassir Y., Adir N., Boger-Nadjar E., Raviv N. G., Rubin-Bejerano I., Sagee S.,
Shenhar G., 2003   Transcriptional regulation of meiosis in budding yeast. Int.
Rev. Cytol. 224: 111–171.

Lardenois A., Becker E., Walther T., Law M. J., Xie B., Demougin P., Strich R.,
Primig M., 2015   Global alterations of the transcriptional landscape during yeast
growth and development in the absence of Ume6-dependent chromatin
modification. Mol. Genet. Genomics 290: 2031–2046.

Lin C. P.-C., Kim C., Smith S. O., Neiman A. M., 2013   A Highly Redundant Gene
Network Controls Assembly of the Outer Spore Wall in S. cerevisiae. PLoS
Genet 9: e1003700.

Liti G., Carter D. M., Moses A. M., Warringer J., Parts L., James S. A., Davey R. P.,
Roberts I. N., Burt A., Koufopanou V., Tsai I. J., Bergman C. M., Bensasson D.,
O'Kelly M. J. T., van Oudenaarden A., Barton D. B. H., Bailes E., Nguyen A. N.,
Jones M., Quail M. A., Goodhead I., Sims S., Smith F., Blomberg A., Durbin R.,
Louis E. J., 2009   Population genomics of domestic and wild yeasts. Mol Syst
Biol 458: 337–341.

Liu F., van der Lijn F., Schurmann C., Zhu G., Chakravarty M. M., Hysi P. G.,
Wollstein A., Lao O., de Bruijne M., Ikram M. A., van der Lugt A., Rivadeneira
F., Uitterlinden A. G., Hofman A., Niessen W. J., Homuth G., de Zubicaray G.,
McMahon K. L., Thompson P. M., Daboul A., Puls R., Hegenscheid K., Bevan
L., Pausova Z., Medland S. E., Montgomery G. W., Wright M. J., Wicking C.,
Boehringer S., Spector T. D., Paus T., Martin N. G., Biffar R., Kayser M., 2012
A Genome-Wide Association Study Identifies Five Loci Influencing Facial
Morphology in Europeans. PLoS Genet 8: e1002932.

Lorenz K., Cohen B. A., 2014   Causal Variation in Yeast Sporulation Tends to
Reside in a Pathway Bottleneck. PLoS Genet 10: e1004634.

Magni P., Ferrazzi F., Sacchi L., Bellazzi R., 2008   TimeClust: a clustering tool for
gene expression time series. Bioinformatics 24: 430–432.

Mai B., Breeden L., 2000   CLN1 and its repression by Xbp1 are important for
efficient sporulation in budding yeast. Molecular and Cellular Biology 20: 478–
487.

Manolio T. A., Collins F. S., Cox N. J., Goldstein D. B., Hindorff L. A., Hunter D. J.,
McCarthy M. I., Ramos E. M., Cardon L. R., Chakravarti A., Cho J. H.,
Guttmacher A. E., Kong A., Kruglyak L., Mardis E., Rotimi C. N., Slatkin M.,
Valle D., Whittemore A. S., Boehnke M., Clark A. G., Eichler E. E., Gibson G.,
Haines J. L., Mackay T. F. C., McCarroll S. A., Visscher P. M., 2009   Finding
the missing heritability of complex diseases. Nature 461: 747–753.

Marzluf G. A., 1997    Genetic regulation of nitrogen metabolism in the fungi.
Microbiology and Molecular Biology Reviews 61: 17–32.
Neiman A. M., 2005    Ascospore formation in the yeast Saccharomyces cerevisiae.
Microbiology and Molecular Biology Reviews 69: 565–584.
Neiman A. M., 2011    Sporulation in the Budding Yeast Saccharomyces cerevisiae.
Genetics 189: 737–765.
Nelson B., Kurischko C., Horecka J., Mody M., Nair P., Pratt L., Zougman A.,
McBroom L. D. B., Hughes T. R., Boone C., Luca F. C., 2003    RAM: a
conserved signaling network that regulates Ace2p transcriptional activity and
polarized morphogenesis. Mol. Biol. Cell 14: 3782–3803.
Primig M., Williams R. M., Winzeler E. A., Tevzadze G. G., Conway A. R., Hwang
S. Y., Davis R. W., Esposito R. E., 2000    The core meiotic transcriptome in
budding yeasts. Nat. Genet. 26: 415–423.
Saint Pierre A., Génin E., 2014   How important are rare variants in common disease?
Briefings in Functional Genomics 13: 353–361.
Schadt E. E., Friend S. H., Shaywitz D. A., 2009    A network view of disease and
compound screening. Nat Rev Drug Discov 8: 286–295.
Spellman P. T., Sherlock G., Zhang M. Q., Iyer V. R., Anders K., Eisen M. B., Brown
P. O., Botstein D., Futcher B., 1998    Comprehensive identification of cell cycle-
regulated genes of the yeast Saccharomyces cerevisiae by microarray
hybridization. Mol. Biol. Cell 9: 3273–3297.
Stankiewicz P., Lupski J. R., 2010   Structural variation in the human genome and its
role in disease. Annu. Rev. Med. 61: 437–455.
Storey J. D., Xiao W., Leek J. T., Tompkins R. G., Davis R. W., 2005   Significance
analysis of time course microarray experiments. Proceedings of the National
Academy of Sciences 102: 12837–12842.
Tang F., Peng Y., Nau J. J., Kauffman E. J., Weisman L. S., 2006    Vac8p, an
armadillo repeat protein, coordinates vacuole inheritance with multiple vacuolar
processes. Traffic 7: 1368–1377.
Teixeira M. C., Monteiro P. T., Guerreiro J. F., Gonçalves J. P., Mira N. P., Santos
dos S. C., Cabrito T. R., Palma M., Costa C., Francisco A. P., Madeira S. C.,
Oliveira A. L., Freitas A. T., Sá-Correia I., 2014    The YEASTRACT database:
an upgraded information system for the analysis of gene and genomic
transcription regulation in Saccharomyces cerevisiae. Nucleic Acids Research 42:
D161–6.
Tomar P., Bhatia A., Ramdas S., Diao L., Bhanot G., Sinha H., 2013    Sporulation
Genes Associated with Sporulation Efficiency in Natural Isolates of Yeast. PLoS
ONE 8: e69765.
Turcotte B., Liang X. B., Robert F., Soontorngun N., 2009   Transcriptional regulation
of nonfermentable carbon utilization in budding yeast. FEMS Yeast Res 10: 2–13.
Wilkening S., Lin G., Fritsch E. S., Tekkedil M. M., Anders S., Kuehn R., Nguyen
M., Aiyar R. S., Proctor M., Sakhanenko N. A., Galas D. J., Gagneur J.,
Deutschbauer A., Steinmetz L. M., 2014    An evaluation of high-throughput
approaches to QTL mapping in Saccharomyces cerevisiae. Genetics 196: 853–
865.
Xu Z., Wei W., Gagneur J., Perocchi F., Clauder-Münster S., Camblong J., Guffanti
E., Stutz F., Huber W., Steinmetz L. M., 2009   Bidirectional promoters generate
pervasive transcription in yeast. Nature 457: 1033–1037.
Zuk O., Schaffner S. F., Samocha K., Do R., Hechter E., Kathiresan S., Daly M. J.,
Neale B. M., Sunyaev S. R., Lander E. S., 2014    Searching for missing

heritability: designing rare variant association studies. Proc. Natl. Acad. Sci.
U.S.A. 111: E455–64.