Male sex pheromone components in the butterfly *Heliconius melpomene*

Sohini Vanjari[1], Florian Mann[2], Richard M. Merrill[1,3], Stefan Schulz[2], Chris Jiggins[1]

1. Department of Zoology, University of Cambridge, Downing Street, Cambridge, CB2 3EJ, United Kingdom.

2. Institute of Organic Chemistry, Technische Universität Braunschweig, Hagenring 30, 38106 Braunschweig, Germany.

3. Smithsonian Tropical Research Institute, MRC 0580-12, Unit 9100 Box 0948, DPO AA 34002-9998, Panama.

Abstract

Sex specific pheromones are known to play an important role in butterfly courtship, and may influence both individual reproductive success and reproductive isolation between species. Extensive ecological, behavioural and genetic studies of *Heliconius* butterflies have made a substantial contribution to our understanding of speciation. However, although long suspected to play an important role, chemical signals have received relatively little attention in this genus. Here, we combine morphological, chemical and behavioural analyses of a male pheromone in the neotropical butterfly *Heliconius melpomene.* First we identify specialized brush-like scales that are putative androconia, and lie within the shiny grey region found on the hindwing of males. We then describe six putative male sex pheromone compounds, which are largely confined to the androconial region of the hindwing of mature males, but not immature males or females. Finally, behavioural assays reveal subtle, but detectable, differences in female response to models scented with hindwing androconial extracts of mature conspecific males as compared to unscented controls. Collectively, the results describe structures involved in release of the pheromone and a list of potential male sex pheromone compounds triggering a behavioural response in females.

Key words: *Heliconius*, pheromone, androconia, Lepidoptera, sexual selection.

Introduction

Organisms use chemicals for a wide variety of forms of communication. In particular, sex pheromones are species-specific blends of chemical compounds mediating intraspecific communication between males and females, and can play key roles in both determining the reproductive success of individuals within a species, and in reproductive isolation between species (Wyatt, 2003, 2014). Some insects deploy single chemicals as signals, but in many insects, pheromone communication is dependent on complex combinations of chemical components (Grillet, Dartevelle, & Ferveur, 2006; Nieberding *et al.*, 2008; Symonds, Johnson, & Elgar, 2012). This chemical complexity provides the potential to convey sophisticated information (Nieberding *et al.*, 2012). The best studied insect sex pheromones are perhaps the chemicals produced by female moths to attract mating partners, often over long distances (Löfstedt, 1993; Smadja & Butlin, 2008). Male insects can also produce sex pheromones but these have generally received less attention (Wyatt, 2014).

Whilst variation in sex pheromone blend is now known to be a major determinant of reproductive isolation and speciation in many species of moths (Löfstedt, 1993; Smadja & Butlin, 2008; Lassance *et al.*, 2010), to date sex pheromones have been studied in just a few species of butterfly. Acceptance behaviour in the queen butterfly *Danaus berenice* is regulated by a dihydropyrrolizine alkaloid released by the male (Brower & Jones, 1965; Meinwald, Meinwald, & Mazzocchi, 1969; Pliske & Eisner, 1969). Another danaine butterfly, *Idea leuconoe* displays 'hair-pencils' during courtship, which contain a mixture of dihydropyrrolizine alkaloids, aromatics, terpenoids, hydrocarbons and a series of γ-lactones (Nishida *et al.*, 1996). This volatile mixture applied on dummy male butterflies elicits an acceptance posture in females. *Pieris rapae* and *Pieris brassicae* use macrocyclic lactones as a pheromone to induce acceptance in females (Yildizhan *et al.*, 2009). In *Bicyclus anynana* males with reduced amounts of male sex pheromone have decreased mating success implying a direct involvement in reproductive fitness (Nieberding *et al.*, 2008, 2012). Male wing compounds are also known to contribute to reproductive isolation between closely related species of butterflies (Grula, McChesney, & Taylor, 1980; Phelan & Baker, 1987; Bacquet *et al.*, 2015).

Here we focus on the potential role of a male contributed pheromone in *Heliconius* butterflies. *Heliconius* is a diverse neotropical genus, studies of which have contributed greatly to our understanding of speciation (Jiggins, 2008; Merrill *et al.*, 2015). These butterflies are known for Müllerian mimicry, where unrelated species converge on the same warning signal to more efficiently advertise their unpalatability to predators. However, closely related *Heliconius* taxa often differ greatly in colour pattern and divergent selection acting on warning patterns is believed to play an important role in speciation within the genus

(Bates, 1862; Jiggins *et al.*, 2001; Merrill *et al.*, 2011b). In particular, males are known to use

colour pattern to recognize mates (Jiggins *et al.*, 2001; Jiggins, Estrada, & Rodrigues, 2004;

Kronforst *et al.*, 2006; Melo *et al.*, 2009; Merrill *et al.*, 2011a; Merrill, Chia, & Nadeau, 2014;

Finkbeiner, Briscoe, & Reed, 2014).

Some aspects of chemical signaling have been studied in *Heliconius*. For example,

males are known to transfer an antiaphrodisiac pheromone to females during mating (Gilbert,

1976). Once mated, females produce a strong odor that acts to repel approaching males

(Schulz *et al.*, 2008). Studies in *H. melpomene* have shown that abdominal glands of males

contain a complex odor bouquet consisting of the volatile compound (*E*)-β-ocimene together

with some trace components and esters of common C16 – and C18 – fatty acids with alcohols,

where β-ocimene acts as the main antiaphrodisiac pheromone. This bouquet is formed during

first few days of eclosion and transferred during copulation to the females (Schulz *et al.*,

2008). This antiaphrodisiac effect can be observed in several *Heliconius* species, which show

species-specific patterns of scent gland constituents (Gilbert, 1976). In addition, *Heliconius*

may also use green leaf volatiles during mate searching. Six-carbon alcohols and acetates are

released in larger amounts after tissue damage by caterpillars, which males of the pupal

mating species *H. charithonia* then use to find potential mates (Estrada & Gilbert, 2010).

Once males find pupae they also use chemical cues to determine sex (Estrada *et al.*, 2010).

There are also a number of observations that indicate a role for chemical recognition

during adult mating. *H. erato* males can distinguish between wings dissected from conspecific

and heterospecific females that are virtually identical in wing pattern, but this effect

disappears after wings have been washed in hexane (Estrada & Jiggins, 2008). *H. cydno*

males show a preference for their own pattern over that of the closely related *H. melpomene*,

but will court wing pattern models of *H. melpomene*. However, *H. cydno* males have virtually

never been observed mating with *H. melpomene* females, suggesting strong barriers in

addition to wing pattern (Naisbit, Jiggins, & Mallet, 2001). In addition, *H. melpomene*

coexists with visually almost identical co-mimics, notably *H. erato*, which is likely to favour

the evolution of pheromonal recognition signals to avoid confusion in courtship.

In some populations the very closely related species *H. timareta*, is sympatric and

mimetic with *H. melpomene*, but nevertheless displays strong assortative mating (Giraldo *et*

al., 2008). A recent study of sympatric *H. melpomene* and *H. timareta* in Peru has provided

some of the first evidence for a role of chemical signals in species recognition (Mérot *et al.*,

2015). Experiments with perfumed males using abdominal scent glands and wing extracts

shows increased probability of inter-specific mating when males were perfumed with

heterospecific extracts. In addition, chemical analysis of both abdominal glands and whole

wings provided evidence for differences between these closely related species in their
chemical signatures (Mérot *et al.*, 2015)
Here we focus on the wing pheromones of *H. melpomene*. First, we investigate
morphological structures potentially associated with pheromone production. In butterflies, a
variety of species-specific structures including brushes, fans, and differentiated scales on
wings, legs or abdomen are used to expose pheromones produced in associated glands (Wyatt,
2003; Nieberding *et al.*, 2008). In particular, male specific scent glands, termed androconia
are common across the Lepidoptera. In male *Heliconius*, a patch of shiny grey scales is
present on the overlapping region of the hind and forewing (Figure 1). The observed sexual
dimorphism in this trait suggests that these are androconia, and may be associated with a male
sex pheromone (Emsley, 1963). Furthermore, earlier authors have identified brush-like scales
in this region that are the putative site for pheromone production (Müller, 1912; Barth, 1952).
Here we investigate the structure of these scales using Scanning Electron Microscopy.
Second, we complement recently published chemical analysis of whole *H. melpomene* wings
by dissecting wing regions to identify those wing regions that are associated with the
production of compounds and identify potential male sex pheromone compounds isolated
from this region. Finally, using a simple behavioural assay we evaluate female response to
male androconial pheromone extracts.
119
Methods
Individuals used for morphological and chemical analyses were from an outbred stock of
Heliconius melpomene plesseni and *Heliconius melpomene malleti* (sold as *H. m. aglaope*)
maintained at the University of Cambridge insectaries (Figure 1). These two races are from
the region of a hybrid zone in the eastern Andes of Ecuador, and showed signs of inter-racial
hybridization in the stocks, so are treated here as a single population and referred to as the
Ecuador samples. These stocks were established from individuals obtained from a commercial
breeder (*Stratford Butterfly Farm*: www.butterflyfarm.co.uk). Laboratory stocks were
maintained on the larval food plants, *Passiflora menispermifolia* and *Passiflora biflora*. Adult
butterflies were fed on ~10% sugar solution mixed with an amino acid supplement (Critical
Care Formula®). In addition, five individual males of *H. m. rosina* were sampled from the
wild around the town of Gamboa in Colón Province, Panama, and are referred to as the
Panama samples.
133
Morphological analysis
The detailed morphology of androconial scales was determined using Field Emission
Scanning Electron Microscope. Three males and two females of *H. melpomene* from Ecuador

were used for this analysis. The androconial grey scale region was dissected out from both

hind and forewings and attached to aluminium stubs with carbon tabs and subsequently

coated with 20nm of gold using a Quorum/Emitech sputter coater. The gold-coated

androconia were then viewed in an FEI XL30 FEGSEM operated at 5kV. Images were

recorded digitally using XL30 software at 500x magnification.

Characterization of potential male sex pheromone

Wing tissue from ten male (five newly emerged and five 10-day old mature individuals) and

five female (10-day old) individuals from the Ecuador stock was collected between November

2011 and March 2012 for chemical analysis. In addition, wing tissue from five male adult

individuals from Panama was collected in August 2012 for chemical analysis. Wings were

dissected into four parts: forewing androconia, hindwing androconia, forewing non-

androconia and hindwing non-androconia. The 'androconia' regions corresponded to the

grey-brown region shown in Figure 1c, with non-androconia corresponding to the remaining

portion of the wing. In females, a region corresponding in size and extent to the grey-brown

region seen in males was dissected. The dissected sections were then allowed to soak in either

hexane or dichloromethane for 3 hours in a glass vial. The solvent was then transferred to new

vial and stored at -20°C. Initial tests showed no major differences between hexane and

dichloromethane extracts. Therefore, the more polar dichloromethane was used in later

analysis.

Extracts were analyzed by gas chromatography/mass spectrometry (GC/MS) using a

Hewlett-Packard model 5975 mass-selective detector connected to a Hewlett-Packard GC

model 7890A, and equipped with a Hewlett-Packard ALS 7683B autosampler. A HP-5MS

fused silica capillary column (Agilent, 30 m × 0.25 mm, 0.25 µm) was used. Injection was

performed in splitless mode (250°C injector temperature) with helium as the carrier gas

(constant flow of 1.2 ml/min). The temperature programme started at 50°C, was held for 5

min, and then rose to 320°C with a heating rate of 5°C/min. All components were identified

by comparison of mass spectra and gas chromatographic retention index with those of

authentic reference samples, and analysis of mass spectra. The double bond position of

unsaturated compounds were determined by derivatisation with dimethyl disulfide (Buser *et*

al., 1983). The alcohols were synthesised from the corresponding esters by reduction

according to established procedures (Becker & Beckert, 1993, p. 570). The aldehydes were

synthesised by oxidation of the respective alcohols (More & Finney, 2002). Synthesis of the

methyl branched alcohols and aldehydes will be reported elsewhere (F. Mann et al., in

preparation). The samples were quantified by using gas chromatography with flame ionisation

detection with a Hewlett-Packard GC model 7890A equipped with a Hewlett-Packard ALS

7683B autosampler. A BPX-5 fused silica capillary column (SGE, 25 m × 0.22 mm, 0.25 μm)

was used. Injection was performed in splitless mode (250°C injector temperature) with

hydrogen as the carrier gas (constant flow of 1.65 ml/min). The temperature programme

started at 50°C, held for 5 min, and then rose to 320°C with a heating rate of 5°C/min.

Pentadecyl acetate (10.1 ng) or (Z)-4-tridecenyl acetate (1 ng) were used as internal standard.

Only compounds eluting earlier than hexacosane were considered for analysis. Later

compounds were identified as cuticular hydrocarbons, 2,5-dialkyltetrahydrofurans, cholesterol

and artefacts like phthalates or adipates. The variability in the late eluting cuticular

hydrocarbons was low and did not show characteristic differences.

Behavioural experiments

Behavioural assays to determine female response to the putative male sex pheromone were

conducted in the *Heliconius* insectaries at the Smithsonian Tropical Research Institute's

(STRI) in Gamboa, Panama between January 2013 and July 2013. Females were assayed

from an outbred stock of *H. melpomene rosina* established from wild individuals collected in

Gamboa (9°7.4′ N, 79°42.2′ W, elevation 60 m) and the nearby Soberanía National Park,

República de Panamá. Stocks were maintained on the larval food plants, *Passiflora*

menispermifolia and *Passiflora biflora*. Adult butterflies were fed on ~20% sugar solution

with pollen supplements and *Psychotria sp.* and *Psiguiria sp.* serving as an additional pollen

source. Males sacrificed for extracts for behavioural experiments were either from this same

stock or were wild individuals collected in Gamboa or the Soberanía National Park. Male

extracts were obtained by dissecting the androconial region from both hindwings, followed by

soaking in 200 μl of hexane for 2-3 hours before transferring the solvent to a new glass vial.

This vial was sealed and stored at -20°C until required.

Behavioural experiments were performed in cages approximately 2m x 2m x 2m.

Ten virgin females (<5 days after eclosion) were presented with a paper model of *H.*

melpomene rosina attached to a ~30cm length of wire, itself attached to a ~60cm stick on

which a camera (Xdreme HD 1080P Action camera) was mounted to record female

behaviour. Pheromone extracts from individual *H. melpomene* males or a control (solvent

only) were applied to a strip of Whatman filter paper no. 4, which was attached to the paper

butterfly model. The extract of a single male was used for each individual female, so in total

ten virgin females and ten male extracts were used. Each trial proceeded as follows: After

mounting the paper model on the wire, 200 μl of extract/control was applied to the filter

paper. Trials were conducted blind: A third party applied the extract/control so that the

experimenter did not know whether they were presenting the focal female with the extract or

control. The paper model was then presented to the focal female, and the apparatus

manipulated to simulate the hovering flight observed in courting males. Each presentation

lasted one minute, followed by a one-minute break when the model was removed from the

vicinity of the focal female. This was repeated five times before a further five-minute pause in

the trial. The whole procedure was repeated six times, alternating between the two treatments,

so that each female experienced a total of 15 presentations with the male extract and 15

presentations with the control (solvent only).

215   Female behaviours were scored from videos recorded during the trials. To avoid

observer bias, the names and order of the digital video files were randomized and only

reassigned to treatment and individual once scoring was complete. We determined the

behaviours to be recorded from observations of interactions between *H. melpomene* males and

females (Table 1). These were 'Slow and moderate wing flapping', where females opened and

closed their wings slowly at regular intervals displaying their wing colors; 'Flying/Flying

facing towards model'where females tried to chase the paper models for short intervals or

females hovered facing the model; 'Slow rhythmic flight' where females displayed slow flight

with intermittent gliding; 'Wing display with abdomen normal' where females opened their

wings displaying the dorsal wing surface with the abdomen between the wings; 'Quick and

jerky wing flapping' where there was continuous wing fluttering by females that effectively

prevented close contact by the model; 'Flying away from model' where females flew away

from the paper model avoiding any interaction; 'Fast erratic flight' where such flight was

notably very fast, abrupt and apparently directionless and finally 'Wings open with abdomen

erect' where females opened their wings displaying the dorsal wing surface with the abdomen

erect. Each of these behaviours was scored as having occurred or not occurred during each of

the one-minute presentations. This led to a dataset for individual females where the proportion

of one-minute presentations in which each behaviour was observed was calculated (for male

extract and control trials separately). To reduce the number of dependent variables we then

performed a principal component analysis on these data. The principal component scores for

each individual were extracted and were used to test for differences in response between

presentations with the male extract and control (solvent only) in paired tests. Statistical

analyses were performed with *R* (version 3.1.2).

Results

Morphological analysis

In order to investigate structural differences in scale morphology potentially associated with

pheromone production, we observed androconial scales of *H. melpomene rosina* males and

females under the scanning electron microscope (Figure 2). We identified a marked sexual

dimorphism in scale structure. In the central region of the male hindwing androconia along

vein Sc+R1 we identified scales with brush-like structures (Figure 2d) at their distal end

(Figure 2a). These scales were completely absent in females and in the forewing androconia

of the males (Figure 2). These scales were also not detected in any other wing region

examined. The brush-like scales were found in alternating rows with scales with a normal

structure. Moving away from the Sc+R1 wing vein, the density and width of these scales

decreased, with isolated brush-like scales found completely surrounded by normal scales. In

addition, it was found that the base of these brush-like scales was more swollen and glandular

as compared to other scales, perhaps indicating a location for the storage or production of the

pheromone (Figure 3).

Characterization of potential male sex pheromone

Using gas chromatography linked to mass spectrometry (GC/MS) and synthesis, six potential

male sex pheromone components were found in the male wing extracts from Ecuador samples

(Figure 4). These compounds were, (*Z*)-9-octadecenal, octadecanal, henicosane, (*Z*)-11-

icosenal, icosanal and (*Z*)-13-docosenal. Out of these six compounds, only henicosane was

found in all regions of the wing, but (*Z*)-9-octadecenal, octadecanal, (*Z*)-11-icosenal, icosanal

and (*Z*)-13-docosenal were restricted to the androconial scale region of the wing. The

hindwing androconial region tended to contain higher titres of these five compounds as

compared to forewing androconial wing region (Figure 5). Octadecanal was present in higher

amounts in hindwing androconia, while icosanal were only found in trace quantities. (Z)-9-

Octadecenal was found in small amounts only in some of the Ecuador samples. A related

compound, (*Z*)-11-Icosanol, was found in trace amounts in two Ecuador mature males. Except

for henicosane, all five compounds were observed to be age-specific and sex-specific.

Henicosane was present in the hindwing androconia of both 10-day old and newly emerged

males and in females of all ages. In contrast, (*Z*)-9-octadecenal, octadecanal, (*Z*)-11-icosenal,

icosanal and (*Z*)-13-docosenal were only present in extracts from the hindwing androconia of

10-day old males. The chemical analysis of the extracts of the Panama samples showed a

similar composition, although they contained more compounds in slightly higher

concentrations. Major components of both the Panama and Ecuador individuals were

octadecanal, (*Z*)-11-icosenal, icosanal, and (*Z*)-13-docosenal. (*Z*)-11-Icosenol was found in

larger amounts in the Panama samples. They additionally contained high amounts of

octadecanol. Small amounts of nonadecanal, methyl-branched octadecanals and their

respective alcohols occurred as well. Some of them were identified to be 15-, 16-, and 17-

methyloctadecanals and the respective alcohols. These compounds were found only in the

Panama samples with the exception of 17-methyloctadecanal, which was present in two of the

samples from Ecuador. *n*-Alkanes were not present in these extracts.

Behavioural experiments

We tested ten individual virgin females in our behavioural trials, each of which was presented
sequentially with either male wing extract or a control (solvent only) applied to a paper
model. Behavioural scores were analysed using a principal components analysis, with two
principal components retained that together accounted for 80.38% of the variance in the data
(Table 1, see also supplementary file 1 for the raw data). The second axis accounted for
18.3% of the variance. There was no significant difference between male extract and control
(solvent only) trials for PC1 (Figure 6; Wilcoxon signed rank test: $V = 33$, $p > 0.05$);
however, there was a significant difference between male extract and control (solvent only)
trials for PC2 (Figure 6; Wilcoxon signed rank test: $V = 49$, $p < 0.05$).

Discussion

Many previous studies of *Heliconius* butterflies have described a role for visual cues in mate
finding, courtship behaviour, reproductive isolation and speciation (Crane, 1955; Brown, 1981;
Jiggins *et al.*, 2001; Kronforst *et al.*, 2006; Giraldo *et al.*, 2008; Estrada & Jiggins, 2008; Merrill
et al., 2011a). Here, we have identified compounds associated with sexually mature male wings
and described morphological structures putatively involved in pheromone release. Furthermore,
we have shown that wing extracts influence female behaviour.

We identified six potential MSP compounds from male wing extracts, namely, (*Z*)-9-
octadecenal, octadecanal, henicosane, (*Z*)-11-icosenal, icosanal and (*Z*)-13-docosenal. Five of
these are restricted to the androconial region of hindwings of 10-day old males, suggesting a
likely role in courtship. Our results are broadly comparable with another recent analysis of wing
compounds in *Heliconius* (Mérot *et al.*, 2015), although this recent study did not compare
different wing regions, or similarly aged males and females. As the study also did not use
synthesis to identify compounds, our work is highly complementary and extends their results
to confirm localization of compounds to older males. Male *Heliconius* do not become sexually
active until several days after eclosion, so the absence of these compounds from females and
younger males is strongly suggestive of a role in mating behaviours. That these five MSP
compounds are largely restricted to the hindwing androconia of mature males (Figure 5a)
suggests that pheromone storage or production is restricted to the hindwing. Trace amounts on
the forewing androconia may be due to contact in the overlapping portion of the fore- and
hindwings, and both wings may play a role in dispersal of the compounds during courtship. The
wild samples from Panama had both a greater diversity of compounds and higher concentrations
on their wings. Further work will be needed to determine whether this is characteristic of natural

populations, or reflects a difference between geographic populations of *H. melpomene*, or is
perhaps simply a result of natural inter-individual variation.

Scanning electron microscope images of the androconia support storage and/or
production of MSPs in the hindwing. The androconial region of male hindwings are equipped
with special brush-like scales (Figure 2a), which might facilitate the release of the pheromone
during courtship and are completely absent in females and any other region of male wings.
These were located primarily around and along the hindwing vein Sc+R1, similar to the
depiction in Figure 73 of Emsley's previous morphological analysis (Emsley, 1963). Similar
scales have been described from light microscopy in other *Heliconius* species, but not
previously in *H. melpomene* (Müller, 1912; Barth, 1952). The base of these special brush-like
scales was more swollen and glandular as compared to other scales (Figure 3), perhaps
indicating a role in storage or production of pheromones by these scales. Even though
Heliconius do not have the dramatic sexual dimorphism in hair pencils or brush-like androconia
seen in other butterflies, they nonetheless do possess male-specific structures likely associated
with pheromone production.

We were able to detect a significant, though subtle, difference in the behavioural
response of females towards models treated with male wing extracts as compared to an
unscented control. This indicates that females both detect compounds found on the male wings,
and alter their behaviour in response to those compounds. This supports our hypothesis that
these compounds act as a pheromone involved in courtship behaviour. Nonetheless, it remains
unclear exactly what the information is that is conveyed by these signals. The signal may
influence female courtship, although it is likely that these compounds convey complex
information about male species identity, quality, age etc. that are interpreted by females in
combination with visual and tactile cues. Recent experiments carried out in Peru also
demonstrate the role of chemical signals in species recognition between *H. melpomene* and *H.
timareta*, although these do not distinguish between the role of wing and abdominal pheromone
signals (Mérot *et al.*, 2015). Further experiments are necessary to disentangle the details of
inter-specific signaling during courtship in *Heliconius*. We are also currently unable to
determine which of the compounds identified here are biologically active. Experiments with
synthetic blends of putative MSP compounds were largely inconclusive, so are not presented
here.

The use of multiple signals is common in animal communication (Candolin, 2003).
Moths being nocturnal mainly depend upon chemical cues to attract their mates (Ando,
Inomata, & Yamamoto, 2004). On the other hand, butterflies primarily use visual cues to locate
mates (Kemp & Rutowski, 2011). It has been shown in *B. anynana* that in addition to visual
cues, chemical cues also play a role and are equally important in sexual selection by female

choice (Costanzo & Monteiro, 2007). Divergent color patterns and extensive mimicry by
different species of *Heliconius* has resulted in the use of multiple signals to maintain species
specificity. Two sympatric cryptic species that share a wing pattern, *H. melpomene malleti* and
H. timareta florencia, nonetheless show strong assortative mating. This suggests that enhanced
divergence in pheromonal signals acts as an important cue in reproductive isolation in these
species (Giraldo *et al.*, 2008). Exploring the pheromonal signals and morphological structures
involved in the process in *H. melpomene* is our first step towards understanding multimodal
signaling and its role in reproductive isolation.
360

Acknowledgments
SJ is funded by a Manmohan Singh studentship from St John's College. RMM is funded by a
Junior Research Fellowship at King's College, Cambridge. CDJ and RMM are supported by a
European Research Council grant number 339873 SpeciationGenetics. We acknowledge the
support of Caroline Nieberding and Christer Löfstedt who provided early encouragement to
pursue this project.
367

References
Ando T, Inomata S ichi, Yamamoto M. **2004**. Lepidopteran Sex Pheromones. In: Schulz
S, ed. Topics in Current Chemistry. The Chemistry of Pheromones and Other
Semiochemicals I. Springer Berlin Heidelberg, 51–96.

Bacquet PMB, Brattström O, Wang HL, Allen CE, Löfstedt C, Brakefield PM,
Nieberding CM. **2015**. Selection on male sex pheromone composition contributes to
butterfly reproductive isolation. *Proceedings. Biological Sciences / The Royal Society* **282**:
20142734.

Barth R. **1952**. Os órgãos odoriferos masculinos de alguns Heliconiinae do Brasil.
Memórias do Instituto Oswaldo Cruz **50**: 355–367.

Bates HW. **1862**. Contributions to an insect fauna of the Amazon valley. Lepidoptera:
Heliconidae. *Transactions of the Linnean Society of London* **23**: 495–566.

Becker HG, Beckert R. **1993**. *Organikum-Organisch-Chemisches Praktikum*. Wiley-VCH
Weinheim.

Brower LP, Jones MA. **1965**. Precourtship interaction of wing and abdominal sex
glands in male Danaus butterflies. *Proceedings of the Royal Entomological Society of*
London. Series A, General Entomology **40**: 147–151.

Brown KS. **1981**. The Biology of Heliconius and Related Genera. *Annual Review of*
Entomology **26**: 427–456.

Buser HR, Arn H, Guerin P, Rauscher S. **1983**. Determination of double bond position
in mono-unsaturated acetates by mass spectrometry of dimethyl disulfide adducts.
Analytical Chemistry **55**: 818–822.

Candolin U. **2003**. The use of multiple cues in mate choice. *Biological Reviews* **78**: 575–
595.

Costanzo K, Monteiro A. **2007**. The use of chemical and visual cues in female choice in
the butterfly *Bicyclus anynana*. *Proceedings of the Royal Society of London B: Biological*
Sciences **274**: 845–851.

Crane J. **1955**. Imaginal behaviour of a Trinidad butterfly, *Heliconius erato hydara*
Hewitson, with special reference to the social use of color. *Zoologica, New York* **40**: 167–
196.

Emsley MG. **1963**. A morphological study of imagine Heliconiinae (Lep., Nymphalidae)
with a consideration of the evolutionary relationships within the group. *Zoologica, New*
York **48**: 85–130.

Estrada C, Yildizhan S, Schulz S, Gilbert LE. **2010**. Sex-specific chemical cues from
immatures facilitate the evolution of mate guarding in *Heliconius* butterflies. *Proceedings*
of The Royal Society. Biological Sciences **277**: 407–413.

Estrada C, Gilbert LE. **2010**. Host plants and immatures as mate-searching cues in
Heliconius butterflies. *Animal Behaviour* **80**: 231–239.

Estrada C, Jiggins CD. **2008**. Interspecific sexual attraction because of convergence in
warning colouration: is there a conflict between natural and sexual selection in mimetic
species? *Journal of Evolutionary Biology* **21**: 749–60.

Finkbeiner SD, Briscoe AD, Reed RD. **2014**. Warning signals are seductive: Relative
contributions of color and pattern to predator avoidance and mate attraction in
Heliconius butterflies. *Evolution* **68**: 3410–3420.

Gilbert LE. **1976**. Postmating female odor in Heliconius butterflies: a male-contributed
antiaphrodisiac? *Science* **193**: 419–420.

Giraldo N, Salazar C, Jiggins CD, Bermingham E, Linares M. **2008**. Two sisters in the
same dress: *Heliconius* cryptic species. *BMC Evolutionary Biology* **8**: 324.

Grillet M, Dartevelle L, Ferveur JF. **2006**. A Drosophila male pheromone affects female
sexual receptivity. *Proceedings of the Royal Society of London B: Biological Sciences* **273**:
315–323.

Grula JW, McChesney JD, Taylor OR. **1980**. Aphrodisiac pheromones of the sulfur
butterfliesColias eurytheme andC. Philodice (Lepidoptera, Pieridae). *Journal of Chemical*
Ecology **6**: 241–256.

Jiggins CD, Naisbit RE, Coe RL, Mallet J. **2001**. Reproductive isolation caused by colour
pattern mimicry. *Nature* **411**: 302–305.

Jiggins CD. **2008**. Ecological Speciation in Mimetic Butterflies. *BioScience* **58**: 541–548.

Jiggins CD, Estrada C, Rodrigues A. **2004**. Mimicry and the evolution of premating
isolation in *Heliconius melpomene* Linnaeus. *Journal of Evolutionary Biology* **17**: 680–
691.

Kemp DJ, Rutowski RL. **2011**. The Role of Coloration in Mate Choice and Sexual
Interactions in Butterflies. Advances in the Study of Behavior. Elsevier, 55–92.

Kronforst MR, Young LG, Kapan DD, McNeely C, O'Neill RJ, Gilbert LE. **2006**. Linkage
of butterfly mate preference and wing color preference cue at the genomic location of
wingless. *Proceedings of the National Academy of Sciences of the United States of America*
103: 6575–6580.

Lassance JM, Groot AT, Liénard MA, Antony B, Borgwardt C, Andersson F,
Hedenström E, Heckel DG, Löfstedt C. **2010**. Allelic variation in a fatty-acyl reductase
gene causes divergence in moth sex pheromones. *Nature* **466**: 486–489.

Löfstedt C. **1993**. Moth Pheromone Genetics and Evolution. *Philosophical Transactions*
of the Royal Society of London B: Biological Sciences **340**: 167–177.

Meinwald J, Meinwald YC, Mazzocchi PH. **1969**. Sex Pheromone of the Queen
Butterfly: Chemistry. *Science* **164**: 1174–1175.

Melo MC, Salazar C, Jiggins CD, Linares M. **2009**. Assortative mating preferences
among hybrids offers a route to hybrid speciation. *Evolution* **63**: 1660–1665.

Mérot C, Frérot B, Leppik E, Joron M. **2015**. Beyond magic traits: Multimodal mating
cues in *Heliconius* butterflies. *Evolution*: n/a–n/a.

Merrill RM, Gompert Z, Dembeck LM, Kronforst MR, McMillan WO, Jiggins CD.
2011a. Mate preference across the speciation continuum in a clade of mimetic
butterflies. *Evolution* **65**: 1489–1500.

Merrill RM, Van Schooten B, Scott JA, Jiggins CD. **2011b**. Pervasive genetic
associations between traits causing reproductive isolation in Heliconius butterflies.
Proceedings of The Royal Society. Biological Sciences **278**: 511–518.

Merrill RM, Dasmahapatra KK, Davey JW, Dell'Aglio DD, Hanly JJ, Huber B, Jiggins
CD, Joron M, Kozak KM, Llaurens V, *et al.* 2015. The diversification of *Heliconius*
butterflies: what have we learned in 150 years? *Journal of Evolutionary Biology* **28**:
1417–1438.

Merrill RM, Chia A, Nadeau NJ. **2014**. Divergent warning patterns contribute to
assortative mating between incipient *Heliconius* species. *Ecology and Evolution* **4**: 911–
917.

More JD, Finney NS. **2002**. A Simple and Advantageous Protocol for the Oxidation of
Alcohols with o-Iodoxybenzoic Acid (IBX). *Organic Letters* **4**: 3001–3003.

Müller F. **1912**. X. The Scent-scales of the Male 'Maracujá butterflies'. In: Longstaff GB,
ed. Butterfly Hunting in Many Lands. New York: Longmans, Green & Co., 655–659.

Naisbit RE, Jiggins CD, Mallet J. **2001**. Disruptive sexual selection against hybrids
contributes to speciation between *Heliconius cydno* and *Heliconius melpomene*.
Proceedings of the Royal Society of London Series B-Biological Sciences **268**: 1849–1854.

Nieberding CM, de Vos H, Schneider MV, Lassance JM, Estramil N, Andersson J,
Bång J, Hedenström E, Löfstedt C, Brakefield PM. **2008**. The Male Sex Pheromone of
the Butterfly *Bicyclus anynana*: Towards an Evolutionary Analysis. *PLoS ONE* **3**: e2751.

Nieberding CM, Fischer K, Saastamoinen M, Allen CE, Wallin EA, Hedenström E,
Brakefield PM. **2012**. Cracking the olfactory code of a butterfly: the scent of ageing.
Ecology Letters **15**: 415–424.

Nishida R, Schulz S, Kim CS, Fukami H, Kuwahara Y, Honda K, Hayashi N. **1996**.
Male sex pheromone of a giant danaine butterfly, *Idea leuconoe*. *Journal of Chemical*
Ecology **22**: 949–972.

Phelan PL, Baker TC. **1987**. Evolution of male pheromones in moths: reproductive
isolation through sexual selection? *Science (New York, N.Y.)* **235**: 205–207.

Pliske TE, Eisner T. **1969**. Sex Pheromone of the Queen Butterfly: Biology. *Science* **164**:
1170–1172.

Schulz S, Estrada C, Yildizhan S, Boppré M, Gilbert LE. **2008**. An Antiaphrodisiac in
Heliconius melpomene Butterflies. *Journal of Chemical Ecology* **34**: 82–93.

Smadja C, Butlin RK. **2008**. On the scent of speciation: the chemosensory system and
its role in premating isolation. *Heredity* **102**: 77–97.

Symonds MRE, Johnson TL, Elgar MA. **2012**. Pheromone production, male abundance,
body size, and the evolution of elaborate antennae in moths. *Ecology and Evolution* **2**:
227–246.

Wyatt TD. **2003**. *Pheromones and Animal Behaviour: Communication by Smell and Taste*.
Cambridge University Press.

Wyatt TD. **2014**. *Pheromones and Animal Behavior: Chemical Signals And Signatures*.
Cambridge: Cambridge University Press.

Yildizhan S, van Loon J, Sramkova A, Ayasse M, Arsene C, ten Broeke C, Schulz S.
2009. Aphrodisiac Pheromones from the Wings of the Small Cabbage White and Large
Cabbage White Butterflies, *Pieris rapae* and *Pieris brassicae*. *ChemBioChem* **10**: 1666–
1677.

Figure 1. A) *H. melpomene malleti* male (left) and *H. melpomene plesseni* female (right). B)

Dissected wings from specimens of *H. melpomene plesseni* showing sexual dimorphism in the

androconial region, with male (left) and female (right). C) Expanded view of the androconial

region with arrow highlighting the vein Sc+R1. The pale grey-brown region in the male wing

was dissected for chemical analysis.

499
500

Figure 2: SEM image of putative androconial scales of *H. melpomene.* Scales in the region of

the hindwing vein Sc+R1 and forewing vein 1A are shown. (A) male hindwing (B) male

forewing and (C) female forewing at 500x magnification. (D) Magnified view of brush-like

structures of the special scales. Scale bars indicate 50 μm (A-C) and 2 μm (D).

-

Figure 3: SEM image of the base of a scale of hindwing androconia of *H. melpomene* (A)

scale from androconial region of male with brush-like structures. (B) scale from androconial

region of female. Scale bars indicate 10 μm.

511
512

Figure 4: Total ion chromatograms of extracts from the androconial region of an

Ecuadorian *H. melpomene* and a Panamanian *H. melpomene* male hindwing. 1: (*Z*)-9-

octadecenal; 2: octadecanal; 3: 15-, 16-, 17-methyloctadecanals, and additional

methyloctadecanal; 4: 1-octadecanol; 5: henicosane; 6: 15-, 16-, and 17-

methyloctadecan-1-ols, additional methyloctadecanol, and nonadecanal; 7: (*Z*)-11-

icosenal; 8: icosanal; 9: (*Z*)-11-icosenol; 10: tricosane; 11: (*Z*)-13-docosenal. All peaks

eluting later than 44 min are cuticular compounds consisting of larger *n*-alkanes, 2,5-

dialkyltetrahydrofurans, cholesterol or are contaminations.

Figure 5: GC/MS analyses of *H. melpomene* wing extracts showing presence of putative male

sex pheromone compounds (a) Presence of compounds in the hindwing androconial region of

five mature females, five young and five mature males, showing that compounds are largely

restricted to mature males. (b) Presence of compounds in five mature females, five young and

five mature males, showing that compounds are largely restricted to mature males. (c)

Presence of compounds in hindwing androconial region of five wild males from Panama.

A

B

C

Amount (ng/ul)

Pheromone analysis
- (Z)-11-icosenal
- (Z)-11-icosenol
- (Z)-13-docosenal
- 15-methyloctadecanal
- 15-methyloctadecanol
- 16-methyloctadecanal
- 16-methyloctadecanol
- 17-methyloctadecanal
- 17-methyloctadekanol
- icosanal
- nonadecanal
- octadecanal
- octadecanol
- unknown mehyloctadecanal
- unknown methyloctadecanol

old_male

Age and Sex

Figure 6: Differences in behavioural response for *H. melpomene* females presented with

models scented with male androconial extracts and unscented controls. Loading scores for

principal components are given in Table 1.

Table 1: Behavioural events scored for females presented with model butterflies and loading

scores for *H. melpomene* female response to differential treatments. The mean proportion of

trials in which a given behavior was observed is shown for both Control and Extract models.

The full data set is given as a supplementary file.

Observed behaviour	PC1	PC2	Ctrl	Extract
Slow and moderate wing flapping	0.15	-0.08	0.92	0.87
Flying/Flying facing towards model	-0.62	-0.12	0.45	0.41
Slow rhythmic flight	-0.41	-0.26	0.23	0.31
Wing display with abdomen normal	0.25	-0.63	0.09	0.15
Quick and jerky wing flapping	0.60	0.11	0.60	0.53
Flying away from model	-0.05	0.11	0.98	0.96
Fast erratic flight	-0.09	0.70	0.35	0.19
Wings open with abdomen erect	-0.03	0.09	0.04	0.04

545