1   **Phenotypic robustness determines genetic regulation of complex traits**

4   **Anupama Yadav, Kaustubh Dhole, Himanshu Sinha***

6   Department of Biological Sciences, Tata Institute of Fundamental Research,

7   Mumbai, 400005, India

10   *Corresponding author: hsinha@tifr.res.in

13   **Keywords:** phenotypic robustness, canalization, variance QTL, gene-environment

14   interaction, gene-gene interaction, cryptic genetic variation

ABSTRACT

A balance between phenotypic variability and robustness is crucial for populations to adapt to

multiple selection pressures. The plasticity of genetic pathways underlies this balance. We

investigated this plasticity by studying the regulation of phenotypic mean and variance in a

biparental recombinant population of *Saccharomyces cerevisiae* grown in a variety of

environments. We found that the growth of this population was well buffered in most

environments, such that majority of alleles regulated the mean value of phenotype, and only a

subset of these alleles regulated phenotypic variance. This latter class of alleles allowed the

other genetic variants to express a range of phenotypic values around a shifted mean. This

shift depends on the population and the environment, *i.e.* based on the evolutionary history of

a strain, buffering can result in either a superior or an inferior phenotype in an environment

but never both. Interestingly, intricate coupling of the genetic network regulating mean

phenotype and robustness was observed in a few environments, which highlighted the

importance of phenotypic buffering in layout of the genetic architecture. For loci regulating

variance, show a higher tendency of genetic interactions, which not only establishes a genetic

basis of release of variance, but also emphasizes the importance of mapping robustness in

understanding the network topology of complex traits. Our study demonstrates differential

robustness as one of the central mechanisms regulating variation in populations and

underlines its role in identifying missing heritability in complex phenotypes and diseases.

INTRODUCTION

The effect of genetic variation can either remain dormant or manifest itself phenotypically, a

phenomenon referred to as phenotypic capacitance (Masel and Siegal 2009; Félix and

Barkoulas 2015; Hayden *et al.* 2015). The process of revelation of phenotypic variability

(decanalization) has adaptive potential for populations in novel or stressful environments

(Masel 2013). Released mutations, often referred to as cryptic genetic variability (CGV)

(Paaby and Rockman 2014) can exhibit high phenotypic variability across multiple

environments, which provides adaptive diversity (Chevin *et al.* 2010; Geiler-Samerotte *et al.*

2013). Understanding the processes underlying phenotypic capacitance is important for

interpreting evolutionary effects of forces acting on populations, which result in missing

heritability and incomplete penetrance of complex traits and disease. Such processes may

ultimately govern the origin and development of complex genetic and molecular networks

(Siegal and Leu 2014).

Recent advances in systems biology have revealed that interpreting molecular mechanisms of evolution is the key in understanding the highly interconnected genetic map (Wagner *et al.* 2007; Costanzo *et al.* 2010). Certain genes, called hubs, show higher connectivity than others and thus holding the network together (Vidal *et al.* 2011). Deciphering the mechanisms, by which these network hubs modulate the effector genes in an environment- and genetic-background dependent manner to influence the phenotype, is the key to understanding how phenotypic capacitance and variability have created the genotype-phenotype (GP) map.

Despite being proposed over half a century ago (Jamniczky *et al.* 2010), only recently, a few studies have begun to identify possible regulators that buffer this genetic diversity (Félix and Barkoulas 2015). Evidence from various genome-wide deletion, protein-protein and transcriptional network analyses have proposed chromatin regulators as network hubs regulating phenotypic capacitance and variability (Levy and Siegal 2008; Tirosh *et al.* 2010). However, the molecular mechanisms through which these networks mediate the variability in the effector genes and generate diverse phenotypic responses are largely unknown. To date, the mechanistic bases have been identified only in a few cases, *e.g.* Hsp90, a highly conserved protein that regulates the phenotypic variance through altered protein folding (Rutherford and Lindquist 1998; Queitsch *et al.* 2002).

In this study, we have attempted to address the following two questions. How important is regulation of robustness for genetic regulation of the phenotype? And, whether perturbation of this robustness and the subsequent release of CGV are beneficial or detrimental for the population growth?

In order to understand the regulation of phenotypic variability and plasticity, it is necessary to investigate the effect of regulators on diverse mutations, an approach applied so far to only a few genes (Queitsch *et al.* 2012). For a single species, though not achievable through single gene perturbation studies, recombinant populations analyzed in various environments provide an excellent platform to study such effects (Lempe *et al.* 2013). The allelic variants that mediate how phenotypes are regulated have two potential outcomes. First, there is an effect on the 'mean' value of some readout, such as growth in different environments in the case of yeast. Mapping such an effect on population 'mean' values allows the identification of alleles that affect phenotypes independent of the genetic background. In addition, there is a second

consequence, *i.e.* an effect on the 'variance' of the readout (Hall *et al.* 2007; Lee *et al.* 2014),

which is often ignored in many studies. Alleles that cause such differences in population

'variance' have a more subtle effect, whereby genetic diversity is retained in the population

(Figure 1A). In our study, we used a synthesis of both these mapping approaches to discern

patterns of regulation of variation in a recombinant population. We find that the genetic loci,

and their networks, regulating robustness are a subset of the conventionally studied loci,

which affect phenotypic mean. However, in certain environments, based on the evolutionary

history of the parental strains, these two networks are very closely intertwined as

demonstrated by complete loss of buffering in the recombinants. Such environments highlight

the importance of mechanisms regulating robustness in the genetic architecture. Based on our

analysis, we propose differential robustness of alleles as a major contributor to antagonistic

pleiotropy. Finally, we provide evidence for genetic basis of loss of robustness and therefore

propose variance mapping as a methodology to uncover the genetic interactions regulating

complex traits.

RESULTS

High overlap between mean and variance QTL

Using a recombinant population generated from a biparental cross, we carried out linkage

mapping to identify genetic loci regulating both phenotypic mean (QTL) and phenotypic

robustness (variance QTL or vQTL) of colony size variation across 34 diverse environments

ranging from different carbon sources to oxidative and DNA damaging stress (see Methods).

To compare the genetic loci identified by QTL and vQTL mapping, the genome was binned

into 595 bins (see Methods, Tables S1, S2). The distribution of loci showing a difference in

mean only, variance (robustness) only or both, differed across environments (Figure 1B). The

majority of all loci identified affected only mean (70%) with 4% affecting only robustness.

However, a substantial 27% loci were both QTL + vQTL (Figure 1C). For these loci,

pleiotropic hotspots were identified, which regulated mean, robustness or both across

multiple environments (Table S2).

Environment dependent regulation of phenotypic mean and robustness

While there was no association between number of QTL and vQTL identified across

environments, there was a negative correlation ($r^2 = -0.4$) between the number of QTL and

the number of loci which were both QTL + vQTL associated across environments. This

indicated that there is a decrease in the number of loci with independent effects as the number
of loci regulating variance increase. Based on the overlap between mean and variance loci,
we could divide the environments into three categories – (i) No overlap between QTL and
vQTL: this indicated that growth was tightly buffered in the recombinants and while the
genetic variants affected the mean of the phenotype, robustness was maintained. These
environments were enriched in various carbon sources. (ii) Partial overlap between QTL and
vQTL: some loci showed difference in robustness but the majority affected only the mean.
(iii) Almost complete overlap: in these environments, all loci which affected the phenotypic
mean resulted in varying degrees of loss of robustness. In such environments only two states
were possible, *i.e.* either the allele contributed to the buffered state or it resulted in loss of
robustness. Growth in such environments emphasized the role of robustness in regulating
phenotypic variation.
133
Release of phenotypic variance is directional
A high overlap of regulation of mean and robustness indicated that the loci regulating
phenotypic variability were a subset of the loci that have been implicated to be associated
with a shift in the mean upon release of variation. While pleiotropic hotspots regulated the
mean and the variance across multiple environments, no discernible pattern was observed in
the directionality of this effect, except when considering each environment independently
(Figure 2A). In the majority of environments (26/33), we observed a strong correlation
between the effect of the allele on the mean and the variance for the significant loci. One half
of the environments (13/26) showed a strong positive correlation ($r^2 > +0.5$) indicating that
loss of robustness was, on an average, advantageous, and the other half showed a negative
correlation ($r^2 < -0.5$), *i.e.* loss of robustness was detrimental for the population (Figure 2B).
145
Differential regulation of robustness as a major contributor to gene-environment
interaction
Pleiotropic loci often have antagonistic mean effects across environments (Yadav *et al.*
2015). While loci regulating robustness show high environment specificity, we asked if it is
possible for a locus to result in robustness in one environment and decanalization in the other
by comparing covariance of alleles across environmental pairs (Haber and Dworkin 2015).
Forty seven bins were identified that had a significant effect in two or more environments.
Eighteen hotspots showed significant difference in the covariance with 10 being significant
across multiple pairs of environments (Table S3). Unlike mean effects, high consistency was

observed in allelic robustness of a single hotspot across environmental pairs, *i.e.* within a
hotspot the same allele showed a low covariance (high buffering or canalized) across multiple
environmental pairs and consequently the other allele consistently showed a high covariance
(decanalization). This consistency in buffering, when coupled with high environment
specificity of directionality (advantageous or deterimental but not both) of release of
variance, explains the abundant antagonistic pleiotropy observed in mean effects of these
pleiotropic hotspots (Figure S1) (Yadav *et al.* 2015). As suggested by our previous analysis,
the environment would determine whether the accumulated mutations would be beneficial or
detrimental to the phenotype. In conclusion, this covariance analysis shows that differential
regulation of robustness is a major contributor to gene-environment interactions, especially
antagonistic pleiotropy.

Release of variance is genetic in nature
While most of the polymorphisms in this population were neutral, some showed their direct
effects and others were hidden showing their effects only under certain environmental and
genetic perturbations. To estimate the extent of genetic basis of loss of robustness, two-QTL
interaction mapping was performed by considering the effects of only significant QTL and
vQTL loci (see Methods). Amongst the loci showing two-QTL interactions, 10% had only
single QTL effects, whereas 80% were either vQTL or both (Table S4) indicating that vQTL
were involved in a greater fraction of genetic interactions. These interacting loci showed
stronger effects in the decanalized allele than the buffered one (Figure 3B, 3C). In addition to
showing that there is a genetic basis regulating the increase in variability, this observation
also suggests that altered variance is a good predictor of extent of genetic interactions.

In our analysis, two kinds of interaction hubs were identified: *within environment interaction*
hubs, where a locus interacted with multiple loci to regulate phenotypic variation in a given
environment; and *across environment interaction hubs*, where a pleiotropic locus interacted
with different loci across several environments to regulate phenotypic variation (Figure 4A).
Overall, our observations support a modular structure of regulation of phenotypic robustness.
In *within environment hubs*, some regulators controlled multiple effector genes in a single
environment. In the presence of a decanalized allele, this control is disrupted, resulting in a
high variance in the same environment. These hubs were identified in the environments
where most loci independently regulated both mean and robustness. On the other hand, in
across environment hubs, a different set of effectors functionally regulated the phenotype

across distinct environments. This suggests that allelic variations in such hubs will cause
changes in variability and control growth across environments.
191
Network of loci regulate phenotypic buffering
Studies showing high interconnectivity and redundancy in the genetic networks argue for a
crosstalk among multiple regulators of phenotypic robustness (Kafri *et al.* 2009; Li *et al.*
2010; Costanzo *et al.* 2010). At a population level, this would translate to maintenance of
robustness by a network of loci instead of a single locus. To identify such interactions
regulating phenotypic variance, we carried out a two-vQTL interaction mapping using
candidate loci (Table S4). To our knowledge, this is the first time that such an analysis has
been performed. A large number of two-vQTL interactions were identified, many more than
would be expected from proportion of single vQTL analysis, emphasizing the functional
relevance of co-regulating variance in population dynamics (Figure 4B). Interestingly, in
addition to just the differential regulation of variance, this interaction mapping also identified
two-locus pairs which had buffered variance in 3 out of their 4 allelic combinations (Figure
3E, 3F). Indeed, in environments showing high overlap between mean and variance QTL,
multiple such pairs were identified (Figure 4C, 4D), indicating that there was a highly
connected, redundant multi-locus network that maintained phenotypic robustness. We also
observed that it is only under very specific allelic combinations that this network was
disrupted to reveal phenotypic variance.
209
210
DISCUSSION
Populations balance maintaining a robust phenotype unaffected by genetic perturbation and
active accumulation of mutations as potential for evolution. Therefore, it is a reasonable
assumption that regulation and mediation of phenotypic variability under different selection
pressures plays a significant role in determining the networks governing complex traits
(Landry and Rifkin 2010; Geiler-Samerotte *et al.* 2013).
217
Once a population adapts to a particular environment, stabilizing selection acts to maintain
the mean at a specific fitness value. While phenotypically uniform, the population needs to
maintain genetic diversity to maintain adaptive potential in the face of a change in selection
pressure (Hartman *et al.* 2001; Gibson and Dworkin 2004). This requires the presence of
genetic hubs that may sense environmental or genetic perturbations, transducing their

responses to the downstream effectors, and therefore, regulate the release of phenotypic

capacitance (Figure 5) (Carlborg *et al.* 2006; Le Rouzic and Carlborg 2008; Hayden *et al.*

2015). This release of phenotypic diversity can be a result of encountering a novel or a stress

environment as well as a result of a change in a regulator (Figure 5) (McGuigan and Sgrò

2009).

Our study showed that high-resolution QTL mapping could be used to distinguish between

genetic loci which have a 'mean' effect on the phenotype and those which regulate

phenotypic robustness. While various studies have compared loci regulating mean and

variance in different organisms in the past they were limited either by sample size or range of

phenotypes considered, thus showing varying degrees of overlap between loci regulating

mean and variance (Hall *et al.* 2007; Sangster *et al.* 2008; Lee *et al.* 2014). Our study shows

empirical evidence that regulation of robustness forms the basis of genetic networks

regulating a phenotype. Growth in an environment can be regulated in two ways: either most

of the loci affect the mean with only a subset affecting robustness, or almost all loci affect

both robustness and the mean. In addition to the nature of the environment, this

categorization would depend on the evolutionary history of the parental strains.

In our study, presence of majority of alleles differentially affected the phenotype mean

without perturbing the buffered state, indicates that the network maintaining robustness is

extensive and the genetic variation in the population is not sufficient to perturb it.

Alternatively, for the environments which parental strains would have encountered during

evolution and were similarly adapted to, *viz.*, various carbon sources, diverse parental alleles

maintain similar robust state. On the other hand, in a few environments, all alleles that

affected mean also affected robustness. This high coupling indicated that robustness was

perturbed in the segregating population such that every allele that affected the phenotype also

perturbed its robustness. This could be possible if either one of the strains was not exposed to

the environment during its evolution or the strains were buffered through incompatible

independent mechanisms. In such environments, the network topology and role of robustness

in their layout becomes apparent.

Independent of the kind of environment, strong directionality was observed in the release of

cryptic genetic variation. This directionality is likely the frozen-in result of the evolutionary

history of the strains or the phenotype (Taute *et al.* 2014). For a stabilized population in a

stressful or a novel environment, the population has an evolutionary imperative to release phenotypic variability, resulting in a positive association between the mean and the variance (Le Rouzic and Carlborg 2008; McGuigan and Sgrò 2009). On the other hand, in an adapted environment, release of variability will be detrimental and would be visible as a negative association between the mean and the variance (see Figure 2A). We observed that while genetic loci showed antagonism in their mean effects across environments, they showed high consistency in directionality of their buffering abilities. Environment-dependent effect of release of variance as well as high consistency in buffering indicates that a locus that buffers the phenotypic capacitance will, as a result, have an antagonistic effect on the population mean in an environment-dependent manner. Additionally, this consistency of effect on robustness is in accordance with the known molecular mechanisms. An allele that perturbs protein folding or chromatin silencing (Taylor and Ehrenreich 2015) will show similar molecular behaviour across environments, whether the release will be beneficial or detrimental will be determined by the phenotype. In summary, our study indicates that altered buffering is a potential cause of antagonistic pleiotropy. While we cannot comment on the genes mediating this phenotypic buffering, our study shows that in different environments, different genetic loci or their hubs regulate robustness. This along with identification of environments with high overlap between regulation of mean and robustness support a scenario where multiple genes are involved in buffering the phenotype and may possibly play a crucial role in layout of genetic networks.

Our variance mapping using biparental recombinant populations adds to the understanding of fundamental questions like missing heritability (Manolio *et al.* 2009; Eichler *et al.* 2010). High population dependence and incomplete penetrance impede the identification of disease-causing alleles (Zuk *et al.* 2012; Mackay 2014). We propose altered phenotypic buffering as a possible causative mechanism behind missing heritability (Queitsch *et al.* 2012). Effect of a disease-causing locus will be neutralized in the presence of an allele that buffers the phenotypic variance, whereas it will be causative in the decanalized allele (Figure 5). While genome-wide association studies suffer from a lack of power to identify such epistatic interactions, analyzing the variance of a population along with the mean can act as a robust refinement to narrow down the possible interactors of disease-causing alleles.

METHODS

Dataset

The raw growth data analysed in this study was derived from a study by Bloom *et al*. (2013),

in which the experimental procedures are described in detail. The data we used was generated

for 1,008 segregants derived from a cross between *S. cerevisiae* strains BY (a laboratory

strain) and RM11-1a (a wine isolate, indicated as RM). These segregants were grown in 46

different conditions. Of these, we studied 34 conditions (see Table S2, see File S1 for more

information).

QTL and vQTL mapping

The single environment QTL and two-QTL mapping was carried out as described previously

(Bhatia *et al.* 2014).

To estimate the difference in phenotypic variance between the two genotypic groups, *i.e.* to

identify vQTL in each environment, the standard Brown-Forsythe (BF) statistic (Lee *et al.*

2014) and the corresponding LOD score were calculated for each genetic marker in each

environment (see File S1). The BF test is equivalent to an F-test performed on the deviations

of the phenotypic values from their respective genotypic medians (or the means). Hence,

under the alternative hypothesis, the phenotypes of the two alleles reveal a difference in the

variance. As a result, the corresponding LOD scores indicate markers responsible for genetic

canalization defined as variance-QTL (vQTL).

At a particular marker, let z_{ij} be the absolute deviation of segregant i's phenotypic value y_{ij}

from its genotypic mean $\tilde{y}_j$ where j can take two values ($j=1$: BY allele and $j=2$: RM

allele).

$$z_{ij} = \left| y_{ij} - \tilde{y}_j \right|$$

Then BF statistic for that marker can be computed as follows:

$$F = \frac{(N-p)}{(p-1)} \frac{\sum_{j=1}^{p} n_j (\tilde{z}_{.j} - \tilde{z}_{..})^2}{\sum_{j=1}^{p} \sum_{i=1}^{n_j} (z_{ij} - \tilde{z}_{.j})^2}$$

Here, N is the total number of segregants, n_1 and n_2 are the number of segregants having

the BY and RM allele respectively ($p = 2$). In order to estimate the effects of vQTL in the

same order as in QTL, LOD scores were computed as described previously (Broman and Sen

2009).

To establish the statistical significance of the putative QTL and vQTL, P-values were

computed using a genome-wide permutation test of 1,000 permutations, where the null

distribution consisted of the highest genome-wide LOD score obtained from each

permutation. A LOD cut off of 3.0 and a P-value cut off of 0.01 was considered.

To estimate pleiotropy, we divided the genome into 20kb non-overlapping bins (Table S1).

Bins containing two or more QTL or vQTL significant (P-value < 0.01) in different

environments were considered as pleiotropic bins. The first markers of each of these

pleiotropic bins, used as representative of the bins were collated to represent the set of

pleiotropic markers (Table S1, S2).

Apart from the standard two-QTL mapping described previously (Bhatia *et al.* 2014), we

mapped variance-controlled interactions, *i.e.* a two-vQTL interaction, which occurs when the

phenotypic variance at one locus depends on the genotype at another locus. To increase

power to identify two-QTL and two-vQTL interactions, for environment specific, targeted

multi-QTL mapping, genetic loci significant in either QTL or vQTL or both mappings were

collated for each environment (Table S2). Additionally the size of the bin was increased from

20kb to 50kb for the same. This collated set of environment-specific loci was tested for both,

two-QTL and two-vQTL interactions in their respective environments. A total of 47 two-

QTL interactions each significant (P-value < 0.001) in at least one of the 34 environments

were obtained. The P-values were computed using a permutation test of 10,000 permutations

with the phenotype data shuffled relative to the genotype data.

Covariance across environmental pairs

To assess the differential covariance of a locus across multiple environments, we considered

the collated set of pleiotropic markers for our study (Table S3). To quantify the differential

covariance across a pair of environments, a Deming regression was calculated between the

phenotype values of the chosen pair of environments for each allele, using R package 'mcr'.

Deming regression, which minimizes errors in multiple dimensions simultaneously, served as
a suitable measurement error model for assessing buffering across two or more environments.
For every possible environment pair, a t-test was performed between the deviations of the
phenotypic values from the Deming fit of the BY and RM allele (P-value < 0.05).
357
358

359 **ACKNOWLEDGEMENTS**
360 We thank Gyan Bhanot for critical reading of the manuscript. This research was supported by
361 Tata Institute of Fundamental Research intramural funds (HS). The funders had no role in
362 study design, data collection and analysis, decision to publish, or preparation of the
363 manuscript.
364
365
366 **AUTHOR CONTRIBUTIONS**
367 Conceived and designed analysis: AY HS. Analysed data: KD AY HS. Wrote the paper: AY
368 HS.

369
370

FIGURE LEGENDS
Figure 1: QTL and vQTL
(A) Schematic showing three categories of QTL mapped. QTL has significantly different
allelic mean but non-significant different allelic variance; vQTL has non-significant mean
difference but significantly different variance; QTL+vQTL has both significant different
allelic mean and variance.
(B) Distribution of QTL mapped in each environment as QTL (red), vQTL (blue) and
QTL+vQTL (black) in the segregating population. The y-axis is proportion of QTL in each
category (see Results). The x-axis is arranged by increasing proportion of QTL and the left
group has the highest proportion of QTL + vQTL. See Table S2 for details.
(C) Total distribution of QTL mapped across 34 environments as QTL (red), vQTL (blue)
and QTL+vQTL (black) in the segregating population.
383

Figure 2: Directionality of alleles in QTL+vQTL category
(A) Representative frequency distributions of three QTL markers showing directionality of
variance release. Blue distribution is of segregants with BY allele and red is for RM

segregants. 4NQO [chrXII (207,340)] marker shows a negative release of variance of BY

allele (with RM mean > BY mean); Lactose [chrXIV (376,315)] marker shows equal variance

of the two alleles; $MgSO_4$ [chrVII (187,538)] marker shows a positive release of variance of

BY allele (with RM mean < BY mean). The QTL are indicated as chromosome number

followed by marker position in bp within brackets. Color bar on the top of each distribution

represents the three classes of correlations between mean and variance.

(B) Correlation between mean and variance of all QTL in each environment. Correlations less

than -0.5 (red) represent environments that have a negative release of variance of BY allele.

Correlations more than +0.5 (yellow) represent environments that have a positive release of

variance of BY allele. Environments with no directional release of variance (correlations

between ±0.5) are represented as blue. The y-axis is correlation between mean and variance;

dashed lines show ±0.5 correlation value.

Figure 3: Representative covariance and two-QTL interactions

(A) Covariance of normalized growth phenotype of BY (blue) and RM (red) segregants for

chrXIV (368,185) marker in 4-HBA and Galactose. Mean and variance of each allele in each

environment is indicated in the box.

(B) Two-QTL interaction between chrXIV (368,185)-chrXIII (46,758) in 4-HBA.

(C) Two-QTL interaction between chrXIV (368,165)-chrXV (555,452) in Galactose.

(D) Covariance of normalized growth phenotype of BY (blue) and RM (red) segregants for

chrXII (649,260) marker in $MgSO_4$ and Cu (copper). Mean and variance of each allele in

each environment is indicated in the box.

(E) Two-QTL interaction between chrXII (649,260)-chrVII (187,538) in $MgSO_4$.

(F) Two-QTL interaction between chrXII (649,260)-chrVIII (208,560) in Cu.

The QTL is indicated as a chromosome number followed by the marker position in bp within

brackets (a convention followed in all QTL plots). For (A, D), the axes are normalized

growth of segregants in the two environments indicated. For (B, C, E, F), the x-axis is

normalized growth of segregants in the environment and the y-axis of number of segregants.

Dash lines in segregant distributions indicate the means of the distributions. The biallelic

marker segregant distributions (in the QTL marker order written above the plots) are

indicated as BB (light blue), BR (light brown), RB (dark green) and RR (purple). Inset plots

show the average distributions of the first marker (BY (blue) and RM (red) alleles). See Table

S3, S4 for details.

Figure 4: Two-QTL and two-vQTL interactions

(A) Two-QTL and two-vQTL interactions between various markers shown as connected

links. The chrII (245,879) marker (red) has multiple two-QTL interactions (deep red) for

growth in same environment (Congo red). The chrXIV (466,590) marker (green) has 6

environment-specific two-QTL interactions (4NQO= dark green, Formamide= deep purple,

IAA= light blue, Lithium chloride= deep orange, Trehalose= deep yellow, Xylose= orange).

Other two-QTL interactions are indicated as light grey links.

(B) All two-QTL (red), two-vQTL (blue) and two-QTL+vQTL (black) interactions across all

environments.

The figures were made using Circos (Krzywinski *et al.* 2009). See Table S4 for data.

(C) Scatter plots showing examples of two-QTL interactions of four markers [chrIII

(191,928), chrIV (997,621), chrVIII (101,016), chrXIV (466,105)] in Indoleacetic acid. The

biallelic marker segregant distributions (in the QTL marker order written above the plots) are

indicated as BB (red), BR (yellow), RB (green) and RR (blue) on x-axis. Mean and variance

of each allelic pair is indicated in the box with allelic pair with most variance indicated in red.

The y-axis is normalized growth phenotype.

(D) Schematic representation of network of four loci (indicated in (C) above) maintaining

phenotypic robustness in Indoleacetic acid.

Figure 5: A schematic showing environment and genetic background dependent loss of

buffering

A regulator (red node) of robustness buffers genetic variability (blue nodes), which results in

a robust population phenotype. This buffering is lost in either a differential allele (green

node) of the regulator or a novel environment (red distribution), thus releasing cryptic genetic

variability (yellow node), which in favourable cases results in better adaptation in the novel

environment, or in unfavourable cases manifests as a detrimental phenotype or a disease.

SUPPORTING INFORMATION

File S1: Scripts and datasets for QTL, vQTL, and two-QTL, two-vQTL interaction mapping.

Table S1: Classification of chromosomal markers into bins.

Table S2: List of markers significant for QTL and vQTL mapping. F and BF statistics along

with their P-values along with the categorization as only vQTL, only QTL and both.

Table S3: Covariance analysis of selected bins. T-test of Deming regression between the two

alleles across environmental pairs along with its mean and variance for each allele, mean and

variance of each allele independently in both environments.

Table S4: List of markers significant for two-QTL and two-vQTL mapping.

Figure S1: Covariance of normalized growth phenotype of BY (red) and RM (blue) allele

chrXIII (45,801) marker in Paraquat and Copper. Mean and variance of each allele in each

environment is indicated in the box.

REFERENCES

Bhatia A., Yadav A., Gagneur J., Zhu C., Steinmetz L. M., Bhanot G., Sinha H., 2014 Yeast
Growth Plasticity Is Regulated by Environment-Specific Multi-QTL Interactions. G3
(Bethesda) **4**: 769–777.

Bloom J. S., Ehrenreich I. M., Loo W. T., Lite T.-L. V., Kruglyak L., 2013 Finding the
sources of missing heritability in a yeast cross. Nature **494**: 234–237.

Broman K. W., Sen S., 2009 *A Guide to QTL Mapping with R/qtl*. Spinger, New York.

Carlborg O., Jacobsson L., Ahgren P., Siegel P., Andersson L., 2006 Epistasis and the release
of genetic variation during long-term selection. Nat Genet **38**: 418–420.

Chevin L.-M., Lande R., Mace G. M., 2010 Adaptation, Plasticity, and Extinction in a
Changing Environment: Towards a Predictive Theory (JG Kingsolver, Ed.). PLoS Biol **8**:
e1000357.

Costanzo M., Baryshnikova A., Bellay J., Kim Y., Spear E. D., Sevier C. S., Ding H., Koh J.
L. Y., Toufighi K., Mostafavi S., Prinz J., St Onge R. P., VanderSluis B., Makhnevych
T., Vizeacoumar F. J., Alizadeh S., Bahr S., Brost R. L., Chen Y., Cokol M., Deshpande
R., Li Z., Lin Z.-Y., Liang W., Marback M., Paw J., San Luis B.-J., Shuteriqi E., Tong A.
H. Y., van Dyk N., Wallace I. M., Whitney J. A., Weirauch M. T., Zhong G., Zhu H.,
Houry W. A., Brudno M., Ragibizadeh S., Papp B., Pal C., Roth F. P., Giaever G.,
Nislow C., Troyanskaya O. G., Bussey H., Bader G. D., Gingras A.-C., Morris Q. D.,
Kim P. M., Kaiser C. A., Myers C. L., Andrews B. J., Boone C., 2010 The genetic
landscape of a cell. Science (New York, NY) **327**: 425–431.

Eichler E. E., Flint J., Gibson G., Kong A., Leal S. M., Moore J. H., Nadeau J. H., 2010
Missing heritability and strategies for finding the underlying causes of complex disease.
Nat Rev Genet **11**: 446–450.

Félix M.-A., Barkoulas M., 2015 Pervasive robustness in biological systems. Nat Rev Genet
16: 483–496.

Geiler-Samerotte K. A., Bauer C. R., Li S., Ziv N., Gresham D., Siegal M. L., 2013 The
details in the distributions: why and how to study phenotypic variability. Curr Opin

Biotechnol **24**: 752–759.

Gibson G., Dworkin I., 2004 Uncovering cryptic genetic variation. Nat Rev Genet **5**: 681–
690.

Haber A., Dworkin I., 2015 *Dis-integrating the fly: A mutational perspective on phenotypic*
integration and covariation. Cold Spring Harbor Labs Journals.

Hall M. C., Dworkin I., Ungerer M. C., Purugganan M., 2007 Genetics of
microenvironmental canalization in Arabidopsis thaliana. Proc Natl Acad Sci U S A **104**:
13717–13722.

Hartman J. L., Garvik B., Hartwell L., 2001 Principles for the buffering of genetic variation.
Science (New York, NY) **291**: 1001–1004.

Hayden E. J., Bendixsen D. P., Wagner A., 2015 Intramolecular phenotypic capacitance in a
modular RNA molecule. Proc Natl Acad Sci USA **112**: 12444–12449.

Jamniczky H. A., Boughner J. C., Rolian C., Gonzalez P. N., Powell C. D., Schmidt E. J.,
Parsons T. E., Bookstein F. L., Hallgrímsson B., 2010 Rediscovering Waddington in the
post-genomic age. Bioessays **32**: 553–558.

Kafri R., Springer M., Pilpel Y., 2009 Genetic Redundancy: New Tricks for Old Genes.
CELL **136**: 389–392.

Krzywinski M. I., Schein J. E., Birol I., Connors J., Gascoyne R., Horsman D., Jones S. J.,
Marra M. A., 2009 Circos: An information aesthetic for comparative genomics. Genome
Res **19**: 1639–1645.

Landry C. R., Rifkin S. A., 2010 Chromatin regulators shape the genotype–phenotype
map. Mol Syst Biol **6**: 1–2.

Le Rouzic A., Carlborg O., 2008 Evolutionary potential of hidden genetic variation. Trends
in Ecology & Evolution **23**: 33–37.

Lee C.-R., Anderson J. T., Mitchell-Olds T., 2014 Unifying Genetic Canalization, Genetic
Constraint, and Genotype-by-Environment Interaction: QTL by Genomic Background by
Environment Interaction of Flowering Time in Boechera stricta (K Bomblies, Ed.). PLoS
Genet **10**: e1004727.

Lempe J., Lachowiec J., Sullivan A. M., Queitsch C., 2013 Molecular mechanisms of
robustness in plants. Current Opinion in Plant Biology **16**: 62–69.

Levy S., Siegal M., 2008 Network Hubs Buffer Environmental Variation in Saccharomyces
cerevisiae (A Levchenko, Ed.). PLoS Biol **6**: e264.

Li J., Yuan Z., Zhang Z., 2010 The cellular robustness by genetic redundancy in budding
yeast (J Zhang, Ed.). PLoS Genet **6**: e1001187.

Mackay T. F. C., 2014 Epistasis and quantitative traits: using model organisms to study gene-
gene interactions. Nat Rev Genet **15**: 22–33.

Manolio T. A., Collins F. S., Cox N. J., Goldstein D. B., Hindorff L. A., Hunter D. J.,
McCarthy M. I., Ramos E. M., Cardon L. R., Chakravarti A., Cho J. H., Guttmacher A.
E., Kong A., Kruglyak L., Mardis E., Rotimi C. N., Slatkin M., Valle D., Whittemore A.
S., Boehnke M., Clark A. G., Eichler E. E., Gibson G., Haines J. L., Mackay T. F. C.,
Mccarroll S. A., Visscher P. M., 2009 Finding the missing heritability of complex
diseases. Nature **461**: 747–753.

Masel J., 2013 Q&A: Evolutionary capacitance. BMC Biology **11**: 103.

Masel J., Siegal M. L., 2009 Robustness: mechanisms and consequences. Trends Genet **25**:
395–403.

McGuigan K., Sgrò C. M., 2009 Evolutionary consequences of cryptic genetic variation.
Trends in Ecology & Evolution **24**: 305–311.

Paaby A. B., Rockman M. V., 2014 Cryptic genetic variation:evolution's hidden substrate.
Nat Rev Genet **15**: 247–258.

Queitsch C., Carlson K. D., Girirajan S., 2012 Lessons from Model Organisms: Phenotypic
Robustness and Missing Heritability in Complex Disease (SM Rosenberg, Ed.). PLoS
Genet **8**: e1003041.

Queitsch C., Sangster T. A., Lindquist S., 2002 Hsp90 as a capacitor of phenotypic variation.
Nature **417**: 618–624.

Rutherford S. L., Lindquist S., 1998 Hsp90 as a capacitor for morphological evolution.
Nature **396**: 336–342.

Sangster T., Salathia N., Undurraga S., Milo R., Schellenberg K., Lindquist S., Queitsch C.,
2008 HSP90 affects the expression of genetic variation and developmental stability in
quantitative traits. Proc Natl Acad Sci USA **105**: –2968.

Siegal M. L., Leu J.-Y., 2014 On the Nature and Evolutionary Impact of Phenotypic
Robustness Mechanisms. Annu. Rev. Ecol. Evol. Syst. **45**: 495–517.

Taute K. M., Gude S., Nghe P., Tans S. J., 2014 Evolutionary constraints in variable
environments, from proteins to networks. Trends Genet **30**: 192–198.

Taylor M. B., Ehrenreich I. M., 2015 Transcriptional Derepression Uncovers Cryptic Higher-
Order Genetic Interactions. (P Wittkopp, Ed.). PLoS Genet **11**: e1005606.

Tirosh I., Reikhav S., Sigal N., Assia Y., Barkai N., 2010 Chromatin regulators as capacitors
of interspecies variations in gene expression. Mol Syst Biol **6**: 435.

Vidal M., Cusick M. E., Barabási A.-L., 2011 Interactome networks and human disease.
CELL **144**: 986–998.

Wagner G. P., Pavlicev M., Cheverud J. M., 2007 The road to modularity. Nat Rev Genet **8**:
921–931.

Yadav A., Radhakrishnan A., Bhanot G., Sinha H., 2015 Differential Regulation of
Antagonistic Pleiotropy in Synthetic and Natural Populations Suggests Its Role in

Adaptation. G3 (Bethesda) **5**: 699–709.

Zuk O., Hechter E., Sunyaev S. R., Lander E. S., 2012 The mystery of missing heritability:
Genetic interactions create phantom heritability. Proc Natl Acad Sci USA **109**: 1193–
1198.

Figure 1

Figure 2

A

4NQO [chrXII (207,340)]

Lactose [chrXIV (376,315)]

MgSO₄ [chrVII (187,538)]

B

Figure 3

Figure 4

Figure 5

Figure S1

chrXIII (45,801) marker

BY allele

$\bar{x}_{BY}(Paraquat) = -0.2$
$\sigma^2_{BY}(Paraquat) = 0.82$
$\bar{x}_{BY}(Cu) = 0.34$
$\sigma^2_{BY}(Cu) = 0.52$

RM allele

$\bar{x}_{RM}(Paraquat) = 0.2$
$\sigma^2_{RM}(Paraquat) = 1.08$
$\bar{x}_{RM}(Cu) = -0.33$
$\sigma^2_{RM}(Cu) = 1.26$

Normalized growth in Paraquat

Normalized growth in Cu