1 BIOLOGICAL SCIENCES: Biophysics and Computational Biology

Measuring and modeling diffuse scattering in protein X-ray crystallography

short title: Diffuse scattering in protein X-ray crystallography

Andrew H. Van Benschoten[1], Lin Liu[1], Ana Gonzalez[2], Aaron S. Brewster[3], Nicholas K. Sauter[3],
James S. Fraser[1,*], Michael E. Wall[4,*]
1 - Dept. of Bioengineering and Therapeutic Sciences, University of California, San Francisco, San
Francisco, CA 11 94158
2 - Stanford Synchrotron Radiation Lightsource, SLAC National Accelerator Laboratory, Menlo Park, CA
94025
3 - Molecular Biophysics & Integrated Bioimaging Division, Lawrence Berkeley National Laboratory,
Berkeley, CA 94720
4 - Computer, Computational, and Statistical Sciences Division, Los Alamos National Laboratory, Los
Alamos, NM, 87545

18
19 *correspondence: jfraser@fraserlab.com, mewall@lanl.gov

21 Keywords: protein dynamics, normal modes, structural biology, diffuse scattering

24
25
26

Abstract

X-ray diffraction has the potential to provide rich information about the structural dynamics of macromolecules. To realize this potential, both Bragg scattering, which is currently used to derive macromolecular structures, and diffuse scattering, which reports on correlations in charge density variations must be measured. Until now measurement of diffuse scattering from protein crystals has been scarce, due to the extra effort of collecting diffuse data. Here, we present three-dimensional measurements of diffuse intensity collected from crystals of the enzymes cyclophilin A and trypsin. The measurements were obtained from the same X-ray diffraction images as the Bragg data, using best practices for standard data collection. To model the underlying dynamics in a practical way that could be used during structure refinement, we tested Translation-Libration-Screw (TLS), Liquid-Like Motions (LLM), and coarse-grained Normal Modes (NM) models of protein motions. The LLM model provides a global picture of motions and were refined against the diffuse data, while the TLS and NM models provide more detailed and distinct descriptions of atom displacements, and only used information from the Bragg data. Whereas different TLS groupings yielded similar Bragg intensities, they yielded different diffuse intensities, none of which agreed well with the data. In contrast, both the LLM and NM models agreed substantially with the diffuse data. These results demonstrate a realistic path to increase the number of diffuse datasets available to the wider biosciences community and indicate that NM-based refinement can generate dynamics-inspired structural models that simultaneously agree with both Bragg and diffuse scattering.

Significance

The structural details of protein motions are critical to understanding many biological processes, but they are often hidden to conventional biophysical techniques. Diffuse X-ray scattering can reveal details of the correlated movements between atoms; however, the data collection historically has required extra effort and dedicated experimental protocols. We have measured three-dimensional diffuse intensities in X-ray diffraction from CypA and trypsin crystals using standard crystallographic data collection techniques. Analysis of the resulting data is consistent with the protein motions resembling diffusion in a liquid or vibrations of a soft solid. Our results show that using diffuse scattering to model protein motions can become a component of routine crystallographic analysis through the extension of commonplace methods.

\body

Introduction:

X-ray crystallography can be a key tool for elucidating the structural basis of protein motions that play critical roles in enzymatic reactions, protein-protein interactions and signaling cascades (van den Bedem and Fraser, 2015). X-ray diffraction yields an ensemble-averaged picture of the protein structure: each photon simultaneously probes multiple unit cells that can vary due to internal rearrangements or changes to the crystal lattice. Bragg analysis of X-ray diffraction only yields the mean charge density of the unit cell, however, which fundamentally limits the information that can be obtained about protein dynamics (Clarage and Phillips, 1997; Keen and Goodwin, 2015).

A key limitation inherent in Bragg analysis is that alternative models with different correlations between atomic motions can yield the same mean charge density (Kuzmanic et al., 2011). The traditional approach to modeling atom movement is to assume a single structural model with individual atomic displacement parameters (B factors). Given sufficient data, anisotropic displacement factors can be modeled, yielding directional insights into motions that might cause variations in the crystal. When the data are more limited, Translation-Libration-Screw (TLS) structural refinement, in which motions are described using rigid body segments of the molecule (Schomaker and Trueblood, 1968), has emerged as a common tool to model protein domain movements in crystallography and has been used by 22% of PDB depositions (Painter and Merritt, 2005, 2006). However, TLS refinements that vary in the rigid body definitions can predict very different motions while maintaining equivalent agreement to Bragg X-ray diffraction data (Urzhumtsev et al., 2015; Van Benschoten et al., 2015).

Additional sources of information have been used to overcome the inherent limitations of Bragg analysis in identifying collective protein motion. Patterns of steric clashes between alternative local conformations (van den Bedem et al., 2013) or time-averaged ensemble refinement (Burnley et al., 2012) can be used to suggest certain modes of concerted motion. However, the atomistic details of these correlated motions may only be reliably (yet indirectly) identified at high resolution, and time-averaged ensemble refinement is additionally complicated by the use of an underlying TLS model to account for crystal packing variations (Burnley et al., 2012). Alternative methods such as solid-state NMR experiments (Ma et al., 2015) or long time scale molecular dynamics simulations (Janowski et al., 2013; Janowski et al., 2015; Wall et al., 2014b) can be used to probe the structural basis of crystal packing variations and internal protein motions.

Complementary information about internal protein motions also can be obtained in the X-ray crystallography experiment itself by analysis of diffuse scattering. Diffuse scattering arises when deviations away from a perfect crystal cause X-rays to be diffracted away from Bragg reflections. When the deviations are due to crystal vibrations, they can be described using textbook temperature diffuse scattering theory (see, e.g. (James, 1948)). When each unit cell varies independently, the diffuse intensity is proportional to the variance in the unit cell structure factor (Guinier, 1963) which is equivalent to the Fourier transform of the Patterson function of the charge density variations. The approximation of independent unit cells can break down when correlations extend across unit cell boundaries; however, motions with long correlation

104 lengths result in diffuse intensity concentrated in the immediate neighborhood of Bragg peaks.
105 When analyzing the more broadly distributed diffuse intensity that corresponds to small
106 correlation lengths (Caspar et al., 1988; Clarage et al., 1992; Wall et al., 1997a; Wall et al.,
107 1997b) the contribution of inter-unit cell atom pairs is a small fraction of the total signal, which is
108 therefore dominated by internal protein motions.
109
Several approaches have been used to connect macromolecular diffuse scattering data to
models of protein motion and lattice disorder. Notably, Peter Moore has emphasized the need to
validate TLS models using diffuse scattering (Moore, 2009), as has been performed in a limited
number of cases (Doucet and Benoit, 1987; Perez et al., 1996; Van Benschoten et al., 2015).
Good agreement with the data has previously been observed for liquid-like motions (LLM)
models (Caspar et al., 1988; Clarage et al., 1992; Wall et al., 1997a; Wall et al., 1997b). In the
LLM model, the atoms in the protein are assumed to move randomly, like in a homogeneous
medium; the motions were termed "liquid-like" by Caspar et al (Caspar et al., 1988) because the
correlations in the displacements were assumed to fall off exponentially with the distance
between atoms.
120
Like the LLM model, normal modes (NM) models treat the protein as a softer substance than
the TLS model while still treating it as a solid. Unlike the LLM model, however, normal-mode
analysis (NMA) provides a much more detailed picture of the conformational ensemble,
enabling a more direct connection to putative mechanisms of protein function (Yang et al.,
2007). In addition, like TLS refinement, the normal modes refinement methods that have been
developed for Bragg analysis use few additional parameters (Gniewek et al., 2012; Kidera et al.,
1994; Lu and Ma, 2008; Ni et al., 2009; Poon et al., 2007). Unfortunately these programs are
not currently available in the standard builds of the major refinement software. Reasonable
qualitative agreement previously been seen using normal modes to model diffuse intensity in
individual diffraction images (Faure et al., 1994; Mizuguchi et al., 1994), and, more recently, the
fit of alternative coarse-grained elastic network models to diffuse scattering data of
staphylococcal nuclease has been investigated (Riccardi et al., 2010).
133
There is also a longstanding interest both in using diffuse scattering to validate improvements in
MD simulations and in using MD to derive a structural basis for the protein motions that give rise
to diffuse scattering (Clarage et al., 1995; Faure et al., 1994; Héry et al., 1998; Meinhold et al.,
2007; Meinhold and Smith, 2005a, 2005b, 2007; Wall et al., 2014b). Recent advances in
computing now enable microsecond duration simulations (Wall et al., 2014b) that can overcome
past barriers to accurate calculations seen using 10 ns or shorter MD trajectories (Clarage et al.,
1995; Meinhold and Smith, 2005a).
141
Despite the fact that diffuse scattering analysis is relatively well developed in small-molecule
crystallography (Welberry, 2004) and materials science (Keen and Goodwin, 2015), it has been
underutilized in protein crystallography. There are relatively few examples of diffuse data
analyzed using individual diffraction images from protein crystallography experiments, including
studies of tropomyosin (Chacko and Phillips, 1992; Phillips et al., 1980), 6-phosphogluconate
dehydrogenase (Helliwell et al., 1986), yeast initiator tRNA (Kolatkar et al., 1994), insulin

(Caspar et al., 1988), lysozyme (Clarage et al., 1992; Doucet and Benoit, 1987; Faure et al.,
1994; Mizuguchi et al., 1994; Perez et al., 1996), myoglobin (Clarage et al., 1995), Gag protein
(Welberry et al., 2011), and the 70s ribosome subunit (Polikanov and Moore, 2015). Moreover,
there are an even smaller number of examples involving complete three-dimensional diffuse
data sets; these include studies of staphylococcal nuclease (Wall et al., 1997b), and calmodulin
(Wall et al., 1997a).
154
To exploit the increased information that is potentially available from diffuse scattering, there is
a pressing need to increase the number of proteins for which complete three-dimensional
diffuse datasets have been experimentally measured. Conventional data collection procedures
use oscillation exposures to estimate the full Bragg intensities. In contrast, the complete three-
dimensional datasets measured by Wall *et al.* (Wall et al., 1997a; Wall et al., 1997b) used
specialized methods for integrating three-dimensional diffuse data from still diffraction images.
Similar methods now can be generalized and applied to other systems using modern beamlines
and X-ray detectors. In particular, the recent commercial development of pixel-array detectors
(PADs), which possess tight point-spread functions and single-photon sensitivity (Gruner, 2012),
have created new opportunities for measuring diffuse scattering as a routine tool in protein
crystallography experiments using more conventional data collection protocols.
166
Here, we present diffuse scattering datasets for the human proline isomerase cyclophilin A
(CypA) and the bovine serine protease trypsin. These datasets substantially increase the
amount of experimental three-dimensional diffuse scattering data available to the
macromolecular crystallography community, providing a necessary foundation for further
advancement of the field (Wall et al., 2014a). To assess the potential for routine collection of
diffuse datasets in crystallography, rather than expending a great deal of effort in optimizing the
diffuse data and collecting still images (Wall et al., 1997a; Wall et al., 1997b), we used
oscillation images obtained using best practices for high-quality Bragg data collection. The
resulting datasets are of sufficient quality that the diffuse scattering can discriminate among
alternative TLS refinements (Van Benschoten et al., 2015), Liquid-Like Motion (LLM) models
(Caspar et al., 1988; Clarage et al., 1992), and Normal Modes (NM) models ((Meinhold and
Smith, 2007; Mizuguchi et al., 1994; Riccardi et al., 2010)). Moreover, the agreement of the NM
models with both Bragg and diffuse scattering data suggests a path forward for anisotropic
refinement of atomic displacements, while maintaining a small set of parameters, using both
data sources simultaneously. Our results demonstrate that diffuse intensity can, and should, be
measured in a typical X-ray crystallography experiment and indicate that diffuse X-ray scattering
can be applied broadly as a tool to understand the conformational dynamics of macromolecules.
184
185

Results:

Experimental diffuse data show crystallographic symmetry

Figure 1. Steps in diffuse data integration. (A) Raw CypA diffraction images are processed (B) to remove Bragg peaks and enable direct comparisons of pixel values to models. (C) Pixels in diffraction images are mapped to reciprocal space and values of diffuse intensity are accumulated on a three-dimensional lattice; each diffraction image produces measurements of diffuse intensity on the surface of an Ewald sphere. (D) The data from individual images is combined and symmetrized to yield a nearly complete dataset (isosurface at a value of 65 photon counts in the total intensity, before subtracting the isotropic component).

The symmetrized anisotropic diffuse datasets processed by LUNUS (**Figure 1**) are shown in **Figure 2A** (CypA) and **Figure 2D** (trypsin) and are available as supplementary material. The CypA dataset is 98% complete to a resolution of 1.4 Å, while the trypsin map is 95% complete to 1.25 Å resolution. We used the Friedel symmetry and Laue group symmetry to quantify the level of crystallographic symmetry in each anisotropic map. To evaluate the degree Friedel symmetry, we averaged intensities between Friedel pairs to create a symmetrized map $I_{Friedel}$ and calculated the Pearson Correlation Coefficient (PCC) between the symmetrized and unsymmetrized data to obtain the statistic $CC_{Friedel}$. For CypA and trypsin, $CC_{Friedel}$ = 0.90 and 0.95 respectively, demonstrating that diffuse intensities obey Friedel symmetry. To assess the degree of Laue group symmetry, we averaged P222-related reflections (the Laue symmetry corresponding to the P 21 21 21 space group of both CypA and trypsin crystals) to produce the symmetrized intensities, I_{P222}. The linear correlation CC_{Sym} was then computed between the symmetrized and unsymmetrized intensities. The correlations were substantial for both CypA (CC_{Sym} = 0.70) and trypsin (CC_{Sym} = 0.69). Thus, our data are consistent with the diffuse intensity following the Bragg peak symmetry. The trypsin data were integrated using one degree oscillation frames, while the CypA data were integrated using 0.5 degree oscillation frames. The comparable degree of symmetry in the CypA and trypsin data suggests that the measurement of diffuse intensity is robust with respect to this difference in data collection.

218
Figure 2. Visualization of anisotropic diffuse intensities. (A) CypA experimental data with isosurfaces
shown using wireframes at a level of 2 photon counts in the resolution range 4.16 Å – 2.97 Å. Positive
intensity is rendered in green, negative in red. (B) Isosurfaces for diffuse scattering predicted by the
CypA LLM model. (C) Residual diffuse scattering (experimental data (A) minus LLM (B)). (D) Trypsin
experimental data with isosurfaces shown using wireframes at a level of 3 photon counts in the resolution
range 4.53 Å – 3.26. (E) Isosurfaces for diffuse scattering predicted by the Trypsin LLM model. (F)
Residual diffuse scattering (experimental data (D) minus LLM (E)).

TLS models yield low correlation with diffuse scattering data

To investigate how well TLS models agree with the molecular motions in the CypA crystal, we
compared the experimental diffuse data to intensities calculated from three alternative TLS
models: *phenix*, *tlsmd* and *whole molecule* (**Figure 3A-D**). Although all three models predict
different motions, the R-factors are very similar: R,R-free = 16.4%,18.1% for the *whole*
molecule and *Phenix* models; and 16.2%,18.1% for the TLSMD model. The correlations
between the calculated diffuse intensity for these models and the anisotropic experimental data
are low: 0.03 for the *phenix* model; 0.04 for the *TLSMD* model; and 0.14 for the *whole molecule*
model. In addition, the pairwise correlations of the calculated diffuse intensities are low: 0.066
for *whole molecule/TLSMD*; 0.116 for *whole molecule/Phenix*; and 0.220 for *Phenix/TLSMD*.

239
240 **Figure 3**. Rigid body domain definitions used for TLS models. CypA and Trypsin TLS groups shown on
241 the tertiary structure for whole molecule (A, E), Phenix (B, F), and TLSMD (C, G) and shown on the
242 primary sequence (D, H).

243
Like CypA, the three trypsin TLS models (**Figure 3E-H**) yielded very similar R,R-free values:
15.1%,16.7% for the *whole molecule* model; 15.3%,16.6% for the Phenix model; and
15.2%,16.6% for the TLSMD model. Correlations between the calculated and experimental
diffuse intensities are again low: 0.02 for the *Phenix* and *TLSMD* models, and 0.08 for the
Whole molecule model. Comparisons of the calculated anisotropic diffuse intensity show that
the *Whole molecule* motion is dissimilar to both the *Phenix* and *TLSMD* predictions (PCC = 0.03
and 0.05, respectively). In contrast, the *Phenix* and *TLSMD* models yield much more similar
diffuse intensities (PCC = 0.515). The relatively high correlation between these models is
consistent with the similarity in the TLS groups (**Figure 3F-H)**.
253
There are several possible explanations for the low correlation between the TLS model and
diffuse data for CypA and trypsin. First, TLS domain groupings other than those identified here
might yield higher agreement with the data. Second, the method used for generating ensembles
(Van Benschoten, 2015) assumes that TLS domains vary independently; it is possible that
accounting for correlations among the domains would more accurately describe the variations.
Lastly, similar to the rigid body motions model of Doucet & Benoit (Doucet and Benoit, 1987),
the correlations among TLS domains might lead to substantial correlations across unit cell
boundaries, which would produce small scale diffuse features in the immediate neighborhood of
Bragg peaks. The data integration methods used here cannot resolve these features, as the
measurements are mapped to a Bragg lattice. Methods to integrate the small-scale features in
protein crystallography onto a finer three-dimensional reciprocal space grid do exist (Wall et al.,
1997a) and might be used to address this last possibility in the future. In any case, the low
correlation of TLS models with the diffuse intensity for CypA and trypsin suggests that the
variations in the protein crystal might not be best explained by motions of relatively large, rigid
domains, and instead might involve motions that are correlated on a shorter length scale than
accounted for by these models.
270
271
272   **Liquid-like motions models yield substantial correlation with diffuse scattering**
273   **data**
274
One model that accounts for short-range correlations is Liquid-Like Motions (LLM) (Caspar et
al., 1988; Clarage et al., 1992). The LLM model assumes that atomic displacements are
uncorrelated between different unit cells, but are correlated within the unit cells. The correlation
in the displacements is assumed to decay exponentially as $f(x) = e^{-x/\gamma}$, where x is the
separation of the atoms, and **γ** is the length scale of the correlation. The displacements of all
atoms are assigned a standard deviation of **σ**. The LLM model previously has been refined
against three-dimensional diffuse intensities obtained from crystalline staphylococcal nuclease
(Wall et al., 1997b) and calmodulin (Wall et al., 1997a), yielding insights into correlated motions.
283
We refined isotropic LLM models of motions in CypA and trypsin against the experimental
diffuse intensities (**Figure 2B, E and Methods**). The CypA model was refined using data in the
resolution range 31.2 Å – 1.45 Å, and the trypsin model using 68 Å – 1.46 Å data. For CypA, the

refinement yielded **γ** = 7.1 Å and **σ** = 0.38 Å with a correlation of 0.518 between the calculated
and experimental anisotropic intensities. The highest correlation between data and experiment
occurs in the range 3.67 Å – 3.28 Å, where the value is 0.74 **(Figure 4A)**.   For the trypsin
dataset, the refinement yielded **γ** = 8.35 Å and **σ** = 0.32Å with a correlation of 0.44, which is
lower than for CypA. The peak value is 0.72 in the resolution range 4.53 Å -4.00 Å (**Figure 4B**).
292

293
294
295 **Figure 4.** Agreement of models of protein motions with diffuse and Bragg data. (A, B) Linear correlation
296 coefficients (CC) between diffuse data and LLM (red bars) or NM models (blue bars) computed by
297 resolution shell for (A) CypA and (B) Trypsin. (C, D) Correlations and R-factors between Bragg data and
298 NM models computed by resolution shell for (C) CypA and (D) Trypsin. Agreement factors for the diffuse
299 and Bragg data were computed using LUNUS (Wall, 2009) and *Phenix* (Adams et al., 2010), respectively.
300
The refined LLM models also were compared to the data using simulated diffraction images.
Images corresponding to frame number 67 of the CypA data were obtained using the LLM
model (Fig. 5A) and the integrated diffuse data (Fig. 5B). The main bright features above and
below the origin are similar between the two. Many of the weaker features also appear to be
similar, both at high and low resolution. The similarity is diminished but still apparent for images
obtained for frame number 45 of trypsin (Supplementary Fig. S1). These simulations provide a
visual confirmation of the substantial correlations obtained for the three-dimensional diffuse
intensity.
309

Figure 5. Simulated diffraction images for CypA frame 67 obtained using: (A) liquid-like motions model; (B) integrated 3D diffuse data; (C) elastic network model. Lighter colors correspond to stronger intensity. White regions correspond to pixel values where there are missing values in the corresponding three-dimensional lattice (Methods).

The substantial correlation of the LLM model with the diffuse data for CypA and trypsin indicates that the variations in the protein crystal can be approximately described using a model of the protein as a soft, homogeneous medium. The model implies that the motions of atoms separated by more than 7-8 Å are relatively independent, and that atoms that are closer to each other move in a more concerted way.

Normal-modes can model both diffuse and Bragg scattering data

To assess the potential of NMA to be developed for diffuse scattering studies, we developed coarse-grained elastic network models of the CypA and trypsin unit cells (Methods). The C_α coordinates and B-factors for the NM models are by definition identical to those derived from the Bragg data (Methods). To assess the agreement of specific NM-derived conformational variations with the Bragg data, we generated 50 member ensembles from the Hessian (Methods). We applied a single isotropic B-factor across all atoms ensuring that all deviations in the individual magnitude and anisotropy originated from the normal modes (Fig. S2). The correlations were high across resolution shells (Fig. 4 C, D), yielding overall R-factors of 38% (CypA) and 31% (Trypsin) (Tables S1 and S2). We also calculated the predicted diffuse intensity from the NM models: the correlation of the CypA model with the data is 0.41 in the resolution range 31.2 Å – 1.45 Å, and the correlation of the trypsin model with the data is 0.38 in the resolution range 68 Å – 1.46 Å. The agreement with the data is substantial within individual resolution shells (Fig. 4). The NM simulated diffraction image for CypA (Fig. 5C) shows bright features that are found in the data (Figs. 5B). The relative strength at high versus low-resolution is higher than in the data, however, suggesting that this NM model is too rigid; this discrepancy might be addressed by softening the intra-residue interactions and optimizing the model against the diffuse scattering data directly. The comparisons of simulated diffraction images for trypsin are consistent with the findings for CypA (Fig. S1).

Overall the agreement of the NM models with the data assessed using either 3D diffuse scattering datasets (Fig. 4) or simulated diffraction images (Figs. 5, S1) is substantial but slightly

less than for the LLM models. However, it is important to interpret this comparison in light of the
fact that the covariance matrices of the NM models were normalized to agree with the Bragg
data and not parameterized against the diffuse data (Methods), while the LLM model is
parameterized against the diffuse data. The agreement with the Bragg data is currently limited
by the fact that the parameter optimization used only the refined C_α positions and B-factors to
agree with the Bragg data and that heteroatoms, such as solvent, were not included in the
calculations. Collectively, these results point to the potential for normal modes to be refined
jointly against Bragg and diffuse scattering data as an alternative atomic displacement model,
replacing TLS or individual B-factors.

Discussion:

Diffuse X-ray scattering is a potentially valuable yet little exploited source of information about
macromolecular dynamics. Diffuse intensities can double the total number of measured data
points in the crystallographic experiment while providing a parallel dataset against which
structural dynamical models can be refined or validated. Until now measurement of three-
dimensional diffuse scattering data only has been pursued in dedicated efforts requiring extra
still diffraction images and substantial optimization of experimental design. The present
collection of two new datasets obtained using oscillation images using best current practices in
room-temperature protein crystallography (Fraser et al., 2011), and the use of the data in
evaluating TLS, LLM, and NM models, illustrates the potential for using diffuse scattering to
increase understanding of protein structure variations in any X-ray crystallography experiment,
representing a significant step towards moving diffuse scattering analysis into the mainstream of
structural biology.

Diffuse data obtained for CypA and trypsin can distinguish among the TLS, LLM, and NM
models of motions. However, the agreement with the data is somewhat lower than in previous
LLM models of three-dimensional diffuse scattering (Wall et al., 1997a; Wall et al., 1997b). In
this study, the correlation of the LLM model with the data was 0.518 in the range 31.2 Å – 1.45
Å for CypA, and 0.44 in the range 68 Å – 1.46 Å for trypsin; in comparison, the correlation was
0.595 in the range 10 Å – 2.5 Å for staphylococcal nuclease (Wall et al., 1997b) and 0.55 in the
range 7.5 Å – 2.1 Å for calmodulin (Wall et al., 1997a). Some possible explanations for the
lower agreement for CypA and trypsin include: the use of higher resolution data in the present
studies; that LLM might be a better description of motions in staphylococcal nuclease and
calmodulin than in CypA and trypsin; and that the measurements might have been more
accurate in the past experiments, as the data collection was tailored for diffuse scattering. The
apparent alignment of the residual intensity distribution with the unit cell axes (**Figures 2C, 2F**)
also suggests that an anisotropic LLM model might be more appropriate than an isotropic LLM
model for CypA and trypsin.

The agreement of the LLM models with three-dimensional experimental diffuse data across
multiple systems warrants further consideration for using diffuse scattering in model refinement
and validation. A key finding is that the agreement of the LLM models with the diffuse data is
higher than the TLS models, which currently are used widely in protein crystallography.

Interestingly, the 7-8 Å length scale of the correlations is comparable to the size of the TLS
domains; however, compared to the sharp domains of the TLS model, the exponential form of
the correlations indicates that there is a smooth spatial transition between the correlated and
uncorrelated atoms in the LLM. The smooth transition might be key to the increased agreement
of the LLM with the diffuse data compared to the rigidly defined regions of the TLS model.
394
Our findings also support the use of NM models in combination with diffuse scattering for model
refinement and validation. The exploratory work here, which did not use diffuse data for
parameterization, indicates that NM models contain features that can capture aspects of the
diffuse scattering data, and motivates further work to incorporate NM in both Bragg and diffuse
refinement. In the coarse-grained NM model in Eq. (2), the residues are treated as rigid, which
artificially increases anisotropic features at high resolution. We performed a limited exploration
of models with decreased intra-residue atom correlations: so far these models have led to lower
correlations with the data; however, in principle such models should be more accurate. Future
work will focus on developing computationally efficient methods for optimizing the accuracy of
the coarse-grained NM models and for using all-atom NM to model diffuse scattering.
405
Overall, the three-dimensional diffuse scattering data obtained here for CypA and trypsin, and
previously for staphylococcal nuclease (Wall et al., 1997b) and calmodulin (Wall et al., 1997a)
suggest that the protein structure varies more like a soft material than like a collection of
independent rigid domains. An important consideration in developing these new refinement
methods is to maintain a key advantage of TLS refinement at lower resolutions: the introduction
of relatively few new parameters for refinement. This requirement also would be satisfied by
NMA, which has a low computational cost and general applicability, making it a promising model
for integrating diffuse scattering into crystallographic model building and refinement (Wall et al.,
2014a). Whether this pursuit is well-motivated hinges on whether new biological insights can be
gained from atomic displacements generated by NM models refined against Bragg and diffuse
data. Indeed, although use of TLS in model refinement is now widespread, it scarcely has been
used to generate biological hypotheses (for exceptions, see: (Chaudhry et al., 2004; Henzler-
Wildman et al., 2007)). In contrast to TLS models, elastic network NM models have been widely
used to draw functional inferences (Bahar et al., 2010). Both the encouraging agreement of the
NM models with the diffuse scattering and the potential for NM models to yield new insights
about the importance of conformational dynamics in protein function provide a strong motivation
for further developing NM models for protein X-ray crystallography.
423
Diffuse scattering also can be used to validate models of molecular motions other than those
considered here, including models produced by ensemble refinement (Burnley et al., 2012);
multiconformer modeling performed by discrete (Keedy et al., 2015; van den Bedem et al.,
2009) or continuous (Burling and Brünger, 1994; Kuriyan et al., 1991; Wall et al., 1997a)
conformational sampling; and molecular dynamics simulations (Clarage and Phillips, 1994;
Clarage et al., 1995; Faure et al., 1994; Héry et al., 1998; Janowski et al., 2013; Janowski et al.,
2015; Meinhold et al., 2007; Meinhold and Smith, 2005a, 2005b; Wall et al., 2014b). In
particular, molecular dynamics simulations now provide sufficient sampling to yield robust
calculations of diffuse intensity (Wall et al., 2014b), and these can be used to consider a myriad

of intramolecular motions (e.g., loop openings and side chain flips) (Wilson, 2013) and lattice
dynamics. Polikanov and Moore (Polikanov and Moore, 2015) recently have demonstrated the
importance of lattice vibrations in explaining experimental diffuse scattering measurements of
ribosome crystals, which indicates that models should simultaneously account for correlations
that are coupled both within and across unit cell boundaries (Clarage et al., 1992; Wall et al.,
1997a); accounting for lattice vibrations more accurately also might yield improved Bragg
integration (Wall et al., 2014a). Moreover, comparisons of crystal simulations and diffuse
scattering can provide a new observable for benchmarking improvements in energy functions
and sampling schemes (Janowski et al., 2015).
442
Although the initial successes of dynamics-based models of diffuse scattering indicates that
crystal defects can play a secondary role in contributing to the diffuse signal, at least in some
cases, consideration of crystal defects might become important to achieve the highest model
accuracy and most general applicability of diffuse scattering in crystallography. Additionally, as
more X-ray data from both brighter conventional and XFEL light sources, accounting for all
sources of Bragg and diffuse scattering will be necessary to model the total scattering needed
for innovative phasing applications (Gaffney and Chapman, 2007). In summary, the new
datasets presented here demonstrate that diffuse scattering can now be routinely collected and
that using these data will help us obtain an increasingly realistic picture of motion in protein
crystals, including integrated descriptions of intramolecular motions, lattice vibrations, and
crystal defects.
454

Methods:

455
456

457   **Protein purification and crystallization**
458

Trypsin crystals were obtained according to the method of Liebschner *et.al* (Liebschner et al.,
2013). Lyophilized bovine pancreas trypsin was purchased from Sigma-Aldrich (T1005) and
dissolved at a concentration of 30 mg/mL into 30mM HEPES pH 7.0, 5 mg/mL benzamidine and
3mM $CaCl_2$. Crystals were obtained from a solution of 200mM Ammonium sulfate, 100mM Na
cacodylate pH 6.5, 20% PEG 8000 and 15% glycerol. CypA was purified and crystallized as
previously described (Fraser et al., 2009). Briefly, the protein was concentrated to 60 mg/mL in
20mM HEPES pH 7.5, 100mM NaCl and 500mM TCEP. Trays were set with a precipitant
solution of 100mM HEPES pH 7.5, 22% PEG 3350 and 5mM TCEP. Both crystal forms were
obtained using the hanging-drop method.
468

469   **Crystallographic data collection**
470

Diffraction data were collected on beamline 11-1 at the Stanford Synchrotron Radiation
Lightsource (Menlo Park, CA). X-ray diffraction images were obtained using a Dectris PILATUS
6M Pixel Array Detector (PAD). Each dataset was collected from a single crystal at an ambient
temperature of 273K. To prevent dehydration, crystals were coated in a thin film of paratone
with minimal surrounding mother liquor. For CypA, a single set of 0.5 degree oscillation images
were collected and used for both Bragg and diffuse data processing. A total of 360 images were

collected across a 180 degree phi rotation. The Trypsin diffraction data consisted of one degree
oscillations across a 135 degree phi rotation; this dataset was similarly used for both Bragg and
diffuse data analysis. Both datasets were collected to optimize the Bragg signal, not the diffuse
signal. Although not used here, we note that data collection using a PAD with fine phi slicing
should be especially well suited for simultaneous collection of Bragg and diffuse data, as it
would enable integration of diffuse intensity at a tunable level of detail in reciprocal space.
483
Bragg data processing
Bragg diffraction data were processed using XDS and XSCALE (Kabsch, 2010) within the *xia2*
software package (Winter et al., 2013). Molecular replacement solutions were found using
Phaser (McCoy et al., 2007) within the *Phenix* software suite (Adams et al., 2010). The PDB
search models were 4I8G for trypsin, and 2CPL for CypA. Initial structural refinement was
performed using *phenix.refine* (Afonine et al., 2012). The strategy included refinement of
individual atomic coordinates and water picking. Both the X-ray/atomic displacement
parameters and X-ray/stereochemistry weights were optimized. Isotropic B-factors were chosen
for the initial structures to allow for non-negligible R-factor optimization by subsequent TLS
refinement strategies. All structures were refined for a total of 5 macrocycles. Statistics for these
initial crystal structure models are shown in **Table 1**.
495

496 **Table 1.** Refinement statistics for CypA and trypsin models, before TLS modeling is applied.
497

	CypA	Trypsin
Resolution range, Å	38.66-1.4	23.29-1.25
Space group	P 21 21 21	P 21 21 21
Unit cell, Å	42.91, 52.44, 89.12	54.81, 58.51, 67.42
Completeness (%)	98	95
R_{work} (%)	17.88	15.9
R_{free} (%)	19.5	17.41
RMS (bonds, Å)	0.007	0.013
RMS (angles, degrees)	1.16	1.61
Ramachandran favored %	97	98
Ramachandran allowed %	3	2
Ramachandran outliers %	0	0
Clashscore	0.79	2.59
Average B-factor, $Å^2$	21.42	14.57

498
499 Diffuse data integration
500

An overview of the diffuse data integration process is presented in **Figure 1**. Image processing
was performed using the LUNUS collection of diffuse scattering tools (Wall, 2009). Pixels
corresponding to the beam stop and image edges were masked using the *punchim* and *windim*
methods. To focus on the diffuse intensity, which compared to Bragg peaks has low individual
pixel values (while, being more broadly distributed in reciprocal space, having comparable total
integrated intensity), pixel values outside of the range 1-10,000 photon counts were masked
using *threshim*. The beam polarization was determined by analyzing the first frame to determine
the azimuthal intensity profile within a 100 pixel wide annulus about the origin, and by fitting the
resulting profile to the theoretical profile (Wall, 1996). Pixel values then were corrected for beam
polarization using *polarim*. A solid-angle normalization (*normim*) correction was also applied.
Mode filtering was used to remove Bragg peaks from diffraction images. This was accomplished
using *modeim*, with a mask width of 20 pixels and a single bin for each photon count increment.
These steps produced diffraction images in which pixel values could be directly compared to
model diffuse intensities. This procedure, originally developed for experiments on

staphylococcal nuclease (Wall et al., 1997b), is similar to the steps used by Polikanov and
Moore (Polikanov and Moore, 2015) to process individual ribosome diffraction images for
analysis of diffuse scattering data.
518
The Lunus processed frames were used to integrate the diffuse data onto a 3D lattice. The
integration was performed using a python script that calls DIALS methods within the
Computational Crystallography Toolbox (CCTBX; (Grosse-Kunstleve et al., 2002; Parkhurst et
al., 2014)). The script obtains an indexing solution using the *real_space_grid_search* method
and uses the results to map each pixel in each diffraction image to fractional Miller indices *h'k'l'*
in reciprocal space. It sums the intensities from pixels in the neighborhood of each integer Miller
index *hkl* and tracks the corresponding pixel counts, while ignoring pixels that fall within a ½ x ½
x ½ region about *hkl*. It writes the intensity sums and pixel counts for each frame on a grid,
populated on an Ewald sphere that varies according to the crystal orientation for each image
(**Figure 1C**). A radial scattering vector intensity profile was calculated for each frame using the
Lunus *avgrim* method and was used to scale diffuse frames across the entire dataset. The
Lunus *sumlt* and *divlt* methods were used to compute the mean diffuse intensity at each grid
point using the scaled sums and pixel counts from all of the frames.
532
Because the model diffuse intensities were computed without considering solvent, experimental
and model diffuse intensities were compared using just the anisotropic component of the signal,
which is primarily due to the protein (Wall et al., 2014b). The Lunus *avgrlt* and *subrflt* routines
were applied to subtract the radial average and obtain the anisotropic signal. Signal intensities
were then symmetrized using *phenix.reflection_file_converter* to obtain a dataset for comparison
to models. Datasets were compared to each other and to models using linear correlations
computed using the *phenix.reflection_statistics* tool.
540
541 All images are available on SBGrid Data Grid (https://data.sbgrid.org/dataset/68/ for CypA;
542 https://data.sbgrid.org/dataset/201/ for Trypsin) and the integrated diffuse scattering maps are
543 available as Supplementary Material.
544
545 **Simulated diffraction images**
546
Diffuse scattering images were simulated using methods similar to those for data integration in
the previous section. After finding an indexing solution, a frame corresponding to the desired
simulated image was selected from the data set. This frame was used as a template for
obtaining a mapping of each pixel to fractional Miller indices. The new value of each pixel was
obtained by linear interpolation of the values of diffuse intensity between the nearest-neighbor
integer points *hkl* for which diffuse intensity was either measured (as in the previous section) or
calculated (as in below sections on liquid-like motions and normal modes models). In the case
of the synthetic images computed from the diffuse data, the images greatly enhanced the ability
to visualize diffuse features compared to the original diffraction images (Supplementary Figs.
S3, S4); the enhancement is due to the improved statistics obtained by averaging many pixel
values to obtain a measurement at each value of *hkl*. The simulated images were processed to
enhance visualization of diffuse features: the minimum pixel value was computed within each

pixel-width annulus about the beam center, and was subtracted from each pixel value within the
annulus. Images were displayed using Adxv (Arvai, 2012), with display parameters selected for
meaningful comparison of the diffuse features.
562
563 **TLS structure refinement and diffuse scattering model**
564
Three independent TLS refinements were performed for CypA (**Figure 3A-D**). The *Whole*
molecule selection consists of the entire molecule as a single TLS group. The *Phenix* selection
consists of the 8 groups (residues 2-14, 15-41, 42-64, 65-84, 85-122, 123-135, 136-145 and
146-165) identified by *phenix.find_tls_groups.* The *TLSMD* selection consists of 8 groups
(residues 2-15, 16-55, 56-80, 81-85, 86-91, 92-124, 125-143 and 144-165) identified by the TLS
Motion Determination web server (Painter and Merritt, 2005, 2006). All TLS refinement was
performed within *phenix.refine* through 5 macrocycles. Aside from the inclusion of TLS
refinement, these macrocycles were identical to the initial structure refinement described above.
573
574 Similarly, for trypsin, we selected Whole *Molecule, Phenix,* and *TLSMD* TLS refinement
575 strategies as described above (**Figure 3E-H**). The *Phenix* selection consists of 7 TLS groups:
576 residues 16-54, 55-103, 104-123, 124-140, 141-155, 156-225 and 226-245. The *TLSMD*
577 selection consists of 9 groups: residues 16-52, 53-98, 99-115, 116-144, 145-171, 172-220, 221-
578 224, 225-237 and 238-245.
579
Structural ensembles of the CypA and trypsin TLS motions were generated through
the *Phenix.tls_as_xyz* method (Urzhumtsev et al., 2015). Each ensemble consisted of 1,000
random samples of the underlying TLS atomic displacement distributions, assuming
independent distributions for each domain. Diffuse scattering models were calculated from the
TLS ensembles using *Phenix.diffuse* (Van Benschoten et al., 2015). CypA and trypsin models
were generated to a final resolution of 1.2 Å and 1.4 Å respectively, to match the resolution of
the experimental data.
587
588 **Liquid-like motions model**
589
We computed Liquid-like motions (LLM) models of diffuse scattering using the structures refined
prior to the TLS refinements (CypA: PDB ; Trypsin: PDB). For both CypA and trypsin, the
temperature factors for all atoms were set to zero and squared calculated structure factors
$I_0(hkl)$ were computed using the *structure_factors*, *as_intensity_array*, and *expand_to_p1*
methods in CCTBX (Grosse-Kunstleve et al., 2002; Parkhurst et al., 2014). The Lunus *symlt*
method was used to fill in missing values in reciprocal space using the appropriate P222 Laue
symmetry.
597
598 Given a correlation length γ and amplitude of motion σ, the diffuse intensity predicted by the
599 LLM model was calculated as
600

$$D_{LLM}(\mathbf{s}) = 4\pi^2 s^2 \sigma^2 e^{-4\pi^2 s^2 \sigma^2} I_0(\mathbf{s}) * \Gamma_\gamma(\mathbf{s}) \tag{1}$$

$$\Gamma_\gamma(\mathbf{s}) = \frac{8\pi\gamma^3}{(1 + 4\pi^2 s^2 \gamma^2)}$$

Fourier methods in Lunus (*fftlt*) were used to compute the convolution. The agreement with the
data was quantified by computing a linear correlation as a target function, using the anisotropic
intensities (**Diffuse data integration**). Optimization of the target with respect to γ and σ was
performed in a python script using *scipy.optimize* (www.scipy.org) with the Powell minimization
method.

Normal modes model

The diffuse intensity was computed using a normal modes (NM) model of correlated atom
displacements, using methods similar to Riccardi et al. (Riccardi et al., 2010). Atomic
coordinates and isotropic displacement parameters were obtained from PDB entries 5F66
(CypA) and 5F6M (trypsin) and were parsed and expanded to the P1 unit cell using the
iotbx.pdb methods in CCTBX (Grosse-Kunstleve et al., 2002). The Hessian matrix **H** was
defined using a modified anisotropic elastic network model (Atilgan et al., 2001), with springs
between C_α atoms (*i,j*) within a cutoff radius of 25 Å. The spring force constants were computed
as $ke^{-r_{ij}/\lambda}$, where r_{ij} is the closest distance between atoms *i* and *j*, either in the same unit cell
or in neighboring unit cells; λ = 10.5 Å; and *k* = 1 for $r_{ij} < 25$ Å and *k* = 0 otherwise (the
nonzero value of *k* is arbitrary due to the normalization employed below). Covariances of atom
pair displacements $v_{ij} = \langle \mathbf{r}_i \cdot \mathbf{r}_j \rangle$ were obtained using the pseudoinverse of **H** as described in
(Atilgan et al., 2001). The values of v_{ij} were renormalized to $\phi_{ij} = v_{ij}\sigma_i\sigma_j/\left(v_{ii}v_{jj}\right)^{1/2}$ using the
isotropic displacement parameters σ_i of the i^{th} C_α atom from the Bragg refinement; the model
was thus consistent with the refined crystal structure.

The diffuse intensity was computed as

$$D_{NM}(\mathbf{s}) = \sum_i \sum_j f_i f_j^* e^{-4\pi^2\left(\sigma_i^2 + \sigma_j^2\right)s^2} \left(e^{-4\pi^2 s^2 \phi_{ij}} - 1\right) \qquad\qquad (2)$$

where f_i is the structure factor of the combined atoms in the residue associated with the i^{th} C_α
atom. Structure factors were computed using a two-gaussian approximation of atomic form
factors; the parameters were obtained using the eltbx.xray_scattering methods in CCTBX
(Grosse-Kunstleve et al., 2002); phase factors were applied using the atomic coordinates.

The Bragg intensities were computed from ensembles generated by using the first 10 nonzero
eigenvectors of **H** with corresponding inverse eigenvalues as their weights. Because the overall
scale of the spring constant was arbitrary in the NM model (see above), the amplitudes of
motion were too large using the absolute eigenvalues; they were therefore scaled to maintain
the connectivity of the backbones. 50 member ensemble models were generated by Normal
Mode Wizard (*NMWiz*) (Bakan et al., 2011), which is a *VMD* (Humphrey et al., 1996) plugin. A
single B-factor of 10 was applied to all atoms in the ensemble. Structure factors were
generated using *phenix.fmodel* and compared to the experimental data using
phenix.reflection_statistics (Adams et al., 2010).

Acknowledgments

We thank Pavel Afonine for computational assistance in converting and comparing structure
factors. We are grateful to the UC Office of the President, Multicampus Research Programs and
Initiatives grant MR-15-338599 and the Program for Breakthrough Biomedical Research, which
is partially funded by the Sandler Foundation. Use of the Stanford Synchrotron Radiation
Lightsource, SLAC National Accelerator Laboratory, is supported by the U.S. Department of
Energy, Office of Science, Office of Basic Energy Sciences under Contract No. DE-AC02-
76SF00515. The SSRL Structural Molecular Biology Program is supported by the DOE Office of
Biological and Environmental Research, and by the National Institutes of Health, National
Institute of General Medical Sciences (including P41GM103393). N.K.S. was supported by NIH
grant GM095887. J.S.F. was supported by a Searle Scholar Award from the Kinship
Foundation, a Pew Scholar Award from the Pew Charitable Trusts, a Packard Fellowship from
the David and Lucile Packard Foundation, NIH OD009180, NIH GM110580, and NSF STC-
1231306. M.E.W. was supported by the US Department of Energy under Contract DE-AC52-
06NA25396 through the Laboratory-Directed Research and Development Program at Los
Alamos National Laboratory (LANL). The LANL technical release number is LA-UR-15-28934.

References

Adams, P.D., Afonine, P.V., Bunkoczi, G., Chen, V.B., Davis, I.W., Echols, N., Headd, J.J.,
Hung, L.W., Kapral, G.J., Grosse-Kunstleve, R.W.*, et al.* (2010). PHENIX: a comprehensive
Python-based system for macromolecular structure solution. Acta Crystallogr D Biol Crystallogr
66, 213-221.

Afonine, P.V., Grosse-Kunstleve, R.W., Echols, N., Headd, J.J., Moriarty, N.W., Mustyakimov,
M., Terwilliger, T.C., Urzhumtsev, A., Zwart, P.H., and Adams, P.D. (2012). Towards automated
crystallographic structure refinement with phenix.refine. Acta Crystallogr D Biol Crystallogr *68*,
352-367.

Arvai, A. (2012). ADXV - a program to display X-ray diffraction images.

Atilgan, A.R., Durell, S.R., Jernigan, R.L., Demirel, M.C., Keskin, O., and Bahar, I. (2001).
Anisotropy of fluctuation dynamics of proteins with an elastic network model. Biophys J *80*, 505-
515.

Bahar, I., Lezon, T.R., Yang, L.W., and Eyal, E. (2010). Global dynamics of proteins: bridging
between structure and function. Annu Rev Biophys *39*, 23-42.

Bakan, A., Meireles, L.M., and Bahar, I. (2011). ProDy: protein dynamics inferred from theory
and experiments. Bioinformatics *27*, 1575-1577.

Burling, F.T., and Brünger, A.T. (1994). Thermal motions and conformational disorder in protein
crystal structures: comparison of multi-conformer and time-averaging models. Israeli J Chem
34, 165-175.

Burnley, B.T., Afonine, P.V., Adams, P.D., and Gros, P. (2012). Modelling dynamics in protein
crystal structures by ensemble refinement. Elife *1*, e00311.

Caspar, D.L., Clarage, J., Salunke, D.M., and Clarage, M. (1988). Liquid-like movements in
crystalline insulin. Nature *332*, 659-662.

Chacko, S., and Phillips, G.N., Jr. (1992). Diffuse x-ray scattering from tropomyosin crystals.
Biophys J *61*, 1256-1266.

Chaudhry, C., Horwich, A.L., Brunger, A.T., and Adams, P.D. (2004). Exploring the structural
dynamics of the E.coli chaperonin GroEL using translation-libration-screw crystallographic
refinement of intermediate states. J Mol Biol *342*, 229-245.

Clarage, J.B., Clarage, M.S., Phillips, W.C., Sweet, R.M., and Caspar, D.L. (1992). Correlations
of atomic movements in lysozyme crystals. Proteins *12*, 145-157.

Clarage, J.B., and Phillips, G.N., Jr. (1994). Cross-validation tests of time-averaged molecular
dynamics refinements for determination of protein structures by X-ray crystallography. Acta
Crystallogr D Biol Crystallogr *50*, 24-36.

Clarage, J.B., and Phillips, G.N., Jr. (1997). Analysis of diffuse scattering and relation to
molecular motion. Methods Enzymol *277*, 407-432.

Clarage, J.B., Romo, T., Andrews, B.K., Pettitt, B.M., and Phillips, G.N., Jr. (1995). A sampling
problem in molecular dynamics simulations of macromolecules. Proc Natl Acad Sci U S A *92*,
3288-3292.

Doucet, J., and Benoit, J.P. (1987). Molecular dynamics studied by analysis of the X-ray diffuse
scattering from lysozyme crystals. Nature *325*, 643-646.

Faure, P., Micu, A., Perahia, D., Doucet, J., Smith, J.C., and Benoit, J.P. (1994). Correlated
intramolecular motions and diffuse X-ray scattering in lysozyme. Nat Struct Biol *1*, 124-128.

Fraser, J.S., Clarkson, M.W., Degnan, S.C., Erion, R., Kern, D., and Alber, T. (2009). Hidden
alternative structures of proline isomerase essential for catalysis. Nature *462*, 669-673.

Fraser, J.S., van den Bedem, H., Samelson, A.J., Lang, P.T., Holton, J.M., Echols, N., and
Alber, T. (2011). Accessing protein conformational ensembles using room-temperature X-ray
crystallography. Proc Natl Acad Sci U S A *108*, 16247-16252.

Gaffney, K.J., and Chapman, H.N. (2007). Imaging atomic structure and dynamics with ultrafast
x-ray scattering. Science *316*, 1444-1448.

Gniewek, P., Kolinski, A., Jernigan, R.L., and Kloczkowski, A. (2012). Elastic network normal
modes provide a basis for protein structure refinement. J Chem Phys *136*, 195101.

Grosse-Kunstleve, R.W., Sauter, N.K., Moriarty, N.W., and Adams, P.D. (2002). The
Computational Crystallography Toolbox: crystallographic algorithms in a reusable software
framework. Journal of Applied Crystallography *35*, 126-136.

Gruner, S.M. (2012). X-ray imaging detectors. Physics Today *65*, 29-34.

Guinier, A. (1963). X-ray diffraction in crystals, imperfect crystals, and amorphous bodies
(Courier Dover Publications).

Helliwell, J., Glover, I., Jones, A., Pantos, E., and Moss, D. (1986). Protein dynamics: use of
computer graphics and protein crystal diffuse scattering recorded with synchrotron X-radiation.
Biochemical Society Transactions *14*, 653-655.

Henzler-Wildman, K.A., Thai, V., Lei, M., Ott, M., Wolf-Watz, M., Fenn, T., Pozharski, E.,
Wilson, M.A., Petsko, G.A., Karplus, M.*, et al.* (2007). Intrinsic motions along an enzymatic
reaction trajectory. Nature *450*, 838-844.

Héry, S., Genest, D., and Smith, J.C. (1998). X-ray diffuse scattering and rigid-body motion in
crystalline lysozyme probed by molecular dynamics simulation. J Mol Biol *279*, 303-319.

Humphrey, W., Dalke, A., and Schulten, K. (1996). VMD: visual molecular dynamics. J Mol
Graph *14*, 33-38, 27-38.

James, R. (1948). The Optical Principles of the Diffraction of X-Rays (London: Bell).

Janowski, P.A., Cerutti, D.S., Holton, J., and Case, D.A. (2013). Peptide crystal simulations
reveal hidden dynamics. J Am Chem Soc *135*, 7938-7948.

Janowski, P.A., Liu, C., Deckman, J., and Case, D.A. (2015). Molecular dynamics simulation of
triclinic lysozyme in a crystal lattice. Protein Sci.

Kabsch, W. (2010). Xds. Acta Crystallogr D Biol Crystallogr *66*, 125-132.

Keedy, D.A., Fraser, J.S., and van den Bedem, H. (2015). Exposing Hidden Alternative
Backbone Conformations in X-ray Crystallography Using qFit. PLoS Comput Biol *11*, e1004507.

Keen, D.A., and Goodwin, A.L. (2015). The crystallography of correlated disorder. Nature *521*,
303-309.

Kidera, A., Matsushima, M., and Go, N. (1994). Dynamic structure of human lysozyme derived
from X-ray crystallography: normal mode refinement. Biophys Chem *50*, 25-31.

Kolatkar, A.R., Clarage, J.B., and Phillips, G.N., Jr. (1994). Analysis of diffuse scattering from
yeast initiator tRNA crystals. Acta Crystallogr D Biol Crystallogr *50*, 210-218.

Kuriyan, J., Osapay, K., Burley, S.K., Brunger, A.T., Hendrickson, W.A., and Karplus, M. (1991).
Exploration of disorder in protein structures by X-ray restrained molecular dynamics. Proteins
10, 340-358.

Kuzmanic, A., Kruschel, D., van Gunsteren, W.F., Pannu, N.S., and Zagrovic, B. (2011).
Dynamics may significantly influence the estimation of interatomic distances in biomolecular X-
ray structures. J Mol Biol *411*, 286-297.

Liebschner, D., Dauter, M., Brzuszkiewicz, A., and Dauter, Z. (2013). On the reproducibility of
protein crystal structures: five atomic resolution structures of trypsin. Acta Crystallogr D Biol
Crystallogr *69*, 1447-1462.

Lu, M., and Ma, J. (2008). A minimalist network model for coarse-grained normal mode analysis
and its application to biomolecular x-ray crystallography. Proc Natl Acad Sci U S A *105*, 15358-
15363.

Ma, P., Xue, Y., Coquelle, N., Haller, J.D., Yuwen, T., Ayala, I., Mikhailovskii, O., Willbold, D.,
Colletier, J.P., Skrynnikov, N.R.*, et al.* (2015). Observing the overall rocking motion of a protein
in a crystal. Nat Commun *6*, 8361.

McCoy, A.J., Grosse-Kunstleve, R.W., Adams, P.D., Winn, M.D., Storoni, L.C., and Read, R.J.
(2007). Phaser crystallographic software. J Appl Crystallogr *40*, 658-674.

Meinhold, L., Merzel, F., and Smith, J.C. (2007). Lattice dynamics of a protein crystal. Phys Rev
Lett *99*, 138101.

Meinhold, L., and Smith, J.C. (2005a). Correlated dynamics determining x-ray diffuse scattering
from a crystalline protein revealed by molecular dynamics simulation. Phys Rev Lett *95*,
218103.

Meinhold, L., and Smith, J.C. (2005b). Fluctuations and correlations in crystalline protein
dynamics: a simulation analysis of staphylococcal nuclease. Biophys J *88*, 2554-2563.

Meinhold, L., and Smith, J.C. (2007). Protein dynamics from X-ray crystallography: anisotropic,
global motion in diffuse scattering patterns. Proteins *66*, 941-953.

Mizuguchi, K., Kidera, A., and Go, N. (1994). Collective motions in proteins investigated by X-
ray diffuse scattering. Proteins *18*, 34-48.

Moore, P.B. (2009). On the relationship between diffraction patterns and motions in
macromolecular crystals. Structure *17*, 1307-1315.

Ni, F., Poon, B.K., Wang, Q., and Ma, J. (2009). Application of normal-mode refinement to X-ray
crystal structures at the lower resolution limit. Acta Crystallogr D Biol Crystallogr *65*, 633-643.

Painter, J., and Merritt, E.A. (2005). A molecular viewer for the analysis of TLS rigid-body
motion in macromolecules. Acta Crystallogr D Biol Crystallogr *61*, 465-471.

Painter, J., and Merritt, E.A. (2006). Optimal description of a protein structure in terms of
multiple groups undergoing TLS motion. Acta Crystallogr D Biol Crystallogr *62*, 439-450.

Parkhurst, J.M., Brewster, A.S., Fuentes-Montero, L., Waterman, D.G., Hattne, J., Ashton, A.W.,
Echols, N., Evans, G., Sauter, N.K., and Winter, G. (2014). : the diffraction experiment toolbox.
J Appl Crystallogr *47*, 1459-1465.

Perez, J., Faure, P., and Benoit, J.P. (1996). Molecular rigid-body displacements in a tetragonal
lysozyme crystal confirmed by X-ray diffuse scattering. Acta Crystallogr D Biol Crystallogr *52*,
722-729.

Phillips, G.N., Jr., Fillers, J.P., and Cohen, C. (1980). Motions of tropomyosin. Crystal as
metaphor. Biophys J *32*, 485-502.

Polikanov, Y.S., and Moore, P.B. (2015). Acoustic vibrations contribute to the diffuse scatter
produced by ribosome crystals. Acta Crystallogr D Biol Crystallogr *71*, 2021-2031.

Poon, B.K., Chen, X., Lu, M., Vyas, N.K., Quiocho, F.A., Wang, Q., and Ma, J. (2007). Normal
mode refinement of anisotropic thermal parameters for a supramolecular complex at 3.42-A
crystallographic resolution. Proc Natl Acad Sci U S A *104*, 7869-7874.

Riccardi, D., Cui, Q., and Phillips, G.N., Jr. (2010). Evaluating elastic network models of
crystalline biological molecules with temperature factors, correlated motions, and diffuse x-ray
scattering. Biophys J *99*, 2616-2625.

Schomaker, V., and Trueblood, K.N. (1968). On the rigid-body motion of molecules in crystals.
Acta Crystallographica Section B *24*, 63-76.

Urzhumtsev, A., Afonine, P.V., Van Benschoten, A.H., Fraser, J.S., and Adams, P.D. (2015).
From deep TLS validation to ensembles of atomic models built from elemental motions. Acta
Crystallogr D Biol Crystallogr *71*, 1668-1683.

Van Benschoten, A.H., Afonine, P.V., Terwilliger, T.C., Wall, M.E., Jackson, C.J., Sauter, N.K.,
Adams, P.D., Urzhumtsev, A., and Fraser, J.S. (2015). Predicting X-ray diffuse scattering from
translation-libration-screw structural ensembles. Acta Crystallogr D Biol Crystallogr *71*, 1657-
1667.

van den Bedem, H., Bhabha, G., Yang, K., Wright, P.E., and Fraser, J.S. (2013). Automated
identification of functional dynamic contact networks from X-ray crystallography. Nat Methods
10, 896-902.

van den Bedem, H., Dhanik, A., Latombe, J.C., and Deacon, A.M. (2009). Modeling discrete
heterogeneity in X-ray diffraction data by fitting multi-conformers. Acta Crystallogr D Biol
Crystallogr *65*, 1107-1117.

van den Bedem, H., and Fraser, J.S. (2015). Integrative, dynamic structural biology at atomic
resolution--it's about time. Nat Methods *12*, 307-318.

Wall, M.E. (2009). Methods and software for diffuse X-ray scattering from protein crystals.
Methods Mol Biol *544*, 269-279.

Wall, M.E., Adams, P.D., Fraser, J.S., and Sauter, N.K. (2014a). Diffuse X-ray scattering to
model protein motions. Structure *22*, 182-184.

Wall, M.E., Clarage, J.B., and Phillips, G.N. (1997a). Motions of calmodulin characterized using
both Bragg and diffuse X-ray scattering. Structure *5*, 1599-1612.

Wall, M.E., Ealick, S.E., and Gruner, S.M. (1997b). Three-dimensional diffuse x-ray scattering
from crystals of Staphylococcal nuclease. Proc Natl Acad Sci U S A *94*, 6180-6184.

Wall, M.E., Van Benschoten, A.H., Sauter, N.K., Adams, P.D., Fraser, J.S., and Terwilliger, T.C.
(2014b). Conformational dynamics of a crystalline protein from microsecond-scale molecular
dynamics simulations and diffuse X-ray scattering. Proc Natl Acad Sci U S A *111*, 17887-17892.

Welberry, T.R. (2004). Diffuse X-Ray Scattering and Models of Disorder (Oxford: Oxford
University Press).

Welberry, T.R., Heerdegen, A.P., Goldstone, D.C., and Taylor, I.A. (2011). Diffuse scattering
resulting from macromolecular frustration. Acta Crystallogr B *67*, 516-524.

Wilson, M.A. (2013). Visualizing networks of mobility in proteins. Nat Methods *10*, 835-837.

Winter, G., Lobley, C.M., and Prince, S.M. (2013). Decision making in xia2. Acta Crystallogr D
Biol Crystallogr *69*, 1260-1273.

Yang, L., Song, G., and Jernigan, R.L. (2007). How well can we understand large-scale protein
motions using normal modes of elastic network models? Biophys J *93*, 920-929.
835