<u>Measuring and modeling diffuse scattering in protein X-ray crystallography</u>

Andrew H. Van Benschoten[1], Ana Gonzalez[2], Aaron S. Brewster[3], Nicholas K. Sauter[3], James
S. Fraser[1,*], Michael E. Wall[4,*]
1 - Dept. of Bioengineering and Therapeutic Sciences, University of California, San Francisco, San
Francisco, CA 11 94158
2 - Stanford Synchrotron Radiation Lightsource, SLAC National Accelerator Laboratory, Menlo Park, CA
94025
3 - Physical Biosciences Division, Lawrence Berkeley National Laboratory, Berkeley, CA 94720
4 - Computer, Computational, and Statistical Sciences Division, Los Alamos National Laboratory, Los
Alamos, NM, 87545
*correspondence: jfraser@fraserlab.com, mewall@lanl.gov

## Abstract:

X-ray diffraction has the potential to provide rich information about the structural dynamics of
macromolecules. To realize this potential fully, it will be necessary to measure and model both
Bragg scattering, which is currently used to derive macromolecular structures, and diffuse
scattering, which reports on correlations in charge density variations. However, due to the extra
effort of collecting diffuse data, experimental measurement of diffuse scattering from protein
crystals has been limited to a small number of systems. Here, we present three-dimensional
measurements of diffuse intensity collected from crystals of the model enzymes cyclophilin A
and trypsin. The measurements were obtained from the same X-ray diffraction images as the
Bragg data, using best practices for data collection in protein crystallography. To model the
underlying dynamics in a practical way that can be used during structure refinement, we tested
both Translation-Libration-Screw (TLS) and Liquid-Like Motions (LLM) models of protein
motions. Whereas different TLS groupings yielded similar Bragg intensities, the diffuse
intensities were clearly different. The agreement of the LLM models with the diffuse data was
much stronger than for TLS. These results demonstrate a path to substantially increase the
number of diffuse datasets available to the wider biosciences community. These and future
datasets will be useful to benchmark computational tools for modeling correlated motions in
macromolecules and to improve refinement of dynamics-inspired structural models emerging in
X-ray crystallography.

## Significance:

The structural details of protein motions are critical to many biological processes, but they are
hidden to conventional biophysical techniques. Diffuse X-ray scattering can reveal these details
by measuring the correlated movements between atoms. However, diffuse scattering data
collection historically has required extra effort and dedicated experimental protocols. We have
measured three-dimensional diffuse intensities in X-ray diffraction from CypA and trypsin
crystals using standard crystallographic data collection techniques. Analysis of the resulting
data is consistent with the presence of liquid-like motions (LLM) in both crystals.  Our results
show that using diffuse scattering to model protein motions can become a component of routine
crystallographic analysis through the extension of commonplace methods.

## Introduction:

**Introduction:**
X-ray crystallography can be a key tool for elucidating the structural basis of protein motions
that play critical roles in enzymatic reactions, protein-protein interactions and signaling
cascades (van den Bedem and Fraser, 2015). X-ray diffraction yields an ensemble-averaged
picture of the protein structure: each photon simultaneously probes multiple unit cells that can
vary due to internal rearrangements or changes to the crystal lattice. Bragg analysis of X-ray
diffraction only yields the mean charge density of the unit cell, however, which fundamentally
limits the information that can be obtained about protein dynamics (Clarage and Phillips, 1997;
Keen and Goodwin, 2015).
51
A key limitation inherent in Bragg analysis is that alternative models with different correlations
between atomic motions can yield the same mean charge density (Kuzmanic et al., 2011). The
traditional approach to modeling atom movement is to assume a single structural model with
individual atomic displacement parameters (B factors). Given sufficient data, anisotropic
displacement factors can be modeled, yielding directional insights into motions that might cause
variations in the crystal. When the data are more limited, Translation-Libration-Screw (TLS)
structural refinement, in which motions are described using rigid body segments of the molecule
(Schomaker and Trueblood, 1968), has emerged as a tool to model protein domain movements
in crystallography (Painter and Merritt, 2005, 2006). However, TLS refinements that vary in the
rigid body definitions can predict very different motions while maintaining equivalent agreement
to Bragg X-ray diffraction data (Urzhumtsev et al., 2015; Van Benschoten et al., 2015).
63
Additional sources of information have been used to overcome the inherent limitations of Bragg
analysis in identifying collective protein motion. Patterns of steric clashes between alternative
local conformations (van den Bedem et al., 2013) or time-averaged ensemble refinement
(Burnley et al., 2012) can be used to suggest certain modes of concerted motion. However, the
atomistic details of these correlated motions may only be reliably (yet indirectly) identified at
high resolution, and time-averaged ensemble refinement is additionally complicated by the use
of an underlying TLS model to account for crystal packing variations (Burnley et al., 2012).
Alternative methods such as solid-state NMR experiments (Ma et al., 2015) or long time scale
molecular dynamics simulations (Janowski et al., 2013; Janowski et al., 2015; Wall et al.,
2014b) can be used to probe the structural basis of crystal packing variations and internal
protein motions.
75
Complementary information about internal protein motions also can be obtained in the X-ray
crystallography experiment itself by analysis of diffuse scattering. Diffuse scattering arises when
deviations away from a perfect crystal cause X-rays to be diffracted away from Bragg
reflections. When the deviations are due to crystal vibrations, they can be described using
textbook temperature diffuse scattering theory (see, e.g. (James, 1948)). When each unit cell
varies independently, the diffuse intensity is proportional to the variance in the unit cell structure
factor (Guinier, 1963) which is equivalent to the Fourier transform of the Patterson function of
the charge density variations. The approximation of independent unit cells can break down
when correlations extend across unit cell boundaries; however, motions with long correlation
lengths result in diffuse intensity concentrated in the immediate neighborhood of Bragg peaks.

When analyzing the more broadly distributed diffuse intensity that corresponds to small
correlation lengths (Caspar et al., 1988; Clarage et al., 1992; Wall et al., 1997a; Wall et al.,
1997b) the contribution of inter-unit cell atom pairs is a small fraction of the total signal, which is
therefore dominated by internal protein motions.
90
Several approaches have been used to connect macromolecular diffuse scattering data to
models of protein motion and lattice disorder. Notably, Peter Moore has emphasized the need to
validate TLS models using diffuse scattering (Moore, 2009), as has been performed in a limited
number of cases (Doucet and Benoit, 1987; Perez et al., 1996; Van Benschoten et al., 2015).
Good agreement with the data has previously been observed for liquid-like motions (LLM)
models (Caspar et al., 1988; Clarage et al., 1992; Wall et al., 1997a; Wall et al., 1997b). In the
LLM model, the atoms in the protein are assumed to move randomly, like in a homogeneous
medium; the motions were termed "liquid-like" by Caspar et al (Caspar et al., 1988) because the
correlations in the displacements were assumed to fall off exponentially with the distance
between atoms. There is also a longstanding interest both in using diffuse scattering to validate
improvements in MD simulations and in using MD to derive a structural basis for the protein
motions that give rise to diffuse scattering (Clarage et al., 1995; Faure et al., 1994; Héry et al.,
1998; Meinhold et al., 2007; Meinhold and Smith, 2005a, b, 2007; Wall et al., 2014b). Recent
advances in computing now enable microsecond duration simulations (Wall et al., 2014b) that
can overcome past barriers to accurate calculations seen using 10 ns or shorter MD trajectories
(Clarage et al., 1995; Meinhold and Smith, 2005a).
107
Despite the fact that diffuse scattering analysis is relatively well developed in small-molecule
crystallography (Welberry, 2004) and materials science (Keen and Goodwin, 2015), it has been
underutilized in protein crystallography. There are relatively few examples of diffuse data
analyzed using individual diffraction images from protein crystallography experiments, including
studies of tropomyosin (Chacko and Phillips, 1992; Phillips et al., 1980), 6-phosphogluconate
dehydrogenase (Helliwell et al., 1986), yeast initiator tRNA (Kolatkar et al., 1994), insulin
(Caspar et al., 1988), lysozyme (Clarage et al., 1992; Doucet and Benoit, 1987; Faure et al.,
1994; Mizuguchi et al., 1994; Perez et al., 1996), myoglobin (Clarage et al., 1995), Gag protein
(Welberry et al., 2011), and the 70s ribosome subunit (Polikanov and Moore, 2015). Moreover,
there are an even smaller number of examples involving complete three-dimensional diffuse
data sets; these include studies of staphylococcal nuclease (Wall et al., 1997b), and calmodulin
(Wall et al., 1997a).
120
To exploit the increased information that is potentially available from diffuse scattering, there is
a pressing need to increase the number of proteins for which complete three-dimensional
diffuse datasets have been experimentally measured. Conventional data collection procedures
use oscillation exposures to estimate the full Bragg intensities. In contrast, the complete three-
dimensional datasets measured by Wall *et al.* (Wall et al., 1997a; Wall et al., 1997b) used
specialized methods for integrating three-dimensional diffuse data from still diffraction images.
Similar methods now can be generalized and applied to other systems using modern beamlines
and X-ray detectors. In particular, the recent commercial development of pixel-array detectors
(PADs), which possess tight point-spread functions and single-photon sensitivity (Gruner, 2012),

have created new opportunities for measuring diffuse scattering as a routine tool in protein
crystallography experiments using more conventional data collection protocols.
132
Here, we present diffuse scattering datasets for the human proline isomerase cyclophilin A
(CypA) and the bovine serine protease trypsin. These datasets substantially increase the
amount of experimental three-dimensional diffuse scattering data available to the
macromolecular crystallography community, providing a necessary foundation for further
advancement of the field (Wall et al., 2014a).  To assess the potential for routine collection of
diffuse datasets in crystallography, rather than expending a great deal of effort in optimizing the
diffuse data and collecting still images (Wall et al., 1997a; Wall et al., 1997b), we used
oscillation images obtained using best practices for high-quality Bragg data collection. The
resulting datasets are of sufficient quality that the diffuse scattering can discriminate among
alternative TLS refinements (Van Benschoten et al., 2015) and Liquid-Like Motion (LLM) models
(Caspar et al., 1988; Clarage et al., 1992).  Our results demonstrate that diffuse intensity can,
and should, be measured in a typical X-ray crystallography experiment and indicate that diffuse
X-ray scattering can be applied broadly as a tool to understand the conformational dynamics of
macromolecules.
147
148    ## Results:
149
150    **Experimental diffuse data show crystallographic symmetry**
151

152
**Figure 1.** Steps in diffuse data integration. (A) Raw CypA diffraction images are processed (B) to remove
Bragg peaks and enable direct comparisons of pixel values to models. (C) Pixels in diffraction images are
mapped to reciprocal space and values of diffuse intensity are accumulated on a three-dimensional
lattice; each diffraction image produces measurements of diffuse intensity on the surface of an Ewald
sphere. (D) The data from individual images is combined and symmetrized to yield a complete dataset
(isosurface at a value of 65 photon counts in the total intensity, before subtracting the isotropic
component).
160
161

The symmetrized anisotropic diffuse datasets processed by LUNUS (**Figure 1**) are shown in
**Figure 2A** (CypA) and **Figure 2D** (trypsin) and are available as supplementary material. The
CypA dataset is 98% complete to a resolution of 1.4 Å, while the trypsin map is 95% complete
to 1.25 Å resolution. We used the Friedel symmetry and Laue group symmetry to quantify the
level of crystallographic symmetry in each anisotropic map. To evaluate the degree Friedel
symmetry, we averaged intensities between Friedel pairs to create a symmetrized map $I_{Friedel}$
and calculated the Pearson Correlation Coefficient (PCC) between the symmetrized and
unsymmetrized data to obtain the statistic $CC_{Friedel}$. For CypA and trypsin, $CC_{Friedel}$ = 0.90 and
0.95 respectively, demonstrating that diffuse intensities obey Friedel symmetry. To assess the
degree of Laue group symmetry, we averaged P222-related reflections (the Laue symmetry
corresponding to the P 21 21 21 space group of both CypA and trypsin crystals) to produce the
symmetrized intensities, $I_{P222}$. The linear correlation $CC_{Sym}$ was then computed between the
symmetrized and unsymmetrized intensities. The correlations were substantial for both CypA
($CC_{Sym}$ = 0.70) and trypsin ($CC_{Sym}$ = 0.69). Thus, our data are consistent with the diffuse
intensity following the Bragg peak symmetry. The trypsin data were integrated using one degree
oscillation frames, while the CypA data were integrated using 0.5 degree oscillation frames. The
comparable degree of symmetry in the CypA and trypsin data suggests that the measurement
of diffuse intensity is robust with respect to this difference in data collection.

182
**Figure 2.** Visualization of anisotropic diffuse intensities. (A) CypA experimental data with isosurfaces
shown using wireframes at a level of 2 photon counts in the resolution range 4.16 Å – 2.97 Å. Positive
intensity is rendered in green, negative in red.  (B) Isosurfaces for diffuse scattering predicted by the
CypA LLM model. (C) Residual diffuse scattering (experimental data (A) minus LLM (B). (D) Trypsin
experimental data with isosurfaces shown using wireframes at a level of 3 photon counts in the resolution
range 4.53 Å – 3.26. (E) Isosurfaces for diffuse scattering predicted by the Trypsin LLM model. (F)
Residual diffuse scattering (experimental data (D) minus LLM (E).

**TLS models yield low correlation with diffuse scattering data**

To investigate how well TLS models agree with the molecular motions in the CypA crystal, we
compared the experimental diffuse data to intensities calculated from three alternative TLS
models: *phenix*, *tlsmd* and *whole molecule* (**Figure 3A-D**). Although all three models predict

different motions, the R-factors are very similar:  R,R-free = 16.4%,18.1% for the *whole*
*molecule* and *Phenix* models; and 16.2%,18.1% for the TLSMD model. The correlations
between the calculated diffuse intensity for these models and the anisotropic experimental data
are low: 0.03 for the *phenix* model; 0.04 for the *TLSMD* model; and 0.14 for the *whole molecule*
model. In addition, the pairwise correlations of the calculated diffuse intensities are low:  0.066
for *whole molecule/TLSMD*; 0.116 for *whole molecule/Phenix*; and 0.220 for *Phenix/TLSMD*.
202

203
204  **Figure 3**. Rigid body domain definitions used for TLS models. CypA and Trypsin TLS groups shown on
205  the tertiary structure for whole molecule (A, E), Phenix (B, F), and TLSMD (C, G) and shown on the
206  primary sequence (D, H).
207
Like CypA, the three trypsin TLS models (**Figure 3E-H**) yielded very similar R,R-free values:
15.1%,16.7% for the *whole molecule* model; 15.3%,16.6% for the Phenix model; and
15.2%,16.6% for the TLSMD model.  Correlations between the calculated and experimental
diffuse intensities are again low: 0.02 for the *Phenix* and *TLSMD* models, and 0.08 for the
*Whole molecule* model. Comparisons of the calculated anisotropic diffuse intensity show that
the *Whole molecule* motion is dissimilar to both the *Phenix* and *TLSMD* predictions (PCC = 0.03
and 0.05, respectively). In contrast, the *Phenix* and *TLSMD* models yield much more similar
diffuse intensities (PCC = 0.515). The relatively high correlation between these models is
consistent with the similarity in the TLS groups (**Figure 3F-H)**.
217
There are several possible explanations for the low correlation between the TLS model and
diffuse data for CypA and trypsin. First, TLS domain groupings other than those identified here
might yield higher agreement with the data. Second, the method used for generating ensembles
(Van Benschoten, 2015) assumes that TLS domains vary independently; it is possible that
accounting for correlations among the domains would more accurately describe the variations.
Lastly, similar to the rigid body motions model of Doucet & Benoit (Doucet and Benoit, 1987),
the correlations among TLS domains might lead to substantial correlations across unit cell
boundaries, which would produce small scale diffuse features in the immediate neighborhood of
Bragg peaks. The data integration methods used here cannot resolve these features, as the
measurements are mapped to a Bragg lattice. Methods to integrate the small-scale features in
protein crystallography onto a finer three-dimensional reciprocal space grid do exist (Wall et al.,
1997a) and might be used to address this last possibility in the future. In any case, the low
correlation of TLS models with the diffuse intensity for CypA and trypsin suggests that the

variations in the protein crystal might not be best explained by motions of relatively large, rigid
domains, and instead might involve motions that are correlated on a shorter length scale than
accounted for by these models.
234
235

236 **Liquid-like motions models yield substantial correlation with diffuse scattering**
237 **data**
238
One model that accounts for short-range correlations is Liquid-Like Motions (LLM) (Caspar et
al., 1988; Clarage et al., 1992). The LLM model assumes that atomic displacements are
uncorrelated between different unit cells, but are correlated within the unit cells. The correlation
in the displacements is assumed to decay exponentially as $f(x) = e^{-x/\gamma}$, where x is the
separation of the atoms, and **γ** is the length scale of the correlation.  The displacements of all
atoms are assigned a standard deviation of **σ**. The LLM model previously has been refined
against three-dimensional diffuse intensities obtained from crystalline staphylococcal nuclease
(Wall et al., 1997b) and calmodulin (Wall et al., 1997a), yielding insights into correlated motions.
247
We refined isotropic LLM models of motions in CypA and trypsin against the experimental
diffuse intensities (**Figure 2B, E and Methods**). The CypA model was refined using data in the
resolution range 31.2 Å – 1.45 Å, and the trypsin model using 68 Å – 1.46 Å data. For CypA, the
refinement yielded **γ** = 7.1 Å and **σ** = 0.38 Å with a correlation of 0.518 between the calculated
and experimental anisotropic intensities. The highest correlation between data and experiment
occurs in the range 3.67 Å – 3.28 Å, where the value is 0.74 **(Figure 4A)**.   For the trypsin
dataset, the refinement yielded **γ** = 8.35 Å and **σ** = 0.32Å with a correlation of 0.44, which is
lower than for CypA. The peak value is 0.72 in the resolution range 4.53 Å -4.00 Å (**Figure 4B**).
256

257
258
259 **Figure 4.** Correlations between LLM models and diffuse data computed by resolution shell for (A) CypA
260 and (B) Trypsin.
261
262 The substantial correlation of the LLM model with the diffuse data for CypA and trypsin indicates
263 that the variations in the protein crystal can be approximately described using a model of the
264 protein as a soft, homogeneous medium. The model implies that the motions of atoms
265 separated by more than 7-8 Å are relatively independent, and that atoms that are closer to each
266 other move in a more concerted way.
267
268

## Discussion:

Diffuse X-ray scattering is a potentially valuable yet little exploited source of information about

macromolecular dynamics. Diffuse intensities can double the total number of measured data

points in the crystallographic experiment while providing a parallel dataset against which

structural dynamical models can be refined or validated. Until now measurement of three-

dimensional diffuse scattering data only has been pursued in dedicated efforts requiring extra

still diffraction images and substantial optimization of experimental design. The present

collection of two new datasets obtained using oscillation images using best current practices in

room-temperature protein crystallography (Fraser et al., 2011), and the use of the data in

evaluating TLS and LLM models, illustrates the potential for using diffuse scattering to increase

understanding of protein structure variations in any X-ray crystallography experiment, and

represents a significant step towards moving diffuse scattering analysis into the mainstream of

structural biology.

Diffuse data obtained for CypA and trypsin can distinguish between the TLS and LLM models of

motions. However, the agreement with the data is somewhat lower than in previous LLM models

of three-dimensional diffuse scattering (Wall et al., 1997a; Wall et al., 1997b). In this study, the

correlation of the LLM model with the data was 0.518 in the range 31.2 Å – 1.45 Å for CypA,

and 0.44 in the range 68 Å – 1.46 Å for trypsin; in comparison, the correlation  was 0.595 in the

range 10 Å – 2.5 Å for staphylococcal nuclease (Wall et al., 1997b) and 0.55 in the range 7.5 Å

– 2.1 Å for calmodulin (Wall et al., 1997a). Some possible explanations for the lower agreement

for CypA and trypsin include: the use of higher resolution data in the present studies; that LLM

might be a better description of motions in staphylococcal nuclease and calmodulin than in

CypA and trypsin; and that the measurements might have been more accurate in the past

experiments, as the data collection was tailored for diffuse scattering. The apparent alignment of

the residual intensity distribution with the unit cell axes (**Figures 2C, 2F**) also suggests that an

anisotropic LLM model might be more appropriate than an isotropic LLM model for CypA and

trypsin.

The agreement of the LLM models with three-dimensional experimental diffuse data across

multiple systems warrants further consideration for using diffuse scattering in model refinement

and validation. A key finding is that the agreement of the LLM models with the diffuse data is

higher than the TLS models, which currently are used widely in protein crystallography.

Interestingly, the 7-8 Å length scale of the correlations is comparable to the size of the TLS

domains; however, compared to the sharp domains of the TLS model, the exponential form of

the correlations indicates that there is a smooth spatial transition between the correlated and

uncorrelated atoms in the LLM. The smooth transition might be key to the increased agreement

of the LLM with the diffuse data compared to the rigidly defined regions of the TLS model.

Overall, the three-dimensional diffuse scattering data obtained here for CypA and trypsin, and

previously for staphylococcal nuclease (Wall et al., 1997b) and calmodulin (Wall et al., 1997a)

suggest that the protein structure varies more like a soft material than like a collection of

independent rigid domains. A normal modes model would be similarly soft, suggesting that

normal modes should be further investigated for analysis of protein motions using three-
dimensional diffuse datasets. Indeed, reasonable qualitative agreement already has been seen
using normal modes to model diffuse intensity in individual diffraction images (Faure et al.,
1994; Mizuguchi et al., 1994), and, like TLS refinement, normal modes refinement methods
have been developed for Bragg analysis (Gniewek et al., 2012; Kidera et al., 1994; Lu and Ma,
2008). An important consideration in developing these new refinement methods is to maintain a
key advantage of TLS refinement at lower resolutions: the introduction of relatively few new
parameters for refinement. This requirement also would be satisfied by the LLM, which has a
low computational cost and general applicability, making it a promising model for integrating
diffuse scattering into crystallographic model building and refinement (Wall et al., 2014a).
323
Diffuse scattering also can be used to validate more detailed models of molecular motions than
were considered here, including models produced by normal mode refinement (Ni et al., 2009),
as mentioned above; ensemble refinement (Burnley et al., 2012); multiconformer modeling
performed by discrete (Keedy et al., 2015a; van den Bedem et al., 2009) or continuous (Burling
and Brünger, 1994; Kuriyan et al., 1991; Wall et al., 1997a) conformational sampling; and
molecular dynamics simulations  (Clarage and Phillips, 1994; Clarage et al., 1995; Faure et al.,
1994; Héry et al., 1998; Janowski et al., 2013; Janowski et al., 2015; Meinhold et al., 2007;
Meinhold and Smith, 2005a, b; Wall et al., 2014b). In particular, molecular dynamics simulations
now provide sufficient sampling to yield robust calculations of diffuse intensity (Wall et al.,
2014b), and these can be used to consider a myriad of intramolecular motions (e.g., loop
openings and side chain flips) (Wilson, 2013) and lattice dynamics. Polikanov and Moore
(Polikanov and Moore, 2015) recently have demonstrated the importance of lattice vibrations in
explaining experimental diffuse scattering measurements of ribosome crystals, which indicates
that models should simultaneously account for correlations that are coupled both within and
across unit cell boundaries (Clarage et al., 1992; Wall et al., 1997a); accounting for lattice
vibrations more accurately also might yield improved Bragg integration (Wall et al., 2014a).
Although the initial successes of dynamics-based models of diffuse scattering indicates that
crystal defects can play a secondary role in contributing to the diffuse signal, at least in some
cases, consideration of crystal defects might become important to achieve the highest model
accuracy and most general applicability of diffuse scattering in crystallography. Additionally, as
more X-ray data from both brighter conventional and XFEL light sources, accounting for all
sources of Bragg and diffuse scattering will be necessary to model the total scattering needed
for innovative phasing applications (Gaffney and Chapman, 2007). In summary, the new
datasets presented here demonstrate that diffuse scattering can now be routinely collected and
that using these data will help us obtain an increasingly realistic picture of motion in protein
crystals, including integrated descriptions of intramolecular motions, lattice vibrations, and
crystal defects.
351
352

# Methods:

**Protein purification and crystallization**

Trypsin crystals were obtained according to the method of Liebschner *et.al* (Liebschner et al., 2013). Lyophilized bovine pancreas trypsin was purchased from Sigma-Aldrich (T1005) and dissolved at a concentration of 30 mg/mL into 30mM HEPES pH 7.0, 5 mg/mL benzamidine and 3mM $CaCl_2$. Crystals were obtained from a solution of 200mM Ammonium sulfate, 100mM Na cacodylate pH 6.5, 20% PEG 8000 and 15% glycerol. CypA was purified and crystallized as previously described (Fraser et al., 2009). Briefly, the protein was concentrated to 60 mg/mL in 20mM HEPES pH 7.5, 100mM NaCl and 500mM TCEP. Trays were set with a precipitant solution of 100mM HEPES pH 7.5, 22% PEG 3350 and 5mM TCEP. Both crystal forms were obtained using the hanging-drop method.

**Crystallographic data collection**

Diffraction data were collected on beamline 11-1 at the Stanford Synchrotron Radiation Lightsource (Menlo Park, CA). X-ray diffraction images were obtained using a Dectris PILATUS 6M Pixel Array Detector (PAD). Each dataset was collected from a single crystal at an ambient temperature of 273K. To prevent dehydration, crystals were coated in a thin film of paratone with minimal surrounding mother liquor. For CypA, a single set of 0.5 degree oscillation images were collected and used for both Bragg (Keedy et al., 2015b) and diffuse data processing. A total of 360 images were collected across a 180 degree phi rotation. The Trypsin diffraction data consisted of one degree oscillations across a 135 degree phi rotation; this dataset was similarly used for both Bragg and diffuse data analysis. Both datasets were collected to optimize the Bragg signal, not the diffuse signal. Although not used here, we note that data collection using a PAD with fine phi slicing should be especially well suited for simultaneous collection of Bragg and diffuse data, as it would enable integration of diffuse intensity at a tunable level of detail in reciprocal space.

**Bragg data processing**

Bragg diffraction data were processed using XDS and XSCALE (Kabsch, 2010) within the *xia2* software package (Winter et al., 2013). Molecular replacement solutions were found using Phaser (McCoy et al., 2007) within the *Phenix* software suite (Adams et al., 2010). The PDB search models were 4I8G for trypsin, and 2CPL for CypA. Initial structural refinement was performed using *phenix.refine* (Afonine et al., 2012). The strategy included refinement of individual atomic coordinates and water picking. Both the X-ray/atomic displacement parameters and X-ray/stereochemistry weights were optimized. Isotropic B-factors were chosen for the initial structures to allow for non-negligible R-factor optimization by subsequent TLS refinement strategies. All structures were refined for a total of 5 macrocycles. Statistics for these initial crystal structure models are shown in **Table 1**.

396    **Table 1.** Refinement statistics for CypA and trypsin models, before TLS modeling is applied.
397

| | CypA | Trypsin |
|---|---|---|
| Resolution range | 38.66-1.4 | 23.29-1.25 |
| Space group | P 21 21 21 | P 21 21 21 |
| Unit cell | 42.91, 52.44, 89.12 | 54.81, 58.51, 67.42 |
| Completeness (%) | 98 | 95 |
| $R_{work}$ (%) | 17.88 | 15.9 |
| $R_{free}$ (%) | 19.5 | 17.41 |
| RMS (bonds, Å) | 0.007 | 0.013 |
| RMS (angles, degrees) | 1.16 | 1.61 |
| Ramachandran favored % | 97 | 98 |
| Ramachandran allowed % | 3 | 2 |
| Ramachandran outliers % | 0 | 0 |
| Clashscore | 0.79 | 2.59 |
| Average B-factor, $Å^2$ | 21.42 | 14.57 |
| PDB ID | 5F66 | 5F6M |
| SBGrid Data Grid ID | 68 | 201 |

398
399    **Diffuse data integration**
400
An overview of the diffuse data integration process is presented in **Figure 1**. Image processing
was performed using the LUNUS collection of diffuse scattering tools (Wall, 2009). Pixels
corresponding to the beam stop and image edges were masked using the *punchim* and *windim*
methods. To focus on the diffuse intensity, which compared to Bragg peaks has low individual
pixel values (while, being more broadly distributed in reciprocal space, having comparable total
integrated intensity), pixel values outside of the range 1-10,000 photon counts were masked
using *threshim*. The beam polarization was determined by analyzing the first frame to determine
the azimuthal intensity profile within a 100 pixel wide annulus about the origin, and by fitting the
resulting profile to the theoretical profile (Wall, 1996). Pixel values then were corrected for beam
polarization using *polarim*. A solid-angle normalization (*normim*) correction was also applied.

Mode filtering was used to remove Bragg peaks from diffraction images. This was accomplished
using *modeim*, with a mask width of 20 pixels and a single bin for each photon count increment.
These steps produced diffraction images in which pixel values could be directly compared to
model diffuse intensities. This procedure is similar to the steps used by Polikanov and Moore
(Polikanov and Moore, 2015) to process individual ribosome diffraction images for analysis of
diffuse scattering data.
417
The Lunus processed frames were used to integrate the diffuse data onto a 3D lattice. The
integration was performed using a python script that calls DIALS methods within the
Computational Crystallography Toolbox (Parkhurst et al., 2014). The script obtains an indexing
solution using the *real_space_grid_search* method and uses the results to map each pixel in
each diffraction image to fractional Miller indices *h'k'l'* in reciprocal space. It sums the intensities
from pixels in the neighborhood of each integer Miller index *hkl* and tracks the corresponding
pixel counts, while ignoring pixels that fall within a ½ x ½ x ½ region about *hkl*. It writes the
intensity sums and pixel counts for each frame on a grid, populated on an Ewald sphere that
varies according to the crystal orientation for each image (**Figure 1C**). A radial scattering vector
intensity profile was calculated for each frame using the Lunus *avgrim* method and was used to
scale diffuse frames across the entire dataset. The Lunus *sumlt* and *divlt* methods were used to
compute the mean diffuse intensity at each grid point using the scaled sums and pixel counts
from all of the frames.
431
Because the model diffuse intensities were computed without considering solvent, experimental
and model diffuse intensities were compared using just the anisotropic component of the signal,
which is primarily due to the protein (Wall et al., 2014b). The Lunus *avgrlt* and *subrflt* routines
were applied to subtract the radial average and obtain the anisotropic signal. Signal intensities
were then symmetrized using *phenix.reflection_file_converter* to obtain a dataset for comparison
to models. Datasets were compared to each other and to models using linear correlations
computed using the *phenix.reflection_statistics* tool.
439
440 All images are available on SBGrid Data Grid (https://data.sbgrid.org/dataset/68/ for CypA;
441 https://data.sbgrid.org/dataset/201/ for Trypsin) and the integrated diffuse scattering maps are
442 available as Supplementary Material.
443
444 **TLS structure refinement and diffuse scattering model**
445
Three independent TLS refinements were performed for CypA (**Figure 3A-D**). The *Whole*
*molecule* selection consists of the entire molecule as a single TLS group. The *Phenix* selection
consists of the 8 groups (residues 2-14, 15-41, 42-64, 65-84, 85-122, 123-135, 136-145 and
146-165) identified by *phenix.find_tls_groups.* The *TLSMD* selection consists of 8 groups
(residues 2-15, 16-55, 56-80, 81-85, 86-91, 92-124, 125-143 and 144-165) identified by the TLS
Motion Determination web server (Painter and Merritt, 2005, 2006). All TLS refinement was
performed within *phenix.refine* through 5 macrocycles. Aside from the inclusion of TLS
refinement, these macrocycles were identical to the initial structure refinement described above.
454

455    Similarly, for trypsin, we selected Whole *Molecule, Phenix,* and *TLSMD* TLS refinement
456    strategies as described above (**Figure 3E-H**). The *Phenix* selection consists of 7 TLS groups:
457    residues 16-54, 55-103, 104-123, 124-140, 141-155, 156-225 and 226-245. The *TLSMD*
458    selection consists of 9 groups: residues 16-52, 53-98, 99-115, 116-144, 145-171, 172-220, 221-
459    224, 225-237 and 238-245.
460
Structural ensembles of the CypA and trypsin TLS motions were generated through
the *Phenix.tls_as_xyz* method (Urzhumtsev et al., 2015). Each ensemble consisted of 1,000
random samples of the underlying TLS atomic displacement distributions, assuming
independent distributions for each domain.  Diffuse scattering models were calculated from the
TLS ensembles using *Phenix.diffuse* (Van Benschoten et al., 2015).  CypA and trypsin models
were generated to a final resolution of 1.2 Å and 1.4 Å respectively, to match the resolution of
the experimental data.
468
469    **Liquid-like motions model**
470
We computed Liquid-like motions (LLM) models of diffuse scattering using the structures refined
prior to the TLS refinements (CypA: PDB 5F66; Trypsin: PDB 5F6M). For both CypA and
trypsin, the temperature factors for all atoms were set to zero and squared calculated structure
factors $I_0(hkl)$ were computed using the *structure_factors*, *as_intensity_array*, and *expand_to_p1*
methods in CCTBX. The Lunus *symlt* method was used to fill in missing values in reciprocal
space using the appropriate P222 Laue symmetry.
477
478    Given a correlation length $\gamma$ and amplitude of motion $\sigma$, the diffuse intensity predicted by the
479    LLM model was calculated as
480

$$D(\mathbf{s}) = 4\pi^2 s^2 \sigma^2 e^{-4\pi^2 s^2 \sigma^2} I_0(\mathbf{s}) * \Gamma_\gamma(\mathbf{s})$$

$$\Gamma_\gamma(\mathbf{s}) = \frac{8\pi\gamma^3}{(1 + 4\pi^2 s^2 \gamma^2)} \tag{1}$$

481
482    Fourier methods in Lunus (*fftlt*) were used to compute the convolution. The agreement with the
483    data was quantified by computing a linear correlation as a target function, using the anisotropic
484    intensities (**Diffuse data integration**). Optimization of the target with respect to $\gamma$ and $\sigma$ was
485    performed in a python script using *scipy.optimize* (www.scipy.org) with the Powell minimization
486    method.
487

488    **Acknowledgments**
489
We are grateful to the UC Office of the President, Multicampus Research Programs and
Initiatives grant MR-15-338599 and the Program for Breakthrough Biomedical Research, which
is partially funded by the Sandler Foundation. Use of the Stanford Synchrotron Radiation
Lightsource, SLAC National Accelerator Laboratory, is supported by the U.S. Department of
Energy, Office of Science, Office of Basic Energy Sciences under Contract No. DE-AC02-
76SF00515. The SSRL Structural Molecular Biology Program is supported by the DOE Office of

Biological and Environmental Research, and by the National Institutes of Health, National Institute of General Medical Sciences (including P41GM103393). N.K.S. was supported by NIH grant GM095887. J.S.F. was supported by a Searle Scholar Award from the Kinship Foundation, a Pew Scholar Award from the Pew Charitable Trusts, a Packard Fellowship from the David and Lucile Packard Foundation, NIH OD009180, NIH GM110580, and NSF STC-1231306. M.E.W. was supported by the US Department of Energy under Contract DE-AC52-06NA25396 through the Laboratory-Directed Research and Development Program at Los Alamos National Laboratory. This document is LANL technical release #LA-UR-15-28934.

**References**

Adams, P.D., Afonine, P.V., Bunkoczi, G., Chen, V.B., Davis, I.W., Echols, N., Headd, J.J., Hung, L.W., Kapral, G.J., Grosse-Kunstleve, R.W.*, et al.* (2010). PHENIX: a comprehensive Python-based system for macromolecular structure solution. Acta Crystallogr D Biol Crystallogr *66*, 213-221.

Afonine, P.V., Grosse-Kunstleve, R.W., Echols, N., Headd, J.J., Moriarty, N.W., Mustyakimov, M., Terwilliger, T.C., Urzhumtsev, A., Zwart, P.H., and Adams, P.D. (2012). Towards automated crystallographic structure refinement with phenix.refine. Acta Crystallogr D Biol Crystallogr *68*, 352-367.

Burling, F.T., and Brünger, A.T. (1994). Thermal motions and conformational disorder in protein crystal structures: comparison of multi-conformer and time-averaging models. Israeli J Chem *34*, 165-175.

Burnley, B.T., Afonine, P.V., Adams, P.D., and Gros, P. (2012). Modelling dynamics in protein crystal structures by ensemble refinement. Elife *1*, e00311.

Caspar, D.L., Clarage, J., Salunke, D.M., and Clarage, M. (1988). Liquid-like movements in crystalline insulin. Nature *332*, 659-662.

Chacko, S., and Phillips, G.N., Jr. (1992). Diffuse x-ray scattering from tropomyosin crystals. Biophys J *61*, 1256-1266.

Clarage, J.B., Clarage, M.S., Phillips, W.C., Sweet, R.M., and Caspar, D.L. (1992). Correlations of atomic movements in lysozyme crystals. Proteins *12*, 145-157.

Clarage, J.B., and Phillips, G.N., Jr. (1994). Cross-validation tests of time-averaged molecular dynamics refinements for determination of protein structures by X-ray crystallography. Acta crystallographica Section D, Biological crystallography *50*, 24-36.

Clarage, J.B., and Phillips, G.N., Jr. (1997). Analysis of diffuse scattering and relation to molecular motion. Methods Enzymol *277*, 407-432.

Clarage, J.B., Romo, T., Andrews, B.K., Pettitt, B.M., and Phillips, G.N., Jr. (1995). A sampling problem in molecular dynamics simulations of macromolecules. Proc Natl Acad Sci U S A *92*, 3288-3292.

Doucet, J., and Benoit, J.P. (1987). Molecular dynamics studied by analysis of the X-ray diffuse
scattering from lysozyme crystals. Nature *325*, 643-646.

Faure, P., Micu, A., Perahia, D., Doucet, J., Smith, J.C., and Benoit, J.P. (1994). Correlated
intramolecular motions and diffuse X-ray scattering in lysozyme. Nat Struct Biol *1*, 124-128.

Fraser, J.S., Clarkson, M.W., Degnan, S.C., Erion, R., Kern, D., and Alber, T. (2009). Hidden
alternative structures of proline isomerase essential for catalysis. Nature *462*, 669-673.

Fraser, J.S., van den Bedem, H., Samelson, A.J., Lang, P.T., Holton, J.M., Echols, N., and
Alber, T. (2011). Accessing protein conformational ensembles using room-temperature X-ray
crystallography. Proc Natl Acad Sci U S A *108*, 16247-16252.

Gaffney, K.J., and Chapman, H.N. (2007). Imaging atomic structure and dynamics with ultrafast
x-ray scattering. Science *316*, 1444-1448.

Gniewek, P., Kolinski, A., Jernigan, R.L., and Kloczkowski, A. (2012). Elastic network normal
modes provide a basis for protein structure refinement. J Chem Phys *136*, 195101.

Gruner, S.M. (2012). X-ray imaging detectors. Physics Today *65*, 29-34.

Guinier, A. (1963). X-ray diffraction in crystals, imperfect crystals, and amorphous bodies
(Courier Dover Publications).

Helliwell, J., Glover, I., Jones, A., Pantos, E., and Moss, D. (1986). Protein dynamics: use of
computer graphics and protein crystal diffuse scattering recorded with synchrotron X-radiation.
Biochemical Society Transactions *14*, 653-655.

Héry, S., Genest, D., and Smith, J.C. (1998). X-ray diffuse scattering and rigid-body motion in
crystalline lysozyme probed by molecular dynamics simulation. Journal of molecular biology
*279*, 303-319.

James, R. (1948). The Optical Principles of the Diffraction of X-Rays (London: Bell).

Janowski, P.A., Cerutti, D.S., Holton, J., and Case, D.A. (2013). Peptide crystal simulations
reveal hidden dynamics. J Am Chem Soc *135*, 7938-7948.

Janowski, P.A., Liu, C., Deckman, J., and Case, D.A. (2015). Molecular dynamics simulation of
triclinic lysozyme in a crystal lattice. Protein Sci.

Kabsch, W. (2010). Xds. Acta Crystallogr D Biol Crystallogr *66*, 125-132.

Keedy, D.A., Fraser, J.S., and van den Bedem, H. (2015a). Exposing Hidden Alternative
Backbone Conformations in X-ray Crystallography Using qFit. PLoS Comput Biol *11*, e1004507.

Keedy, D.A., Kenner, L.R., Warkentin, M., Woldeyes, R.A., Hopkins, J.B., Thompson, M.C.,
Brewster, A.S., Van Benschoten, A.H., Baxter, E.L., Uervirojnangkoorn, M.*, et al.* (2015b).
Mapping the conformational landscape of a dynamic enzyme by multitemperature and XFEL
crystallography. Elife *4*.

Keen, D.A., and Goodwin, A.L. (2015). The crystallography of correlated disorder. Nature *521*,
303-309.

Kidera, A., Matsushima, M., and Go, N. (1994). Dynamic structure of human lysozyme derived
from X-ray crystallography: normal mode refinement. Biophys Chem *50*, 25-31.

Kolatkar, A.R., Clarage, J.B., and Phillips, G.N., Jr. (1994). Analysis of diffuse scattering from
yeast initiator tRNA crystals. Acta Crystallogr D Biol Crystallogr *50*, 210-218.

Kuriyan, J., Osapay, K., Burley, S.K., Brunger, A.T., Hendrickson, W.A., and Karplus, M. (1991).
Exploration of disorder in protein structures by X-ray restrained molecular dynamics. Proteins
*10*, 340-358.

Kuzmanic, A., Kruschel, D., van Gunsteren, W.F., Pannu, N.S., and Zagrovic, B. (2011).
Dynamics may significantly influence the estimation of interatomic distances in biomolecular X-
ray structures. J Mol Biol *411*, 286-297.

Liebschner, D., Dauter, M., Brzuszkiewicz, A., and Dauter, Z. (2013). On the reproducibility of
protein crystal structures: five atomic resolution structures of trypsin. Acta Crystallogr D Biol
Crystallogr *69*, 1447-1462.

Lu, M., and Ma, J. (2008). A minimalist network model for coarse-grained normal mode analysis
and its application to biomolecular x-ray crystallography. Proc Natl Acad Sci U S A *105*, 15358-
15363.

Ma, P., Xue, Y., Coquelle, N., Haller, J.D., Yuwen, T., Ayala, I., Mikhailovskii, O., Willbold, D.,
Colletier, J.P., Skrynnikov, N.R.*, et al.* (2015). Observing the overall rocking motion of a protein
in a crystal. Nat Commun *6*, 8361.

McCoy, A.J., Grosse-Kunstleve, R.W., Adams, P.D., Winn, M.D., Storoni, L.C., and Read, R.J.
(2007). Phaser crystallographic software. J Appl Crystallogr *40*, 658-674.

Meinhold, L., Merzel, F., and Smith, J.C. (2007). Lattice dynamics of a protein crystal. Phys Rev
Lett *99*, 138101.

Meinhold, L., and Smith, J.C. (2005a). Correlated dynamics determining x-ray diffuse scattering
from a crystalline protein revealed by molecular dynamics simulation. Phys Rev Lett *95*,
218103.

Meinhold, L., and Smith, J.C. (2005b). Fluctuations and correlations in crystalline protein
dynamics: a simulation analysis of staphylococcal nuclease. Biophys J *88*, 2554-2563.

Meinhold, L., and Smith, J.C. (2007). Protein dynamics from X-ray crystallography: anisotropic,
global motion in diffuse scattering patterns. Proteins *66*, 941-953.

Mizuguchi, K., Kidera, A., and Go, N. (1994). Collective motions in proteins investigated by X-
ray diffuse scattering. Proteins *18*, 34-48.

Moore, P.B. (2009). On the relationship between diffraction patterns and motions in
macromolecular crystals. Structure *17*, 1307-1315.

Ni, F., Poon, B.K., Wang, Q., and Ma, J. (2009). Application of normal-mode refinement to X-ray
crystal structures at the lower resolution limit. Acta Crystallogr D Biol Crystallogr *65*, 633-643.

Painter, J., and Merritt, E.A. (2005). A molecular viewer for the analysis of TLS rigid-body
motion in macromolecules. Acta Crystallogr D Biol Crystallogr *61*, 465-471.

Painter, J., and Merritt, E.A. (2006). Optimal description of a protein structure in terms of
multiple groups undergoing TLS motion. Acta Crystallogr D Biol Crystallogr *62*, 439-450.

Parkhurst, J.M., Brewster, A.S., Fuentes-Montero, L., Waterman, D.G., Hattne, J., Ashton, A.W.,
Echols, N., Evans, G., Sauter, N.K., and Winter, G. (2014). : the diffraction experiment toolbox.
J Appl Crystallogr *47*, 1459-1465.

Perez, J., Faure, P., and Benoit, J.P. (1996). Molecular rigid-body displacements in a tetragonal
lysozyme crystal confirmed by X-ray diffuse scattering. Acta Crystallogr D Biol Crystallogr *52*,
722-729.

Phillips, G.N., Jr., Fillers, J.P., and Cohen, C. (1980). Motions of tropomyosin. Crystal as
metaphor. Biophys J *32*, 485-502.

Polikanov, Y.S., and Moore, P.B. (2015). Acoustic vibrations contribute to the diffuse scatter
produced by ribosome crystals. Acta Crystallogr D Biol Crystallogr *71*, 2021-2031.

Schomaker, V., and Trueblood, K.N. (1968). On the rigid-body motion of molecules in crystals.
Acta Crystallographica Section B *24*, 63-76.

Urzhumtsev, A., Afonine, P.V., Van Benschoten, A.H., Fraser, J.S., and Adams, P.D. (2015).
From deep TLS validation to ensembles of atomic models built from elemental motions. Acta
Crystallogr D Biol Crystallogr *71*, 1668-1683.

Van Benschoten, A.H., Afonine, P.V., Terwilliger, T.C., Wall, M.E., Jackson, C.J., Sauter, N.K.,
Adams, P.D., Urzhumtsev, A., and Fraser, J.S. (2015). Predicting X-ray diffuse scattering from
translation-libration-screw structural ensembles. Acta Crystallogr D Biol Crystallogr *71*, 1657-
1667.

van den Bedem, H., Bhabha, G., Yang, K., Wright, P.E., and Fraser, J.S. (2013). Automated
identification of functional dynamic contact networks from X-ray crystallography. Nat Methods
*10*, 896-902.

van den Bedem, H., Dhanik, A., Latombe, J.C., and Deacon, A.M. (2009). Modeling discrete
heterogeneity in X-ray diffraction data by fitting multi-conformers. Acta Crystallogr D Biol
Crystallogr *65*, 1107-1117.

van den Bedem, H., and Fraser, J.S. (2015). Integrative, dynamic structural biology at atomic
resolution--it's about time. Nat Methods *12*, 307-318.

Wall, M.E. (2009). Methods and software for diffuse X-ray scattering from protein crystals.
Methods Mol Biol *544*, 269-279.

Wall, M.E., Adams, P.D., Fraser, J.S., and Sauter, N.K. (2014a). Diffuse X-ray scattering to
model protein motions. Structure *22*, 182-184.

Wall, M.E., Clarage, J.B., and Phillips, G.N. (1997a). Motions of calmodulin characterized using
both Bragg and diffuse X-ray scattering. Structure *5*, 1599-1612.

Wall, M.E., Ealick, S.E., and Gruner, S.M. (1997b). Three-dimensional diffuse x-ray scattering
from crystals of Staphylococcal nuclease. Proceedings of the National Academy of Sciences of
the United States of America *94*, 6180-6184.

Wall, M.E., Van Benschoten, A.H., Sauter, N.K., Adams, P.D., Fraser, J.S., and Terwilliger, T.C.
(2014b). Conformational dynamics of a crystalline protein from microsecond-scale molecular
dynamics simulations and diffuse X-ray scattering. Proc Natl Acad Sci U S A *111*, 17887-17892.

Welberry, T.R. (2004). Diffuse X-Ray Scattering and Models of Disorder (Oxford: Oxford
University Press).

Welberry, T.R., Heerdegen, A.P., Goldstone, D.C., and Taylor, I.A. (2011). Diffuse scattering
resulting from macromolecular frustration. Acta Crystallogr B *67*, 516-524.

Wilson, M.A. (2013). Visualizing networks of mobility in proteins. Nat Methods *10*, 835-837.

Winter, G., Lobley, C.M., and Prince, S.M. (2013). Decision making in xia2. Acta Crystallogr D
Biol Crystallogr *69*, 1260-1273.
657