1

Genomic analysis reveals major determinants of *cis*-regulatory variation in
Capsella grandiflora

Kim A. Steige[1,2,§], Benjamin Laenen[2,§,*], Johan Reimegård[3], Douglas G. Scofield[1,4],
Tanja Slotte[1,2,*]

[1]Dept. of Ecology and Genetics, Evolutionary Biology Centre, Uppsala University,
Norbyv. 18D, 75236 Uppsala, SWEDEN
[2]Science for Life Laboratory, Dept. of Ecology, Environment and Plant Sciences,
Stockholm University, Lilla Frescati, SE-10691 Stockholm, SWEDEN
[3]Science for Life Laboratory, Dept. of Cell and Molecular Biology, Uppsala
University, Box 596, 75124 Uppsala, SWEDEN
[4]Uppsala Multidisciplinary Center for Advanced Computational Science, Department
of Information Technology, Uppsala University, Box 137, Uppsala 751 05, SWEDEN

[§]These authors contributed equally to the work.

*Authors for correspondence, email: Benjamin Laenen; Benjamin.Laenen@su.se;
Tanja Slotte; Tanja.Slotte@su.se

Keywords
allele-specific expression; distribution of fitness effects; purifying selection; positive
selection; transposable elements; gene-body methylation

Understanding the causes of *cis*-regulatory variation is a long-standing aim in

evolutionary biology. Although *cis*-regulatory variation has long been considered

important for adaptation, we still have a limited understanding of the selective

importance and genomic determinants of standing *cis*-regulatory variation. To

address these questions, we studied the prevalence, genomic determinants and

selective forces shaping *cis*-regulatory variation in the outcrossing plant *Capsella

grandiflora*. We first identified a set of 1,010 genes with common *cis*-regulatory

variation using analyses of allele-specific expression (ASE). Population genomic

analyses of whole-genome sequences from 32 individuals showed that genes with

common *cis*-regulatory variation are 1) under weaker purifying selection and 2)

undergo less frequent positive selection than other genes. We further identified

genomic determinants of *cis*-regulatory variation. Gene-body methylation (gbM)

was a major factor constraining *cis*-regulatory variation, whereas presence of

nearby TEs and tissue specificity of expression increased the odds of ASE. Our

results suggest that most common *cis*-regulatory variation in *C. grandiflora* is

under weak purifying selection, and that gene-specific functional constraints are

more important for the maintenance of *cis*-regulatory variation than genome-

scale variation in the intensity of selection. Our results agree with previous

findings that suggest TE silencing affects nearby gene expression, and provide

novel evidence for a link between gbM and *cis*-regulatory constraint, possibly

reflecting greater dosage-sensitivity of body-methylated genes. Given the

extensive conservation of gene-body methylation in flowering plants, this

suggests that gene-body methylation could be an important predictor of *cis*-

regulatory variation in a wide range of plant species.

Significance

Despite long-standing interest in the contribution of *cis*-regulatory changes to adaptation, we still have a limited understanding of the selective importance and genomic determinants of *cis*-regulatory variation in natural populations. Using a combination of analyses of allele-specific expression and population genomic analyses, we investigate the selective forces and genomic determinants of *cis*-regulatory variation in the outcrossing plant species *Capsella grandiflora*. We conclude that gene-specific functional constraints shape *cis*-regulatory variation and that genes with *cis*-regulatory variation are under relaxed purifying selection compared to other genes. Finally, we identify a novel link between gene-body methylation and the extent of *cis*-regulatory constraint in natural populations.

Introduction

Understanding the causes of regulatory variation is of major importance for many areas of biology and medicine (1). Much interest has centered on *cis*-regulatory variation, which has long been thought to be particularly important for adaptation (2-5). Like other quantitative traits, *cis*-regulatory variation is expected to be shaped by the interplay of mutation, selection, and drift. However, the relative importance of these forces remains unclear in most species.

Recently, prospects for quantifying *cis*-regulatory variation have greatly improved, and as a result, ample heritable cis-regulatory variation has been identified in many species (reviewed in (6)). This is resulting in a growing consensus that a large amount of standing *cis*-regulatory variation is under weak purifying selection (7-9). Clarifying why the impact of purifying selection varies across the genome is therefore important to understand the maintenance of *cis*-regulatory variation.

Variation in the intensity of purifying selection across the genome can result from differences in selective constraint that are due to the specific functions of the genes involved. For example, according to the dosage balance hypothesis, genes that encode interacting proteins are expected to experience stronger constraint than other genes (10). In yeast, there is empirical evidence that purifying selection on expression noise constrains regulatory evolution of dosage-sensitive genes (11-13) and in plants, dosage-sensitivity affects the retention of duplicate genes following whole-genome duplication (14). However, many other genomic features, including expression level,

85   tissue specificity and gene-body methylation (gbM), are also known to be associated

86   with constraint (15-18) and could affect *cis*-regulatory variation.

87   Variation in purifying selection can also result from broad, genome-scale

forces that affect genes mainly as a result of their genomic environment, and not due

to their specific function. For instance, in the self-fertilizing species *Caenorhabditis*

elegans, variation in the impact of background selection across the genome had a

major effect on the distribution of *cis*-regulatory variation across the genome (8). If

background selection is important, then one might generally expect levels of *cis*-

regulatory variation to be associated with recombination rate and/or gene density (19).

At present however, the relative importance of gene-level constraint vs. genome-scale

evolutionary forces for the distribution of *cis*-regulatory variation remains unclear in

most species.

97   In this study, we have investigated the selective importance and genomic

correlates of common *cis*-regulatory variation in the outcrossing crucifer species

Capsella grandiflora. This species is particularly well suited for studying differences

in the impact of selection across the genome, as it has relatively low population

structure (20) and a large, stable effective population size (21, 22). Indeed, selection

on both protein-coding (23) and regulatory regions (18) is highly efficient in *C.*

grandiflora, and high levels of polymorphism enhance the power to detect *cis*-

regulatory variation and quantify selection. Genomic studies are facilitated by the

close relationship between *C. grandiflora* and the selfing species *Capsella rubella*, for

which a genome sequence is available (22).

107   Here, we identified genes with common *cis*-regulatory variation in *C.*

grandiflora based on analyses of allele-specific expression (ASE) in deep

transcriptome sequencing data. To quantify the impact of positive and purifying

selection on genes with *cis*-regulatory variation, we conducted population genomic

analyses of high-coverage whole genome resequencing data from 32 *C. grandiflora*

individuals. Finally, we identified genomic predictors of *cis*-regulatory variation. Our

results show that there is pervasive *cis*-regulatory variation in *C. grandiflora*, and

genes that harbor *cis*-regulatory variation are under weaker purifying selection and

undergo less frequent positive selection than other genes. We find no evidence for a

role of recombination rate or gene density in shaping *cis*-regulatory variation,

suggesting that gene-specific variation in functional constraint is more important in

this species. We further identify gbM as a major factor constraining *cis*-regulatory

variation, whereas presence of nearby TEs and tissue specificity of expression

increase the odds of ASE. Our results provide novel evidence for a link between gbM

and *cis*-regulatory constraint, possibly reflecting greater dosage-sensitivity of body-

methylated genes.

Results

Widespread cis-regulatory variation in C. grandiflora

To identify genes with *cis*-regulatory variation, we quantified allele-specific

expression based on deep whole transcriptome sequencing data (total 93.5 Gbp with

Q≥30) from flower buds and leaves of three *C. grandiflora* F1s (Supplementary Table

S1). Each F1 harbored an average of about 235,700 high-confidence heterozygous

coding SNPs, which were phased prior to analyses of ASE. After filtering, on average

approximately 13,400 genes per F1 were amenable to ASE analyses (Table 1).

We assessed ASE using a Bayesian method (24), accounting for technical

variation in allelic counts using high-coverage whole genome resequencing data for

each F1 (mean coverage of 40x, total 26.6 Gbp with Q≥30; Supplementary Table S2).

We estimated that a mean of 35% (range 33-39%) of analyzed genes show ASE in

individual *C. grandiflora* F1s (Table 1). Similar proportions of genes had ASE in both

leaves and flower buds (Table 1) and allelic expression biases were moderate for most

genes with ASE, with strong allelic expression biases ($0.2 \leq$ ASE ratio ≥ 0.8) shown

by an average of 5.1% of genes (Figure 1, Supplementary Figures S1 and S2).

Out of a total of 11,532 genes that were amenable to analysis of ASE in all

F1s, there were 1,010 genes that showed ASE in either leaves or flower buds, 313

genes showed ASE in flower buds but not leaves, 404 genes showed ASE in leaf

samples but not flower buds, and 293 genes had ASE in both flower buds and leaves

of all F1s (Supplementary Figure S3). Among the 1,010 genes with ASE leaves or

flower buds of all F1s, one GO category, GO:0006952, "defense response" was

significantly enriched at FDR ≤ 0.01. This was likely driven by genes with ASE in

leaves, as there was no significant enrichment of Gene Ontology (GO) terms among

genes with ASE in flower buds, whereas six biological process GO terms associated

with photosynthesis and defense responses were significantly enriched (FDR ≤ 0.01)

among genes with ASE in leaves (Supplementary Table S3). Among control genes,

there was a nominally significant enrichment of genes in only two GO terms, protein

binding (GO:0005515) and zinc ion binding (GO:0008270) (Weighted Fisher $P \leq$

0.01), but this was not significant at FDR ≤ 0.01.

Lower intensity of purifying selection on genes with cis-regulatory variation

To assess the impact of selection on genes showing *cis*-regulatory variation in *C.*

grandiflora, we sequenced the genomes of 21 individuals from one population in the

Zagory region of Greece (the 'population sample') as well as 12 individuals from

separate populations across the species range (the 'range-wide sample') using 233.2

Gbp of high-quality (Q≥30) paired-end 100 bp Illumina reads and a mean coverage of

25x per individual (Supplementary Table S2). We called variants using GATK best

practices and filtered genomic regions as previously described (25) to identify a total

of 6,492,075 high-quality SNPs, most of which (5,240,485) were also segregating in

the population sample.

We compared levels of polymorphism at genes that show ASE in all of our

F1s (1,010 genes; 'ASE genes'), using as a control set the 10,552 genes that were

amenable to ASE analyses in all F1s but did not show significant ASE (termed

'control genes') (Supplementary Figure S3). To reduce bias resulting from the

requirement of expressed polymorphisms for analyses of ASE, all population genetic

analyses were conducted only on these paired gene sets, and genes that were not

amenable to analysis of ASE were not included. ASE genes had elevated

polymorphism levels compared to the control at all investigated site classes, as well as

an elevated ratio of nonsynonymous to synonymous polymorphism (Table 2;

Supplementary Table S4), suggesting that the impact of purifying selection might

differ between ASE and control gene sets (Table 2; Supplementary Table S4).

To quantify the impact of purifying selection on ASE genes and control genes,

we used the DFE-alpha method (26, 27), which allows estimation of a gamma-

distribution of negative fitness effects based on site frequency spectra (SFS) at

putatively neutral and selected sites. We found that ASE genes have a significantly

higher proportion of nearly neutral nonsynonymous mutations than control genes, as

well as a significantly reduced proportion of nonsynonymous mutations under strong

purifying selection (strength of purifying selection N_es>10) (Figure 2). This result

applies broadly, both for the population and the range-wide samples, and when

assuming a constant population size as well as after correcting for population size

change (Supplementary Figure S4). The result also holds after controlling for

differences in the expression level among genes with and without ASE

(Supplementary Figure S5), and when classifying genes based on a single F1

individual (Supplementary Figure S6), suggesting that the results hold broadly for

common *cis*-regulatory variation. Our results further remain unchanged after

removing defense-response genes (GO:0006952) with ASE (Supplementary Figure

S7) prior to DFE-alpha analyses, and thus strong balancing selection on these genes

does not drive the patterns we observe.

In contrast to the clear evidence for weaker purifying selection at

nonsynonymous sites for genes with ASE, there were no significant differences in the

DFE depending on ASE status at 5'-UTRs (Supplementary Figure S8). For introns,

results were inconsistent, with some but not all analyses pointing to weaker purifying

selection on control genes (Figure 2, Supplementary Figure S9, Supplementary Table

S5). This could suggest that patterns of selection differ among coding and noncoding

regions. However, at other noncoding regions than introns, such as promoter regions

500 bp upstream of the TSS and at 3'-UTRs, there was some evidence for relaxed

purifying selection at ASE genes (Figure 2; Supplementary Figures S10-S11,

Supplementary Table S5). These results held only under the 1-epoch model, which

could in part be due to a lack of power, as regulatory motifs are expected to make up a

small fraction of the analyzed sites. Consistent with this, we infer weaker purifying

selection on upstream regions and UTRs than on nonsynonymous mutations

(Supplementary Figures S8-S11; Supplementary Table S5).

Genes with cis-regulatory variation undergo less frequent adaptive evolution

To investigate the impact of positive selection on genes with and without ASE we

obtained estimates of ω_a, the rate of adaptive substitutions relative to neutral

divergence (28) in DFE-alpha. For this purpose, we relied on genome-wide

divergence between *Capsella* and *Arabidopsis*, with 4-fold synonymous sites

considered to be evolving mainly neutrally (see Methods for details). Using this

method, we find that ASE genes show a significantly lower proportion of adaptive

nonsynonymous substitutions than control genes (Figure 3). In contrast, we found no

significant differences in ω_a among ASE genes or control genes for UTRs or regions

500 bp upstream of the TSS (Supplementary Table S5). Second, we estimated α, the

proportion of adaptive fixations in the selected site class, based on the approximate

method of (29), designed to yield accurate estimates in the presence of linked

selection. Results generated with this method were consistent with DFE-alpha, with a

significantly lower estimate of the proportion of adaptive nonsynonymous

substitutions at genes with *cis*-regulatory variation than at control genes in *C.*

grandiflora (Figure 3).

Determinants of cis-regulatory variation in C. grandiflora

To identify genomic factors and potential drivers of *cis*-regulatory variation, we

conducted logistic regression analyses with presence/absence of ASE as the response

variable. We included a total of 12 predictor variables, chosen to include proxies for

variation in mutation rate, recombination rate, gene density, expression level and

degree of constraint, which could be expected to affect levels of *cis*-regulatory

variation (see Methods for details). The best-fit model retained eight of these

predictor variables (Table 3). In this model, gbM had the greatest effect on *cis*-

regulatory variation, resulting in a reduction of 49% in the odds of observing ASE

(Table 3), whereas the presence of polymorphic TEs within 1 kb of the gene also had

a substantial effect, increasing the odds of ASE by 38%, followed in turn by tissue

specificity of expression, promoter diversity, expression level, gene length and

nonsynonymous/synonymous polymorphism, all of which increased the odds of ASE

(Table 3). Including network connectivity improved model fit, although the effect was

not individually significant (Table 3). Notably, gene density and recombination rate,

which affect the intensity of linked selection, were not included in the best-fit model.

Similar results were obtained in an analysis that followed the approach of (30) to

ensure orthogonality of predictors by using principal components of all continuous

predictors in logistic regression analyses (Supplementary Tables S6 and S7). These

analyses suggest that variation in gene-specific constraint are important for shaping

the distribution of *cis*-regulatory variation across the *C. grandiflora* genome, and that

gbM and presence of nearby TEs are strong predictors of *cis*-regulatory constraint.

Discussion

Our results show that genes that harbor common *cis*-regulatory variation in *C.*

grandiflora are under weaker purifying selection, and experience less frequent

positive selection than other genes. We further find that gene-specific features that are

likely to reflect the degree of functional constraint and mutational input are better

predictors of *cis*-regulatory variation than those that are expected to shape the broad

impact of linked selection across the genome. These functional constraints do not

appear to limit the potential for adaptation at coding sequences, as positive selection

had a greater impact on coding divergence at genes that did not exhibit common *cis*-

regulatory variation in *C. grandiflora*.

Our findings support the view that most standing *cis*-regulatory variation in

natural populations is weakly deleterious (7), and our robust inference of relaxed

purifying selection on genes with common *cis*-regulatory variation agrees well with

those of a recent eQTL mapping study in *C. grandiflora* (9). Our inference of relaxed

purifying selection on genes with common *cis*-regulatory variation do not appear to be

driven by balancing selection or conditional neutrality affecting a subset of defense-

related genes that show ASE, as our results remain unchanged after removing such

genes.

The major association between gbM and *cis*-regulatory constraint that we

detected is particularly interesting, because the function of gbM is currently unclear

(31, 32). The conservation of gbM of orthologs in very distantly related plant species

suggests that gbM has functional importance, but intriguingly, some plants lack gbM

(31-33). Body-methylated genes tend to be longer than other genes, expressed at

intermediate levels, evolve slowly at the sequence level (17, 34, 35), and are stably

expressed under different conditions (36). A recent study found that *A. thaliana* from

northern Sweden show elevated gene body methylation, mainly due to *trans*-acting

loci (36), but as far as we are aware, no study has directly linked gbM to *cis*-

regulatory variation in natural plant populations.

It is possible that the associations we detected between genomic features and

cis-regulatory variation are caused by underlying drivers that were not directly

measured. One natural candidate is gene essentiality. However, while gbM is

significantly associated with predicted gene essentiality (37) (Fisher exact test

P<0.001), our results do not appear to be driven by essentiality, which was not

retained in our best-fit logistic regression model for *cis*-regulatory variation. Instead,

we hypothesize that selection for increased stability of expression of dosage-sensitive

genes could underlie several of the associations we observe. Dosage-sensitive genes

exhibit less expression noise (12, 38), show less variation in expression among tissues,

and are expected to be part of larger regulatory network modules(10, 12). In our study,

reduced tissue-specificity of expression and increased network connectivity were

associated with a reduced likelihood of ASE. Furthermore, expression variation

among three biological replicates of a *C. rubella* genotype (25) that likely represents

mainly noise, is significantly lower for genes with no ASE than for those with ASE

(median CV of FPKM=0.28 for genes with ASE, 0.18 for control genes, Wilcoxon

rank-sum test, P-value<10^{-5}). Finally, defense-related genes, which are thought to be

dosage-insensitive in plants (39), were significantly enriched among genes with *cis*-

regulatory variation in our study, whereas protein-binding genes were nominally

enriched among control genes without ASE. Both promoter polymorphism and TE

insertions, which can impact expression in several ways (40), might be expected to be

more likely to be tolerated near dosage-insensitive genes. Our results are therefore

consistent with dosage sensitivity causing strong constraint on *cis*-regulatory variation

and shaping the impact of positive and purifying selection on coding variation. Thus,

similar functional constraints that shape duplicate gene retention after whole genome

duplication (14) may also be key for the genomic distribution of *cis*-regulatory

variation in natural plant populations. Future studies should explore the connection

between dosage-sensitivity, gbM, and *cis*-regulatory variation in greater detail across

a wider range of plant species.

Materials and Methods

Plant Material

For analyses of ASE, we generated three intraspecific *C. grandiflora* F1s by crossing

six individuals sampled across the range of *C. grandiflora* (Supplementary Table S8).

For population genomic analyses of *C. grandiflora*, we grew a single offspring from

field-collected seeds of each of 32 plants ('the population genomic sample';

Supplementary Table S9), representing 21 plants from one population from Greece

(the 'population sample'), and 11 additional plants from 11 separate Greek

populations covering the species' range. Together with an individual from the

population sample, these represent a 12-plant 'range-wide sample'. We grew plants at

standard long-day conditions and collected leaf and mixed stage flower bud samples

for RNA sequencing, and leaf samples for whole genome sequencing as previously

described (25).

Sample preparation and sequencing

We extracted total RNA from the intraspecific F1s using a Qiagen RNEasy Plant Mini

Kit (Qiagen, Hilden, Germany). RNAseq libraries were constructed using the TruSeq

RNA v2 kit. For genomic resequencing, we extracted genomic DNA using a modified

CTAB extraction method. Whole genome sequencing libraries with an insert size of

300-400 bp were prepared using the TruSeq DNA v2 protocol. Sequencing of 100bp

paired-end reads was performed on an Illumina HiSeq 2000 instrument (Illumina, San

Diego, CA, USA). All sequence data has been submitted to the European

Bioinformatics Institute (www.ebi.ac.uk), with study accession numbers:

PRJEB12070 and PRJEB12072.

Sequence quality and trimming

RNA and DNA reads from the F1s were trimmed as previously described (25).

Adapters and low quality sequence were trimmed using CutAdapt 1.3. We analyzed

genome coverage using BEDTools v.2.17.0 (41) and removed potential PCR

duplicates using Picard v.1.92 (http://picard.sourceforge.net).

Read mapping, variant calling and filtering

We mapped RNAseq reads from the F1s to the v1.0 reference *C. rubella* assembly

(22) using STAR v.2.3.0.1 (42) with default parameters. For genomic reads from F1s,

we mapped reads with STAR as in (25). Genomic reads from the population genomic

sample were mapped using BWA-MEM v.0.7.12 (43) using default parameters and

the –M flag.

Variant calling was done using GATK v. 2.5-2 UnifiedGenotyper (44)

according to GATK best practices (45, 46). We conducted duplicate marking, local

realignment around indels and recalibrated base quality scores using a set of

1,538,085 SNPs identified in *C. grandiflora* (18) as known variants, and retained only

SNPs considered high quality by GATK.

We removed centromeric and pericentromeric regions where we have low

confidence in our variant calls, and prior to ASE analysis, we conducted additional

filtering of SNPs as in (25). Using this procedure, we identified an average of 235,719

heterozygous coding SNPs in 17,973 genes in each F1. For population genomic

analyses, we further filtered all genomic regions annotated as repeats using

RepeatMasker 4.0.1, and removed sites with extreme coverage (DP < 15 or DP > 200)

and too many missing individuals (≥20%) using VCFtools (47). Indels and non-

biallelic SNP were also pruned prior to analysis.

Phasing

Prior to ASE analysis, we conducted read-backed phasing of genomic variants in F1s

using GATK v. 2.5-2 ReadBackPhasing (-phaseQualityThresh 10). RNAseq data

from all F1s were subsequently phased by reference to the phased genomic variants.

Read counts for all phased fragments were obtained using Samtools mpileup. This

resulted in a mean number of 31,313 contiguous phased fragments per F1 (Table 1).

To validate our phasing procedure, we compared the phased fragments, based

on reads, with the phased chromosomes, based on heritage, in three interspecific *C.*

grandiflora x *C. rubella* F1s from (25). For most genes, over 95% of SNPs were

correctly phased in the interspecific F1s, demonstrating that our phasing procedure is

reliable (Supplementary Figure S13; Supplementary Figure S14).

Analyses of allele-specific expression

We analyzed ASE using a hierarchical Bayesian method which requires phased data,

in the form of read counts at heterozygous SNPs for both genomic and transcriptomic

data (24). Genomic read counts are used to obtain an empirical estimate of technical

variation which is then used in analyses of the RNAseq data. We used this method to

obtain estimates of the posterior probability and degree of ASE, for the longest phased

fragment per gene with at least three transcribed SNPs. We analyzed ~14,000 reliably

expressed genes for ASE in flower buds, and ~13,400 genes in leaves (Table 1). All

analyses were run in triplicate, and we checked MCMC convergence by comparing

parameter estimates from independent runs with different starting points, and by

assessing mixing. Runs were completed using the pqR version of R (http://www.pqr-

project.org) for 200,000 generations or a maximum runtime of 10 days, with the first

10% of each run discarded as burn-in.

Population genomic analyses

To assess whether patterns of polymorphism differ among ASE and control genes, we

tested for a difference in median levels of polymorphism and Tajima's *D* in the *C.*

grandiflora population sample, using Mann-Whitney U-tests, with Benjamini-

Hochberg correction for multiple comparisons. Estimates of nucleotide diversity (π),

Watterson's theta (θ_W) and Tajima's D (D_T) were obtained using custom R scripts by

BL. Separate estimates were obtained for 6 classes of sites: 4-fold degenerate sites, 0-

fold degenerate sites, 3'- and 5'-untranslated regions (UTRs), introns, and intergenic

regions 500 bp upstream of the transcription start site (TSS).

Selection on genes with ASE

To test whether there was evidence for a difference in the strength and direction of

natural selection on ASE and control genes, we first estimated the distribution of

fitness effects (DFE) as in (26), and the proportion of adaptive substitutions relative to

the total number of synonymous substitutions (ω_a) using the method of (28). The DFE

was estimated under a constant population size model and under a model with

stepwise population size change. We obtained confidence intervals for our estimates

of three bins of the DFE ($0<N_es<1$; $1<N_es<10$; $10<N_es$) and for α and ω_a by

resampling genes in 200 bootstrap replicates and tested for a difference in the DFE,

and ω_a among sets of genes with ASE and control genes, as in (27). Separate

estimates were obtained for 0-fold degenerate sites, 3'- and 5'-untranslated regions

(UTRs), introns, and promoter regions 500 bp upstream of the TSS likely enriched for

regulatory elements, using 4-fold degenerate sites as neutral standard. For estimates of

α and ω_a, we relied on divergence to *Arabidopsis*; specifically, we generated a whole

genome alignment using lastz v. 1.03.54, with chaining of *C. rubella*, *A. thaliana* and

A. lyrata as described in (48), and counted divergence differences and sites as in (18).

DFE-alpha analyses were run using Method I (27).

To assess the effect of expression level on our DFE-alpha inference, we

selected genes among the control set of genes to match the distribution of expression

level of ASE gene by resampling the control genes to match the distribution of

expression levels in the ASE gene set. Purifying selection and positive selection were

then re-evaluated in DFE-alpha, using the resampled control gene set and the ASE set.

To assess whether our results were robust to the sampling strategy for ASE analyses,

we based our classification of ASE and control genes based on a single F1 individual

and repeated the DFE analyses. To investigate whether our results could be driven by

the inclusion of defense-related genes, we removed genes annotated as defense-

response genes (GO:0006952), and repeated the DFE-alpha analyses.

Genomic determinants of cis-regulatory variation

We assessed the relative importance of a number of genomic features for

presence/absence of ASE using logistic regression, on a set of genes that was

restricted to those for which we could assess ASE. We included the following

genomic features that may affect linked selection: recombination rate and gene

density (in 50 kb windows). Gene density were based on the annotation of *C. rubella*

v1.0 reference genome (22). We obtained recombination rates per 50kb windows

based on 878 markers from (49) by fitting a smooth spline. We further included gene

length, tissue specificity (*t*;(16)), expression level (log FPKM values), and as a proxy

for mutation rate variation, we included 4-fold synonymous divergence to

Arabidopsis (d_S). Because promoter polymorphism may cause *cis*-regulatory variation,

we included nucleotide diversity (π) for the region 500bp upstream of the TSS. We

included nonsynonymous/synonymous nucleotide diversity (π_N/π_S) to reflect the level

of constraint at the coding sequence level. According to the dosage balance

hypothesis, genes in smaller co-expression modules may be under reduced regulatory

constraint. We therefore included information on *A. thaliana* co-expression module

size (37) in our analyses. We further included information on the presence of retained

paralogs from the Brassicaceae α whole genome duplication or the β and γ whole

genome duplication (37). We identified a set of genes with gbM in both *C. rubella*

(33) and *A. thaliana* (17), which are highly likely to also harbor gbM in *C.*

grandiflora. Finally, we included information on polymorphic TEs within 1 kb of

genes in the range-wide sample. We identified TE insertions in our range-wide

sample as in (25), except that we required a minimum of 5 reads to call a TE insertion.

All continuous variables were centered and scaled prior to logistic regression, with

model selection using a stepwise AIC procedure with backward and forward selection

of variables to find the best-fit model (Table 3). We repeated the analysis using an

analysis strategy which is superior to partial correlation analysis and robust in the

presence of noisy genomic data and multicollinearity of predictor variables (30). We

used a set of orthogonal predictor variables obtained by identifying principal

components for a data set including all the continuous variables using the "pls"

package in R, well as gene body methylation and presence of heterozygous TEs as

binary factors, and conducted model selection as described above.

Acknowledgements

We thank Lauren McIntyre and Stephen Wright for valuable discussion, Daniel Halligan for sharing scripts for DFE-alpha analyses and Daniel Skelly for advice on ASE analyses. We thank Veronika Scholz and Michael Nowak for bioinformatic assistance, and Julia Dankanich and Cindy Canton for assistance with experiments and lab work. Sequencing was performed by the SNP&SEQ Technology Platform in Uppsala. The computations were performed on resources provided by SNIC through Uppsala Multidisciplinary Center for Advanced Computational Science (UPPMAX) under projects b2012122 and b2012190. This study was funded by grants from the Swedish Research Council, the Nilsson-Ehle foundation, the Magnus Bergvall foundation, and the Erik Philip-Sörensen foundation to T.S.

References

1. Albert FW, Kruglyak L (2015) The role of regulatory variation in complex traits and disease. *Nat Rev Genet* 16(4):197–212.

2. King MC, Wilson AC (1975) Evolution at two levels in humans and chimpanzees. *Science* 188(4184):107-116

3. Wray GA (2007) The evolutionary significance of cis-regulatory mutations. *Nat Rev Genet* 8(3):206–216.

4. Carroll SB (2008) Evo-devo and an expanding evolutionary synthesis: a genetic theory of morphological evolution. *Cell* 134(1):25–36.

5. Wittkopp PJ, Kalay G (2012) Cis-regulatory elements: molecular mechanisms and evolutionary processes underlying divergence. *Nat Rev Genet* 13(1):59–69.

6. Fraser HB (2011) Genome-wide approaches to the study of adaptive gene expression evolution. *Bioessays* 33(6):469–477.

7. Fay JC, Wittkopp PJ (2008) Evaluating the role of natural selection in the evolution of gene regulation. *Heredity* 100(2):191–199.

8. Rockman MV, Skrovanek SS, Kruglyak L (2010) Selection at linked sites shapes heritable phenotypic variation in C. elegans. *Science* 330(6002):372–376.

9. Josephs EB, Lee YW, Stinchcombe JR, Wright SI (2015) Association mapping reveals the role of purifying selection in the maintenance of genomic variation in gene expression. *Proc Natl Acad Sci USA* 112(50):15390-15395.

10. Birchler JA, Veitia RA (2012) Gene balance hypothesis: connecting issues of dosage sensitivity across biological disciplines. *Proc Natl Acad Sci USA*

491   109(37):14746–14753.

11. Lemos B, Meiklejohn CD, Hartl DL (2004) Regulatory evolution across the protein interaction network. *Nat Genet* 36(10):1059–1060.

12. Lehner B (2008) Selection to minimise noise in living systems and its implications for the evolution of gene expression. *Mol Syst Biol* 4(1):170.

13. Metzger BPH, Yuan DC, Gruber JD, Duveau F, Wittkopp PJ (2015) Selection on noise constrains variation in a eukaryotic promoter. *Nature* 521(7552):344-347.

14. Li Z, et al. (2016) Gene duplicability of core genes is highly consistent across all angiosperms. *Plant Cell* 28(2):326–344.

15. Rocha EPC (2006) The quest for the universals of protein evolution. *Trends Genet* 22(8):412–416.

16. Slotte T, et al. (2011) Genomic determinants of protein evolution and polymorphism in Arabidopsis. *Genome Biol Evol* 3:1210–1219.

17. Takuno S, Gaut BS (2012) Body-methylated genes in *Arabidopsis thaliana* are functionally important and evolve slowly. *Mol Biol Evol* 29(1):219–227.

18. Williamson RJ, et al. (2014) Evidence for widespread positive and negative selection in coding and conserved noncoding regions of *Capsella grandiflora*. *PLoS Genet* 10(9):e1004622.

19. Slotte T (2014) The impact of linked selection on plant genomic variation. *Brief Funct Genomics* 13(4):268–275.

20. St Onge KR, Källman T, Slotte T, Lascoux M, Palmé AE (2011) Contrasting demographic history and population structure in *Capsella rubella* and *Capsella grandiflora*, two closely related species with different mating systems. *Mol Ecol* 20(16):3306–3320.

21. Foxe JP, et al. (2009) Recent speciation associated with the evolution of selfing in *Capsella*. *Proc Natl Acad Sci USA* 106(13):5241–5245.

22. Slotte T, et al. (2013) The *Capsella rubella* genome and the genomic consequences of rapid mating system evolution. *Nat Genet*. 45:831-835.

23. Slotte T, Foxe JP, Hazzouri KM, Wright SI (2010) Genome-wide evidence for efficient positive and purifying selection in *Capsella grandiflora*, a plant species with a large effective population size. *Mol Biol Evol* 27(8):1813–1821.

24. Skelly DA, Johansson M, Madeoy J, Wakefield J, Akey JM (2011) A powerful and flexible statistical framework for testing hypotheses of allele-specific gene

expression from RNA-seq data. *Genome Res* 21(10):1728–1737.

25. Steige KA, Reimegård J, Koenig D, Scofield DG, Slotte T (2015) *Cis*-
regulatory changes associated with a recent mating system shift and floral
adaptation in *Capsella*. *Mol Biol Evol* 32(10):2501–2514.

26. Keightley PD, Eyre-Walker A (2007) Joint inference of the distribution of
fitness effects of deleterious mutations and population demography based on
nucleotide polymorphism frequencies. *Genetics* 177(4):2251–2261.

27. Eyre-Walker A, Keightley PD (2009) Estimating the rate of adaptive molecular
evolution in the presence of slightly deleterious mutations and population size
change. *Mol Biol Evol* 26(9):2097–2108.

28. Gossmann TI, et al. (2010) Genome wide analyses reveal little evidence for
adaptive evolution in many plant species. *Mol Biol Evol* 27(8):1822–1832.

29. Messer PW, Petrov DA (2013) Frequent adaptation and the McDonald-
Kreitman test. *Proc Natl Acad Sci USA* 110(21):8615-8620.

30. Drummond DA, Raval A, Wilke CO (2006) A single determinant dominates
the rate of yeast protein evolution. *Mol Biol Evol* 23(2):327–337.

31. Takuno S, Ran J-H, Gaut BS (2016) Evolutionary patterns of genic DNA
methylation vary across land plants. *Nature Plants* 2(2):15222.

32. Bewick AJ, et al. (2016) On the origin and evolutionary consequences of gene
body DNA methylation *Proc Natl Acad Sci USA* doi:
10.1073/pnas.1604666113

33. Niederhuth CE, et al. (2016) Widespread natural variation of DNA methylation
within angiosperms *bioRxiv* doi:10.1101/045880.

34. Takuno S, Gaut BS (2013) Gene body methylation is conserved between plant
orthologs and is of evolutionary consequence. *Proc Natl Acad Sci USA*
110(5):1797–1802.

35. Seymour DK, Koenig D, Hagmann J, Becker C, Weigel D (2014) Evolution of
DNA methylation patterns in the Brassicaceae is driven by differences in
genome organization. *PLoS Genet* 10(11):e1004785.

36. Dubin MJ, et al. (2015) DNA methylation in Arabidopsis has a genetic basis
and shows evidence of local adaptation. *Elife* 4:e05255.

37. Lloyd JP, Seddon AE, Moghe GD, Simenc MC, Shiu S-H (2015)
Characteristics of plant essential genes allow for within- and between-species
prediction of lethal mutant phenotypes. *Plant Cell* 27(8):2133–2147.

38. Fraser HB, Hirsh AE, Giaever G, Kumm J, Eisen MB (2004) Noise
minimization in eukaryotic gene expression. *Plos Biol* 2(6):e137.

39. Coate JE, Song MJ, Bombarely A, Doyle JJ (2016) Expression-level support
for gene dosage sensitivity in three *Glycine* subgenus *Glycine* polyploids and
their diploid progenitors. *New Phytol*. doi:10.1111/nph.14090.

40. Feschotte C (2008) Transposable elements and the evolution of regulatory
networks. *Nat Rev Genet* 9(5):397–405.

41. Quinlan AR, Hall IM (2010) BEDTools: a flexible suite of utilities for
comparing genomic features. *Bioinformatics* 26(6):841–842.

42. Dobin A, et al. (2013) STAR: ultrafast universal RNA-seq aligner.
Bioinformatics 29(1):15–21.

43. Li H (2013) Aligning sequence reads, clone sequences and assembly contigs
with BWA-MEM. *arXiv* 1303.3997v2.

44. McKenna A, et al. (2010) The Genome Analysis Toolkit: a MapReduce
framework for analyzing next-generation DNA sequencing data. *Genome Res*
20(9):1297–1303.

45. DePristo MA, et al. (2011) A framework for variation discovery and
genotyping using next-generation DNA sequencing data. *Nat Genet* 43(5):491–
498.

46. Van der Auwera GA, et al. (2013) From FastQ data to high confidence variant
calls: the Genome Analysis Toolkit best practices pipeline. *Curr Protoc*
Bioinformatics 11(1110):11.10.1–11.10.33.

47. Danecek P, et al. (2011) The variant call format and VCFtools. *Bioinformatics*
27(15):2156–2158.

48. Haudry A, et al. (2013) An atlas of over 90,000 conserved noncoding
sequences provides insight into crucifer regulatory regions. *Nat Genet*
45(8):891–898.

49. Slotte T, Hazzouri KM, Stern D, Andolfatto P, Wright SI (2012) Genetic
architecture and adaptive significance of the selfing syndrome in *Capsella*.
Evolution 66(5):1360–1374.

Figure Legends

Figure 1. The extent of ASE in leaves (A, B) and flower buds (C, D) of one

representative *C. grandiflora* F1. Panels A and C show the deviation from equal

expression for all assayed genes. Genes with strong evidence for ASE (posterior

probability of ASE ≥ 0.95) show stronger deviations from equal expression (B and D).

Figure 2. The impact of purifying selection differs between genes with and without

ASE in *C. grandiflora.* The estimated proportion of mutations in each $N_e s$ bin of the

distribution of negative fitness effects (DFE) is shown, with whiskers corresponding

to 95% confidence intervals. The panels show the DFE for nonsynonymous sites (0-

fold degenerate sites) (A), for introns (B), for promoter regions 500 bp upstream of

the transcription start site (C) and for 3'-UTRs (D). Significance levels for

comparisons of ASE and control genes, that were also amenable to analysis of ASE,

are indicated by asterisks (*: P-value ≤ 0.05, ** P-value ≤ 0.01). These results are

based on the population sample and the one-epoch model.

Figure 3. A lower proportion of adaptive nonsynonymous fixations at genes with ASE.

(A) Estimation of α using the asymptotic method of Messer and Petrov (2013), which

fits an exponential function to estimates of α based on polymorphisms at different

frequencies. Orange dots show values for control genes, and green dots show values

for genes with ASE. The grey shaded area indicates 95% confidence intervals. The

point estimate for genes with and without ASE is 0.06 vs 0.28, respectively. (B) The

estimated proportion of adaptive fixations relative to 4-fold synonymous substitutions

(ω_a) for genes with and without ASE. Whiskers correspond to 95% confidence

intervals, and significance levels for comparisons of ASE and control genes are

indicated by asterisks (*: P-value ≤ 0.05, ** P-value ≤ 0.01).

Tables

Table 1. Genes amenable to analysis of ASE in flower buds and leaves, and ASE

results.

Sample type	F1	Analyzed[1]	ASE genes[2]	ASE prop.[3]	FDR[4]
Flower buds	6.3	13521	3065	0.33	0.00134
	7.2	14390	3829	0.36	0.00240
	8.2	14232	3601	0.35	0.00198
Leaves	6.3	12390	3425	0.34	0.00182
	7.2	13074	3749	0.39	0.00242
	8.2	12796	3550	0.34	0.00195

[1]Genes with expression data for at least one replicate, and with a phased fragment

containing at least three transcribed SNPs after filtering.

[2]Genes with posterior probability of ASE $\geq$ 0.95.

[3]Estimated proportion of genes with ASE

[4]False Discovery Rate

Table 2. Population genetic summary statistics and divergence estimates for the

different site classes, separately for ASE and control genes.

Site class[1]	Gene set	Mean π	Mean θ_W	$\pi_{\text{siteclass}}/\pi_{\text{4-fold}}$[2]	d
4-fold	ASE	0.029	0.029	NA	0.16
	control	0.023	0.024	NA	0.15
0-fold	ASE	0.009	0.011	0.32	0.04
	control	0.005	0.007	0.23	0.03
3'UTR	ASE	0.018	0.021	0.62	0.13
	control	0.014	0.018	0.62	0.12
5'UTR	ASE	0.016	0.016	0.55	0.12
	control	0.012	0.012	0.54	0.12
500 bp upstream	ASE	0.017	0.020	0.6	0.16
	control	0.015	0.019	0.68	0.15
intron	ASE	0.020	0.022	0.69	0.15
	control	0.018	0.020	0.79	0.14

[1]Class of sites investigated, including 4-fold degenerate sites (4-fold), 0-fold

degenerate sites (0-fold), 5'UTRs, 3'UTRs, 500 bp upstream of the TSS (500 bp

upstream) and introns.

[2] Ratio of nucleotide diversity at focal site class to nucleotide diversity at 4-fold

synonymous sites.

Table 3. The best-fit logistic regression model (AIC=3086.9) predicting ASE from
genomic features. Regression coefficients and their standard error, z-statistics and
associated P-values, and odds ratios (OR) are shown. Coexpression module size and
gene length were included in the best-fit model, but did not have individually
significant effects.

Model parameter	Coeff. (SE)	z value	P-value	OR
Gene-body methylation	-0.67 (0.20)	-3.41	$<10^{-3}$	0.51
π_N/π_S	0.08 (0.04)	2.25	0.024	1.09
Expression level	0.20 (0.06)	3.31	<0.001	1.22
Promoter polymorphism	0.21 (0.05)	4.45	$<10^{-3}$	1.23
Tissue specificity	0.30 (0.06)	5.03	$<10^{-3}$	1.35
TE within 1 kb	0.32 (0.13)	2.50	0.013	1.38
Coexpression module size	-0.08 (0.05)	-1.59	NS	0.92
Gene length	0.08 (0.06)	1.49	NS	1.09
Intercept	-2.60 (0.06)	-42.91	$<10^{-3}$	0.07