In silico identification of metabolic enzyme drug targets in *Burkholderia pseudomallei*

Jean F. Challacombe [1]*

[1] Bioscience Division, Los Alamos National Laboratory, Los Alamos, NM 87545; jchalla@lanl.gov

* Correspondence: jchalla@lanl.gov; Tel.: +1-505-665-1485

Abstract: The intracellular pathogen *Burkholderia pseudomallei,* which is endemic to parts of southeast Asia and northern Australia, causes the disease melioidosis. Although acute infections can be treated with antibiotics, melioidosis is difficult to cure, and some patients develop chronic infections or a recrudescence of the disease months or years after treatment of the initial infection. *B. pseudomallei* strains have a high level of natural resistance to a variety of antibiotics, and with limited options for new antibiotics on the horizon, new alternatives are needed. The aim of the present study was to characterize the metabolic capabilities of *B. pseudomallei*, identify metabolites crucial for pathogen survival, understand the metabolic interactions that occur between pathogen and host cells, and determine if metabolic enzymes produced by the pathogen might be potential antibacterial targets. This aim was accomplished through genome scale metabolic modeling under different external conditions: 1) including all nutrients that could be consumed by the model, and 2) providing only the nutrients available in culture media. Using this approach, candidate chokepoint enzymes were identified, then knocked out *in silico* under the different nutrient conditions. The effect of each knockout on the metabolic network was examined. When five of the candidate chokepoints were knocked out *in silico*, the flux through the *B. pseudomallei* network was decreased, depending on the nutrient conditions. These results demonstrate the utility of genome-scale metabolic modeling methods for drug target identification in *B. pseudomallei*.

Keywords: Burkholderia, pathogen, metabolism, metabolic network, nutrients, infection

1. Introduction

The intracellular pathogen *Burkholderia pseudomallei,* which causes the disease melioidosis, is acquired from the environment in parts of southeast Asia and northern Australia [1,2]. Although acute infections can be treated with antibiotics, melioidosis is difficult to cure, requiring lengthy treatment in two phases for a duration of ~20 weeks [3,4]. Despite antibiotic therapy, some patients have persistent cases that develop into chronic infections, and others experience a recrudescence of the disease months or years after treatment of the initial infection with antibiotics [5]. *B. pseudomallei* strains have a high level of natural resistance to a variety of antibiotics [6-8], and with limited options for new antibiotics on the horizon, alternatives are desperately needed.

The availability of many *B. pseudomallei* genomes and advances in computational analysis methods make possible the rapid identification of novel antibacterial targets by selecting the most likely targets from complete sets of protein coding genes. Previous studies have demonstrated that essential genes present in pathogen genomes, but not in the host, make the best therapeutic targets [9]. Many known antibacterial compounds are enzyme inhibitors [10,11], so metabolic enzymes specific to pathogenic bacteria represent promising drug targets [9].

Enzyme targets in key metabolic pathways have been identified in *B. pseudomallei* and other bacterial pathogens; these pathways include fatty acid biosynthesis [12-14], the glyoxalate shunt [15,16], the chorismate pathway for biosynthesis of aromatic amino acids [17], purine, histidine, 4-aminobenzoate, and lipoate biosynthesis [18,19], leucine, threonine, p-aminobenzoic acid, aromatic

compound biosynthesis [20], branched chain amino acid biosynthesis [21], purine metabolism [22].
Other enzyme targets have been identified that are not in pathways - alanine racemase (interconverts
L- and D-alanine) [23], superoxide dismutase [24] and cyclic di-GMP phosphodiesterase [25].

Drugs acting on pathogen targets that are not present in the host should not cause significant
side effects. However, before the human genome was available, the process of identifying bacterial
pathogen-specific drug targets was labor intensive, involving comparison of candidate pathogen
targets against all known eukaryotic sequences to filter out targets likely to occur in the human [9].
Since then, various software tools have made the process of *in silico* target identification in pathogen
genomes easier. Available *in silico* tools encompass various cheminformatic [26] and bioinformatic
[27,28] approaches to identify new protein targets. Among the bioinformatic tools, metabolic
pathway/metabolic network analysis has emerged as an efficient *in silico* method to identify
candidate metabolic enzyme targets in pathogen genomes.

Several software packages are available to facilitate genome-scale metabolic network analyses
[29]. Starting with an annotated pathogen genome, the components of metabolic pathways are
identified, curated and refined [30]. The resulting genome-scale metabolic model can be used to
integrate omics datasets and to perform various analyses to determine the most likely drug targets
[10]. Perhaps the most important task with respect to finding good candidate targets is metabolic
chokepoint identification. By definition, a chokepoint enzyme either consumes a unique substrate or
produces a unique product in the pathogen metabolic network [31]. Inhibition of chokepoint
enzymes may disrupt crucial metabolic processes in the pathogen, so chokepoints that are essential
to the pathogen represent good potential drug targets [32-34].

The aim of the present study was to characterize the metabolic capabilities of *B. pseudomallei*,
identify metabolites and aspects of the metabolic network crucial for pathogen survival, understand
the metabolic interactions that occur between pathogen and host cells, and determine if any of the
metabolic enzymes produced by the pathogen might be potential antibacterial targets. This aim was
accomplished through genome scale metabolic modeling of *B. pseudomallei* under different external
conditions, including all nutrients that could be consumed by the model and only the nutrients
available in culture media. Using this approach, candidate chokepoints were identified, then knocked
out the genes encoding chokepoint enzymes *in silico* under the different nutrient conditions, and
examined the effect of each knockout on the metabolic network. The result of this analysis was five
candidate antibacterial targets, demonstrating the utility of genome-scale metabolic modeling
methods for *in silico* studies of pathogen metabolism and for drug target identification in *B.*
pseudomallei.

2. Materials and Methods

2.1 Metabolic pathway reconstructions and annotation curation
Pathway genome databases (PGDBs) for *B. pseudomallei* strains MSHR668 and K96243 were
obtained through the Pathway Tools software (version 18.5) from the PGDB registry [35]. We found
that the original annotation of *B. pseudomallei* K96243 identified many fewer coding sequences than
that of MSHR668, so we re-annotated the original complete genome sequences of both strains using
the RAST system [36] to better compare them [37]. The RAST-annotated genome sequences were
loaded into Pathway Tools, using the PathoLogic component to predict the metabolic pathways [38].
For each genome, the set of protein coding sequences from the original annotation was compared to
those from the RAST annotation, using blast to identify coding sequences in common between the
two annotations, and to identify coding sequences that were missing from each annotation. For the
MSHR668 genome, RAST annotation identified 247 fewer coding sequences than the original
annotation. However, there were some predicted coding sequences with annotated functions in the
RAST annotation that were not present in the original, so these were added to the original PGDB for
MSHR668. The RAST annotation of the K96243 genome identified 1,317 more protein coding
sequences than the original annotation, so the PGDB created from the RAST annotation was used as
the starting point. Coding sequences from the original annotation that were not present in the RAST
annotation of K96243 were added to the PGDB that was created from the RAST annotated genome.

97
98   *2.2 Chokepoint Identification*
99   Chokepoints were identified in each PGDB using the chokepoint reaction finder in Pathway
100   Tools. All reactions were included except those found in human.
101
2.3 Flux balance analysis
For each metabolic network reconstruction, flux balance analysis (FBA) was performed using
the MetaFlux module within the Pathway Tools [39]. Development FBA models were constructed
iteratively to determine the compounds that each model could use and produce. This was
accomplished by trying all compounds in the PGDB as biomass metabolites, nutrients and secretions
in the various compartments (cytosol, periplasmic space and extracellular). Each model was refined
iteratively, first identifying the specific biomass components that could be produced, then trying all
compounds as nutrients and secretions, then specifying the biomass metabolites and nutrients and
trying all compounds as secretions.
Once the nutrients, secretions and biomass components that could be consumed or produced by
the metabolic networks were determined, the log file produced by MetaFlux was examined and
problematic reactions were fixed, if possible. Most of the problematic reactions were unbalanced due
to missing chemical formulas of one or more metabolites. A few of these reactions were corrected by
copying the missing structures from other PGDBs. For example, the structures of D-ribose, D-
glucuronate, D-glucose and some other compounds were copied from the more highly curated
Escherichia coli K12 substr. MG1655 PGDB. Many of the reactions with compounds that were lacking
chemical formulas were generic so no suitable chemical structure could be found or created.
Reactions involving starch, glycogen, and glucans with variable lengths and non-numeric
stoichiometries, could not be balanced. Other reactions were missing H^+ or H_2O on one side or the
other, and the addition of the missing compound balanced the reaction. However, there were a small
number of reactions that could not be fixed and these were marked as unbalanced. Once all reactions
that could be fixed were corrected in the PGDB, MetaFlux was run again in development mode to
identify additional biomass metabolites, nutrients, and secretions. Once the set of biomass
metabolites was constant, the nutrients that could be consumed by the model were determined,
followed by identification of the secretions produced by the model. The result of this process was a
final unconstrained FBA model.
To mimic the nutrient conditions in culture, only the estimated set of ingredients present in LB
medium were included as nutrients in the FBA model. LB medium includes as its main ingredients
tryptone [40] and yeast extract [41]. Tryptone provides peptides and peptones, which are good
sources   of   amino   acids   (http://www.sigmaaldrich.com/analytical-
chromatography/microbiology/basic-ingredients/protein-sources.html); yeast extract provides
vitamins, nitrogenous compounds, carbon, sulfur, trace elements and minerals [42]. The LB media
composition reported in a *Bacillus subtilis* modeling study [43] was used as a starting point for this
study. Starting with the nutrient set that mimicked LB media [43], all compounds were included in
the try-biomass set for the cytosol and periplasmic cellular compartments to see which biomass
metabolites could be produced given only the nutrients present in LB media. Once a stable set of
nutrients was determined, iterations were performed to determine the stable set of biomass
metabolites that could be produced. Then all compounds were tried as secretions in the cytosol and
periplasmic compartments to determine the compounds that could be secreted by the model.
A list of nutrients that might be available to *B. pseudomallei* while residing inside host cells during
infection was compiled by searching the literature for infection studies involving *B. pseudomallei* and
host cells. Gene expression studies of other intracellular pathogens and host cells during infection
were also considered. A development FBA model mimicking infection conditions was constructed as
described above for the LB media model.
146
2.4 Essential gene and candidate drug target identification

To reduce the set of potential drug targets, candidate chokepoint enzymes were compared
against the list of essential genes and all of the drug targets in DrugBank. Essential genes in the
MSHR668 genome were determined by blasting all protein coding sequences (amino acid format)
against the Database of Essential Genes [44] using the blastp program with an E-value cut-off of 1e -
10 and 70% identity as thresholds. Candidate drug target sequences were determined by blasting all
protein coding sequences in the MSHR668 genome (as both nucleotide and amino acid formats)
against the complete set of DrugBank targets [45], using an E-value cutoff of 0.005 and a threshold
identity of 70% to select likely targets.
156
157 *2.5 In silico knockout experiments*
158   Knockout experiments were performed *in silico* through the Pathway Tools MetaFlux module
159 [39]; the chokepoint genes were knocked out one at a time and the effects on total biomass flux
160 through the metabolic network were noted.
161
2.6 Network visualization
For each PGDB, an .sbml file was exported from the Pathway Tools and loaded into Cytoscape [46]
version 3.1.1 for visualization and comparison of network features.

3. Results

3.1. General features of B. pseudomallei genomes and metabolic networks

The general characteristics of each PGDB and metabolic network for B. pseudomallei MSHR668 and
K96243 are listed in Table 1.

Table 1. Features of *B. pseudomallei* PGDBs and metabolic networks

Genome (Pathway Tools PGDB)	Curated MSHR668 (original + RAST)	Curated K96243 (RAST + original)
Coding sequences	7133	7045
Pathways	339	387
Enzymatic reactions	1870	1995
Transport reactions	144	82
Enzymes	1666	1685
Transporters	292	38
Compounds	1397	1578
Metabolic Network		
Nodes (metabolites)	4295	4219
Edges (reaction steps)	10860	9747
Chokepoints		
Producing	444	506
Consuming	419	498
Candidate	479	348
Dead-end reactants	166	217
Dead-end products	72	118

The *B. pseudomallei* MSHR668 and K96243 genomes contained similar numbers of coding
sequences, pathways, enzymatic reactions and enzymes. Differences between the PGDBs were noted
in the numbers of transporters, transport reactions and compounds. The K96243 PGDB contained
fewer transporters and transport reactions and more compounds than MSHR668. In terms of the
metabolic network characteristics, both networks contained similar numbers of nodes (representing
metabolites), while the MSHR668 network had more edges (reaction steps) than K96243. This is likely

due to the more extensive curation of the MSHR668 network that was performed during refinement
of the metabolic network models (see Methods).

In a metabolic network, chokepoints are reactions that either uniquely produce or uniquely
consume a metabolite. Inhibiting an enzyme that consumes a unique substrate may cause that
metabolite to accumulate, and it may be toxic to the cell; inhibiting an enzyme that produces a unique
product may starve the cell of an essential metabolite [31]. Identifying chokepoint enzymes in
pathogens is a promising *in silico* approach to recognize potential metabolic drug targets. For
example, analysis of *Plasmodium falciparum* metabolism revealed that 87.5% of proposed drug targets
supported by evidence are chokepoint reactions [31]. However, to be a valid chokepoint, the
metabolite in question must be balanced by a producing or consuming reaction and not be a dead-
end metabolite [31]. Table 1 compares the numbers of chokepoint reactions and dead-end metabolites
that were identified in each PGDB. Overall the numbers were similar between the two PGDBs, and
the lower numbers of dead-ends in the MSHR668 database were likely due to the more extensive
curation performed on the MSHR668 PGDB (see Materials and Methods).

3.2. Flux balance analysis (FBA)

3.2.1. Unconstrained FBA model

Given a set of nutrients for consumption, along with secretions and metabolites that can be
produced, a FBA model predicts the steady-state flux rates of the metabolic reactions in an organism,
and provides an estimate of the overall biomass flux. FBA was conducted on each metabolic network
as described in the Materials and Methods section. It took several cycles of refinement to solve an
initial unconstrained MSHR668 model. Even after several iterations, the K96243 unconstrained model
did not reach a stable solution, likely because there were missing transporter-encoding genes (and
possibly other genes) in the PGDB, indicating that the annotation needed more curation. Since the
unconstrained MSHR668 model reached a stable solution after performing the initial network
curation steps suggested by MetaFlux, only this model was analyzed further. The initial
unconstrained FBA model of the MSHR668 metabolic network (Table 2) included all possible biomass
compounds that could be produced, all nutrients that could be consumed, and had no weights
imposed on the biomass metabolites and no constraints imposed on the nutrients. [Supplementary
materials,   S1_Final_unconstrained_model_inputs.pdf,
S2_Final_unconstrained_model_solution.pdf].

3.2.2. LB media FBA model

To mimic the conditions that *B. pseudomallei* experiences in culture, only the nutrients present in
LB media [43] plus glycerol were included as inputs to the LB media FBA model (Table 2). In this
model, constraints were included on some of the nutrients (ADP, Pi, proton and glycerol).
[Supplementary materials, S3_Final_LB_model_inputs.pdf and S4_Final_LB_model_solution.pdf].

Table 2. Characteristics of the MSHR668 unconstrained and LB media models

Model	Total rxns	Rxns carrying flux	Biomass metabolites produced	Nutrients consumed	Secre-tions	Biomass flux
Unconstrained	3193	1619	1403	667	10	15000.00
LB media	3213	1025	282	47	0	0.079412

3.2.3. Host cell infection model

In addition to the unconstrained and LB media models, the original plan for this study included the development of a model of *B. pseudomallei* metabolism that mimics infection conditions. However, there was very little information available on the growth requirements of *B. pseudomallei* inside human macrophages. In addition, no comprehensive studies have been performed to identify the complete list of host cell nutrients that are available to *B. pseudomallei* during infection. Most studies of the nutritional requirements of intracellular pathogens growing inside host cells have been performed on *Legionella pneumophila* [47,48], which can grow and replicate similarly in human macrophages and amoebae [49]. Growing in both human macrophages and amoebae, *L. pneumophila* utilizes amino acids as its main sources of carbon, nitrogen and energy; *L. pneumophila* obtains amino acids from the host through proteasomal degradation [48]. However, glucose is also used to feed central metabolism under both culture and infection conditions [50].

Comparing the nutrients provided by the LB media model to the nutrients used by *L. pneumophila* in culture and during infection of amoebae [47], the only difference was the carbohydrate carbon source: glycerol (*B. pseudomallei* LB media [51]) vs. glucose (*L. pneumophila* AYE media [50] and in amoebae [47,50]). When glycerol was replaced by glucose in the LB media model of *B. pseudomallei*, no FBA solution was found [data not shown]. Several possible explanations for this result are presented in the Discussion.

While the specific carbon requirements of *B. pseudomallei* in either human macrophages or amoebae have not been determined, one study produced whole-genome tiling array expression data to assess *B. pseudomallei* transcriptional responses under 82 different conditions, including infection [52]. From their supplemental table S2, a list of metabolic genes expressed in the infection conditions was used to infer the potential nutrients consumed by *B. pseudomallei* during infection. Additional candidate host cell nutrients were identified from the literature, focusing on studies of intracellular pathogen-mammalian host infections. The nutrients identified as potential carbon sources for intracellular survival of various pathogens included aromatic compounds, such as benzoate and phenylacetic acid and related derivatives [53], sugar acids like gluconate, galactonate, glucuronate, and galacturonate [54], ribo- and deoxyribonucleosides, hexuronates [55], glutathione [56], glucose 6-phosphate [57,58], glycerol-3-phosphate [59]. The complete list of potential host cell nutrients is in the [Supplementary material, S5_Nutrients_infection_model.pdf] file.

FBA was performed for a *B. pseudomallei* MSHR668 model that included the list of candidate host cell nutrients identified as above. However, some of the nutrients were not present in any reactions in the MSHR668 PGDB. When the rest of the compounds were included as nutrients in addition to glycerol, none of them were consumed by the model, but glycerol was consumed and biomass was produced. When glycerol was excluded from the nutrient list, none of the other nutrients were consumed and the model was not solvable [data not shown].

3.3. Metabolic chokepoints and candidate drug targets

To narrow down the list of metabolic chokepoints, which represented candidate drug targets, essential genes and genes with sequence homology to existing DrugBank targets were identified in the MSHR668 genome. This analysis identified 34 chokepoint genes that were also essential genes and DrugBank targets (Figure 1, Table 3).

Figure 1. Process for identifying candidate metabolic enzyme drug targets (chokepoints) in the *B. pseudomallei* MSHR668 genome.

Table 3. Chokepoint genes that encode candidate metabolic enzyme drug targets in *B. pseudomallei* MSHR668

Locus_tag	Gene symbol	Enzyme name	E.C. number	DrugBank target (drug IDs)	Human target?	Choke-point in K96243?	Bp Mutant(s) exist?
BURPS668 _0305	argG+	argininosucc inate synthase	6.3.4.5	P0A6E4 Argininosuccina te synthase (DB00536; DB04077)	no	yes	no*
BURPS668 _0328	folB+	dihydroneo pterin aldolase	4.1.2.25	P56740 Dihydroneopteri n aldolase (DB01778; DB01906; DB02119; DB02489; DB03231; DB03571; DB04168; DB04400; DB04425; DB06906)	no	yes	no
BURPS668 _0567	pth+	peptidyl-tRNA hydrolase	3.1.1.29	No	yes	yes	no
BURPS668 _0675	aspS+	aspartyl-tRNA synthase	6.1.1.12	P36419 Aspartate— tRNA ligase (DB01895)	no	no	no

BURPS668_0810	RecA+	RecA protein	5.99.1.-	P62219 RecA (DB01660; DB03222; DB04444; DB04395)	no	yes	no*
BURPS668_0964	dut+	deoxyuridine5'-triphosphate nucleotidohydrolase	3.6.1.19/ 3.6.1.23	P06968 Deoxyuridine 5'-triphosphate nucleotidohydrolase (DB01965; DB02333; DB03413; DB03800)	no	yes	no
BURPS668_0967	ileS+	isoleucyl-tRNA synthetase	6.1.1.5	Q9NSE4 Isoleucine--tRNA ligase (DB00167)	yes	no	no
BURPS668_1446	pckG+	phosphoenolpyruvate carboxykinase	4.1.1.32	P35558 Phosphoenolpyruvate carboxykinase, cytosolic [GTP] (DB01819; DB02008; DB03267; DB03725)	yes	yes	no
BURPS668_1465	dnaQ+	DNA polymerase III subunit epsilon	2.7.7.7	P03007 DNA polymerase III subunit epsilon (DB01643)	no	yes	no
BURPS668_1544	NA+	alpha-glucosidase	3.2.1.20	O33830 Alpha-glucosidase (DG01769; DB03323)	yes lysosomal	no	no
BURPS668_1669	valS+	valyl-tRNA synthetase	6.1.1.9	P26640 Valine--tRNA ligase (DB00161)	yes	no	no
BURPS668_1712	thrS+	threonyl-tRNA synthetase	6.1.1.3	P0A8M5 Threonine--tRNA ligase (DB03355; DB03869; DB04024)	no	no	no
BURPS668_1750	sucA+	2-oxoglutarate dehydrogenase E1	1.2.4.2	Q02218 2-oxoglutarate dehydrogenase, mitochondrial (DB00157; DB00313)	yes	yes	no
BURPS668_1752	lpdA+	dihydrolipoamide	1.8.1.4	P14218 Dihydrolipoyl	no	yes	no

		dehydrogen ase		dehydrogenase (DB03147)			
BURPS668 _2178	purA+	adenylosucc inate synthetase	6.3.4.4	Q83P33 Adenylosuccinat e synthetase (DB02954; DB04160; DB04566)	no	yes	no
BURPS668 _2189	hisS+	histidyl- tRNA synthetase	6.1.1.21	P60906 Histidine--tRNA ligase (DB03811; DB04201)	no	no	no
BURPS668 _2308	pgi+	glucose-6- phosphate isomerase	5.3.1.9	P06744 Glucose- 6-phosphate isomerase (DB02007; DB02076; DB02093; DB02548; DB03042; DB03581; DB03937; DB04493)	yes	no	no
BURPS668 _2426	lpxD+	UDP-3-O-[3- hydroxymyr istoyl] glucosamine N- acyltransfera se	2.3.1.191	O67648 UDP-3- O-[3- hydroxymyristo yl] N- acetylglucosami ne deacetylase (DB01991; DB04257; DB04399; DB07355; DB07536; DB08231; DB07861)	no	yes	no
BURPS668 _2433	uppS+	undecapren yl diphosphate synthase	2.5.1.31	P60472 Ditrans,polycis- undecaprenyl- diphosphate synthase ((2E,6E)- farnesyl- diphosphate specific) (DB04695; DB04714; DB07404; DB07409; DB07410; DB07426; DB07780)	no	yes	no

BURPS668_2610	lpdA+	pyruvate dehydrogenase complex E3 component, dihydrolipoamide dehydrogenase	1.8.1.4	P10802 Dihydrolipoyllysine-residue acetyltransferase component of pyruvate dehydrogenase complex (DB01846; DB01992; DB08120)	no	yes	no
BURPS668_2788	fabF+	3-oxoacyl-ACP synthase	2.3.1.41	Q02054 Actinorhodin polyketide synthase acyl carrier protein (DB08585; DB08586)	no	no	no
BURPS668_3103	rfbA+	glucose-1-phosphate thymidylyltransferase	2.7.7.24	Q9HU22 Glucose-1-phosphate thymidylyltransferase (DB01643; DB02452; DB02843; DB03723; DB03751; DB04272; DB04485; DB02452; DB04355)	no	yes	no
BURPS668_3328	tgt+	queuine tRNA-ribosyltransferase	2.4.2.29	P28720 Queuine tRNA-ribosyltransferase (DB01825; DB02041; DB02441; DB02599; DB03074; DB03304; DB03505; DB03780; DB04004; DB04169; DB04239; DB04543; DB07012; DB07452; DB07481; DB07564; DB07704; DB08267;	no	yes	no

				DB08268; DB08511; DB08512; DB08514)			
BURPS668 _3366	ruvB+	Holliday junction DNA helicase RuvB	3.1.22.4	Q5SL87 Holliday junction ATP-dependent DNA helicase RuvB (DB00173)	no	no	no
BURPS668 _3464	aroQ+	3-dehydroquinate dehydratase	4.2.1.10	P15474 3-dehydroquinate dehydratase (DB02786; DB02801; DB04347; DB04656; DB08485)	no	yes	no
BURPS668 _3525	murG+	undecaprenyldiphospho-muramoylpentapeptide beta-N-acetylglucosaminyltransferase	2.4.1.227	P17443 UDP-N-acetylglucosamine--N-acetylmuramyl-(pentapeptide)pyrophosphoryl-undecaprenol N-acetylglucosamine transferase (DB02196)	no	yes	no
BURPS668 _3561	ung+	uracil-DNA glycosylase	3.2.2.27	Q8X444 Uracil-DNA glycosylase (DB03419)	no	yes	no
BURPS668 _3668	murA+	UDP-N-acetylglucosamine 1-carboxyvinyltransferase		P33038 UDP-N-acetylglucosamine 1-carboxyvinyltransferase (DB01879; DB02435; DB02995; DB03089; DB04174; DB04474)	no	yes	no
BURPS668 _A0384	acnD+	aconitate hydratase	4.2.1.3	Q99798 Aconitate hydratase, mitochondrial (DB01727; DB03964; DB04072; DB04351;	yes	yes	no

				DB04562); P36683 Aconitate hydratase 2 (DB04351)			
BURPS668 _A1869	ileS+	isoleucyl- tRNA synthetase	6.1.1.5	P41972 isoleucyl-tRNA synthetase (DB00410)	no	no	no
BURPS668 _A2053	NA+	putative acetyl-CoA carboxylase, biotin carboxylase	6.3.4.14	P24182 Biotin carboxylase (DB08074; DB08075; DB08076; DB08144; DB08145; DB08146; DB08314; DB08315; DB08316; DB08317; DB08318)	no	yes	no
BURPS668 _A2451	leuB+	3- isopropylma late dehydrogen ase	1.1.1.85	Q56268 3- isopropylmalate dehydrogenase (DB04279)	no	yes	no
BURPS668 _A2546	polA+	DNA polymerase I	2.7.7.7	P00582 DNA polymerase I (DB00548; DB08432)	no	yes	no
BURPS668 _A3190	lpdA+	dihydrolipo amide dehydrogen ase	1.8.1.4	P09063 Dihydrolipoyl dehydrogenase (DB03147)	no	yes	no

* mutants exist in other *Burkholderia* species, +essential gene in *B. pseudomallei* K96243 and MSHR668

Eight of the targets in Table 3 were also DrugBank targets in human; DrugBank was searched for the
remaining twenty-six targets, showing that they also occur in other bacteria.

In silico knockout experiments were performed with the MetaFlux module of Pathway Tools to
test the effect of inhibiting each chokepoint enzyme on *B. pseudomallei* growth in both the
unconstrained and LB media models. Results (Table 4) show that knockout of BURPS668_3328 (*tgt*)
and BURPS668_A2451 (*leuB*), eliminated the biomass flux in the unconstrained model, while
knockout of BURPS668_2426 (*lpxD*), BURPS668_2433 (*uppS*), and BURPS668_3525 (*murG*) eliminated
the biomass flux in the LB media model. Knockout of BURPS668_2433 (*uppS*), and BURPS668_3525
(*murG*) decreased the biomass flux in the unconstrained model, but did not eliminate it.

Table 4. Results of *in silico* knockout experiments on twenty-six chokepoint reactions in *B.*
pseudomallei

locus_tag	Gene name	Reaction(s)	Biomass flux (unconstrained model)	Biomass flux (LB media model)
BURPS668_0305	argG	ARGSUCCINSYN-RXN	15000.000000	0.079412
BURPS668_0328	folB	H2NEOPTERINALDOL-RXN	15000.000000	0.079412
BURPS668_0675	aspS	ASPARTATE--TRNA-LIGASE-RXN-ASP-tRNAs/L-ASPARTATE/ATP/PROTON//Charged-ASP-tRNAs/AMP/PPI.60	15000.000000	0.079412
BURPS668_0810	recA	RXN0-5100	15000.000000	0.079412
BURPS668_0964	dut	DUTP-PYROP-RXN	15000.000000	0.079412
BURPS668_1465	dnaQ	DNA-DIRECTED-DNA-POLYMERASE-RXN	15000.000000	0.079412
BURPS668_1712	thrS	THREONINE--TRNA-LIGASE-RXN-THR-tRNAs/THR/ATP/PROTON//Charged-THR-tRNAs/AMP/PPI.52	15000.000000	0.079412
BURPS668_1752	lpdA	RXN-9718	15000.000000	0.079412
BURPS668_2178	purA	ADENYLOSUCCINATE-SYNTHASE-RXN	15000.000000	0.079412
BURPS668_2189	hisS	HISTIDINE--TRNA-LIGASE-RXN-HIS-tRNAs/HIS/ATP/PROTON//Charged-HIS-tRNAs/AMP/PPI.52	15000.000000	0.079412
BURPS668_2426	lpxD	UDPHYDROXYMYRGLUCOSAMNACETYLTRANS-RXN-R-3-hydroxymyristoyl-ACPs/UDP-OHMYR-GLUCOSAMINE//OH-MYRISTOYL/ACP/PROTON.73	15000.000000	0.000000
BURPS668_2433	uppS	RXN-8999	14545.454545	0.000000
BURPS668_2610	lpdA	1.8.1.4-RXN	15000.000000	0.079412
BURPS668_2788	fabF	3-OXOACYL-ACP-SYNTH-RXN /3-OXOACYL-ACP-SYNTH-BASE-RXN /2.3.1.41-RXN / many other reactions	15000.000000	0.079412
BURPS668_3103	rfbA	DTDPGLUCOSEPP-RXN	15000.000000	0.079412
BURPS668_3328	tgt	QUEUOSINE-TRNA-RIBOSYLTRANSFERASE-RXN; RXN0-1321	0.000000	0.079412
BURPS668_3366	ruvB	3.1.22.4-RXN	15000.000000	0.079412
BURPS668_3464	aroQ	3-DEHYDROQUINATE-DEHYDRATASE-RXN	15000.000000	0.079412
BURPS668_3525	murG	RXN-11346, RXN-8976, NACGLCTRANS-RXN, RXN-11029	10000.000000	0.000000
BURPS668_3561	ung	RXN0-2584	15000.000000	0.079412

BURPS668_3668	murA	UDPNACETYLGLUCOSAMENOL PYRTRANS-RXN	15000.000000	0.079412
BURPS668_A1869	ileS	ISOLEUCINE--TRNA-LIGASE-RXN	15000.000000	0.079412
BURPS668_A2053	acc	BIOTIN-CARBOXYL-RXN	15000.000000	0.079412
BURPS668_A2451	leuB	RXN-13158; 3-ISOPROPYLMALDEHYDROG-RXN	0.000000	0.079412
BURPS668_A2546	polA	DNA-DIRECTED-DNA-POLYMERASE-RXN	15000.000000	0.079412
BURPS668_A3190	lpdA	1.8.1.4-RXN/RXN0-1132/RXN-8629	15000.000000	0.079412

282 The overall biomass fluxes were different in the two models. The unconstrained model had a much
283 greater total biomass flux than the LB media model. This is likely because the unconstrained model
284 included more nutrient inputs than the LB media model.

4. Discussion

4.1 Links between metabolism and virulence

In order to colonize the host, establish an infection, and proliferate, pathogens employ various
strategies, often involving links between metabolic pathways and virulence genes. Although current
knowledge regarding the connections between metabolism and virulence is limited [60], this topic is
becoming an increasing focus for host-pathogen studies. Some general links between metabolism and
virulence include regulatory connections between specific metabolites and virulence gene expression
[61-65], metabolic requirements for adaptation of the pathogen to the host niche [59,60,66,67], and
carbon catabolite repression [68]. More detailed information on the topic of metabolism and virulence
can be found in reviews [59,60,66,67,69-74].
Inside host cells, the survival of pathogenic bacteria depends on their acquisition of nutrients
and carbon sources, such as carbohydrates, lipids, glycolipids, dicarboxylic acids and amino acids,
from their host environment [59,60,66,67,71,75]. Preferred carbon sources vary among intracellular
pathogens, and the types of nutrients available in the host cell cytosol may determine the cell-type
specificities of different intracellular pathogens [76]. For example, many bacteria prefer hexoses, like
glucose, as sources of carbon and energy. These sugars are catabolized through glycolysis, the
pentose phosphate and Entner-Doudoroff pathways [75]. Some bacteria lack the glycolysis pathway
and preferentially metabolize glucose via the Entner-Doudoroff pathway [77], while others lack both
glycolysis and Entner-Doudoroff pathways and live on pyruvate that they obtain from the host cell
cytosol [78].

4.2 Metabolic potential of B. pseudomallei MSHR668

We previously reported that *B. pseudomallei* MSHR668, K96243 and 1106a have abundant
capabilities to metabolize hexoses, including the complete sets of genes encoding the glycolysis,
pentose phosphate cycle, and Entner-Doudoroff pathways. They also have several pathways for
metabolism of pyruvate to acetyl-CoA, acetate and ethanol [37]. In general, *B. pseudomallei* as a species
has a very diverse set of metabolic capabilities, likely a reflection of its ability to live both in the
natural environment and in hosts.
The metabolic power of *B. pseudomallei* has been targeted in mutation studies to address the roles
of various metabolic genes in virulence. Mutation of various metabolic genes that affect cell growth
results in attenuation of *B. pseudomallei* virulence; these genes include

phosphoribosylformylglycinamidine cyclo-ligase (*purM*) [79], aspartate-*β*-semialdehyde dehydrogenase (*asd*) [80], acetolactate synthase (*ilvI*) [21], dehydroquinate synthase (*aroB*) [20], chorismate synthase (*aroC*) [17], phosphoserine aminotransferase (*serC*) [81], phosphoribosylglycinamide formyltransferase 1 (*purN*) and phosphoribosylformylglycinamide cyclo-ligase (*purM*) [18], two phospholipase C enzymes [82], disulfide oxidoreductase (*dsbA*) [83]. Targeted mutation of *purM*, which encodes aminoimidazole ribotide, a precursor of de novo adenine and thiamine biosynthesis, predictably causes a deficiency in adenine and thiamine biosynthesis [79]. Aspartate-*β*-semialdehyde dehydrogenase (*asd*) mutants cannot synthesize diaminopimelate for cell wall biosynthesis [80], while acetolactate synthase (*ilvI*) mutants cannot synthesize the branched chain amino acids isoleucine, valine, and leucine [21]. Dehydroquinate synthase (*aroB*) mutants are defective in the shikimate pathway for chorismate biosynthesis, and addition of the aromatic compounds tyrosine, tryptophan, phenylalanine, PABA, and 2,3-dihydroxybenzoate is required to restore the growth in minimal medium [20]. Chorismate synthase (*aroC*) mutants are also defective in aromatic compound synthesis and cannot grow without the addition of aromatic compounds to the media [17]. Phosphoserine aminotransferase (*serC*) mutants are defective in serine and pyridoxal 5-phosphate biosynthesis and require minimal medium supplemented with serine for normal growth [81].

4.3 Vaccine studies

Using live attenuated bacteria as vaccines can be effective in preventing disease, as attenuated bacteria may still be able to replicate in the host and may contain immune-stimulatory epitopes that are not found in subunit or heat-inactivated vaccines [84]. Many of the *B. pseudomallei* mutants described above have already been tried as attenuated vaccines with mixed results. Vaccination with the attenuated *asd* mutant protected BALB/c mice against acute melioidosis, but did not protect against chronic melioidosis [80]. Vaccination with the attenuated *ilvI* mutant of *B. pseudomallei* protected BALB/c mice against a challenge with a virulent strain [21]. In mice vaccinated with the *aroB* mutant, the time to death following challenge with the virulent K96243 strain was a bit longer than in unvaccinated mice, but all mice eventually died [20]. The *aroC* mutant was unable to persist in vaccinated BALB/c mice long enough to elicit protective immunity, however C57BL/6 mice were protected against challenge with a virulent strain [17]. Intraperitoneal vaccination of BALB/c mice with a *serC* mutant resulted in higher levels of survival after challenge with K96243 virulent strain [81]. While immunization of mice with attenuated *B. pseudomallei* mutants has resulted in the induction of protective immunity in some cases, sterile immunity was rarely reported (reviewed by [85]). Also, the live attenuated vaccine model may not be the best solution for the prevention of melioidosis, because an attenuated mutant might revert to virulence, and might also establish a latent infection [85].

4.4 Identification of antimicrobial therapeutics

An alternative avenue to combat melioidosis is through the development of novel antimicrobial therapeutics. Regardless of the specific metabolic capabilities possessed by a pathogen, essential nutrient acquisition and utilization mechanisms are proving to be good potential therapeutic targets, as inhibition of these targets might deprive the pathogen of needed substrates for growth and replication inside host cells [31]. There are currently 699 *B. pseudomallei* genomes available at the National Center for Biotechnology Information (NCBI, http://www.ncbi.nlm.nih.gov/genome/476). While some of these entries represent re-annotations of previous submissions, and some genomes represent colony morphology variants of the same strain, an impressive number of individual genomes are available to use with computational approaches to identify new therapeutic targets. With this large number of available genomes, a core genome approach [86,87] could be used to identify potential metabolic enzyme targets that are present in all sequenced *B. pseudomallei* genomes.

In silico methods for the identification of therapeutic targets in bacterial pathogens include comparative genomics-based approaches, such as identification of essential genes specific to the

pathogen, and techniques based on metabolic pathway analysis and metabolic network modeling.
The more robust approaches use a combination of comparative genomics and metabolic pathway
analysis. These approaches have been used to identify essential gene targets in *Mycoplasma genitalium*
[88] and *Mycobacterium ulcerans* [89]. Another method, subtractive target identification, involves
identification of enzymes in the metabolic pathways of the pathogen, and comparing them to human
proteins to identify pathogen enzymes that are not found in human. A list of likely targets is compiled
by focusing on enzymes in pathways that are usually essential for pathogen growth and survival,
like lipid metabolism, carbohydrate metabolism, amino acid metabolism, energy metabolism,
vitamin and cofactor biosynthetic pathways and nucleotide metabolism. This approach has been used
to identify putative targets in *M. tuberculosis* [90], MRSA [91-93], as well as a collection of other
bacterial pathogens [94].

Methods for therapeutic target discovery based on metabolic pathway analysis and metabolic
network modeling have become very popular in the last ten years. Numerous studies have identified
candidate drug targets in various bacterial [27,34,95-106], fungal [106] and protosoan [31,106-109]
pathogens using a variety of methods to analyze metabolic pathways and networks. One method
employs chokepoint analysis to identify metabolic enzymes that are critical to the pathogen, because
they uniquely consume and/or produce certain metabolites. Chokepoint analysis has been used to
identify candidate metabolic enzyme targets in various pathogen genomes
[27,31,34,99,107,108,110,111]. However, no studies to date have used this approach to identify
potential drug targets in *B. pseudomallei* metabolic networks.

4.4.1 Identification of metabolic chokepoints

For this study, metabolic chokepoints were identified in the curated *B. pseudomallei* MSHR668
metabolic network using the Pathway Tools software [35]. Table 3 lists the chokepoint enzymes
identified in the metabolic networks of *B. pseudomallei* MSHR668 and K96243. Twenty-four of the *B.*
pseudomallei chokepoints were not indicated as human targets in DrugBank, and therefore
represented good candidate therapeutic targets against melioidosis. Six of the chokepoints in Table 3
were aminoacyl-tRNA synthetases, which are likely good targets as they are critical enzymes
involved in protein translation. These chokepoints included aspartyl-, threonyl-, histidyl-, valyl- and
isoleucyl-tRNA synthetase (2 copies). Aspartyl-tRNA synthase (*aspS*) is an essential gene target in *M.*
tuberculosis [112]. Threonyl-tRNA synthetase (*thrS*) inhibitors have been identified [113] and shown
to have anti-malarial activity against *Plasmodium falciparum* [114]. Isoleucyl-tRNA synthetase (*ileS*) is
a well-documented bacterial target [115-117]. The antimicrobial drug, mupirocin (pseudomonic acid),
selectively inhibits bacterial isoleucyl-tRNA synthetase without inhibiting its human homolog [114].
However, resistance is seen in bacteria that possess an isoleucyl-tRNA synthetase that is similar to
eukaryotic versions [118]. Histidyl-tRNA synthetase (*hisS*) has been explored as a target in
Trypanosoma cruzi [119]. The chokepoint enzyme queuine tRNA-ribosyltransferase (*tgt*) incorporates
the wobble base queuine into tRNA, and is also a target in *Zymomonas mobilis* and *Shigella* [120,121]

Several *B. pseudomallei* chokepoint enzymes are likely involved in DNA-related processes. Two
of these enzymes, encoded by the *recA* and *dnaQ* genes, are involved in the SOS pathway [122], which
mediates the bacterial response to DNA damage. Activation of the SOS response by ciprofloxacin
induces mutations, which can lead to fluoriquinolone resistance [123]. The RecA protein is a target
for antibacterial drug discovery in *M. tuberculosis* [124] and *Mycoplasma hyopneumoniae* [96], and has
been proposed as a specific target for reducing the evolution of antimicrobial resistance [125]. The
chokepoint enzyme deoxyuridine 5'-triphosphate nucleotidohydrolase (dUTPase) prevents
incorporation of uracil into DNA and is important for DNA integrity [126]. dUTPase is a potential
antimalarial drug target against *P. falciparum* [127,128]. Holliday junction DNA helicase (*ruvB*)
participates in homologous recombination and repair of replication forks, and is therefore essential
for bacterial growth. Holliday junction processing components were previously identified as targets
for antimicrobials in *E. coli* [129] and *Neisseria gonorrhoeae* [130]. The chokepoint enzyme uracil-DNA
glycosylase (*ung*) has a role in uracil excision repair, and is a candidate anti-malarial drug target [131],
as well as a potential target to control growth of GC-rich bacteria such as *Pseudomonas aeruginosa* and

Mycobacterium smegmatis [132]. DNA polymerase I was identified as a chokepoint in *B. pseudomallei*.
Putative inhibitors of DNA polymerase I (*polA*), and subsequently DNA synthesis, have been
explored as possible antimicrobials [133,134].
Some chokepoint enzymes in *B. pseudomallei* have annotated functions in the biosynthesis of cell
wall components. One of these, UDP-3-O-[3-hydroxymyristoyl] glucosamine N-acyltransferase
(*lpxD*), catalyzes the third step in the lipid A biosynthesis pathway [135]. The lipid A component of
bacterial LPS is of particular interest because it is essential for cell viability and is highly conserved
[136]. This pathway is a target for new antibacterial therapeutics in *Escherichia coli* [137]. Another
chokepoint involved in cell wall synthesis is undecaprenyl diphosphate synthase (*uppS*), which
catalyzes the synthesis of a polyisoprenoid essential for both peptidoglycan and cell wall teichoic
acid synthesis. UppS is a critical enzyme required for bacterial survival, and is an antibacterial target
in *Staphylococcus aureus* [138], *Bacteroides fragilis*, *Vibrio vulnificus*, *E. coli* [139] and *H. pylori* [140].
Several classes of compounds that inhibit UppS function have been discovered [138,141,142]. Two
additional *B. pseudomallei* chokepoints, undecaprenyldiphospho-muramoylpentapeptide beta-N-
acetylglucosaminyltransferase (*murG*), and UDP-N-acetylglucosamine 1-carboxyvinyltransferase
(*murA*) are likely involved in peptidoglycan biosynthesis. MurG is the target of the antibiotic
ramoplanin in *Staphylococcus aureus* [143]. Other potential inhibitors of MurG have been identified by
high throughput screening [144]. A small molecule inhibitor of MurG that augments the activity of
β-lactams against methicillin-resistant *Staphylococcus aureus* was recently identified [145]. MurA has
been a popular target for the design of novel antibiotics, and several inhibitors of MurA have been
identified that are active against various bacterial species [91,146-151]. The chokepoint enzyme
glucose-1-phosphate thymidylyltransferase (*rfbA/rmlA*), involved in O antigen biosynthesis, is also a
target in *Streptococcus pneumoniae* [152] and *Pseudomonas aeruginosa* [153].
The rest of the chokepoint enzymes in Table 3 are components of various biosynthesis pathways.
These enzymes include argininosuccinate synthase (*argG*), which catalyzes the second to last step in
L-arginine biosynthesis, and is associated with pathogenesis in the parasite *Leishmania donovani* [154],
Streptococcus pneumoniae [155], and *B. cenocepacia* [156]. The chokepoint enzyme 3-isopropylmalate
dehydrogenase (*leuB*) is the third enzyme specific to leucine biosynthesis in microorganisms [157],
and has been investigated as an antibacterial target in *M. tuberculosis* [158]. The *B. pseudomallei*
chokepoint dihydroneopterin aldolase (*folB*) is part of the tetrahydrofolate biosynthesis process, and
is essential for growth and biomass production in *Acinetobacter baylyi*, *Bacillus anthracis*, *Francisella*
tularensis, *F. tularensis* subsp. novicida strain U112, *Mycobacterium tuberculosis*, *Helicobacter pylori*,
Pseudomonas aeruginosa, *Salmonella enterica* serovar Typhi and *Yersinia pestis* [95]. Three of the
chokepoint genes identified in *B. pseudomallei* encoded lipoamide dehydrogenase, a component of the
pyruvate dehydrogenase complex, which converts pyruvate to acetyl-CoA as part of central
metabolism [159]. Lipoamide dehydrogenase is also a target in *M. tuberculosis*, where deletion
drastically impaired the pathogen's ability to establish infection in the mouse [160]. The
mycobacterial version has only 36% identity with the human homolog. Lipoamide dehydrogenase is
a target of drugs against trypanosomal infections [161]. Adenylosuccinate synthetase (*purA*) is also a
chokepoint enzyme and potential therapeutic target that is involved in purine salvage in *Leishmania*
donovani [162]. Chokepoint enzyme 3-dehydroquinate dehydratase (*aroQ*) is a component of the
shikimate pathway for chorismate biosynthesis and is a target of known inhibitors in *M. tuberculosis*,
Enterococcus faecalis and *Streptomyces coelicolor* [163-167].
The *B. pseudomallei* chokepoint enzyme 3-oxoacyl-ACP synthase (*fabF*), involved in fatty acid
synthesis, is already an antibacterial target in *E. coli*, and a specific inhibitor, cerulenin, has been
identified [168,169]. Another fatty acid synthesis chokepoint enzyme in *B. pseudomallei* was a biotin-
dependent acetyl-CoA carbosylase. Biotin dependent carboxylases comprise a large group of
enzymes that participate in a variety of cellular processes, including fatty acid metabolism, amino
acid metabolism, carbohydrate metabolism, polyketide biosynthesis, urea utilization, etc. (reviewed
by [170]). Acetyl-CoA carboxylase is comprised of two enzymes, biotin carboxylase and
carboxyltransferase, and catalyzes the first committed step in fatty acid synthesis [171]. Acetyl-CoA
carboxylase is an antimicrobial target in *M. tuberculosis* [172], *E. coli* [173,174], other bacteria and most

living organisms (reviewed by [175]). All of the chokepoints in Table 3 are essential genes in *B.*
pseudomallei MSHR668 and K96243, as determined by blasting the chokepoint enzyme sequences
against essential gene sequences in the Database of Essential Genes [176] and by comparing to the list
of essential genes previously identified in K96243 [177].

To determine if *B. pseudomallei* deletion mutants were available for each of the chokepoints in
Table 3, searches of the internet, PubMed, and the Burkholderia Genome Database
(http://burkholderia.com) were performed. Based on these searches, none of the chokepoint enzymes
in Table 3 had a mutant available; however, a *B. cenocepacia argG* mutant has attenuated virulence
[178], and *recA* mutants have been identified in *B. cepacia* [179].

Additional metabolic enzymes, not identified as chokepoints in this study, and pathways critical
for bacterial growth and survival have been mentioned with respect to target identification. These
include anaplerotic pathways that turned on by limiting carbon sources [180], the glyxoalate shunt
enzyme isocitrate lyase [16,181,182], involved in the metabolism of fatty acids [15,181], enoyl-ACP
reductase (FabI) in the type II fatty acid biosynthesis pathway [183], and alanine racemase [23].

4.4.2 Flux balance analysis

To gain an understanding of the metabolic processes in *B. pseudomallei* MSHR668 that are active
under different environmental conditions, and to test the effect of deletion of each chokepoint enzyme
on the growth of *B. pseudomallei in silico*, metabolic network models were constructed and FBA was
performed. The first FBA model, of the unconstrained network in MSHR668, included all possible
biomass compounds that could be produced, all nutrients that could be consumed, had no weights
imposed on the biomass metabolites and no constraints imposed on the nutrients. [Supplementary
material, S1_Final_unconstrained_model_inputs.pdf, S2_Final_unconstrained_model_solution.pdf].
This model likely represents the metabolic potential of *B. pseudomallei* in a soil or water environment
where abundant carbon and nitrogen sources are available. To mimic the conditions that *B.*
pseudomallei experiences in culture, a separate model was constructed that provided only the
nutrients present in LB media plus glycerol [43]. Constraints were included on some of the nutrients
in this model (ADP, Pi, proton and glycerol). [Supplementary material,
S3_Final_LB_model_inputs.pdf, S4_Final_LB_model_solution.pdf]. A third model was attempted, to
mimic infection conditions using only the nutrients present in the host cell cytosol. However, no
comprehensive studies have identified a complete list of host cell nutrients that are available to *B.*
pseudomallei during infection. Also, the specific carbon requirements of *B. pseudomallei* in either
human macrophages or amoebae have not been determined. After trying to compile a list of nutrients
that mimic the content of the host cytosol, from the literature and from gene expression studies of *B.*
pseudomallei during infection, this model did not produce a solution so it was abandoned. However,
the LB media model contained a similar set of nutrients to those used by the intracellular pathogen
L. pneumophila [47,48,50],. so it may in fact be somewhat representative of infection conditions.

In silico knockout experiments were performed, where each chokepoint enzyme was knocked
out, one at a time, to assess the effect on the total flux through the unconstrained and LB media
models. Five chokepoint enzymes, when knocked out, had an effect on the models (Table 4).
Specifically, knocking out BURPS668_3328 (*tgt*) and BURPS668_A2451 (*leuB*), eliminated the biomass
flux in the unconstrained model, while knockout of BURPS668_2426 (*lpxD*), BURPS668_2433 (*uppS*),
and BURPS668_3525 (*murG*) eliminated the biomass flux in the LB media model. Knockout of
BURPS668_2433 (*uppS*), and BURPS668_3525 (*murG*) decreased the biomass flux in the unconstrained
model, but did not eliminate it. The metabolic pathways that these enzymes belong to are shown in
Figure 2.

Figure 2. Metabolic pathways in *B. pseudomallei* MSHR668 that show reduced flux when chokepoint enzymes (indicated by pink arrows) are deleted. A. The mono-trans, poly-cis decaprenyl phosphate biosynthesis pathway that contains the chokepoint enzyme undecaprenyl diphosphate synthase (*uppS*). B. The two chokepoints UDP-N-acetylglucosamine--N-acetylmuramyl-(pentapeptide) pyrophosphoryl-decaprenol N-acetylglucosamine transferase (*murG*) and UDP-3-O-(R-3-hydroxymyristoyl)-glucosamine N-acyltransferase (*lpxD*), involved in peptidoglycan and lipid A biosynthesis, respectively. C. The chokepoint enzyme tRNA-guanine transglycosylase (*tgt*), involved in queosine biosynthesis. D. The 3-isopropylmalate dehydrogenase (*leuB*) chokepoint enzyme performs the third step in leucine biosynthesis. *In silico* deletion of UDP-3-O-(R-3-hydroxymyristoyl)-glucosamine N-acyltransferase (*lpxD*) reduced flux through the *B. pseudomallei* metabolic network in the LB media model, deletion of undecaprenyl diphosphate synthase (*uppS*) reduced flux through both unconstrained and LB media models, and deletion of tRNA-guanine transglycosylase (*tgt*) and 3-isopropylmalate dehydrogenase (*leuB*) reduced flux in the unconstrained model. These pathways were rendered by the Cellular Overview feature of Pathway Tools.

In terms of carbon sources, glucose was utilized as a nutrient in the unconstrained model of *B. pseudomallei* MSHR668. However, when glycerol was replaced by glucose in the LB media model, no FBA solution was found [data not shown]. This was a somewhat unexpected result, as *B. pseudomallei* can utilize glucose as a carbon source in culture [184]. One possible explanation for this result is that additional nutrients required for glucose utilization were missing from the input nutrients list. Co-metabolism of more than one carbon substrate is a metabolic strategy employed by intracellular bacteria replicating inside host cells to provide carbon for energy and biosynthesis [59]. For example, *Listeria monocytogenes* can use both glycerol and lactate as carbon sources, [57,185-187]. It has been suggested that during infection by *Listeria*, the host cell may not contain enough glucose to activate bacterial PTS glucose transporters, so alternative carbon sources are important for survival and virulence of the pathogen [57]. *M. tuberculosis* also relies on glycerol and fatty acids as carbon sources

in the macrophage environment [188] As the unconstrained *B. pseudomallei* model included a much
longer list of nutrients than the LB model and also could use glucose as a nutrient, this could be the
case. Another explanation for glucose not being utilized by the LB media model is that intracellular
bacteria seem to prefer other substrates over glucose during infection [189], and glycerol may be a
major carbon source for intracellular bacteria during infection [189,190]. This may also be the
situation for *B. pseudomallei* inside host cells. Glycerol feeds into the second half of the
glycolysis/gluconeogenesis pathway through its conversion to dihydroxyacetone phosphate
[www.metacyc.org; [191]], bypassing the first four steps of glycolysis. We previously determined that
the *B. pseudomallei* MSHR668 genome has the full set of genes to perform this conversion [37]. Two
studies of *L. monocytogenes* infection support the idea that intracellular pathogens generally may use
glycerol rather than glucose as a main carbon source while inside host cells. Transcription profiles of
L. monocytogenes grown in mouse macrophages showed reduced expression of genes encoding some
of the enzymes involved in glycolysis, in particular phosphoglucose isomerase (*pgi*), which converts
glucose-6-phosphate into fructose-6-phosphate, and the five steps involved in the conversion of
glyceraldehyde-3-phosphate to pyruvate [189]. Similar transcription profiles were seen during *L.*
monocytogenes infection of Caco-2 epithelial cells [192]. Both studies showed increased expression of
genes involved in the uptake and utilization of glycerol [189,192]: these genes were *glpF, glpK, glpD*
and *dhaK*.

To date, no study has determined precisely which carbon substrates are utilized by *B.*
pseudomallei during infection of host cells. In addition to glycerol, there is evidence that *B. pseudomallei*
may utilize aromatic carbon compounds such as benzoate and phenylacetic acid as carbon sources
for intracellular survival [53]. One study showed that in *B. pseudomallei* 1026b, glycolytic pathway
and TCA cycle genes were down-regulated during infection of hamster [193], supporting the idea
that *B. pseudomallei* may prefer carbon sources other than glucose while inside host cells. Other studies
examined genes induced by hypoxia, which is a condition present in infected macrophages [194] and
changes in *B. pseudomallei* gene expression during infection of rat lungs {van Schaik, 2008 #194.

Complicating the situation even more, the complete nutrient content of a representative
mammalian host cell cytosol has not been determined yet, so a consensus set of nutrients present in
the cytosol of different host cell types is still out of reach {Eisenreich, 2013 #40}. This is largely due to
the challenges in designing appropriate infection models and robust analytical approaches to
measure metabolic changes occurring in host cells during infection. Because of these limitations on
both the pathogen and host sides, it is difficult to predict which carbon sources pathogens can use to
grow inside host cells. While we don't know the exact biochemical composition of a mammalian cell
cytosol, we do know some details about mammalian cells in general. For example, the cytosol of a
typical cell has low magnesium, sodium and calcium concentrations, and a high potassium
concentration at neutral pH [195]. In addition, mammalian cells contain small amounts of amino
acids, plus significant amounts of TCA cycle intermediates [196,197]. Once inside host cells,
intracellular bacteria may stimulate host cell responses to produce needed nutrients [190]. However,
host-pathogen interactions during infection are complicated, as some host defense responses are
aimed at inhibiting pathogen survival and proliferation, for instance by decreasing metabolic
activities that provide nutrients to the pathogen [180].

5. Conclusions

This work is the first to use genome scale metabolic modeling to address *B. pseudomallei*
metabolism as a source of new drug targets. While identifying the nutrients available to *B.*
pseudomallei inside host cells was difficult, the effort described here identified a set of twenty-six
chokepoint enzyme drug targets; *in silico* deletion of five of these target enzymes reduced the total
biomass flux through the *B. pseudomallei* metabolic network. While a genome-based approach like
this can streamline the initial steps of antibacterial target identification, the true utility of this process
will be demonstrated when the targets are experimentally verified by performing knockout
experiments in culture, followed by efficacy testing of candidate drugs in culture and in animal
models of infection.

Supplementary Materials: The following are available online at www.mdpi.com/link, S1_Final_unconstrained_model_inputs.pdf,S2_Final_unconstrained_model_solution.pdf, S3_Final_LB_model_inputs.pdf, S4_Final_LB_model_solution.pdf, S5_Nutrients_infection_model.pdf

Acknowledgments: This project was supported by the Defense Threat Reduction Agency (JSTO-CBD Proposal # CBCALL12-IS1-1-0283).

Conflicts of Interest: The author declares no conflict of interest. The founding sponsors had no role in the design of the study; in the collection, analyses, or interpretation of data; in the writing of the manuscript, and in the decision to publish the results.

References

1. White, N.J. Melioidosis. *Lancet* **2003**, *361*, 1715-1722.

2. Currie, B.J. Melioidosis: An important cause of pneumonia in residents of and travellers returned from endemic regions. *Eur Respir J* **2003**, *22*, 542-550.

3. Schweizer, H.P. Mechanisms of antibiotic resistance in *Burkholderia pseudomallei*: Implications for treatment of melioidosis. *Future Microbiol.* **2012**, *7*, 1389-1399.

4. Dance, D. Treatment and prophylaxis of melioidosis. *Int J Antimicrob Agents.* **2014**, *43*, 310-318.

5. Mays, E.E.; Ricketts, E.A. Melioidosis: Recrudescence associated with bronchogenic carcinoma twenty-six years following initial geographic exposure. *Chest* **1975**, *68*.

6. Thibault, F.M.; Hernandez, E.; Vidal, D.R.; Girardet, M.; Cavallo, J.D. Antibiotic susceptibility of 65 isolates of *Burkholderia pseudomallei* and *Burkholderia mallei* to 35 antimicrobial agents. *J. Antimicrob. Chemother.* **2004**, *54*, 1134-1138.

7. Jenney, A.W.; Lum, G.; Fisher, D.A.; Currie, B.J. Antibiotic susceptibility of *Burkholderia pseudomallei* from tropical northern australia and implications for therapy of melioidosis. *Int. J. Antimicrob. Agents* **2001**, *17*, 109-113.

8. Dance, D.A.; Wuthiekanun, V.; Chaowagul, W.; White, N.J. The antimicrobial susceptibility of *Pseudomonas pseudomallei*. Emergence of resistance in vitro and during treatment. *J Antimicrob Chemother.* **1989**, *24*, 295-309.

9. Galperin, M.Y.; Koonin, E.V. Searching for drug targets in microbial genomes. *Curr Opin Biotechnol.* **1999**, *10*, 571-578.

10. Oberhardt, M.A.; Yizhak, K.; Ruppin, E. Metabolically re-modeling the drug pipeline. *Curr Opin Pharmacol.* **2013**, *13*, 778-785.

11. Silver, L.L. Challenges of antibacterial discovery. *Clin Microbiol Rev.* **2011**, *24*, 71-109.

12. Cummings, J.E.; Kingry, L.C.; Rholl, D.A.; Schweizer, H.P.; Tonge, P.J.; Slayden, R.A. The *Burkholderia pseudomallei* enoyl-acyl carrier protein reductase fabi1 is essential for in vivo growth and is the target of a novel chemotherapeutic with efficacy. *Antimicrob Agents Chemother.* **2014**, *58*, 931-935.

13. Lu, H.; Tonge, P.J. Inhibitors of fabi, an enzyme drug target in the bacterial fatty acid biosynthesis pathway. *Acc Chem Res.* **2008**, *41*, 11-20.

14. Tong, L.; Harwood, H.J.J. Acetyl-coenzyme a carboxylases: Versatile targets for drug discovery. *J Cell Biochem* **2006**, *99*, 1478-1488.

15. Muñoz-Elías, E.J.; McKinney, J.D. *Mycobacterium tuberculosis* isocitrate lyases 1 and 2 are jointly required for in vivo growth and virulence. *Nat Med.* **2005**, *11*, 638-644.

16. van Schaik, E.J.; Tom, M.; Woods, D.E. *Burkholderia pseudomallei* isocitrate lyase is a
persistence factor in pulmonary melioidosis: Implications for the development of isocitrate
lyase inhibitors as novel antimicrobials. *Infect Immun.* **2009**, *77*, 4275-4283.

17. Srilunchang, T.; Proungvitaya, T.; Wongratanacheewin, S.; Strugnell, R.; Homchampa, P.
Construction and characterization of an unmarked aroc deletion mutant of *Burkholderia*
pseudomallei strain a2. *Southeast Asian J Trop Med Public Health* **2009**, *40*, 123-130.

18. Breitbach, K.; Köhler, J.; Steinmetz, I. Induction of protective immunity against *Burkholderia*
pseudomallei using attenuated mutants with defects in the intracellular life cycle. *Trans R Soc*
Trop Med Hyg **2008**, *102 Suppl 1*, S89-94.

19. Pilatz, S.; Breitbach, K.; Hein, N.; Fehlhaber, B.; Schulze, J.; Brenneke, B.; Eberl, L.; Steinmetz,
I. Identification of *Burkholderia pseudomallei* genes required for the intracellular life cycle and
in vivo virulence. *Infect Immun* **2006**, *74*, 3576-3586.

20. Cuccui, J.; Easton, A.; Chu, K.K.; Bancroft, G.J.; Oyston, P.C.; Titball, R.W.; Wren, B.W.
Development of signature-tagged mutagenesis in *Burkholderia pseudomallei* to identify genes
important in survival and pathogenesis. *Infect Immun* **2007**, *75*, 1186-1195.

21. Atkins, T.; Prior, R.G.; Mack, K.; Russell, P.; Nelson, M.; Oyston, P.C.; Dougan, G.; Titball,
R.W. A mutant of *Burkholderia pseudomallei*, auxotrophic in the branched chain amino acid
biosynthetic pathway, is attenuated and protective in a murine model of melioidosis. *Infect*
Immun **2002**, *70*, 5290-5294.

22. Levine, H.B.; Maurer, R.L. Immunization with an induced avirulent auxotrophic mutant of
Pseudomonas pseudomallei. *J Immunol* **1958**, *81*, 433-438.

23. Zajdowicz, S.L.; Jones-Carson, J.; Vazquez-Torres, A.; Jobling, M.G.; Gill, R.E.; Holmes, R.K.
Alanine racemase mutants of *Burkholderia pseudomallei* and *Burkholderia mallei* and use of
alanine racemase as a non-antibiotic-based selectable marker. *PLoS One* **2011**, *6*, e21523.

24. Vanaporn, M.; Wand, M.; Michell, S.L.; Sarkar-Tyson, M.; Ireland, P.; Goldman, S.;
Kewcharoenwong, C.; Rinchai, D.; Lertmemongkolchai, G.; Titball, R.W. Superoxide
dismutase c is required for intracellular survival and virulence of *Burkholderia pseudomallei*.
Microbiology **2011**, *157*, 2392-2400.

25. Lee, H.S.; Gu, F.; Ching, S.M.; Lam, Y.; Chua, K.L. Cdpa is a *Burkholderia pseudomallei* cyclic
di-gmp phosphodiesterase involved in autoaggregation, flagellum synthesis, motility,
biofilm formation, cell invasion, and cytotoxicity. *Infect Immun* **2010**, *78*, 1832-1840.

26. McPhillie, M.J.; Cain, R.M.; Narramore, S.; Fishwick, C.W.; Simmons, K.J. Computational
methods to identify new antibacterial targets. *Chem Biol Drug Des.* **2015**, *85*, 22-29.

27. Chung, B.K.; Dick, T.; Lee, D.Y. In silico analyses for the discovery of tuberculosis drug
targets. *J Antimicrob Chemother.* **2013**, *68*, 2701-2709.

28. Mobegi, F.M.; van Hijum, S.A.; Burghout, P.; Bootsma, H.J.; de Vries, S.P.; van der Gaast-de
Jongh, C.E.; Simonetti, E.; Langereis, J.D.; Hermans, P.W.; de Jonge, M.I., *et al.* From microbial
gene essentiality to novel antimicrobial drug targets. *BMC Genomics* **2014**, *15*, 958.

29. Hamilton, J.J.; Reed, J.L. Software platforms to facilitate reconstructing genome-scale
metabolic networks. *Environ Microbiol.* **2014**, *16*, 49-59.

30. Thiele, I.; Palsson, B.Ø. A protocol for generating a high-quality genome-scale metabolic
reconstruction. *Nat. Protoc.* **2010**, *5*, 93-121.

31.   Yeh, I.; Hanekamp, T.; Tsoka, S.; Karp, P.D.; Altman, R.B. Computational analysis of
Plasmodium falciparum metabolism: Organizing genomic information to facilitate drug
discovery. *Genome Res* **2004**, *14*, 917-924.

32.   Duffield, M.; Cooper, I.; McAlister, E.; Bayliss, M.; Ford, D.; Oyston, P. Predicting conserved
essential genes in bacteria: In silico identification of putative drug targets. *Mol Biosyst.* **2010**,
6, 2482-2489.

33.   Sakharkar, K.R.; Sakharkar, M.K.; Chow, V.T. A novel genomics approach for the
identification of drug targets in pathogens, with special reference to *Pseudomonas aeruginosa.*
In Silico Biol. **2004**, *4*, 355-360.

34.   Singh, S.; Malik, B.K.; Sharma, D.K. Metabolic pathway analysis of *s. Pneumoniae*: An in silico
approach towards drug-design. *J Bioinform Comput Biol* **2007**, *5*, 135-153.

35.   Karp, P.D.; Paley, S.; Romero, P. The pathway tools software. *Bioinformatics* **2002**, *18 Suppl 1*,
S225-S232.

36.   Aziz, R.K.; Bartels, D.; Best, A.A.; DeJongh, M.; Disz, T.; Edwards, R.A.; Formsma, K.; Gerdes,
S.; Glass, E.M.; Kubal, M., *et al.* The rast server: Rapid annotations using subsystems
technology. *BMC Genomics* **2008**, *9*, 75.

37.   Challacombe, J.F.; Stubben, C.J.; Klimko, C.P.; Welkos, S.L.; Kern, S.J.; Bozue, J.A.; Worsham,
P.L.; Cote, C.K.; Wolfe, D.N. Interrogation of the *Burkholderia pseudomallei* genome to address
differential virulence among isolates. *PLoS One* **2014**, *9*, e115951.

38.   Karp, P.D.; Latendresse, M.; Caspi, R. The pathway tools pathway prediction algorithm.
Stand Genomic Sci **2011**, *5*, 424-429.

39.   Latendresse, M.; Krummenacker, M.; Trupp, M.; Karp, P.D. Construction and completion of
flux balance models from pathway databases. *Bioinformatics* **2102**, *28*, 388-396.

40.   Fraser, D.; Powell, R.E. The kinetics of trypsin digestion. *J Biol Chem.* **1950**, *187*, 803-820.

41.   Bertani, G. Studies on lysogenesis. I. The mode of phage liberation by lysogenic *Escherichia*
coli. *J Bacteriol.* **1951**, *62*, 293-300.

42.   Grant, C.L.; Pramer, D. Minor element composition of yeast extract. *J Bacteriol.* **1962**, *84*, 869-
870.

43.   Oh, Y.K.; Palsson, B.O.; Park, S.M.; Schilling, C.H.; Mahadevan, R. Genome-scale
reconstruction of metabolic network in *Bacillus subtilis* based on high-throughput
phenotyping and gene essentiality data. *J Biol Chem.* **2007**, *282*, 28791-28799.

44.   Zhang, R.; Lin, Y. Deg 5.0, a database of essential genes in both prokaryotes and eukaryotes.
Nucleic Acids Res. **2009**, *37(Database issue)*, D455-458.

45.   Wishart, D.S.; Knox, C.; Guo, A.C.; Cheng, D.; Shrivastava, S.; Tzur, D.; Gautam, B.;
Hassanali, M. Drugbank: A knowledgebase for drugs, drug actions and drug targets. *Nucleic*
Acids Res. **2008**, *36(Database issue)*, D901-906.

46.   Shannon, P.; Markiel, A.; Ozier, O.; Baliga, N.S.; Wang, J.T.; Ramage, D.; Amin, N.;
Schwikowski, B.; Ideker, T. Cytoscape: A software environment for integrated models of
biomolecular interaction networks. *Genome Res* **2003**, *13*, 2498-2504.

47.   Price, C.T.; Richards, A.M.; Abu Kwaik, Y. Nutrient generation and retrieval from the host
cell cytosol by intra-vacuolar *Legionella pneumophila. Front Cell Infect Microbiol* **2014**, *4*, 111.

48.   Price, C.T.; Abu Kwaik, Y. Amoebae and mammals deliver protein-rich atkins diet meals to
Legionella cells. *Microbe* **2012**, *7*, 506-513.

49.   Gao, L.Y.; Harb, O.S.; Abu Kwaik, Y. Utilization of similar mechanisms by *Legionella*
719   *pneumophila* to parasitize two evolutionarily distant host cells, mammalian macrophages and
720   protozoa. *Infect. Immun.* **1997**, *65*, 4738-4746.

721   50.   Eylert, E.; Herrmann, V.; Jules, M.; Gillmaier, N.; Lautner, M.; Buchrieser, C.; Eisenreich, W.;
722   Heuner, K. Isotopologue profiling of *Legionella pneumophila*: Role of serine and glucose as
723   carbon substrates. *J Biol Chem.* **2010**, *285*, 22232-22243.

724   51.   DeShazer, D. Genomic diversity of *Burkholderia pseudomallei* clinical isolates: Subtractive
725   hybridization reveals a *Burkholderia mallei*-specific prophage in *B. pseudomallei* 1026b. *J.*
726   *Bacteriol.* **2004**, *186*, 3938-3950.

727   52.   Ooi, W.F.; Ong, C.; Nandi, T.; Kreisberg, J.F.; Chua, H.H.; Sun, G.; Chen, Y.; Mueller, C.;
728   Conejero, L.; Eshaghi, M., *et al.* The condition-dependent transcriptional landscape of
729   *Burkholderia pseudomallei.* PLoS Genet. **2013**, *9*, e1003795.

730   53.   Chieng, S.; Carreto, L.; Nathan, S. *Burkholderia pseudomallei* transcriptional adaptation in
731   macrophages. *BMC Genomics* **2012**, *13*, 328.

732   54.   Eriksson, S.; Lucchini, S.; Thompson, A.; Rhen, M.; Hinton, J.C. Unravelling the biology of
733   macrophage infection by gene expression profiling of intracellular *Salmonella enterica. Mol*
734   *Microbiol* **2003**, *47*, 103-118.

735   55.   Zdziarski, J.; Brzuszkiewicz, E.; Wullt, B.; Liesegang, H.; Biran, D.; Voigt, B.; Grönberg-
736   Hernandez, J.; Ragnarsdottir, B.; Hecker, M.; Ron, E.Z., *et al.* Host imprints on bacterial
737   genomes--rapid, divergent evolution in individual patients. *PLoS Pathog.* **2010**, *6*, e1001078.

738   56.   Reniere, M.L.; Whiteley, A.T.; Hamilton, K.L.; John, S.M.; Lauer, P.; Brennan, R.G.; Portnoy,
739   D.A. Glutathione activates virulence gene expression of an intracellular pathogen. *Nature*
740   **2015**, *517*, 70-31.

741   57.   Eylert, E.; Schär, J.; Mertins, S.; Stoll, R.; Bacher, A.; Goebel, W.; Eisenreich, W. Carbon
742   metabolism of *Listeria monocytogenes* growing inside macrophages. *Mol Microbiol.* **2008**, *69*,
743   1008-1017.

744   58.   Chico-Calero, I.; Suárez, M.; González-Zorn, B.; Scortti, M.; Slaghuis, J.; Goebel, W.; Vázquez-
745   Boland, J.A.; Consortium, E.L.G. Hpt, a bacterial homolog of the microsomal glucose- 6-
746   phosphate translocase, mediates rapid intracellular proliferation in *Listeria. Proc Natl Acad Sci*
747   *U S A* **2002**, *99*, 431-436.

748   59.   Eisenreich, W.; Dandekar, T.; Heesemann, J.; Goebel, W. Carbon metabolism of intracellular
749   bacterial pathogens and possible links to virulence. *Nat Rev Microbiol* **2010**, *8*, 401-412.

750   60.   Fuchs, T.M.; Eisenreich, W.; Heesemann, J.; Goebel, W. Metabolic adaptation of human
751   pathogenic and related nonpathogenic bacteria to extra- and intracellular habitats. *FEMS*
752   *Microbiol Rev.* **2012**, *36*, 435-462.

753   61.   Gore, A.L.; Payne, S.M. Csra and cra influence *Shigella flexneri* pathogenesis. *Infect Immun.*
754   **2010**, *78*, 4674-4682.

755   62.   Litwin, C.M.; Calderwood, S.B. Role of iron in regulation of virulence genes. *Clin Microbiol*
756   *Rev.* **1993**, *6*, 137-149.

757   63.   Mellies, J.L.; Barron, A.M.; Carmona, A.M. Enteropathogenic and enterohemorrhagic
758   *Escherichia coli* virulence gene regulation. *Infect Immun* **2007**, *75*, 4199-4210.

759   64.   Njoroge, J.W.; Nguyen, Y.; Curtis, M.M.; Moreira, C.G.; Sperandio, V. Virulence meets
760   metabolism: Cra and kdpe gene regulation in enterohemorrhagic *Escherichia coli*. *MBio* **2012**,
761   *3*, e00280-00212.

762   65.   Porcheron, G.; Dozois, C.M. Interplay between iron homeostasis and virulence: Fur and ryhb
763   as major regulators of bacterial pathogenicity. *Vet Microbiol* **2015**, *179*, 2-14.

764   66.   Ray, K.; Marteyn, B.; Sansonetti, P.J.; Tang, C.M. Life on the inside: The intracellular lifestyle
765   of cytosolic bacteria. *Nat Rev Microbiol* **2009**, *7*, 333-340.

766   67.   Rohmer, L.; Hocquet, D.; Miller, S.I. Are pathogenic bacteria just looking for food?
767   Metabolism and microbial pathogenesis. *Trends Microbiol.* **2011**, *19*, 341-348.

768   68.   Gorke, B.; Stulke, J. Carbon catabolite repression in bacteria: Many ways to make the most
769   out of nutrients. *Nat Rev Microbiol* **2008**, *6*, 613-624.

770   69.   Barel, M.; Charbit, A. *Francisella tularensis* intracellular survival: To eat or to die. *Microbes
771   Infect* **2013**, *pii*, S1286-4579(1213)00206-00202.

772   70.   Kwaik, Y.A. Nutrition-based evolution of intracellular pathogens. *Environmental Microbiology
773   Reports* **2015**, *7*, 2-3.

774   71.   Kwaik, Y.A.; Bumann, D. Microbial quest for food in vivo: 'Nutritional virulence' as an
775   emerging paradigm. *Cell Microbiol.* **2013**, *15*, 882-890.

776   72.   Meibom, K.L.; Charbit, A. *Francisella tularensi*s metabolism and its relation to virulence. *Front
777   Microbiol.* **2010**, *1*, 140.

778   73.   Olive, A.J.; Sassetti, C.M. Metabolic crosstalk between host and pathogen: Sensing, adapting
779   and competing. *Nature Reviews Microbiology* **2016**, *14*, 221-234.

780   74.   Santic, M.; Kwaik, Y.A. Nutritional virulence of *Francisella tularensis*. *Front Cell Infect Microbiol*
781   **2013**, *3*, 112.

782   75.   Muñoz-Elías, E.J.; McKinney, J.D. Carbon metabolism of intracellular bacteria. *Cell Microbiol.*
783   **2006**, *8*, 10-22.

784   76.   Brown, S.A.; Palmer, K.L.; Whiteley, M. Revisiting the host as a growth medium. *Nat Rev
785   Microbiol.* **2008**, *6*, 657-666.

786   77.   Fuhrer, T.; Fischer, E.; Sauer, U. Experimental identification and quantification of glucose
787   metabolism in seven bacterial species. *J Bacteriol.* **2005**, *187*, 1581-1590.

788   78.   Renesto, P.; Ogata, H.; Audic, S.; Claverie, J.M.; Raoult, D. Some lessons from *Rickettsia*
789   genomics. *FEMS Microbiol Rev* **2005**, *29*, 99-117.

790   79.   Propst, K.L.; Mima, T.; Choi, K.H.; Dow, S.W.; Schweizer, H.P. A *Burkholderia pseudomallei*
791   deltapurm mutant is avirulent in immunocompetent and immunodeficient animals:
792   Candidate strain for exclusion from select-agent lists. *Infect Immun* **2010**, *78*, 3136-3143.

793   80.   Norris, M.H.; Propst, K.L.; Kang, Y.; Dow, S.W.; Schweizer, H.P.; Hoang, T.T. The
794   *Burkholderia pseudomallei* δasd mutant exhibits attenuated intracellular infectivity and imparts
795   protection against acute inhalation melioidosis in mice. *Infect Immun* **2011**, *79*, 4010-4018.

796   81.   Rodrigues, F.; Sarkar-Tyson, M.; Harding, S.V.; Sim, S.H.; Chua, H.H.; Lin, C.H.; Han, X.;
797   Karuturi, R.K.; Sung, K.; Yu, K., *et al.* Global map of growth-regulated gene expression in
798   *Burkholderia pseudomallei*, the causative agent of melioidosis. *J Bacteriol* **2006**, *188*, 8178-8188.

799   82.   Korbsrisate, S.; Tomaras, A.P.; Damnin, S.; Ckumdee, J.; Srinon, V.; Lengwehasatit, I.; Vasil,
800   M.L.; Suparak, S. Characterization of two distinct phospholipase c enzymes from *Burkholderia
801   pseudomallei*. *Microbiology* **2007**, *153*, 1907-1915.

802   83.   Ireland, P.M.; McMahon, R.M.; Marshall, L.E.; Halili, M.; Furlong, E.; Tay, S.; Martin, J.L.;
803   Sarkar-Tyson, M. Disarming *Burkholderia pseudomallei:* Structural and functional
804   characterization of a disulfide oxidoreductase (DsbA) required for virulence in vivo. *Antioxid*
805   *Redox Signal* **2014**, *20*, 606-617.

806   84.   Drabner, B.; Guzmán, C.A. Elicitation of predictable immune responses by using live
807   bacterial vectors. *Biomol Eng* **2001**, *17*, 75-82.

808   85.   Peacock, S.J.; Limmathurotsakul, D.; Lubell, Y.; Koh, G.C.; White, L.J.; Day, N.P.; Titball, R.W.
809   Melioidosis vaccines: A systematic review and appraisal of the potential to exploit biodefense
810   vaccines for public health purposes. *PLoS Negl Trop Dis* **2012**, *6*, e1488.

811   86.   Sim SH, Y.Y., Lin CH, Karuturi RKM, Wuthiekanun V, Tuanyok A. The core and accessory
812   genomes of *Burkholderia pseudomallei*: Implications for human melioidosis. *PLoS Pathog* **2008**,
813   *4*, e1000178.

814   87.   Sahl, J.W.; Vazquez, A.J.; Hall, C.M.; Busch, J.D.; Tuanyok, A.; Mayo, M.; Schupp, J.M.;
815   Lummis, M.; Pearson, T.; Shippy, K.*, et al.* The effects of signal erosion and core genome
816   reduction on the identification of diagnostic markers. *mBio* **2016**, *7*, e00846-00816.

817   88.   Butt, A.M.; Tahir, S.; Nasrullah, I.; Idrees, M.; Lu, J.; Tong, Y. *Mycoplasma genitalium*: A
818   comparative genomics study of metabolic pathways for the identification of drug and vaccine
819   targets. *Infect Genet Evol* **2012**, *12*, 53-62.

820   89.   Butt, A.M.; Nasrullah, I.; Tahir, S.; Tong, Y. Comparative genomics analysis of *Mycobacterium*
821   *ulcerans* for the identification of putative essential genes and therapeutic candidates. *PLoS*
822   *One* **2012**, *7*, e43080.

823   90.   Anishetty, S.; Pulimi, M.; Pennathur, G. Potential drug targets in *Mycobacterium tuberculosis*
824   through metabolic pathway analysis. *Comput Biol Chem* **2005**, *29*, 368-378.

825   91.   Yadav, P.K.; Singh, G.; Singh, S.; Gautam, B.; Saad, E.I. Potential therapeutic drug target
826   identification in community acquired-methicillin resistant *Staphylococcus aureus* (ca-mrsa)
827   using computational analysis. *Bioinformation* **2012**, *8*, 664-672.

828   92.   Uddin, R.; Saeed, K. Identification and characterization of potential drug targets by
829   subtractive genome analyses of methicillin resistant *Staphylococcus aureus*. *Comput Biol Chem*
830   **2014**, *48*, 55-63.

831   93.   Uddin, R.; Saeed, K.; Khan, W.; Azam, S.S.; Wadood, A. Metabolic pathway analysis
832   approach: Identification of novel therapeutic target against methicillin resistant
833   *Staphylococcus aureus*. *Gene* **2015**, *556*, 213-226.

834   94.   Barh, D.; Tiwari, S.; Jain, N.; Ali, A.; Santos, A.R.; Misra, A.N.; Azevedo, V.; Kumar, A. In
835   silico subtractive genomics for target identification in human bacterial pathogens. *Drug Dev.*
836   *Res.* **2011**, *72*, 162-177.

837   95.   Ahn, Y.Y.; Lee, D.S.; Burd, H.; Blank, W.; Kapatral, V. Metabolic network analysis-based
838   identification of antimicrobial drug targets in category a bioterrorism agents. *PLoS One* **2014**,
839   *9*, e85195.

840   96.   Damte, D.; Suh, J.W.; Lee, S.J.; Yohannes, S.B.; Hossain, M.A.; Park, S.C. Putative drug and
841   vaccine target protein identification using comparative genomic analysis of kegg annotated
842   metabolic pathways of *Mycoplasma hyopneumoniae*. *Genomics* **2013**, *102*, 47-56.

97.   Fang, K.; Zhao, H.; Sun, C.; Lam, C.M.; Chang, S.; Zhang, K.; Panda, G.; Godinho, M.; Martins
dos Santos, V.A.; Wang, J. Exploring the metabolic network of the epidemic pathogen
Burkholderia cenocepacia j2315 via genome-scale reconstruction. *BMC Syst Biol.* **2011**, *5*, 83.

98.   Lee, D.S.; Burd, H.; Liu, J.; Almaas, E.; Wiest, O.; Barabási, A.L.; Oltvai, Z.N.; Kapatral, V.
Comparative genome-scale metabolic reconstruction and flux balance analysis of multiple
Staphylococcus aureus genomes identify novel antimicrobial drug targets. *J Bacteriol.* **2009**, *191*,
4015-4024.

99.   Rahman, S.A.; Schomburg, D. Observing local and global properties of metabolic pathways:
'Load points' and 'choke points' in the metabolic networks. *Bioinformatics* **2006**, *22*, 1767-1774.

100. Baths, V.; Roy, U.; Singh, T. Disruption of cell wall fatty acid biosynthesis in *Mycobacterium*
tuberculosis using a graph theoretic approach. *Theor Biol Med Model* **2011**, *8*, 5.

101. Chaudhury, S.; Abdulhameed, M.D.; Singh, N.; Tawa, G.J.; D'haeseleer, P.M.; Zemla, A.T.;
Navid, A.; Zhou, C.E.; Franklin, M.C.; Cheung, J., *et al.* Rapid countermeasure discovery
against *Francisella tularensis* based on a metabolic network reconstruction. *PLoS One* **2013**, *8*,
e63369.

102. Heinemann, M.; Kümmel, A.; Ruinatscha, R.; Panke, S. In silico genome-scale reconstruction
and validation of the *Staphylococcus aureus* metabolic network. *Biotechnol Bioeng* **2005**, *92*, 850-
864.

103. Herrgård, M.J.; Fong, S.S.; Palsson, B.Ø. Identification of genome-scale metabolic network
models using experimentally measured flux profiles. *PLoS Comput Biol* **2006**, *2*, e72.

104. Kim, H.U.; Kim, S.Y.; Jeong, H.; Kim, T.Y.; Kim, J.J.; Choy, H.E.; Yi, K.Y.; Rhee, J.H.; Lee, S.Y.
Integrative genome-scale metabolic analysis of *Vibrio vulnificus* for drug targeting and
discovery. *Mol Syst Biol* **2011**, *7*, 460.

105. Raghunathan, A.; Shin, S.; Daefler, S. Systems approach to investigating host-pathogen
interactions in infections with the biothreat agent Francisella. Constraints-based model of
Francisella tularensis. BMC Syst Biol **2010**, *4*, 118.

106. Chavali, A.K.; D'Auria, K.M.; Hewlett, E.L.; Pearson, R.D.; Papin, J.A. A metabolic network
approach for the identification and prioritization of antimicrobial drug targets. *Trends*
Microbiol. **2012**, *20*, 113-123.

107. Fatumo, S.; Plaimas, K.; Mallm, J.P.; Schramm, G.; Adebiyi, E.; Oswald, M.; Eils, R.; König, R.
Estimating novel potential drug targets of *Plasmodium falciparum* by analysing the metabolic
network of knock-out strains *in silico. Infect Genet Evol* **2009**, *9*, 351-358.

108. Guimerà, R.; Sales-Pardo, M.; Amaral, L.A. A network-based method for target selection in
metabolic networks. *Bioinformatics* **2007**, *23*, 1616-1622.

109. Chavali, A.K.; Whittemore, J.D.; Eddy, J.A.; Williams, K.T.; Papin, J.A. Systems analysis of
metabolism in the pathogenic trypanosomatid *Leishmania major. Mol Syst Biol* **2008**, *4*, 177.

110. Perumal, D.; Lim, C.S.; Sakharkar, M.K. A comparative study of metabolic network topology
between a pathogenic and a non-pathogenic bacterium for potential drug target
identification. *Summit on Translat Bioinforma.* **2009**, *2009*, 100-104.

111. Taylor, C.M.; Wang, Q.; Rosa, B.A.; Huang, S.C.; Powell, K.; Schedl, T.; Pearce, E.J.;
Abubucker, S.; Mitreva, M. Discovery of anthelmintic drug targets and drugs using
chokepoints in nematode metabolic pathways. *PLoS Pathog* **2013**, *9*, e1003505.

112. Ioerger, T.R.; O'Malley, T.; Liao, R.; Guinn, K.M.; Hickey, M.J.; Mohaideen, N.; Murphy, K.C.;
Boshoff, H.I.; Mizrahi, V.; Rubin, E.J., *et al.* Identification of new drug targets and resistance
mechanisms in *Mycobacterium tuberculosis*. *PLoS One* **2013**, *8*, e75245.

113. Orelle, C.; Szal, T.; Klepacki, D.; Shaw, K.J.; Vázquez-Laslop, N.; Mankin, A.S. Identifying the
targets of aminoacyl-trna synthetase inhibitors by primer extension inhibition. *Nucleic Acids*
Res **2013**, *41*, e144.

114. Novoa, E.M.; Camacho, N.; Tor, A.; Wilkinson, B.; Moss, S.; Marín-García, P.; Azcárate, I.G.;
Bautista, J.M.; Mirando, A.C.; Francklyn, C.S., *et al.* Analogs of natural aminoacyl-trna
synthetase inhibitors clear malaria in vivo. *Proc Natl Acad Sci U S A.* **2014**, *111*, E5508-5517.

115. Schimmel, P.; Tao, J.; Hill, J. Aminoacyl trna synthetases as targets for new anti-infectives.
FASEB J **1998**, *12*, 1599-1609.

116. Kim, S.; Lee, S.W.; Choi, E.C.; Choi, S.Y. Aminoacyl-trna synthetases and their inhibitors as a
novel family of antibiotics. *Appl Microbiol Biotechnol.* **2003**, *61*, 278-288.

117. Hurdle, J.G.; O'Neill, A.J.; Chopra, I. Prospects for aminoacyl-trna synthetase inhibitors as
new antimicrobial agents. *Antimicrob Agents Chemother.* **2005**, *49*, 4821-4833.

118. Thomas, C.M.; Hothersall, J.; Willis, C.L.; Simpson, T.J. Resistance to and synthesis of the
antibiotic mupirocin. *Nat Rev Microbiol* **2010**, *8*, 281-289.

119. Koh, C.Y.; Siddaramaiah, L.K.; Ranade, R.M.; Nguyen, J.; Jian, T.; Zhang, Z.; Gillespie, J.R.;
Buckner, F.S.; Verlinde, C.L.; Fan, E., *et al.* A binding hotspot in *Trypanosoma cruzi* histidyl-
trna synthetase revealed by fragment-based crystallographic cocktail screens. *Acta Crystallogr*
D Biol Crystallogr **2015**, *71*, 1684-1698.

120. Biela, I.; Tidten-Luksch, N.; Immekus, F.; Glinca, S.; Nguyen, T.X.; Gerber, H.D.; Heine, A.;
Klebe, G.; Reuter, K. Investigation of specificity determinants in bacterial trna-guanine
transglycosylase reveals queuine, the substrate of its eucaryotic counterpart, as inhibitor.
PLoS One **2013**, *8*, e64240.

121. Grädler, U.; Gerber, H.D.; Goodenough-Lashua, D.M.; Garcia, G.A.; Ficner, R.; Reuter, K.;
Stubbs, M.T.; Klebe, G. A new target for shigellosis: Rational design and crystallographic
studies of inhibitors of trna-guanine transglycosylase. *J Mol Biol* **2001**, *306*, 455-467.

122. Pohlhaus, J.R.; Long, D.T.; O'Reilly, E.; Kreuzer, K.N. The epsilon subunit of DNA
polymerase iii is involved in the nalidixic acid-induced sos response in E*scherichia coli*. *J*
Bacteriol **2008**, *190*, 5239-5247.

123. Cirz, R.T.; Chin, J.K.; Andes, D.R.; de Crécy-Lagard, V.; Craig, W.A.; Romesberg, F.E.
Inhibition of mutation and combating the evolution of antibiotic resistance. *PLoS Biology*
2005, *3*, e176.

124. Nautiyal, A.; Patil, K.N.; Muniyappa, K. Suramin is a potent and selective inhibitor of
Mycobacterium tuberculosis reca protein and the sos response: Reca as a potential target for
antibacterial drug discovery. *J Antimicrob Chemother* **2014**, *69*, 1834-1843.

125. Culyba, M.J.; Mo, C.Y.; Kohli, R.M. Targets for combating the evolution of acquired antibiotic
resistance. *Biochemistry* **2015**, *54*, 3573-3582.

126. Tye, B.-K.; Lehman, I.R. Excision repair of uracil incorporated in DNA as a result of a defect
in dutpase. *J. Mol. Biol.* **1977**, *117*, 293–306.

127. Nguyen, C.; Kasinathan, G.; Leal-Cortijo, I.; Musso-Buendia, A.; Kaiser, M.; Brun, R.; Ruiz-
Pérez, L.M.; Johansson, N.G.; González-Pacanowska, D.; Gilbert, I.H. Deoxyuridine

triphosphate nucleotidohydrolase as a potential antiparasitic drug target. *J Med Chem* **2005**, *48*, 5942-5954.

128. Hampton, S.E.; Baragaña, B.; Schipani, A.; Bosch-Navarrete, C.; Musso-Buendía, J.A.; Recio, E.; Kaiser, M.; Whittingham, J.L.; Roberts, S.M.; Shevtsov, M., *et al.* Design, synthesis, and evaluation of 5'-diphenyl nucleoside analogues as inhibitors of the *Plasmodium falciparum* dutpase. *ChemMedChem* **2011**, *6*, 1816-1831.

129. Kepple, K.V.; Boldt, J.L.; Segall, A.M. Holliday junction-binding peptides inhibit distinct junction-processing enzymes. *Proc Natl Acad Sci U S A* **2005**, *102*, 6867-6872.

130. Sechman, E.V.; Kline, K.A.; Seifert, S. Loss of both holliday junction processing pathways is synthetically lethal in the presence of gonococcal pilin antigenic variation. *Mol Microbiol* **2006**, *61*, 185-193.

131. Suksangpleng, T.; Leartsakulpanich, U.; Moonsom, S.; Siribal, S.; Boonyuen, U.; Wright, G.E.; Chavalitshewinkoon-Petmitr, P. Molecular characterization of *Plasmodium falciparum* uracil-DNA glycosylase and its potential as a new anti-malarial drug target. *Malar J* **2014**, *13*, 149.

132. Venkatesh, J.; Kumar, P.; Krishna, P.S.; Manjunath, R.; Varshney, U. Importance of uracil DNA glycosylase in *Pseudomonas aeruginosa* and *Mycobacterium smegmatis*, g+c-rich bacteria, in mutation prevention, tolerance to acidified nitrite, and endurance in mouse macrophages. . *J Biol Chem.* **2003**, *278*, 24350-24358.

133. Landini, P.; Corti, E.; Goldstein, B.P.; Denaro, M. Mechanism of action of purpuromycin. *Biochem J* **1992**, *284*, 47-52.

134. Nishio, A.; Uyeki, E.M. Cellular uptake and inhibition of DNA synthesis by dihydroxyanthraquinone and two analogues. *Cancer Res* **1983**, *43*, 1951-1956.

135. Kelly, T.M.; Stachula, S.A.; Raetz, C.R.; Anderson, M.S. The firA gene of *Escherichia coli* encodes udp-3-o-(r-3-hydroxymyristoyl)-glucosamine n-acyltransferase. The third step of endotoxin biosynthesis. *J Biol Chem.* **1993**, *268*, 19866-19874.

136. Raetz, C.R.; Whitfield, C. Lipopolysaccharide endotoxins. *Annu Rev Biochem* **2002**, *71*, 635-700.

137. Emiola, A.; George, J.; Andrews, S.S. A complete pathway model for lipid a biosynthesis in *Escherichia coli*. *PLoS One* **2015**, *10*, e0121216.

138. Farha, M.A.; Czarny, T.L.; Myers, C.L.; Worrall, L.J.; French, S.; Conrady, D.G.; Wang, Y.; Oldfield, E.; Strynadka, N.C.; Brown, E.D. Antagonism screen for inhibitors of bacterial cell wall biogenesis uncovers an inhibitor of undecaprenyl diphosphate synthase. *Proc Natl Acad Sci U S A* **2015**, *112*, 11048-11053.

139. Dodbele, S.; Martinez, C.D.; Troutman, J.M. Species differences in alternative substrate utilization by the antibacterial target undecaprenyl pyrophosphate synthase. *Biochemistry* **2014**, *53*, 5042-5050.

140. Kuo, C.J.; Guo, R.T.; Lu, I.L.; Liu, H.G.; Wu, S.Y.; Ko, T.P.; Wang, A.H.; Liang, P.H. Structure-based inhibitors exhibit differential activities against *Helicobacter pylori* and *Escherichia coli* undecaprenyl pyrophosphate synthases. *J Biomed Biotechnol* **2008**, *2008*, 841312.

141. Zhu, W.; Zhang, Y.; Sinko, W.; Hensler, M.E.; Olson, J.; Molohon, K.J.; Lindert, S.; Cao, R.; Li, K.; Wang, K., *et al.* Antibacterial drug leads targeting isoprenoid biosynthesis. *Proc Natl Acad Sci U S A* **2013**, *110*, 123-128.

142.   Sinko, W.; Wang, Y.; Zhu, W.; Zhang, Y.; Feixas, F.; Cox, C.L.; Mitchell, D.A.; Oldfield, E.;
McCammon, J. Undecaprenyl diphosphate synthase inhibitors: Antibacterial drug leads. *J*
Med Chem **2014**, *57*, 5693-5701.

143.   Walker, S.; Chen, L.; Hu, Y.; Rew, Y.; Shin, D.; Boger, D.L. Chemistry and biology of
ramoplanin: A lipoglycodepsipeptide with potent antibiotic activity. *Chem Rev* **2005**, *105*, 449-
476.

144.   Helm, J.S.; Hu, Y.; Chen, L.; Gross, B.; Walker, S. Identification of active-site inhibitors of
murg using a generalizable, high-throughput glycosyltransferase screen. *J Am Chem Soc* **2003**,
125, 11168-11169.

145.   Mann, P.A.; Müller, A.; Xiao, L.; Pereira, P.M.; Yang, C.; Ho Lee, S.; Wang, H.; Trzeciak, J.;
Schneeweis, J.; Dos Santos, M.M., *et al.* Murgocil is a highly bioactive staphylococcal-specific
inhibitor of the peptidoglycan glycosyltransferase enzyme murg. *ACS Chem Biol* **2013**, *8*,
2442-2451.

146.   Olesen, S.H.; Ingles, D.J.; Yang, Y.; Schönbrunn, E. Differential antibacterial properties of the
mura inhibitors terreic acid and fosfomycin. *J Basic Microbiol* **2014**, *54*, 322-326.

147.   Bensen, D.C.; Rodriguez, S.; Nix, J.; Cunningham, M.L.; Tari, L.W. Structure of mura (udp-n-
acetylglucosamine enolpyruvyl transferase) from *Vibrio fischeri* in complex with substrate
udp-n-acetylglucosamine and the drug fosfomycin. *Acta Crystallogr Sect F Struct Biol Cryst*
Commun **2012**, *68*, 382-385.

148.   Gautam, A.; Rishi, P.; Tewari, R. Udp-n-acetylglucosamine enolpyruvyl transferase as a
potential target for antibacterial chemotherapy: Recent developments. *Appl Microbiol*
Biotechnol. **2011**, *92*, 211-212.

149.   Han, H.; Yang, Y.; Olesen, S.H.; Becker, A.; Betzi, S.; Schönbrunn, E. The fungal product
terreic acid is a covalent inhibitor of the bacterial cell wall biosynthetic enzyme udp-n-
acetylglucosamine 1-carboxyvinyltransferase (mura) *Biochemistry* **2010**, *49*, 4276-4282.

150.   Bachelier, A.; Mayer, R.; Klein, C.D. Sesquiterpene lactones are potent and irreversible
inhibitors of the antibacterial target enzyme mura. *Bioorg Med Chem Lett* **2006**, *16*, 5605-5609.

151.   Eschenburg, S.; Priestman, M.A.; Abdul-Latif, F.A.; Delachaume, C.; Fassy, F.; Schönbrunn,
E. A novel inhibitor that suspends the induced fit mechanism of udp-n-acetylglucosamine
enolpyruvyl transferase (mura). *J Biol Chem.* **2005**, *280*, 14070-14075.

152.   Smithen, D.A.; Forget, S.M.; McCormick, N.E.; Syvitski, R.T.; Jakeman, D.L. Polyphosphate-
containing bisubstrate analogues as inhibitors of a bacterial cell wall thymidylyltransferase.
Org Biomol Chem **2015**, *13*, 3347-3350.

153.   Blankenfeldt, W.; Asuncion, M.; Lam, J.S.; Naismith, J.H. The structural basis of the catalytic
mechanism and regulation of glucose-1-phosphate thymidylyltransferase (rmla). *EMBO J*
2000, *19*, 6652-6663.

154.   Lakhal-Naouar, I.; Jardim A, S., R.; Luo S, K., Y.; Nakhasi, H.L.; Duncan, R.C. *Leishmania*
donovani argininosuccinate synthase is an active enzyme associated with parasite
pathogenesis. *PLoS Negl Trop Dis.* **2012**, *6*, e1849.

155.   Piet, J.R.; Geldhoff, M.; van Schaik, B.D.; Brouwer, M.C.; Valls Seron, M.; Jakobs, M.E.;
Schipper, K.; Pannekoek, Y.; Zwinderman, A.H.; van der Poll, T., *et al. Streptococcus*
pneumoniae arginine synthesis genes promote growth and virulence in pneumococcal
meningitis. *J Infect Dis.* **2014**, *209*, 1781-1791.

156. Somvanshi, V.S.; Viswanathan, P.; Jacobs, J.L.; Mulks, M.H.; Sundin, G.W.; Ciche, T.A. The
type 2 secretion pseudopilin, gspj, is required for multihost pathogenicity of *Burkholderia*
cenocepacia au1054. *Infect Immun.* **2010**, *78*, 4110-4121.

157. Yang, H.L.; Kessler, D.P. Genetic analysis of the leucine region in *Escherichia coli* b-r: Gene-
enzyme assignments. *J Bacteriol* **1974**, *117*, 63-72.

158. Singh, R.K.; Kefala, G.; Janowski, R.; Mueller-Dieckmann, C.; von Kries, J.P.; Weiss, M.S. The
high-resolution structure of leub (rv2995c) from *Mycobacterium tuberculosis. J Mol Biol.* **2005**,
346, 1-11.

159. Pettit, F.H.; Reed, L.J. Alpha-keto acid dehydrogenase complexes. 8. Comparison of
dihydrolipoyl dehydrogenases from pyruvate and alpha-ketoglutarate dehydrogenase
complexes of *Escherichia coli. Proc Natl Acad Sci U S A.* **1967**, *58*, 1126-1130.

160. Bryk, R.; Arango, N.; Maksymiuk, C.; Balakrishnan, A.; Wu, Y.T.; Wong, C.H.; Masquelin, T.;
Hipskind, P.; Lima, C.D.; Nathan, C. Lipoamide channel-binding sulfonamides selectively
inhibit mycobacterial lipoamide dehydrogenase. *Biochemistry* **2013**, *52*, 9375-9384.

161. Krauth-Siegel, R.L.; Schöneck, R. Flavoprotein structure and mechanism. 5. Trypanothione
reductase and lipoamide dehydrogenase as targets for a structure-based drug design. *FASEB*
J **1995**, *9*, 1138-1146.

162. Boitz, J.M.; Strasser, R.; Yates, P.A.; Jardim, A.; Ullman, B. Adenylosuccinate synthetase and
adenylosuccinate lyase deficiencies trigger growth and infectivity deficits in *Leishmania*
donovani. J Biol Chem. **2013**, *288*, 8977-8990.

163. Howard, N.I.; Dias, M.V.; Peyrot, F.; Chen, L.; Schmidt, M.F.; Blundell, T.L.; Abell, C. Design
and structural analysis of aromatic inhibitors of type ii dehydroquinase from *Mycobacterium*
tuberculosis. ChemMedChem **2015**, *10*, 116-133.

164. Cheung, V.W.; Xue, B.; Hernandez-Valladares, M.; Go, M.K.; Tung, A.; Aguda, A.H.;
Robinson, R.C.; Yew, W.S. Identification of polyketide inhibitors targeting 3-dehydroquinate
dehydratase in the shikimate pathway of *Enterococcus faecalis. PLoS One* **2014**, *9*, e103598.

165. Ballester, P.J.; Mangold, M.; Howard, N.I.; Robinson, R.L.; Abell, C.; Blumberger, J.; Mitchell,
J.B. Hierarchical virtual screening for the discovery of new molecular scaffolds in
antibacterial hit identification. *J R Soc Interface* **2012**, *9*, 3196-3207.

166. Kumar, A.; Siddiqi, M.I.; Miertus, S. New molecular scaffolds for the design of *Mycobacterium*
tuberculosis type ii dehydroquinase inhibitors identified using ligand and receptor based
virtual screening. *J Mol Model* **2010**, *16*, 693-712.

167. Prazeres, V.F.; Sánchez-Sixto, C.; Castedo, L.; Lamb, H.; Hawkins, A., R.; Riboldi-Tunnicliffe,
A.; Coggins, J.R.; Lapthorn, A.J.; González-Bello, C. Nanomolar competitive inhibitors of
Mycobacterium tuberculosis and *Streptomyces coelicolor* type ii dehydroquinase. *ChemMedChem*
2007, *2*, 194-207.

168. Siggaard-Andersen, M.; Wissenbach, M.; Chuck, J.A.; Svendsen, I.; Olsen, J.G.; von Wettstein-
Knowles, P. The fabj-encoded beta-ketoacyl-[acyl carrier protein] synthase iv from *Escherichia*
coli is sensitive to cerulenin and specific for short-chain substrates. *Proc Natl Acad Sci U S A.*
1994, *91*, 11027-11031.

169. Magnuson, K.; Carey, M.R.; Cronan, J.E.J. The putative fabJ gene of *escherichia coli* fatty acid
synthesis is the fabF gene. *J Bacteriol* **1995**, *177*, 3593-3595.

170. Tong, L. Structure and function of biotin-dependent carboxylases. *Cell Mol Life Sci* **2013**, *70*, 863-891.

171. Silvers, M.A.; Robertson, G.T.; Taylor, C.M.; Waldrop, G.L. Design, synthesis, and antibacterial properties of dual-ligand inhibitors of acetyl-coa carboxylase. *J Med Chem* **2014**, *57*, 8947-8959.

172. Reddy, M.C.; Breda, A.; Bruning, J.B.; Sherekar, M.; Valluru, S.; Thurman, C.; Ehrenfeld, H.; Sacchettini, J.C. Structure, activity, and inhibition of the carboxyltransferase β-subunit of acetyl coenzyme a carboxylase (accd6) from *Mycobacterium tuberculosis. Antimicrob Agents Chemother* **2014**, *58*, 6122-6132.

173. Levert, K.L.; Waldrop, G.L. A bisubstrate analog inhibitor of the carboxyltransferase component of acetyl-coa carboxylase. *Biochem Biophys Res Commun* **2002**, *291*, 1213-1217.

174. Freiberg, C.; Brunner, N.A.; Schiffer, G.; Lampe, T.; Pohlmann, J.; Brands, M.; Raabe, M.; Häbich, D.; Ziegelbauer, K. Identification and characterization of the first class of potent bacterial acetyl-coA carboxylase inhibitors with antibacterial activity. *J Biol Chem.* **2004**, *279*, 26066-26073.

175. Tong, L. Acetyl-coenzyme a carboxylase: Crucial metabolic enzyme and attractive target for drug discovery. *Cell Mol Life Sci* **2005**, *62*, 1784-1803.

176. Zhang, R.; Ou, H.Y.; Zhang, C.T. Deg: A database of essential genes. *Nucleic Acids Res* **2004**, *32(Database issue)*, D271-272.

177. Moule, M.G.; Hemsley, C.M.; Seet, Q.; Guerra-Assunção, J.A.; Lim, J.; Sarkar-Tyson, M.; Clark, T.G.; Tan, P.B.; Titball, R.W.; Cuccui, J., *et al.* Genome-wide saturation mutagenesis of *Burkholderia pseudomallei* k96243 predicts essential genes and novel targets for antimicrobial development. *MBio* **2014**, *5*, e00926-00913.

178. Somvanshi, V.S.; Viswanathan, P.; Jacobs, J.L.; Mulks, M.H.; Sundin, G.W.; Ciche, T.A. The type 2 secretion pseudopilin, gspj, is required for multihost pathogenicity of *Burkholderia cenocepacia* au1054. *Infect Immun* **2010**, *78*, 4110-4121.

179. van Waasbergen, L.G.; Kidambi, S.P.; Miller, R.V. Construction of a reca mutant of *Burkholderia* (formerly pseudomonas), *cepacia. Appl Microbiol Biotechnol* **1998**, *49*, 59-65.

180. Appleberg, R. Macrophage nutriprive antimicrobial mechanisms. *J Leukocyte Biol* **2006**, *70*, 1117-1128.

181. McKinney, J.D.; Bentrup, K.H.; Muñoz-Elias, E.J.; Miczak, A.; Chen, B.; Chan, W.T.; Swenson, D.; Sacchettini, J.C.; Jacobs Jr., W.R.; Russell, D.G. Persistence of *Mycobacterium tuberculosis* in macrophages and mice requires the glyoxylate shunt enzyme isocitrate lyase. *Nature* **2000**, *406*, 735–738.

182. Fang, F.C.; Libby, S.J.; Castor, M.E.; Fung, A.M. Isocitrate lyase (aceA) is required for salmonella persistence but not for acute lethal infection in mice. *Infect. Immun.* **2005**, *73*, 2547–2549.

183. Liu, N.; Cummings, J.E.; England, K.; Slayden, R.A.; Tonge, P.J. Mechanism and inhibition of the fabi enoyl-acp reductase from *Burkholderia pseudomallei. J Antimicrob Chemother* **2011**, *66*, 564-573.

184. Bokman, A.H.; Levine, H.B.; Lusby, M. Glucose catabolism in *Malleomyces pseudomallei. J Bacteriol* **1957**, *73*, 649-654.

185. Kouassi, Y.; Shelef, L.A. Metabolic activities of *Listeria monocytogenes* in the presence of
sodium propionate, acetate, lactate and citrate. *J Appl Bacteriol* **1996**, *81*, 147-153.

186. Premaratne, R.J.; Lin, W.J.; Johnson, E.A. Development of an improved chemically defined
minimal medium for *Listeria monocytogenes*. *Appl Environ Microbiol.* **1991**, *57*, 3046-3048.

187. Joseph, B.; Przybilla, K.; Stühler, C.; Schauer, K.; Slaghuis, J.; Fuchs, T.M.; Goebel, W.
Identification of *Listeria monocytogenes* genes contributing to intracellular replication by
expression profiling and mutant screening. *J Bacteriol.* **2006**, *188*, 556-568.

188. Bordbar, A.; Lewis, N.E.; Schellenberger, J.; Palsson, B.Ø.; Jamshidi, N. Insight into human
alveolar macrophage and *M. tuberculosis* interactions via metabolic reconstructions. *Mol Syst*
Biol **2010**, *6*, 422.

189. Chatterjee, S.S.; Hossain, H.; Otten, S.; Kuenne, C.; Kuchmina, K.; Machata, S.; Domann, E.;
Chakraborty, T.; Hain, T. Intracellular gene expression profile of *Listeria monocytogenes*. *Infect*
Immun **2006**, *74*, 1323-1338.

190. Eisenreich, W.; Heesemann, J.; Rudel, T.; Goebel, W. Metabolic host responses to infection by
intracellular bacterial pathogens. *Front Cell Infect Microbiol* **2013**, *3*, 24.

191. Caspi, R.; Altman, T.; Dale, J.M.; Dreher, K.; Fulcher, C.A.; Gilham, F.; Kaipa, P.; Karthikeyan,
A.S.; Kothari, A.; Krummenacker, M., *et al.* The metacyc database of metabolic pathways and
enzymes and the biocyc collection of pathway/genome databases. *Nucleic Acids Res.* **2010**,
38(Database issue), D473-479.

192. Joseph, B.; Przybilla, K.; Stühler, C.; Schauer, K.; Slaghuis, J.; Fuchs, T.M.; Goebel, W.
Identification of *Listeria monocytogenes* genes contributing to intracellular replication by
expression profiling and mutant screening. *J Bacteriol.* **2006**, 556-568.

193. Tuanyok, A.; Tom, M.; Dunbar, J.; Woods, D.E. Genome-wide expression analysis of
Burkholderia pseudomallei infection in a hamster model of acute melioidosis. *Infect Immun* **2006**,
74, 5465-5476.

194. Hamad, M.A.; Austin, C.R.; Stewart, A.L.; Higgins, M.; Vázquez-Torres, A.; Voskuil, M.I.
Adaptation and antibiotic tolerance of anaerobic *Burkholderia pseudomallei*. *Antimicrob Agents*
Chemother **2011**, *55*, 3313-3323.

195. Alberts, B.; Johnson, A.; Lewis, J.; Raff, M.; Roberts, K.; Walter, P. *Molecular biology of the cell,*
4th edition. Garland Science: New York, 2002.

196. Myint, K.T.; Uehara, T.; Aoshima, K.; Oda, Y. Polar anionic metabolome analysis by nano-
lc/ms with a metal chelating agent. *Anal Chem* **2009**, *81*, 7766-7772.

197. Lee, I.J.; Hom, K.; Bai, G.; Shapiro, M. Nmr metabolomic analysis of caco-2 cell differentiation.
J Proteome Res **2009**, *8*, 4104-4108.

1131
1132