COGNITIVE ENDOPHENOTYPES OF MODERN AND EXTINCT

HOMININS ASSOCIATED WITH *NTNG* GENE PARALOGS

Pavel Prosselkov[1,2]*, Ryota Hashimoto[3,4], Denis Polygalov[5], Kazutaka Ohi[3,4], Qi Zhang[1],

Thomas J. McHugh[5], Masatoshi Takeda[3,4] and Shigeyoshi Itohara[1]*

[1]Laboratory of Behavioral Genetics, RIKEN Brain Science Institute, Wakoshi, 351-0198

Saitama, Japan

[2]Graduate School, Department of Veterinary Medicine, Faculty of Agriculture, Tokyo

University, 113-8657 Tokyo, Japan

[3]Department of Psychiatry, Osaka University Graduate School of Medicine, Osaka, Japan

[4]Molecular Research Center for Children's Mental Development, United Graduate School of

Child Development, Osaka University, Kanazawa University and Hamamatsu University

School of Medicine, Osaka, Japan

[5]Laboratory of Circuit and Behavior Physiology, RIKEN Brain Science Institute, 351-0198

Saitama, Japan

*corresponding authors:

prosselkov@brain.riken.jp

sitohara@brain.riken.jp

Keywords: evolution, gene duplication, psychogenetics, cognitive endophenotype,

paleogenetics

ABSTRACT

A pair of gene paralogs, *NTNG1* and *NTNG2*, contributes to the Intellectual Quotient

(IQ) test scores in a complementary manner. Single nucleotide polymorphisms (SNPs)

of *NTNG1* are associated with attenuated verbal comprehension (VC) or processing

speed (PS) while *NTNG2* SNPs affect working memory (WM), and perceptual

organization (PO) forming cognitive endophenotypes in healthy and schizophrenia

(SCZ)-affected human subjects. Regions of interest (ROIs), defined as 21 nucleotide

(nu) long loci embedding the IQ-affecting mutation alleles, underwent dramatic

evolutionary changes from mice through primates to hominin gene orthologs, and

resulted in accelerated evolution of VC and WM/PO affecting ROIs. Mutation alleles

associated with the higher VC and WM IQ scores are found in genomes of extinct

hominins of Neolithic times, however, lower WM scores associated allele is detectable in

Mesolithic hunters genomes. Protein sequence of *NTNG1* is 100% conserved among the

archaic and modern extinct hominins while *NTNG2* underwent a recent selection sweep

encoding a primate-specific S371A/V (~50,000 yrs BC), and a modern human (5,300 yrs

BC) T346A substitutions. Hereby, we show that a 500 mln yrs old genomic duplication

of a synapse primordial gene of an urochordate provided a substrate for further

synapse elaborations and its ultimate capacitive expansion of what evolved into a

vertebrate cognitive superior complexity – intelligence.

INTRODUCTION

In 1970 S.Ohno ingeniously proposed that new gene function can result from a gene duplication and following it gene paralogs SF (**Ohno, 1970**). Two gene paralogs, *NTNG1* and *NTNG2*, expressed predominantly in the brain (**Nakashiba et al., 2002**), and encoding Netrin-G1 and Netrin-G2 proteins, respectively, are localised pre-synaptically and segregate in a non-overlapping manner into distinct neuronal circuits (**Nishimura-Akiyoshi et al., 2007; Matsukawa et al., 2014**). They are related to the netrin family of axonal guidance cues (**Sun et al., 2011**) but differ in that they attach to the axonal membrane via a glycosylphosphatidylinositol (GPI) link (**Yu et al., 2013**; **Sevcsik et al., 2015**), a known lipid raft-associated membrane signaling cascade organiser (**Klotzsch and Schutz, 2013; Yu et al., 2013**). The first evolutionary precursor of *NTNG* as a single gene copy can be located in the genome of a primitive vertebrate tunicate/urochordate *Ciona intestinalis* (sea squirt, ENSCING00000024925), reported to be the first organism with the neural crest primordials (**Abitua et al., 2012**), multipotent brain progenitor cells (**Stolfi et al., 2015**), and neurogenic placodes, facilitating the transition from pelagic invertebrate life style to a predatory vertebrate (**Abitua et al., 2015**). The dramatic expansion of human cerebral cortex over the course of evolution (**Wise, 2008**; **Preuss, 2012**; **Geschwind and Rakic, 2013**; **Belmonte et al., 2015**) had provided new niches for accommodating either *de novo* or advancing pre-existed cognitive features and culminating in the positively selected human cognitive functions (**Joshi et al., 2015**).

IQ tests are a surrogate measure of general human cognitive ability characterising intelligence. They are often administered as WAIS-III/IV (**Wechsler, 1958**) and represent a cumulative score of 4 cognitive indices: VC, WM, PO and PS frequently referred as "cognitive domains" (**Deary et al., 2006**). IQ has been validated by factor analyses (**Glascher et al., 2009**), and a common factor (correlate) influencing each of them frequently

referred as *g*, or "general intelligence", proposed by Spearman in 1904 (**Spearman, 1904**),

and recently challenged as the only existing correlate (**Hampshire et al., 2012**). IQ is

affected by several mental disorders including schizophrenia (SCZ) characterized by severe

cognitive deficits in WM and behavioral flexibility and resulting in low performance in

several cognitive tests (**Forbes et al., 2009; Leeson et al., 2009**; **Barnett et al., 2010**). Since

both *NTNG* paralogs have been reported associated with SCZ (**Fukasawa et al., 2004**; **Aoki-**

Suzuki et al., 2005; **JSSLG et al., 2005**; **Eastwood and Harrison, 2008**; **Ohtsuki et al.,**

2008; **Zakharyan et al., 2011**; **Zhu et al., 2011**; **Ayalew et al., 2012**; **Wilcox and Quadri,**

2014) we investigated whether these gene paralogs contribute to human intelligence by

assessing the IQ of human carriers for the *NTNG1* and *NTNG2* SNP alleles against non-

carriers with and without SCZ.

RESULTS

Several *NTNG1* and *NTNG2* SNPs have been previously reported to be associated with SCZ

(e.g. **Aoki-Suzuki et al., 2005**; **Ohtsuki et al., 2008**}. We have found that out of 11 SNPs

tested, five affect the IQ scores and composite domains in human subjects (**Figure 1**,

Supplementary Table 1 (**ST1**)). SCZ patients carrying rs2218404 T allele (**Figure 1A-1**) of

NTNG1 (T/G and T/T genotypes, N = 25 patients) compared with G/G genotype (N = 36

patients) demonstrated attenuated full-scale IQ (FIQ, ANCOVA $p = 0.0057$ (F = 7.80)), and

VIQ ($p = 0.0033$ (F = 8.87)). VC domain score was the main contributor to the VIQ decline

($p = 0.0050$ (F = 8.08), **Figure 1B-1**), with low scores across all parameters except

comprehension (CH): vocabulary (Vc, $p = 0.020$ (F = 5.49)), similarities (SiM, $p = 0.041$ (F

= 4.23)), and information (IF, $p = 0.0067$ (F = 7.50), **Figure 1B-1** lower panel). Thus, a point

mutation in *NTNG1*, rs2218404, is associated with low VC affecting the VIQ via the

attenuated Vc, SiM and IF subscores. The next *NTNG1* SNP found to affect IQ was rs96501,

with attenuated PS in healthy human subjects C allele carriers (N = 45) vs T/T genotypes (N

= 98, $p = 0.028$ (F = 4.89), **Figure 1B-1**) with no effect on SCZ patients. The contributing

affecting score was symbol search (SS, $p = 0.053$ (F = 3.79), **Figure 1B-1**, lower panel) with

digit symbol coding (DSC) being also attenuated but non-significantly ($p = 0.12$ (F = 2.40)).

Three other SNPs mapped to *NTNG2* (**Figure 1A-2**) have been also shown to affect IQ.

Healthy carriers of the *NTNG2* SNP rs1105684 A allele (N = 49) showed a lower FIQ ($p =$

0.018 (F = 5.70)), VIQ ($p = 0.029$ (F = 4.90)), and PIQ ($p = 0.048$ (F = 3.99)) when

compared with the T/T genotypes (N = 96, **Figure 1B-2**). To check for a potential dosage-

dependent effect of a mutation allele on IQ score, *NTNG2* SNP rs2149171 SCZ and healthy

human subject cohorts were each split on 3 genotypes, respectively: C/C (N = 14 and 39),

C/T (N = 29 and 73), and T/T (N = 15 and 30). The presence of the C allele as a single copy

(C/T genotype) was strongly associated with a prominent attenuation in the IQ scores of SCZ

patients and was essentially identical to that produced by the C/C genotype when both are

compared to the T allele carriers (FIQ: $p = 0.014$ (F = 4.35); VIQ: $p = 0.029$ (F = 3.60); PIQ:

$p = 0.035$ (F = 3.42), **Figure 1B-2**). If in case of *NTNG1* located SNP rs2218404 the lower

VIQ score was contributed mainly by the decreased VC domain scores for Vc, SiM, and IF

(**Figure 1B-1**), in the case of *NTNG2* located rs2149171 the CH and WM domain scores

were responsible for the VIQ decline in C allele carriers ($p = 0.012$ (F = 4.54) for CH and $p =$

0.040 (F = 3.27) for WM (N = 12 (C/C); N = 25 (C/T) and N = 14 (T/T)). Similarly to VIQ,

where CH of rs2149171 complements the cognitive endophenotype produced by the T-allele

of rs2218404, the PIQ attenuated score in the case of rs2149171 was due to lower PO score

($p = 0.050$ (F = 3.04), **Figure 1B-2**) in the C allele carriers. The third *NTNG2* located SNP

found to affect the human IQ was rs2274855 (**Figure 1A-2**) with a cognitive endophenotype

associated with the A-allele presence in SCZ patients (N = 33) vs G/G genotypes (N = 26)

and resembling that of the described above C-allele of rs2149171. Accordingly, the

attenuated scores were: FIQ ($p = 0.012$ (F = 6.44)), VIQ ($p = 0.018$ (F = 5.70)), and PIQ ($p =$

0.036 (F = 4.46), **Figure 1B-2**). Similarly to rs2149171 the lower VIQ score was due to

declined CH ($p = 0.035$ (F = 4.49)) but unaffected VC that is contrary (complementary) to the

rs2218404 endophenotype (**Figure 1B-1**). WM was robustly affected by the A-allele

presence (N = 29; $p = 0.023$ (F = 5.28)) contributed by the low DS score ($p = 0.026$ (F =

5.04)) with LNS and AM being unaffected (**Figure 1B-2**, lower panel). The observed PO

score was comprised by the declined matrix reasoning (MR, $p = 0.038$ (F = 4.34)), block

design (BD, $p = 0.041$ (F = 4.23)) with picture completion being unchanged (**Figure 1B-2**,

lower panel). Thus, all three aforementioned *NTNG2* SNPs affect the VIQ and PIQ in human

subjects with the first one contributed by the CH subscore and WM and the latter by the PO

score (**Figure 1C**). Contrary to this, the *NTNG1* located SNP (rs2218404) affects the VIQ

through the lower Vc, SiM and IF scores and affecting the VC domain scores. Another SNP,

rs96501, affects PS domain, though in healthy subjects only. It can be concluded that both

genes (as paralogs) contribute to the cognitive scoring produced upon the implemented IQ

testing but in a cognitive domain-complementary manner, *NTNG1* is responsible for the VC

and PS in human while *NTNG2* for the WM and PO domain scores (**Figure 1C**).

The robust link observed between a single SNP and affected cognitive domain IQ

score (**Figure 1**) can be explained by some global dramatic perturbations caused by the

presence of a mutated allele and/or a functinal importance of its context-dependent

positioning on the gene (**Figure 2A** and **3A**). To determine a potential significance of the

SNP alleles' epistatic environment we compared the nucleotide (nu) sequence within the

immediate vicinity of a SNP allele positioning (50 nu upstream and downstream) in mice,

primates and modern human. We compared all 11 SNPs used for the IQ screening and plotted

the identity percent as a function of distance from the mutated allele position (**Figure 2B** and

3B, see **Supplementary Materials = SM**). We found that the identity percent distribution

over the analysed areas of $\pm$50 nu is not uniform and displays a SNP allele position-centred

dramatic evolutionary changes pointing towards a potential functional significance of the

immediate vicinity of a SNP as short as $\pm$10 nu and not further, referred from here and

beyond as a Region Of Interest (ROI) for each specific SNP allele. We calculated the rates of

evolutionary changes for each ROI as a percent identity change over the lapsed mln yrs of

evolution (**Figure 2C and 3C**). Among the 6 *NTNG1*-located SNP ROIs three of them

display accelerated rates of evolution from marmoset to chimpanzee (rs2218404, rs628117,

rs96501) when compared to the mouse-marmoset rates, and rs2218404 (affecting VC in

human subjects) additionally demonstrates an accelerated rate of evolution on the

chimpanzee to human path (**Figure 2C**). As for the *NTNG2* located ROIs (**Figure 3-B**),

rs1105684 is remarkably consistent at displaying high evolutionary rates around 0.8 and,

together with rs2274855, both have identical rates at the mouse-marmoset and chimpanzee-

human paths, but differ dramatically at the marmoset-chimpanzee point (0.8 vs 0,

respectively). rs2274855 is the only *NTNG2*-nested SNP ROI which underwent an AE from

chimpanzee to human. Next we compared the DNA sequences of all 11 ROIs across mice,

primates and extinct hominins *NTNG* gene paralogs (see **ST2** for the datasets sources used

for the genes reconstruction). T-allele of rs2218404 is detectable in marmoset and in mouse

its position corresponds to adenosine (**Figure 2D**). G-alelle (associated with a higher VC

score comparing to the human T-allele carriers) is found in Mesolithic hunter Loschbour

(8,000 BC) but not in another ancient hunter Motala12 and is also present in other two

hominins belonging to the Neolithic period, Iceman and Eskimos (5,300 and 4,000 yrs BC,

respectively). rs628117 is the only *NTNG1*-related mutation near vicinity of which ($\pm$50 nu)

is located an intra-hominins (Es, Ice, Lo) mutation (**Figure 2B**, low left). The next, PS-

affecting, ROI of rs96501 displays an intricate path of T-allele evolution (associated with a

higher PS score) being anciently conserved from mice to primates but later substituted on the

less efficient (in terms of the generated IQ scores) C allele in Neanderthals, later again

replaced by the T allele in Mesolithic hunters and coming back during the Neolithic times

(**Figure 2D**). The first *NTNG2* SNP rs1105684 is located at the beginning of the gene and

affects WM in healthy human subjects (**Figure 3A**). The origin of the T-allele is evolutionary

bound to marmoset since its position in mice is occupied by another pyrimidine base C

(**Figure 3D**). Next on the gene are two SNP alleles for rs7851893 and rs3824574 which do

not affect IQ and similarly to rs1105684 are surrounded by highly conserved ROIs not only

in hominins and chimpansee (100% identity) but also in marmoset (except 1 mutation for

rs7851893). rs2149171 ROI (affecting the WM and PO scores) similarly to rs3824574 is

100% conserved across the all species (including the mouse) except the allele itself. The

attenuating IQ C-allele position is occupied in mice genome by T but present in Iceman and

Eskimos genes. A distinct evolutionary path is taken by another cognitive endophenotype-

associated and affecting WM and PO scores A-allele of rs2274855 and its ROI (**Figure 3D**).

Its position in mice is likely to be occupied by the C pyrimidine base (the software places a

blank instead of it) which is gradually substituted on purine G in chimpanzee and misplaced

by the lower IQ score-associated A-allele in Mesolithic hominins. And 20 nu downstream of

the centre of ROI is located a modern human-specific point mutation translated into the

T346A protein substitution (as described below).

The distinct picture of evolutionary changes among the *NTNG1* and *NTNG2* nested

SNPs has prompted us to compare evolutionary rates for the full-length proteins encoded by

these gene paralogs, Netrin-G1 and Netrin-G2, respectively (**Nakashiba et al., 2000 and**

2002). Netrin-G1 undergoes only few changes in its amino acid (aa) composition with the

maximum calculated rate of evolution reaching 0.05 when mice and marmoset proteins are

compared, 0.01 among the primates, and 0.03 between chimpanzee and human due to a

single point mutation A81S (**Figure 2E** and **SM:** Netrin-G1), absent in other primates.

Netrin-G2 evolves 2.8 times faster between mouse and marmoset than its paralog Netrin-G1

and continues evolving with a steady rate of 0.05 from primates to human (**Figure 3E** and

SM: Netrin-G2). We have also reconstructed both proteins from ancient (Neanderthals,

Paleolithic time) and extinct hominins (Mesolithic and Neolithic times) and compared them

with primates' and mice' Netrin-G orthologs (**Figure 2F** and **3F**). Netrin-G1 is a highly

conserved protein among the primates and hominins (**Figure 3F**). As for Netrin-G2, a

mutation shared among the Neanderthals' and Mesolithic genomes, primates and mice

(T346A) is absent in the Neolithic Iceman and modern human (the signal for Motala3,

Motala1 and MezmayskayaNea is not clear due to low sequence coverage). Primates

(marmoset and chimpanzee) share another mutation (S371A/V) preserved in mouse and

absent in hominins (further details can be found in the **SM:** Results).

DISCUSSION

NTNG **paralog SNPs and associated cognitive endophenotypes of human subjects.**

Shortcomings of cognitive and information processing are key features of SCZ diagnosis

(**APA, 2013**). They are manifested as impairments in PO, WM, VC and PS (see **Yoon et al.,**

2014 for references) frequently reported as attenuated scores upon IQ tests implementation.

SCZ patients carrying a mutation allele for one of *NTNG* gene paralog SNPs form cognitive

endophenotypes affecting the IQ scores (**Figure 1C**). The given endophenotypic groups

comprise from subjects with either affected VIQ (via attenuated VC by *NTNG1* rs2218404 or

WM by *NTNG2* rs2149171 and rs2274855) or affected PIQ (via attenuated PO by *NTNG2*

rs2149171 and rs2274855). In two extra cases PS is affected by rs96501 of *NTNG1* and WM

by rs1105684 of *NTNG2* (**Figure 1B-1** and **B-2**, respectively) but in healthy human subjects.

Such intriguing non-overlapping effect on the IQ domains prompts us to conclude that *NTNG*

paralogs complement each other function and represent an example of how a synapse-

expressed genes affect the human cognitive abilities, perhaps through the precision of

neuronal connectivity perturbations and concomitant miswirings. The observed phenomena

of the affected WM is the most striking due to its multifaceted constructive nature (**Frydecka**

et al., 2014) underlying many, if not all, cognitive tasks such as comprehension, reasoning

and learning (**Baddeley, 1992**) and historically introduced by Baddeley as the reading span

test (**Mackintosh, 2011**). Lack of the localization effect of *NTNG2* SNP mutation alleles, all

three are located in different parts of the gene (**Figure 1A-2**) but associated with identical

endophenotype (**Figure 1B-2**), points to a uniform nature of the *NTNG2* function distribution

over the entire gene. An obviously non-coding nature of all five IQ-affecting alleles

(rs2149171 despite being exon 4-located encodes a silent F246F mutation) corroborates an

idea that anthropoid trait-associated loci lie outside coding protein areas (**del Rosario et al.,**

2014; Kellis et al., 2014) and hints towards a potential of these alleles to perturb genes

regulatory functions, e.g. mRNA splicing, affecting downstream located pivotally functional

NTNG elements such as Ukd-domain encoding exons 6 and 7 or a unique Netrin-Gs trait –

GPI-link. Alternatively, or simultaneously, the *NTNG* SNP alleles may be embedded into an

epistatic network of other genes influencing human cognitive traits (**Hemani et al., 2014**).

However since it is usual for a SNP effect to be estimated using an additive model (assuming

either independent and cumulative single contribution) to the mean of a trait with the small

effect size the power to detect the epistatic environment drastically declines. Contrary to the

genetic associations with gene expression having large effect sizes (**Hemani et al., 2014**),

cognitive trait-associated effect sizes are reportedly small (**Plomin and Deary, 2015**), e.g.

the largest effect sizes of the variance of intelligence scores accounted for only 0.2%

(**Benyamin et al., 2014**), 0.5% on GWA studies of 1,583 adolescence (**Desrivieres et al.,**

2015) or was predicted to be ~1% on 3,511 adults (**Davies et al., 2011**). Another GWAS of

educational attainment (sharing a moderate correlate with intelligence), which included

126,559 individuals, reports on just 1% of the variance but only 0.02% in a replication

sample (**Rietveld et al., 2013**). Our data support the preexisted conclusion that human

cognitive traits modalities are not described by statistically large effect sizes.

Evolutionary elaborations of the embedding IQ-affecting mutations loci. Eleven

previously published SCZ-associated SNPs were tested for their effect on IQ performance of

human subjects and 5 of them were found to be associated with attenuated IQ cognitive

endophenotypes (**Figure 1B-1** and **B-2**). ROIs of 3 of them (rs2218404, rs1373336 and

rs2274855) underwent an AE from chimpanzee to human when compared to New World

monkeys to apes (marmoset-chimpanzee) path (**Figure 2C** and 3**C**). Two of them affect IQ in

humans (rs2218404 – VC and rs2274855 – WM, **Figure 1B-1** and **B-2**) while being located

within the vicinity of exon 5 (2,275 nu downstream and 15 nu upstream, respectively) – a

part of the lowest percent identity coding DNA among the *NTNG* gene paralogs (**Prosselkov**

et al., in press). Presence of the evolutionary accelerated regions within the *NTNG* genes

non-coding areas underscores them as contributors to the human-specific traits along with

other genes (**Prabhakar et al., 2006**). However, not only ROIs of the IQ-affecting alleles but

the alleles themselves demonstrate several unique evolutionary features (see **SM:** Discussion).

To understand evolutionary forces driving the emergence of cognitive endophenotype-

associated alleles we have deduced a set of rules outlined as follows. 1. An alternative

(mutated) allele evolutionary appearance coincides with the lack of any other mutations

within ROI (a conserved island rule); 2. positioning of the future mutation often represents a

turning point of dramatic changes of an allele ROI (e.g. as seen in marmoset: rs2218404 (50-

90%), rs628117 (30-80%), rs96501 (100-40%)); 3. an AE of ROI often precedes the

emergence of a mutation allele (e.g. rs2218404: chimpanzee to human ($k = 1.59$), appearance

of "G" in Loschbour; rs628117: marmoset to chimpanzee ($k = 1.10$), appearance of "T" in

AltaiNea; rs96501: marmoset to chimpanzee ($k = 0.83$), appearance of "C" in AltaiNea;

rs2274855: chimpanzee to human ($k = 0.79$), appearance of "A" in Motala12; 4. low identity

percent (equivalent to subsequent substantial evolutionary changes) among the evolutionary

species within the allele surrounding proximity of as long as $\pm$50 nu is not sufficient for the

future mutated allele significance as a cognitive endophenotype determinant (as deduced by

the IQ score) as seen for the rs1373336, rs1444042, rs4915045 and rs7851893 (none of them

are IQ-affecting, though associated with SCZ, despite showing (very) low identity in mice).

Rather some dramatic changes within the allele's immediate proximity of $\pm$10 nu (defined as

a ROI) preceded or followed by more stringently conserved DNA are necessary (the

conserved island perturbation rule). Currently we are unable to state that the IQ-associated

alleles ROIs are regulatory loci and an important source of evolutionary innovation

(**Rubinstein and de Souza, 2013**) but they may be the smallest functional blocks of a strong

positive selection exerts its action upon similarly to the 20–30 nu clusters of strongly

conserved non-coding elements (CNEs), transcription factor binding sites (TFBS), RNA

splicing and editing motifs (**Harmston et al., 2013**).

Extinct hominins and IQ-associated mutation alleles. Availability of archaic genomes

allows excavation for the advantageous alleles that modern humans acquired from archaic

extinct hominins such as Neandertals and Denisovans who used to live 230,000-30,000 years

ago (Middle/Upper Paleolithic, Old Stone Age) defined by distinct morphological features

(**Meyer et al., 2012**), and from modern extinct humans (hunters, farmers) from Mesolithic

(Middle Stone Age, ~10,000 yrs BC, **Lazaridis et al., 2014**) and Neolithic (New Stone Age,

~5,000 BC, **Keller et al., 2012**; **Rasmussen et al., 2010**) periods. Though exhibiting several

anatomical features, making archaic hominins different from the modern human, there are

studies challenging the idea that reserve symbolism and abstract thinking was an exclusive

prerogative of modern human (**Appenzeller, 2013**; **Wong, 2015**). The time Neanderthals

used to live in is thought to be associated with the onset of cognitive fluidity involving the

capacity to draw analogies (early paintings), to combine concepts (making tools) and to adapt

ideas for new contexts (**Gabora and Russon, 2011**). Wynn and Coolidge believe that

evolution of WM was central to the evolution of human cognitive traits consisted from few

genetic mutations that led to "enhanced WM" 200,000-40,000 BC (**Balter, 2010**). Our work

partially supports this idea showing the perseverance of higher WM score-associated alleles

across the hominins such as T of rs1105684 and G of rs2274855 (**Figure 3D**) but a Neolithic

appearance of rs2149171 T in Iceman previously found only in mice genome supporting the

conclusion made by **Crabtree (2013**) that modern humans as species "are surprisingly

intellectually fragile and perhaps reached a peak 2,000–6,000 years ago". Mesolithic period

has been always considered as a key gate for the evolution of human languages (**Haak et al.,**

2015) with our data showing that rs2218404 G-allele associated with a higher VC score

(**Figure 1B-1**) emerges for the first time in the Loschbour hunter *NTNG1* gene (**Figure 3D**).

VC as a part of abstract symbol usage is associated with the global network efficiency (as a

part of fronto-parietal network in **Song et al., 2008**; **Glascher et al., 2009**), and global

communication and intellectual performance (**Pamplona et al., 2015**). From this point of

view it is not surprising that appearance of the G allele in Loschbour and Iceman genomes

coincides with the presence of PS enhancing rs96501 T allele (**Figure 3D**). A wealth of data

has been collected characterising possible look and health status of archaic hominins and

modern but extinct humans (for ref. see **Sarkissian et al., 2015** and **SM:** Discussion). Based

on our own data we may also speculate that the extinct hominins may have had a lower VC

comparing to us, and consequently, Neanderthals were unlikely able for a semantic

communication due to a global network inefficiency VC is associated with; they have had

similar to us PS (if they had lived beyond the Mesolithic period), and were likely to have had

identical to modern human WM, corroborating the advanced evolutionary nature of this

important human cognitive domain of a limited capacity and associated with intelligence.

CONCLUSION

Evolution of a novel function relies on enhanced genetic robustness through functional

redundancy potentially provided by a gene duplication event. Further evolutionary outcome

depends on the substrate availability (undergoing its own evolution) upon which the novel

function(s) exerts its action. Nature does not create but tinkers to perfection provided to it

material exploring available evolutionary tools. Half a billion years ago a gene duplication

event had provided a plethora of such substrate thus converting the evolution itself into a

"Creator" of new functions. Here we have described how a pair of twin genes got themselves

involved into the human cognitive functioning believed to be emerged in a primordial state in

primitive vertebrates prior to the first recorded gene duplication. Subsequent process of the

function specialisation made *NTNG* paralogs to subfunctionalise into distinct cognitive

domains in a complementary manner (**Prosselkov et al., in press**).

MATERIALS AND METHODS

Ethics statement. This study was performed in accordance with the World Medical Association's Declaration of Helsinki and approved by the Osaka University Research Ethics Committee. A written informed consent was obtained from all subjects after the procedures had been fully explained.

Subjects. The procedures were performed as per established protocols at Osaka University as described previously (**Ohi et al., 2012**). The subjects consisted from 339 patients with SCZ and 716 healthy controls. The sex ratio did not differ significantly between the groups, but the mean age was significantly different. The subjects were all biologically unrelated Japanese and recruited from both outpatient and inpatient units at Osaka University Hospital and other psychiatric hospitals. Each patient with SCZ had been diagnosed by at least two trained psychiatrists based on unstructured clinical interviews, according to the criteria of the DSM-IV (**APA, 2013**). In case if the diagnosis of the two trained psychiatrists was discordant, it was resolved through the further negotiations on both specialist opinions. In case of unresolved diagnostic disputes, the patient was omitted from the study. Psychiatrically healthy controls were recruited through local advertisements and were evaluated by means of unstructured interviews to exclude individuals with current or past contact with psychiatric services, those who experienced psychiatric medications, or who were not Japanese. Controls for family history of a CD, such as SCZ, BD, or major depressive disorder were not included. Ethnicity was determined by self-report and was not confirmed by genetic analyses. Additionally, subjects were excluded from this study if they had neurologic or medical conditions that could have potentially affect their central nervous system, such as atypical headaches, head trauma with loss of consciousness, chronic lung disease, kidney disease, chronic hepatic disease, thyroid disease, active cancer, cerebrovascular disease, epilepsy, seizures, substance abuse related disorders, or mental retardation.

SNPs selection, genotyping, and genomic sequencing. This study was designed to examine

the association of SCZ patients cognitive performance (through WAIS-III implementation,

Wechsler, 1958) with *NTNG* genes. Venous blood was collected from the subjects. Genomic

DNA was extracted from the whole blood using standard procedures. The SNPs (**Fukasawa**

et al., 2004; **Aoki-Suzuki et al., 2005**; **JSSLG et al., 2005**; **Eastwood and Harrison, 2008**;

Ohtsuki et al., 2008; **Zakharyan et al., 2011**; **Zhu et al., 2011**; **Ayalew et al., 2012**; **Wilcox**

and Quadri, 2014) were genotyped using the TaqMan allelic discrimination assay (Applied

Biosystems, Foster City, CA). No deviations from the Hardy-Weinberg equilibrium in the

examined SNPs were detected ($p > 0.05$).

Statistical analysis. The effects of the diagnosis, genotype and their interaction on cognitive

performances in the WAIS were analyzed by two-way analyses of covariance (ANCOVA).

Diagnosis and genotype statuses were included in the model as independent variables (**ST1**).

FIQ and each WAIS subscale score (VIQ, PIQ, VC, PO, WM, PS, Vc, SiM, IF, CH, AM, DS,

LNS, PC, BD, and MR) were included as dependent variables. Sex, age and years of

education were treated as covariates, as they were possible confounding factors. All p values

are two tailed, and statistical significance was defined as $*p < 0.05$ and $**p < 0.01$.

Identity percent calculations and the definition of ROIs. The complete procedure is

described in the **Figure 4** legend. Stretcher (**McWilliam et al., 2013**) was used for the

alignments (the default values were: gap penalty – 16 (DNA) and 12 (protein), and the extend

penalty – 4 (DNA) and 2 (protein)), for the percent identity calculations and evolutionary

rates. A ROI was selected as a minimal area surrounding a SNP mutation allele incorporating

the outmost evolutionary dramatic changes.

Mice, primates and hominins *NTNG* paralogs DNA and encoded aa sequences

reconstruction. Genomes for mouse (GRC38.p3) and marmoset (C_jacchus3.2.1) were from

Ensemble. Since chimpanzee's genome is based only on a single individual (CHIMP2.1.4,

Clint) and contains several questionable information we have reconstructed a consensus

genome sequence for both *NTNG* genes based on 25 primate sequences of *Pan troglodytes*

(**Prado-Martinez et al., 2013**). All datasets used for the *NTNG* paralogs DNA and encoded

by them proteins reconstruction are listed in **ST2**. For details refer to **SM**.

SUPPLEMENTARY MATERIALS (**SM**)

Contain additional Results and Discussion, Supplementary Methods (ancient and primate

genomes reconstructions), and Supplementary Tables (**ST1** and **ST2**) as a single compiled

pdf file. Also included are human Netrin-G1 and Netrin-G2 alignments, as well as 101 nu

alignments for all 11 ROIs across the all analysed species.

ACKNOWLEDGEMENTS

Authors would like to acknowledge the financial support provided by Funding Program for

World-Leading Innovative R&D on Science and Technology (FIRST Programme) and

KAKENHI 15H04290 by the Japan Society for the Promotion of Science (JSPS).

COMPETING INTERESTS

Authors would like to express a lack of any competing interests associated with the work.

REFERENCES

Abitua PB, Wagner E, Navarrete IA, Levine M. 2012. Identification of a rudimentary neural

crest in a non-vertebrate chordate. *Nature* **492:**104-107. doi: 10.1038/nature11589.

Abitua PB, Gainous TB, Kaczmarczyk AN, Winchell CJ, Hudson C, Kamata K, Nakagawa

M, Tsuda M, Kusakabe, TG, Levine M. 2015. The pre-vertebrate origins of neurogenic

placodes. *Nature* **524:**462-465. doi: 10.1038/nature14657.

Aoki-Suzuki M, Yamada K, Meerabux J, Iwayama-Shigeno Y, Ohba H, Iwamoto K, Takao

H, Toyota T, Suto Y, Nakatani N, Dean B, Nishimura S, Seki K, Kato T, Itohara S,

Nishikawa T, Yoshikawa, T. 2005. A Family-Based Association Study and Gene

Expression Analyses of Netrin-G1 and -G2 Genes in Schizophrenia. *Biological Psychiatry*

57:382–393. doi: 10.1016/j.biopsych.2004.11.022.

APA (American Psychiatric Association). 2013. Diagnostic and Statistical Manual of Mental

Disorders (DSM-V®), Fifth Edition. 991 pp. doi: 10.1176/appi.books.9780890425596.

Appenzeller T. 2013. Old masters. *Nature* **497:** 302-304.

Ayalew M, Le-Niculescu H, Levey DF, Jain N, Changala B, Patel SD, Winiger E, Breier A,

Shekhar A, Amdur R, Koller D, Nurnberger JI, Corvin A, Geyer M, Tsuang MT, Salomon

D, Schork NJ, Fanous AH, O'Donovan MC, Niculescu AB. 2012. Convergent functional

genomics of schizophrenia: from comprehensive understanding to genetic risk prediction.

Molecular Psychiatry **17:**887-905. doi: 10.1038/mp.2012.37.

Baddeley A. 1992. Working memory. *Science* **255:**556-559. doi: 10.1126/science.1736359.

Balter M. 2010. Did Working Memory Spark Creative Culture? *Science* **328:**160-163. doi:

10.1126/science.328.5975.160.

Barnett JH, Robbins TW, Leeson VC, Sahakian BJ, Joyce EM, Blackwell AD. 2010.

Assessing cognitive function in clinical trials of schizophrenia. *Neuroscience and*

Biobehavioral Reviews **34:**1161-1177. doi: 10.1016/j.neubiorev.2010.01.012.

Belmonte JCI, Callaway EM, Churchland P, Caddick SJ, Feng G, Homanics GE, Lee K-F,

425   Leopold DA, Miller CT, Mitchell JF, Mitalipov S, Moutri AR, Movshon JA, Okano H,

426   Reynolds JH, Ringach D, Sejnowski TJ, Silva AC, Strick PL, Wu J, Zhang F. 2015.

427   Brains, Genes, and Primates. *Neuron* **86:**617-631. doi: 10.1016/j.neuron.2015.03.021.

428   **Benyamin B**, Pourcain BSt, Davis OS, Davies G, Hansell NK, Brion M-JA, Kirkpatrick RM,

429   Cents RAM, Franic S, Miller MB, Haworth CMA, Meaburn E, Price TS, Evans DM,

430   Timpson N, Kemp J, Ring S, McArdle W, Medland SE, Yang J, Harris SE, Liewald DC,

431   Scheet P, Xiao X, Hudziak JJ, de Geus EJC, Wellcome Trust Case Control Consortium 2

432   (WTCCC2), Jaddoe VWV, Starr JM, Verhulst FC, Pennell C, Tiemeier H, Iacono WG,

433   Palmer LJ, Montgomery GW, Martin NG, Boomsma DI, Posthuma D, McGue M, Wright

434   MJ, Davey Smith G, Deary IJ, Plomin R, Visscher PM. 2014. Childhood intelligence is

435   heritable, highly polygenic and associated with *FNBP1L. Molecular Psychiatry* **19:**253-

436   258. doi: 10.1038/mp.2012.184.

437   **Crabtree GR.** 2013. Our fragile intellect. Part I. *Trends in Genetics* **29:**1-3. doi:

438   10.1016/j.tig.2012.10.002.

439   **Davies G,** Tenesa A, Payton A, Yang J, Harris SE, Liewald D, Ke X, Hellard SLe,

440   Christoforou A, Luciano M, McGhee K, Lopez L, Gow AJ, Corley J, Redmond P, Fox HC,

441   Haggarty P, Whalley LJ, McNeill G, Goddard ME, Espeseth T, Lundervold AJ, Reinvang

442   I, Pickles A, Steen VM, Ollier W, Porteous DJ, Horan M, Starr JM, Pendleton N, Visscher

443   PM, Deary IJ. 2011. Genome-wide association studies establish that human intelligence is

444   highly   heritable   and   polygenic.   *Molecular   Psychiatry*   **16:**996-1005.   doi:

445   10.1038/mp.2011.85.

446   **Deary IJ,** Spinath FM, Bates TC. 2006. Genetics of intelligence. *European Journal of*

447   *Human Genetics* **14:**690-700. doi: 10.1038/sj.ejhg.5201588.

del Rosario RCH, Rayan NA, Prabhakar S. 2014. Noncoding Origins of Anthropoid Traits

and a New Null Model of Transposon Functionalization. *Genome Research* **24:**1469-1484.

doi: 10.1101/gr.168963.113.

Eastwood SL, Harrison PJ. 2008. Decreased mRNA expression of netrin-G1 and netrin-G2

in the temporal lobe in schizophrenia and bipolar disorder. *Neuropsychopharmacology*

33:933-945. doi: 10.1038/sj.npp.1301457.

Forbes NF, Carrick LA, McIntosh AM, Lawrie SM. 2009. Working memory in

schizophrenia: a meta-analysis. *Psychological Medicine* **39:**889-905. doi:

10.1017/s0033291708004558.

Frydecka D, Eissa AM, Hewedi DH, Ali M, Drapała J, Misiak B, Kłosińska E, Phillips JR,

Moustafa AA. 2014. Impairments of working memory in schizophrenia and bipolar

disorder: the effect of history of psychotic symptoms and different aspects of cognitive

task demands. *Frontiers in Behavioral Neuroscience* **8**:A416. doi:

10.3389/fnbeh.2014.00416.

Fukasawa M, Aoki M, Yamada K, Iwayama-Shigeno Y, Takao H, Meerabux J, Toyota T,

Nishikawa T, Yoshikawa T. 2004. Case-control association study of human netrin G1

gene in Japanese schizophrenia. *Journal of Medical and Dental Sciences* **51:** 121-128.

Gabora L and Russon A. 2011. "The evolution of Intelligence". *In* "The Cambridge

Handbook of Intelligence". Edited by Sternberg RJ, Kaufman SB. p. 328-350.

Geschwind DH and Rakic P. 2013. Cortical Evolution: Judge the Brain by Its Cover.

Neuron **80:**633-647. doi: 10.1016/j.neuron.2013.10.045.

Glascher J, Tranel D, Paul LK, Rudrauf D, Rorden C, Hornaday A, Grabowski T, Damasio

H, Adolphs R. 2009. Lesion Mapping of Cognitive Abilities Linked to Intelligence.

Neuron **61:**681-691. doi: 10.1016/j.neuron.2009.01.026.

Green RE, Krause J, Briggs AW, Maricic T, Stenzel U, Kircher M, Patterson N, Li H, Zhai

W, Fritz MH-Y, Hansen NF, Durand EY, Malaspinas A-S, Jensen JD, Marques-Bonet T,

Alkan C, Prüfer K, Meyer M, Burbano HA, Good JM, Schultz R, Aximu-Petri A, Butthof

A, Höber B, Höffner B, Siegemund M, Weihmann A, Nusbaum C, Lander ES, Russ C,

Novod N, Affourtit J, Egholm M, Verna C, Rudan P, Brajkovic D, Kucan Ž, Gušic I,

Doronichev VB, Golovanova LV, Lalueza-Fox C, de la Rasilla M, Fortea J, Rosas A,

Schmitz RW, Johnson PLF, Eichler EE, Falush D, Birney E, Mullikin JC, Slatkin M,

Nielsen R, Kelso J, Lachmann M, Reich D, Pääbo S. 2010. A Draft Sequence of the

Neandertal Genome. *Science* **328:**710-722. doi: 10.1126/science.1188021.

Haak W, Lazardis I, Patterson N, Rohland N, Mallick S, Llamas B, Brandt G, Nordenfelt S,

Harney E, Stewardson K, Fu Q, Mittnik A, Bánffy E, Economou C, Francken M,

Friederich S, Pena RG, Hallgren F, Khartanovich V, Khokhlov A, Kunst M, Kuznetsov P,

Meller H, Mochalov O, Moiseyev V, Nicklisch N, Pichler SL, Risch R, Guerra MAR,

Roth C, Szécsényi-Nagy A, Wahl J, Meyer M, Krause J, Brown D, Anthony D, Cooper A,

Alt KW, Reich D. 2015. Massive migration from the steppe was a source for Indo-

European languages in Europe. *Nature* **522:**207-211. doi: 10.1038/nature14317.

Hampshire A, Highfield RR, Parkin BL, Owen AM. 2012. Fractionating Human Intelligence.

Neuron **76:**1225-1237. doi: 10.1016/j.neuron.2012.06.022.

Harmston N, Baresic A, Lenhard B. 2013. The mystery of extreme non-coding conservation.

Philosophical Transactions of the Royal Society B-Biological Sciences **368**:A20130021.

doi: 10.1098/rstb.2013.0021.

Hemani G, Shakhbazov K, Westra H-J, Esko T, Henders AK, McRae AF, Yang J, Gibson G,

Martin NG, Metspalu A, Franke L, Montgomery GW, Visscher PM, Powell JE. 2014.

Detection and replication of epistasis influencing transcription in humans. *Nature*

508:249–253. doi: 10.1038/nature13005.

Joshi PK, Esko T, Mattsson H, Eklund N, Gandin I, Nutile T, Jackson AU, Schurmann C,

Smith AV, Zhang W, Okada Y, Stančáková A, Faul JD, Zhao W, Bartz TM, Concas MP,

Franceschini N, Enroth S, Vitart V, Trompet S, Guo X, Chasman DI, O'Connel JR, Corre

T, Nongmaithem SS, Chen Y, Mangino M, Ruggiero D, Traglia M, Farmaki A-E,

Kacprowski T, Bjonnes A, van der Spek A, Wu Y, Giri AK, Yanek LR, Wang L, Hofer E,

Rietveld CA, McLeod O, Cornelis MC, Pattaro C, Verweij N, Baumbach C, Abdellaoui A,

Warren HR, Vuckovic D, Mei H, Bouchard C, Perry JRB, Cappellani S, Mirza SS, Benton

MC, Broeckel U, Medland SE, Lind PA, Malerba G, Drong A, Yengo L, Bielak LF, Zhi D,

van der Most PJ, Shriner D, Maegi R, Hemani G, Karaderi T, Wang Z, Liu T, Demuth I,

Zhao JH, Meng W, Lataniotis L, van der Laan SW, Bradfield JP, Wood AR, Bonnefond

A, Ahluwalia TS, Hall LM, Salvi E, Yazar S, Carstensen L, de Haan HG, Abney M, Afzal

U, Allison MA, Amin N, Asselbergs FW, Bakker SJL, Barr RG, Baumeister SE, Benjamin

DJ, Bergmann S, Boerwinkle E, Bottinger EP, Campbell A, Chakravarti A, Chan Y,

Chanock SJ, Chen C, Chen Y-DI, Collins FS, Connell J, Correa A, Cupples LA, Smith

GD, Davies G, Dörr M, Ehret G, Ellis SB, Feenstra B, Feitosa MF, Ford I, Fox CS,

Frayling TM, Friedrich N, Geller F, Generation Scotland, Gillham-Nasenya I, Gottesman

O, Graff M, Grodstein F, Gu C, Haley C, Hammond CJ, Harris SE, Harris TB, Hastie ND,

Heard-Costa NL, Heikkilä K, Hocking LJ, Homuth G, Hottenga J-J, Huang J, Huffman JE,

Hysi PG, Ikram MA, Ingelsson E, Joensuu A, Johansson Å, Jousilahti P, Jukema JW,

Kähönen M, Kamatani Y, Kanoni S, Kerr SM, Khan NM, Koellinger P, Koistinen HA,

Kooner MK, Kubo M, Kuusisto J, Lahti J, Launer LJ, Lea RA, Lehne B, Lehtimäki T,

Liewald DCM, Lind L, Loh M, Lokki M-L, London SJ, Loomis SJ, Loukola A, Lu Y,

Lumley T, Lundqvist A, Männistö S, Marques-Vidal P, Masciullo C, Matchan A, Mathias

RA, Matsuda K, Meigs JB, Meisinger C, Meitinger T, Menni C, Mentch FD, Mihailov E,

Milani L, Montasser ME, Montgomery GW, Morrison A, Myers RH, Nadukuru R,

Navarro P, Nelis M, Nieminen MS, Nolte IM, O'Connor GT, Ogunniyi A, Padmanabhan S,

Palmas WR, Pankow JS, Patarcic I, Pavani F, Peyser PA, Pietilainen K, Poulter N,

Prokopenko I, Ralhan S, Redmond P, Rich SS, Rissanen H, Robino A, Rose LM, Rose R,

Sala C, Salako B, Salomaa V, Sarin A-P, Saxena R, Schmidt H, Scott LJ, Scott WR,

Sennblad B, Seshadri S, Sever P, Shrestha S, Smith BH, Smith JA, Soranzo N,

Sotoodehnia N, Southam L, Stanton AV, Stathopoulou MG, Strauch K, Strawbridge RJ,

Suderman MJ, Tandon N, Tang S-T, Taylor KD, Tayo BO, Töglhofer AM, Tomaszewski

M, Tšernikova N, Tuomilehto J, Uitterlinden AG, Vaidya D, van Hylckama Vlieg A, van

Setten J, Vasankari T, Vedantam S, Vlachopoulou E, Vozzi D, Vuoksimaa E,

Waldenberger M, Ware EB, Wentworth-Shields W, Whitfield JB, Wild S, Willemsen G,

Yajnik CS, Yao J, Zaza G, Zhu X, The BioBank Japan Project, Salem RM, Melbye M,

Bisgaard H, Samani NJ, Cusi D, Mackey DA, Cooper RS, Froguel P, Pasterkamp G, Grant

SFA, Hakonarson H, Ferrucci L, Scott RA, Morris AD, Palmer CAN, Dedoussis G,

Deloukas P, Bertram L, Lindenberger U, Berndt SI, Lindgren CM, Timpson NJ, Toenjes

A. Munroe PB, Sørensen TIA, Rotimi CN, Arnett DK, Oldehinkel AJ, Kardia SLR,

Balkau B, Gambaro G, Morris AP, Eriksson JG, Wright MJ, Martin NG, Hunt SC, Starr

JM, Deary IJ, Griffiths LR, Tiemeier H, Pirastu N, Kaprio J, Wareham NJ, Pérusse L,

Wilson JG, Girotto G, Caulfield MJ, Raitakari O, Boomsma DI, Gieger C, van der Harst

Pim, Hicks AA, Kraft P, Sinisalo J, Knekt P, Johannesson M, Magnusson PKE, Hamsten

A, Schmidt R, Borecki IB, Vartiainen E, Becker DM, Bharadwaj D, Mohlke KL, Boehnke

M, van Duijn CM, Sanghera DK, Teumer A, Zeggini E, Metspalu A, Gasparini P, Ulivi S,

Ober C, Toniolo D, Rudan I, Porteous DJ, Ciullo M, Spector TD, Hayward C, Dupuis J,

Loos RJF, Wright AF, Chandak GR, Vollenweider P, Shuldiner AR, Ridker PM, Rotter JI,

Sattar N, Gyllensten U, North KE, Pirastu M, Psaty BM, Weir DR, Laakso M, Gudnason

V, Takahashi A, Chambers JC, Kooner JS, Strachan DP, Campbell H, Hirschhorn JN,

Perola M, Polašek O, Wilson JF. 2015. Directional dominance on stature and cognition in

diverse human populations. *Nature.* **523:**459-462. doi: 10.1038/nature14618.

Keller A, Graefen A, Ball M, Matzas M, Boisguerin V, Maixner F, Leidinger P, Backes C,

Khairat R, Forster M, Stade B, Franke A, Mayer J, Spangler J, McLaughlin S, Shah M,

Lee C, Harkins TT, Sartori A, Moreno-Estrada A, Henn B, Sikora M, Semino O, Chiaroni

J, Rootsi S, Myres NM, Cabrera VM, Underhill PA, Bustamante CD, Vigl EE, Samadelli

M, Cipollini G, Haas J, Katus H, O'Connor BD, Carlson MRJ, Meder B, Blin N, Meese E,

Pusch CM, Zink A. 2012. New insights into the Tyrolean Iceman's origin and phenotype

as inferred by whole-genome sequencing. *Nature Communications* **3:**A698. doi:

10.1038/ncomms1701.

Kellis M, Wold B, Snyder MP, Bernstein BE, Kundaje A, Marinov GK, Ward LD, Birney E,

Crawford GE, Dekker J, Dunham I, Elnitski LL, Farnham PJ, Feingold EA, Gerstein M,

Giddings MC, Gilbert DM, Gingeras TR, Green ED, Guigo R, Hubbard T, Kent J, Lieb JD,

Myers RM, Pazin MJ, Ren B, Stamatoyannopoulos JA, Weng Z, White KP, Hardison RC.

2014. Defining functional DNA elements in the human genome. *Proceedings of the*

National Academy of Sciences of the United States of America **111:**6131-6138. doi:

10.1073/pnas.1318948111.

Klotzsch E and Schutz GJ. 2013. A critical survey of methods to detect plasma membrane

rafts. *Philosophical Transactions of the Royal Society B-Biological Sciences*

368:20120033. doi: 10.1098/rstb.2012.0033.

Lazaridis I, Patterson N, Mittnik A, Renaud G, Mallick S, Kirsanow K, Sudmant PH,

Schraiber JG, Castellano S, Lipson M, Berger B, Economou C, Bollongino R, Fu Q, Bos

KI, Nordenfelt S, Li H, de Filippo C, Prüfer K, Sawyer S, Posth C, Haak W, Hallgren F,

Fornander E, Rohland N, Delsate D, Francken M, Guinet J-M, Wahl J, Ayodo G, Babiker

HA, Bailliet G, Balanovska E, Balanovsky O, Barrantes R, Bedoya G, Ben-Ami H, Bene J,

Berrada F, Bravi CM, Brisighelli F, Busby GBJ, Cali F, Churnosov M, Cole David EC,

Corach D, Damba L, van Driem G, Dryomov S, Dugoujon J-M, Fedorova SA, Romero IG,

Gubina M, Hammer M, Henn BM, Hervig T, Hodoglugil U, Jha AR, Karachanak-

Yankova S, Khusainova R, Khusnutdinova E, Kittles R, Kivisild T, Klitz W, Kučinskas V,

Kushniarevich A, Laredj L, Litvinov S, Loukidis T, Mahley RW, Melegh B, Metspalu E,

Molina J, Mountain J, Näkkäläjärvi K, Nesheva D, Nyambo T, Osipova L, Parik J,

Platonov F, Posukh O, Romano V, Rothhammer F, Rudan I, Ruizbakiev R, Sahakyan H,

Sajantila A, Salas A, Starikovskaya EB, Tarekegn A, Toncheva D, Turdikulova S,

Uktveryte I, Utevska O, Vasquez R, Villena M, Voevoda M, Winkler CA, Yepiskoposyan

L, Zalloua P, Zemunik T, Cooper A, Capelli C, Thomas MG, Ruiz-Linares A, Tishkoff

SA, Singh L, Thangaraj K, Villems R, Comas D, Sukernik R, Metspalu M, Meyer M,

Eichler EE, Burger J, Slatkin M, Pääbo S, Kelso J, Reich D, Krause J. 2014. Ancient

human genomes suggest three ancestral populations for present-day Europeans. *Nature*

513:409-413. doi: 10.1038/nature13673.

Leeson VC, Robbins TW, Matheson E, Hutton SB, Ron MA, Barnes TRE, Joyce EM. 2009.

Discrimination Learning, Reversal, and Set-Shifting in First-Episode Schizophrenia:

Stability Over Six Years and Specific Associations with Medication Type and

Disorganization Syndrome. *Biological Psychiatry* **66:**586-593. doi:

10.1016/j.biopsych.2009.05.016.

Lips ES, Cornelisse LN, Toonen RF, Min JL, Hultman CM, the International Schizophrenia

Consortium, Holmans PA, O'Donovan MC, Purcell SM, Smit AB, Verhage M, Sullivan

PF, Visscher PM, Posthuma D. 2012. Functional gene group analysis identifies synaptic

gene groups as risk factor for schizophrenia. *Molecular Psychiatry* **17:**996-1006. doi:

10.1038/mp.2012.37.

Mackintosh NJ. 2011. "History of theories and measurement of Intelligence". *In* "The

Cambridge Handbook of Intelligence". Edited by Sternberg RJ, Kaufman SB. p. 3-19. doi:

10.1017/CBO9780511977244.

Matsukawa H, Akiyoshi-Nishimura S, Zhang Q, Lujan R, Yamaguchi K, Goto H, Yaguchi

K, Hashikawa T, Sano C, Shigemoto R, Nakashiba T, Itohara S. 2014. Netrin-G/NGL

Complexes Encode Functional Synaptic Diversification. *Journal of Neuroscience*

34:15779-15792. doi: 10.1523/jneurosci.1141-14.2014.

McWilliam H, Li W, Uludag M, Squizzato S, Park YM, Buso N, Cowley AP, Lopez R. 2013.

Analysis Tool Web Services from the EMBL-EBI. *Nucleic Acids Research* **41:**597-600.

doi: 10.1093/nar/gkt376.

Meyer M, Kircher M, Gansauge MT, Li H, Racimo F, Mallick S, Schraiber JG, Jay F, Prüfer

K, de Filippo C, Sudmant PH, Alkan C, Fu QM, Do R, Rohland N, Tandon A, Siebauer M,

Green RE, Bryc K, Briggs AW, Stenzel U, Dabney J, Shendure J, Kitzman J, Hammer MF,

Shunkov MV, Derevianko AP, Patterson N, Andrés AM, Eichler EE, Slatkin M, Reich D,

Kelso J, Pääbo S. 2012. A High-Coverage Genome Sequence from an Archaic Denisovan

Individual. *Science* **338:**222-226. doi: 10.1126/science.1224344.

Nakashiba T, Ikeda T, Nishimura S, Tashiro K, Honjo T, Culotti JG, Itohara, S. 2000.

Netrin-G1: a Novel Glycosyl Phosphatidylinositol-Linked Mammalian Netrin That Is

Functionally Divergent from Classical Netrins. *The Journal of Neuroscience* **20:**6540-

6550.

Nakashiba T, Nishimura S, Ikeda T, Itohara S. 2002. Complementary expression and neurite

outgrowth activity of netrin-G subfamily members. *Mechanisms of Development* **111:**47-

60. doi: 10.1016/S0925-4773(01)00600-1.

Nishimura-Akiyoshi S, Niimi K, Nakashiba T, and Itohara S. 2007. Axonal netrin-Gs

transneuronally determine lamina-specific subdendritic segments. *Proceedings of the*

National Academy of Sciences of the United States of America **104:**14801-14806. doi:

10.1073/pnas.0706919104.

Ohi K, Hashimoto R, Nakazawa T, Okada T, Yasuda Y, Yamamori H, Fukumoto M,

624 Umeda-Yano S, Iwase M, Kazui H, Yamamoto T, Kano M, Takeda M. 2012. The

625 p250GAP Gene Is Associated with Risk for Schizophrenia and Schizotypal Personality

626 Traits. PLoS One:e35696. doi: 10.1371/journal.pone.0035696.

627 **Ohno S.** 1970. Evolution by Gene Duplication. Springer, New York.

628 **Ohtsuki T,** Horiuchi Y, Koga M, Ishiguro H, Inada T, Iwata N, Ozaki N, Ujike H, Watanabe

629 Y, Someya T, Arinami T. 2008. Association of polymorphisms in the haplotype block

630 spanning the alternatively spliced exons of the *NTNG1* gene at 1p13.3 with schizophrenia

631 in Japanese populations. *Neuroscience Letters* **435:**194-197. doi:

632 10.1016/j.neulet.2008.02.053.

633 **Pamplona GSP,** Neto GSS, Rosset SRE, Rogers BP, Salmon CEG. 2015. Analyzing the

634 association between functional connectivity of the brain and intellectual performance.

635 *Frontiers in Human Neuroscience* **9:**A61. doi: 10.3389/fnhum.2015.00061.

636 **Plomin R and Deary IJ.** 2015. Genetics and intelligence differences: five special findings.

637 *Molecular Psychiatry* **20:**98-108. doi: 10.1038/mp.2014.105.

638 **Prabhakar S,** Noonan JP, Pääbo S, Rubin EM. 2006. Accelerated evolution of conserved

639 noncoding sequences in humans. *Science* **314:**786-786. doi: 10.1126/science.1130738.

640 **Prado-Martinez J,** Sudmant PH, Kidd JM, Li H, Kelley JL, Lorente-Galdos B, Veeramah

641 KR, Woerner AE, O'Connor TD, Santpere G, Cagan A, Theunert C, Casals F, Laayouni H,

642 Munch K, Hobolth A, Halager AE, Malig M, Hernandez-Rodriguez J, Hernando-Herraez I,

643 Prüfer K, Pybus M, Johnstone L, Lachmann, M, Alkan C, Twigg D, Petit N, Baker C,

644 Hormozdiari F, Fernandez-Callejo M, Dabad M, Wilson ML, Stevison L, Camprubí C,

645 Carvalho T, Ruiz-Herrera A, Vives L, Mele M, Abello T, Kondova I, Bontrop RE, Pusey

646 A, Lankester F, Kiyang JA, Bergl RA, Lonsdorf E, Myers S, Ventura M, Gagneux P,

647 Comas D, Siegismund H, Blanc J, Agueda-Calpena L, Gut M, Fulton L, Tishkoff SA,

648 Mullikin JC, Wilson RK, Gut IG, Gonder MK, Ryder OA, Hahn BH, Navarro A, Akey JM,

Bertranpetit J, Reich D, Mailund T, Schierup MH, Hvilsom C, Andrés AM, Wall JD,

Bustamante CD, Hammer MF, Eichler EE, Marques-Bonet T. 2013. Great ape genetic

diversity and population history. *Nature* **499:**471-475. doi: 10.1038/nature12228.

Preuss TM. 2012. Human brain evolution: From gene discovery to phenotype discovery.

Proceedings of the National Academy of Sciences of the United States of America

109:10709-10716. doi : 10.1073/pnas.1201894109.

Prüfer K, Racimo F, Patterson N, Jay F, Sankararaman S, Sawyer S, Heinze A, Renaud G,

Sudmant PH, de Filippo C, Li H, Mallick S, Dannemann M, Fu Q, Kircher M, Kuhlwilm

M, Lachmann M, Meyer M, Ongyerth M, Siebauer M, Theunert C, Tandon A, Moorjani P,

Pickrell J, Mullikin JC, Vohr SH, Green RE, Hellmann I, Johnson PLF, Blanche H, Cann

H, Kitzman JO, Shendure J, Eichler EE, Lein ES, Bakken TE, Golovanova LV,

Doronichev VB, Shunkov MV, Derevianko AP, Viola B, Slatkin M, Reich D, Kelso J,

Pääbo S. 2013. The complete genome sequence of a Neanderthal from the Altai

Mountains. *Nature* **505:**43–49. doi: 10.1038/nature12886.

Rasmussen M, Li YR, Lindgreen S, Pedersen JS, Albrechtsen A, Moltke I, Metspalu M,

Metspalu E, Kivisild T, Gupta R, Bertalan M, Nielsen K, Gilbert MTP, Wang Y,

Raghavan M, Campos PF, Kamp HM, Wilson AS, Gledhill A, Tridico S, Bunce M,

Lorenzen ED, Binladen J, Guo XS, Zhao J, Zhang XQ, Zhang H, Li Z, Chen MF, Orlando

L, Kristiansen K, Bak M, Tommerup N, Bendixen C, Pierre TL, Grønnow B, Meldgaard

M, Andreasen C, Fedorova SA, Osipova LP, Higham TFG, Ramsey CB, Hansen TVO,

Nielsen FC, Crawford MH, Brunak S, Sicheritz-Pontén T, Villems R, Nielsen R, Krogh A,

Wang J, Willerslev E. 2010. Ancient human genome sequence of an extinct Palaeo-

Eskimo. *Nature* **463:**757-762. doi: 10.1038/nature08835.

Rietveld CA, Medland SE, Derringer J, Yang J, Esko T, Martin NW, Westra HJ,

Shakhbazov K, Abdellaoui A, Agrawal A, Albrecht E, Alizadeh BZ, Amin N, Bamard J,

Baumeister SE, Benke KS, Bielak LF, Boatman JA, Boyle PA, Davies G, de Leeuw C,

Eklund N, Evans DS, Ferhmann R, Fischer K, Gieger C, Gjessing HK, Hägg S, Harris JR,

Hayward C, Holzapfel C, Ibrahim-Verbaas CA, Ingelsson E, Jacobsson B, Joshi PK,

Jugessur A, Kaakinen M, Kanoni S, Karjalainen J, Kolcic I, Kristiansson K, Kutalik Z,

Lahti J, Lee SH, Lin P, Lind PA, Liu YM, Lohman K, Loitfelder M, McMahon G, Vidal

PM, Meirelles O, Milani L, Myhre R, Nuotio ML, Oldmeadow CJ, Petrovic KE, Peyrot

WJ, Polaśek O, Quaye L, Reinmaa E, Rice JP, Rizzi TS, Schmidt H, Schmidt R, Smith

AV, Smith JA, Tanaka T, Terracciano A, van der Loos MJHM, Vitart V, Völzke H,

Wellmann J, Yu L, Zhao W, Allik J, Attia JR, Bandinelli S, Bastardot F, Beauchamp J,

Bennett DA, Berger K, Bierut LJ, Boomsma DI, Bültmann U, Campbell H, Chabris CF,

Cherkas L, Chung MK, Cucca F, de Andrade M, De Jager PL, De Neve J-E, Deary IJ,

Dedoussis GV, Deloukas P, Dimitriou M, Eiríksdóttir G, Elderson MF, Eriksson JG,

Evans DM, Faul JD, Ferrucci L, Garcia ME, Grönberg H, Guonason V, Hall P, Harris JM,

Harris TB, Hastie ND, Heath AC, Hernandez DG, Hoffmann W, Hofman A, Holle R,

Holliday EG, Hottenga JJ, Iacono WG, Illig T, Järvelin MR, Kähönen M, Kaprio J,

Kirkpatrick RM, Kowgier M, Latvala A, Launer LJ, Lawlor DA, Lehtimäki T, Li JM,

Lichtenstein P, Lichtner P, Liewald DC, Madden PA, Magnusson PKE, Mäkinen TE,

Masala M, McGue M, Metspalu A, Mielck A, Miller MB, Montgomery GW, Mukherjee S,

Nyholt DR, Oostra BA, Palmer LJ, Palotie A, Penninx B, Perola M, Peyser PA, Preisig M,

Räikkönen K, Raitakari OT, Realo A, Ring SM, Ripatti S, Rivadeneira F, Rudan I,

Rustichini A, Salomaa V, Sarin AP, Schlessinger D, Scott RJ, Snieder H, St Pourcain B,

Starr JM, Sul JH, Surakka I, Svento R, Teumer A, The LifeLines Cohort Study, Tiemeier

H, van Rooij FJA, Van Wagoner DR, Vartiainen E, Viikari J, Vollenweider P, Vonk JM,

Waeber G, Weir DR, Wichmann H-E, Widen E, Willemsen G, Wilson JF, Wright AF,

Conley D, Davey-Smith G, Franke L, Groenen PJF, Johannesson M, Kardia SLR, Krueger

RF, Laibson D, Martin NG, Meyer MN, Posthuma D, Thurik AR, Timpson NJ,

Uitterlinden AG, van Duijn CM, Visscher PM, Benjamin DJ, Cesarini D, Koellinger PD.

2013. GWAS of 126,559 Individuals Identifies Genetic Variants Associated with

Educational Attainment. *Science* **340:**1467-1471. doi: 10.1126/science.1235488.

Rubinstein M and de Souza FSJ. 2013. Evolution of transcriptional enhancers and animal

diversity. *Philosophical Transactions of the Royal Society B-Biological Sciences*

368:A20130017. doi: 10.1098/rstb.2013.0017.

Sarkissian CD, Allentoft ME, Ávila-Arcos MC, Barnett R, Campos PF, Cappellini E, Ermini

L, Fernández R, da Fonseca R, Ginolhac A, Hansen AJ, Jónsson H, Korneliussen T,

Margaryan A, Martin MD, Moreno-Mayar JV, Raghavan M, Rasmussen M, Velasco MS,

Schroeder H, Schubert M, Seguin-Orlando A, Wales N, Gilbert MTP, Willerslev E,

Orlando L. 2015. Ancient genomics. *Philosophical Transactions of the Royal Society B-*

Biological Sciences **370**:20130387. doi: 10.1098/rstb.2013.0387.

Sevcsik E, Brameshuber M, Foelser M, Weghuber J, Honigmann A, Schuetz GJ. 2015. GPI-

anchored proteins do not reside in ordered domains in the live cell plasma membrane.

Nature Communications **6**:A6969. doi: 10.1038/ncomms7969.

Song M, Zhou Y, Li J, Liu Y, Tian L, Yu C, Tianzi, J. 2008. Brain spontaneous functional

connectivity and intelligence. *Neuroimage* **41:**1168-1176. doi:

10.1016/j.neuroimage.2008.02.036.

Spearman C. 1904. "General intelligence" objectively determined and measured. *American*

Journal of Psychology **15:**201-292. doi: 10.2307/1412107.

Stolfi A, Ryan K, Meinertzhagen IA, Christiaen L. 2015. Migratory neuronal progenitors

arise from the neural plate borders in tunicates. *Nature* **527:**371-374. doi:

10.1038/nature15758.

Sun KLW, Correia JP, Kennedy TE. 2011. Netrins: versatile extracellular cues with diverse

functions. *Development* **138:**2153-2169. doi: 10.1242/dev.044529.

The Japanese Schizophrenia Sib-Pair Linkage Group (JSSLG), Arinami T, Ohtsuki T,

Ishiguro H, Ujike H, Tanaka Y, Morita Y, Mineta M, Takeichi M, Yamada S, Imamura A,

Ohara K, Shibuya H, Ohara, K, Suzuki Y, Muratake T, Kaneko N, Someya T, Inada T,

Yoshikawa T, Toyota T, Yamada K, Kojima T, Takahashi S, Osamu O, Shinkai T,

Nakamura M, Fukuzako H, Hashiguchi T, Niwa S, Ueno T, Tachikawa H, Hori T, Asada T,

Nanko S, Kunugi H, Hashimoto R, Ozaki N, Iwata N, Harano M, Arai H, Ohnuma T,

Kusumi I, Koyama T, Yoneda H, Fukumaki Y, Shibata H, Kaneko S, Higuchi H, Yasui-

Furukori N, Numachi Y, Itokawa M, Okazaki Y. 2005. Genomewide high-density SNP

linkage analysis of 236 Japanese families supports the existence of schizophrenia

susceptibility loci on chromosomes 1p, 14q, and 20p. *American Journal of Human*

Genetics **77:**937-944. doi: 10.1086/498122.

Wechsler D. 1958. *The Measurement And Appraisal Of Adult Intelligence*, 4th edition. The

Williams & Wilkins Company. 324 pp.

Wilcox JA and Quadri S. 2014. Replication of NTNG1 association in schizophrenia.

Psychiatric Genetics **24:**266-268. doi: 10.1097/ypg.0000000000000061.

Wise SP. 2008. Forward frontal fields: phylogeny and fundamental function. *Trends in*

Neurosciences **31:**599-608. doi: 10.1016/j.tins.2008.08.008.

Wong K. 2015. Neandertal Minds. *Scientific American* **312:**36-43. doi:

10.1038/scientificamerican0215-36.

Wu JQ, Wang X, Beveridge NJ, Tooney PA, Scott RJ, Carr VJ, Cairns MJ. 2012.

Transcriptome Sequencing Revealed Significant Alteration of Cortical Promoter Usage

and Splicing in Schizophrenia. *PLoS One* **7:**e36351. doi: 10.1371/journal.pone.0036351.

Yoon JH, Sheremata SL, Rokem A, Silver MA. 2014. Windows to the soul: vision science as

a tool for studying biological mechanisms of information processing deficits in

schizophrenia. *Frontiers in Psychology* **4**:A681. doi: 10.3389/fpsyg.2014.00681.

Yu S, Guo Z, Johnson C, Gu G, Wu Q. 2013. Recent progress in synthetic and biological

studies of GPI anchors and GPI-anchored proteins. *Current Opinion in Chemical Biology*

17: 1006-1013. doi: 10.1016/j.cbpa.2013.09.016.

Zakharyan R, Boyajyan A, Arakelyan A, Gevorgyan A, Mrazek F, Petrek M. 2011.

Functional variants of the genes involved in neurodevelopment and susceptibility to

schizophrenia in an Armenian population. *Human Immunology* **72:**746-748. doi:

10.1016/j.humimm.2011.05.018.

Zhu Y, Yang H, Bi Y, Zhang Y, Zhen C, Xie S, Qin H, He J, Liu L, Liu Y. 2011. Positive

association between *NTNG1* and schizophrenia in Chinese Han population. *Journal of*

Genetics **90:**499-502.

Figure 1. Complementary effect of *NTNG* paralog SNPs on IQ cognitive domains of human subjects as measured by WAIS-III. (A-1, A-2) *NTNG1* and *NTNG2* gene structures with the SNPs location indicated. (B-1, B-2, C) Affected cognitive domains and scores. Red highlights *NTNG1* and green (or blue, in case of heterozygosity) – *NTNG2* located alleles associated with the attenuated IQ scores. The data are presented as a mean±SEM. *$p<0.05$,**$p<0.01$ (two ways ANCOVA (sex, education, and age as co-variates)). The number of human subjects is indicated as N. For statistical and diagnosis details refer to **ST1.**

Figure 2.

Figure 3.

Figures 2 and 3. Accelerated evolution (AE) and definition of a ROI of *NTNG* loci embedding associated with the cognitive endophenotype alleles. Hominins and primate-specific protein amino acid substitutions. (**A**) *NTNG1* and *NTNG2* SNPs' gene locations. Alleles associated with the lower IQ scores are shown either in red (*NTNG1*) or green (*NTNG2*). (**B**) Calculated identity percent of primates and mice to human gene loci as a function of distance from the position of a mutated allele (denoted as a star). Comparison was done in a stepwise manner as +10 nu to the maximum of +50 nu using Stretcher (http://www.ebi.ac.uk/Tools/psa, McWilliam et al., 2013). The areas were compared based on the positioning relative to the point mutation without any manual curation for "the best-fit" alignment, but as per the algorithm output only. Due to low level of identity for mouse and marmoset the initial search of the mutation allele positioning was done aligning against the human corresponding full-length intron, and then second time against the obtained 101 nu query ("-50 nu-SNP+50 nu"). Two dashed lines define an area of -10 nu to +10 nu from the mutation allele position. This area (21 nu in total) is defined as a ROI of the given mutation allele of a representing SNP. An arrow indicates a position of extinct hominins-specific mutations (see below). (**C**) Evolutionary rates for the ROIs calculated as a percent identity change relative to the hg19 over the mln of years of evolution. The spectrum color reflects the mutations' positioning order on a gene, as purple-yellow for *NTNG1*, and yellow-blue for *NTNG2*. (**D**) ROIs DNA sequences across hominins, primates and mice loci. The extinct and ancient hominin's *NTNG* paralogs were reconstructed from the available datasets (see **SM**): Saqqaq-Eskimos (Es: **Rasmussen et al., 2010**); Iceman (Ice, **Keller et al., 2012**); Hunters (Loschbour (Lo), Motala12 (Mo12), Motala3 (Mo3), Motala1: **Lazaridis et al., 2014**); VindijaNea (Vi: **Green et al., 2010**); Denisovan (De), AltaiNea (Al) and MezmaiskayaNea (Me): **Prüfer et al., 2013**; chimpanzee (Chi: reconstructed from **Prado-Martinez et al., 2013**, n=25 animals). Non-available sequences due to poor reads quality are denoted as X. Ensemble was used for the initial *NTNG* paralogs retrieval in marmoset (Ma) and mouse (Mo). Yellow (vertical strip) denotes the position of the SNP-related mutation allele and non-matched to human substitutions. (**E**) Evolution rates for the proteins encoded by the *NTNG1* (Netrin-G1) and *NTNG2* (Netrin-G2). (**F**) Amino acid changes for Netrin-Gs across hominins, primates and mice. For Netrin-G1, there are no common mutations among primates, hominins and modern human. Es contains 5 point mutations absent in other hominins; Chi contains 1 mutation outside the depicted area (A81S); all 4 mutations for marmoset are shown, and 10 more extra mutations for mouse are not shown. For Netrin-G2, all hominins except Ice (relative to hg19) carry a T346A mutation (exon 5-located and known as rs4962173), also detectable in primates and mouse. Chi's Netrin-G2 differs from all hominins by S371A mutation (exon 6 - nested) and present in marmoset as S371V. Non-matched to hg19 amino acids are highlighted as pink. Refer to **SM** for the full alignments.