COGNITIVE ENDOPHENOTYPES OF MODERN AND EXTINCT HOMININS ASSOCIATED WITH *NTNG* GENE PARALOGS

Pavel Prosselkov[1,2]*, Ryota Hashimoto[3,4], Denis Polygalov[5], Kazutaka Ohi[3,4], Qi Zhang[1],

Thomas J. McHugh[5], Masatoshi Takeda[3,4] and Shigeyoshi Itohara[1]*

[1]Laboratory of Behavioral Genetics, RIKEN Brain Science Institute, Wakoshi, 351-0198

Saitama, Japan

[2]Graduate School, Department of Veterinary Medicine, Faculty of Agriculture, Tokyo

University, 113-8657 Tokyo, Japan

[3]Department of Psychiatry, Osaka University Graduate School of Medicine, Osaka, Japan

[4]Molecular Research Center for Children's Mental Development, United Graduate School of

Child Development, Osaka University, Kanazawa University and Hamamatsu University

School of Medicine, Osaka, Japan

[5]Laboratory of Circuit and Behavior Physiology, RIKEN Brain Science Institute, 351-0198

Saitama, Japan

*corresponding authors:

    prosselkov@brain.riken.jp

    sitohara@brain.riken.jp

Keywords: evolution, gene duplication, psychogenetics, cognitive endophenotype,

    paleogenetics

ABSTRACT

A pair of vertebrate-specific and brain-expressed pre-synaptic genes, *NTNG1* and *NTNG2*,

contributes to the Intellectual Quotient (IQ) test scores in a complementary manner. Single

nucleotide polymorphisms (SNPs) of *NTNG1* are associated with attenuated verbal

comprehension (VC) or processing speed (PS) while *NTNG2* SNPs affect working memory

(WM) and perceptual organization (PO), forming cognitive endophenotypes in healthy and

schizophrenia (SCZ)-affected human subjects. Regions of interest (ROIs), defined as 21

nucleotide (nu) long *NTNG* gene loci symmetrically embedding the IQ-affecting mutation

alleles (VC/PS and WM/PO), underwent dramatic evolutionary changes from mice through

primates to hominins, at the accelerated rates. Mutation alleles associated with the higher VC

and WM IQ scores are found in the genomes of extinct hominins of Neolithic times, however,

lower WM scores associated allele is also detectable in Mesolithic hunters genomes. Protein

sequence of *NTNG1* is 100% conserved among the primates, archaic and modern extinct

hominins while *NTNG2* underwent a recent selection sweep encoding a primate-specific

S371A/V (~50,000 yrs BC), and a modern human (5,300 yrs BC) T346A substitutions. We

show that a 500 mln yrs old genomic duplication of a synapse primordial gene provided a

substrate for further synapse elaborations and its ultimate capacitive expansion of what

evolved into a vertebrate cognitive superior complexity – intelligence.

S.Ohno ingeniously proposed that new gene function can result from a gene duplication and

following it gene paralogs SF (1). Two gene paralogs, *NTNG1* and *NTNG2*, expressed

predominantly in the brain (2), and encoding Netrin-G1 and Netrin-G2 proteins, respectively,

are expressed pre-synaptically and segregate in a non-overlapping manner into distinct

neuronal circuits (3, 4). They are related to the netrin family of axonal guidance cues (5) but

differ in that they attach to the axonal membrane via a glycosylphosphatidylinositol (GPI)

link (6, 7), a known lipid raft-associated membrane signaling cascade organiser (6, 8). The

first evolutionary precursor of *NTNG* as a single gene copy can be located in the genome of a

primitive vertebrate tunicate/urochordate *Ciona intestinalis* (sea squirt,

ENSCING00000024925), reported to be the first organism with the neural crest primordials

(9), multipotent brain progenitor cells (10), and neurogenic placodes, facilitating the

transition from pelagic invertebrate life style to a predatory vertebrate (11). The dramatic

expansion of human cerebral cortex over the course of evolution (12-15) had provided new

niches for accommodating either *de novo* or advancing pre-existed cognitive features and

culminating in the positively selected human cognitive functions (16).

IQ tests are a surrogate measure of general human cognitive ability characterising

intelligence. They are often administered as WAIS-III/IV (17) and represent a cumulative

score of 4 cognitive indices: VC, WM, PO and PS frequently referred as "cognitive domains"

(18). IQ has been validated by factor analyses (19), and a common factor (correlate)

influencing each of them frequently referred as *g*, or "general intelligence", proposed by

Spearman (20), but recently challenged as the only existing correlate (21). IQ is affected by

several mental disorders including schizophrenia (SCZ) characterised by severe cognitive

deficits in WM (22), behavioral flexibility and resulting in low performance in cognitive tests

(23-25). Since both *NTNG* paralogs have been reportedly associated with SCZ (26-34) we

investigated whether these gene paralogs contribute to human intelligence by assessing the

IQ of human carriers for the *NTNG1* and *NTNG2* SNP alleles against non-carriers with and

without SCZ.

RESULTS

Several *NTNG1* and *NTNG2* SNPs have been previously reported to be associated with SCZ

(27, 30). We have found that out of 11 SNPs tested, five affect the IQ scores and composite

domains in human subjects (Figure 1, Supplementary Table 1 (ST1)). SCZ patients carrying

rs2218404 T allele (Figure 1A-1) of *NTNG1* (T/G and T/T genotypes, N = 25 patients)

compared with G/G genotype (N = 36 patients) demonstrated attenuated full-scale IQ (FIQ,

ANCOVA $p = 0.0057$ (F = 7.80)), and VIQ ($p = 0.0033$ (F = 8.87)). VC domain score was

the main contributor to the VIQ decline ($p = 0.0050$ (F = 8.08), Figure 1B-1), with low scores

across all parameters except comprehension (CH): vocabulary (Vc, $p = 0.020$ (F = 5.49)),

similarities (SiM, $p = 0.041$ (F = 4.23)), and information (IF, $p = 0.0067$ (F = 7.50), Figure

1B-1 lower panel). Thus, a mutation allele of *NTNG1*, rs2218404, is associated with low VC

and affects the VIQ via the attenuated Vc, SiM and IF subscores. The next *NTNG1* SNP

found to affect IQ was rs96501, with attenuated PS in healthy human subjects C allele

carriers (N = 45) vs T/T genotypes (N = 98, $p = 0.028$ (F = 4.89), Figure 1B-1) with no effect

on SCZ patients. The contributing affecting score was symbol search (SS, $p = 0.053$ (F =

3.79), Figure 1B-1, lower panel) with digit symbol coding (DSC) being also attenuated but

non-significantly ($p = 0.12$ (F = 2.40)). Three other SNPs mapped to *NTNG2* (Figure 1A-2)

have been also shown to affect IQ. Healthy carriers of the *NTNG2* SNP rs1105684 A allele

(N = 49) showed a lower FIQ ($p = 0.018$ (F = 5.70)), VIQ ($p = 0.029$ (F = 4.90)), and PIQ (p

$= 0.048$ (F = 3.99)) when compared with the T/T genotypes (N = 96, Figure 1B-2). To check

for a potential dosage-dependent effect of a mutation allele on IQ score, *NTNG2* SNP

rs2149171 SCZ and healthy human subject cohorts were each split on 3 genotypes,

respectively: C/C (N = 14 and 39), C/T (N = 29 and 73), and T/T (N = 15 and 30). The

presence of the C allele as a single copy (C/T genotype) was strongly associated with a

prominent attenuation in the IQ scores of SCZ patients and was essentially identical to that

produced by the C/C genotype when both are compared to the T allele carriers (FIQ: $p =$

0.014 (F = 4.35); VIQ: p = 0.029 (F = 3.60); PIQ: p = 0.035 (F = 3.42), Figure 1B-2). If in

the case of *NTNG1* located SNP rs2218404 the lower VIQ score was contributed mainly by

the decreased VC domain scores for Vc, SiM, and IF (Figure 1B-1), in the case of *NTNG2*

located rs2149171 the CH and WM domain scores were responsible for the VIQ decline in C

allele carriers (p = 0.012 (F = 4.54) for CH and p = 0.040 (F = 3.27) for WM (N = 12 (C/C);

N = 25 (C/T) and N = 14 (T/T)). Similarly to VIQ, where CH of rs2149171 complements the

cognitive endophenotype produced by the T-allele of rs2218404, the PIQ attenuated score in

the case of rs2149171 was due to lower PO score (p = 0.050 (F = 3.04), Figure 1B-2) in the C

allele carriers. The third *NTNG2* located SNP found to affect the human IQ was rs2274855

(Figure 1A-2) with a cognitive endophenotype associated with the A-allele presence in SCZ

patients (N = 33) vs G/G genotypes (N = 26) and resembling that of the described above C-

allele of rs2149171. Accordingly, the attenuated scores were: FIQ (p = 0.012 (F = 6.44)),

VIQ (p = 0.018 (F = 5.70)), and PIQ (p = 0.036 (F = 4.46), Figure 1B-2). Similarly to

rs2149171 the lower VIQ score was due to declined CH (p = 0.035 (F = 4.49)) but unaffected

VC that is contrary (complementary) to the rs2218404 endophenotype (Figure 1B-1). WM

was robustly affected by the A-allele presence (N = 29; p = 0.023 (F = 5.28)) contributed by

the low DS score (p = 0.026 (F = 5.04)) with LNS and AM being unaffected (Figure 1B-2,

lower panel). The observed PO score was comprised by the declined matrix reasoning (MR, *p*

= 0.038 (F = 4.34)), block design (BD, p = 0.041 (F = 4.23)) with picture completion being

unchanged (Figure 1B-2, lower panel). Thus, all three aforementioned *NTNG2* SNPs affect

the VIQ and PIQ in human subjects with the first one contributed by the CH subscore and

WM and the latter by the PO score (Figure 1C). Contrary to this, the *NTNG1* located SNP

(rs2218404) affects the VIQ through the lower Vc, SiM and IF scores and affecting the VC

domain scores. Another SNP, rs96501, affects PS domain, although in healthy subjects only.

It can be concluded that both genes (as paralogs) contribute to the cognitive scoring produced

upon the implemented IQ testing but in a cognitive domain-complementary manner. *NTNG1*

contributes to the VC and PS while *NTNG2* for the WM and PO domain scores (Figure 1C).

The robust link observed between a single SNP and affected cognitive domain IQ

score (Figure 1) can be explained by some global dramatic perturbations caused by the

presence of a mutated allele and/or a functinal importance of its context-dependent

positioning on the gene (Figure 2A and 3A). To determine a potential significance of the SNP

alleles' epistatic environment we compared the nucleotide (nu) sequence within the

immediate vicinity of a SNP allele positioning (50 nu upstream and downstream) in mice,

primates and modern human. We compared all 11 SNPs used for the IQ screening and plotted

the identity percent as a function of distance from the mutated allele position (Figure 2B and

3B, see Supplementary Materials = SM). We found that the identity percent distribution over

the analysed areas of ±50 nu is not uniform and displays a SNP allele position-centred

dramatic evolutionary changes pointing towards a potential functional significance of the

immediate vicinity of a SNP as short as ±10 nu and not further, referred from here and

beyond as a Region Of Interest (ROI) for each specific SNP allele. We calculated the rates of

evolutionary changes for each ROI as a percent identity change over the lapsed mln yrs of

evolution (Figure 2C and 3C). Among the 6 *NTNG1*-located SNP ROIs three of them display

accelerated rates of evolution from marmoset to chimpanzee (rs2218404, rs628117, rs96501)

when compared to the mouse-marmoset rates, and rs2218404 (affecting VC in human

subjects) additionally demonstrates an accelerated rate of evolution on the chimpanzee to

human path (Figure 2C). As for the *NTNG2* located ROIs (Figure 3-B), rs1105684 is

remarkably consistent at displaying high evolutionary rates around 0.8 and, together with

rs2274855, both have identical rates at the mouse-marmoset and chimpanzee-human paths,

but differ dramatically at the marmoset-chimpanzee point (0.8 vs 0, respectively). rs2274855

is the only *NTNG2*-nested SNP ROI that underwent an AE from chimpanzee to human. Next

we compared the DNA sequences of all 11 ROIs across mice, primates and extinct hominins

NTNG gene paralogs (see ST2 for the datasets sources used for the genes reconstruction). T-

allele of rs2218404 is detectable in marmoset and its position in mouse corresponds to

adenosine (Figure 2D). G-alelle (associated with a higher VC score comparing to the human

T-allele carriers) is found in Mesolithic hunter Loschbour (8,000 BC) but not in another

ancient hunter Motala12 and is also present in other two hominins belonging to the Neolithic

period, Iceman and Eskimos (5,300 and 4,000 yrs BC, respectively). rs628117 is the only

NTNG1-related mutation near vicinity of which ($\pm$50 nu) is located an intra-hominins (Es, Ice,

Lo) mutation (Figure 2B, low left). The next, PS-affecting, ROI of rs96501 displays an

intricate path of T-allele evolution (associated with a higher PS score) being anciently

conserved from mice to primates but later substituted on the less efficient (in terms of the

generated IQ scores) C allele in Neanderthals, later again replaced by the T allele in

Mesolithic hunters and coming back during the Neolithic times (Figure 2D). The first *NTNG2*

SNP rs1105684 is located at the beginning of the gene and affects WM in healthy human

subjects (Figure 3A). The origin of the T-allele is evolutionary bound to marmoset since its

position in mice is occupied by another pyrimidine base C (Figure 3D). Next on the gene are

two SNP alleles for rs7851893 and rs3824574 which do not affect IQ and similarly to

rs1105684 are surrounded by highly conserved ROIs not only in hominins and chimpansee

(100% identity) but also in marmoset (except 1 mutation for rs7851893). rs2149171 ROI

(affecting the WM and PO scores) similarly to rs3824574 is 100% conserved across the all

species (including the mouse) except the allele itself. The attenuating IQ C-allele position is

occupied in mice genome by T but present in Iceman and Eskimos genes. A distinct

evolutionary path is taken by another cognitive endophenotype-associated and affecting WM

and PO scores A-allele of rs2274855 and its ROI (Figure 3D). Its position in mice is likely to

be occupied by the C pyrimidine base (the software places a blank instead of it) which is

gradually substituted on purine G in chimpanzee and misplaced by the lower IQ score-

associated A-allele in Mesolithic hominins. And 20 nu downstream of the centre of ROI is

located a modern human-specific point mutation translated into the T346A protein

substitution (as described below).

The distinct picture of evolutionary changes among the *NTNG1* and *NTNG2* nested

SNPs has prompted us to compare evolutionary rates for the full-length proteins encoded by

these gene paralogs, Netrin-G1 and Netrin-G2, respectively (2, 35). Netrin-G1 undergoes

only few changes in its amino acid composition with the maximum calculated rate of

evolution reaching 0.05 when mice and marmoset proteins are compared, 0.01 among the

primates, and 0.03 between chimpanzee and human due to a single point mutation A81S

(Figure 2E and SM: Netrin-G1), absent in other primates. Netrin-G2 evolves 2.8 times faster

between mouse and marmoset than its paralog, Netrin-G1, and continues evolving with a

steady rate of 0.05 from primates to human (Figure 3E and SM: Netrin-G2). We have also

reconstructed both proteins from ancient (Neanderthals, Paleolithic time) and extinct

hominins (Mesolithic and Neolithic times) and compared them with primates' and mice'

Netrin-G orthologs (Figure 2F and 3F). Netrin-G1 is a highly conserved protein among the

primates and hominins (Figure 3F). As for Netrin-G2, a mutation shared among the

Neanderthals' and Mesolithic genomes, primates and mice (T346A) is absent in the Neolithic

Iceman and modern human (the signal for Motala3, Motala1 and MezmayskayaNea is not

clear due to low sequence coverage). Primates (marmoset and chimpanzee) share another

mutation (S371A/V) preserved in mouse and absent in hominins (further details can be found

in the SM: Results).

DISCUSSION

NTNG paralog SNPs and associated cognitive endophenotypes of human subjects.

Shortcomings of cognitive and information processing are key features of SCZ diagnosis (36).

They are frequently manifested as impairments in PO, WM, VC and PS (see (37) for

references) and reported as attenuated scores upon IQ tests implementation. SCZ patients

carrying a mutation allele for one of *NTNG* gene paralog SNPs form cognitive

endophenotypes affecting the IQ scores (Figure 1C). The term "endophenotype" was

originally coined by (38) and later advanced through the field of psychiatry by Gottesman

and Shields (reviewed in (39)) as a biomarker associated with a specific phenotypic trait (40).

The formed SCZ endophenotypic groups comprise from subjects with either affected VIQ

(via attenuated VC by *NTNG1* rs2218404 or WM by *NTNG2* rs2149171 and rs2274855) or

affected PIQ (via attenuated PO by *NTNG2* rs2149171 and rs2274855). In two extra cases PS

is affected by rs96501 of *NTNG1* and WM by rs1105684 of *NTNG2* (Figure 1B-1 and B-2,

respectively) but in healthy human subjects. Such intriguing non-overlapping effect on the IQ

domains prompts us to conclude that *NTNG* paralogs complement each other function and

represent an example of how a synapse-expressed genes affect the human cognitive abilities,

perhaps through the precision of neuronal connectivity perturbations and concomitant

miswirings. The observed phenomena of the affected WM is the most striking due to its

multifaceted constructive nature (41) underlying many, if not all, cognitive tasks such as

comprehension, reasoning and learning (42) and historically introduced by Baddeley as the

reading span test (43). Lack of the localization effect of *NTNG2* SNP mutation alleles, all

three are located in different parts of the gene (Figure 1A-2) but associated with identical

endophenotype (Figure 1B-2), points to a uniform nature of the *NTNG2* function distribution

over the entire gene. An obviously non-coding nature of all five IQ-affecting alleles

(rs2149171 despite being exon 4-located encodes a silent F246F mutation) corroborates an

idea that anthropoid trait-associated loci lie outside coding protein areas (44, 45). This hints

towards a potential of these alleles to perturb genes regulatory functions, e.g. mRNA splicing,

affecting downstream located pivotally functional *NTNG* elements such as Ukd-domain

encoding exons 6 and 7 or a unique Netrin-Gs trait – GPI-link. Alternatively, or

simultaneously, the *NTNG* SNP alleles may be embedded into an epistatic network of other

genes influencing human cognitive traits (46). However since it is usual for a SNP effect to

be estimated using an additive model (assuming either independent and cumulative single

contribution) to the mean of a trait with the small effect size the power to detect the epistatic

environment drastically declines. Contrary to the genetic associations with gene expression

having large effect sizes (46), cognitive trait-associated effect sizes are reportedly small (47),

e.g. the largest effect sizes of the variance of intelligence scores accounted for only 0.2% (48),

0.5% on GWA studies of 1,583 adolescence (49) or was predicted to be ~1% on 3,511 adults

(50). Another GWAS of educational attainment (sharing a moderate correlate with

intelligence), which included 126,559 individuals, reports on just 1% of the variance but only

0.02% in a replication sample (51). Our data support the preexisted conclusion that human

cognitive traits modalities are not described by statistically large effect sizes.

Evolutionary elaborations of the embedding IQ-affecting mutations loci. Eleven previously

published SCZ-associated SNPs were tested for their effect on IQ performance of human

subjects and 5 of them were found to be associated with attenuated IQ cognitive

endophenotypes (Figure 1B-1 and B-2). ROIs of 3 of them (rs2218404, rs1373336 and

rs2274855) underwent an AE from chimpanzee to human when compared to New World

monkeys to apes (marmoset-chimpanzee) path (Figure 2C and 3C). Two of them affect IQ in

humans (rs2218404 – VC and rs2274855 – WM, Figure 1B-1 and B-2) while being located

within the vicinity of exon 5 (2,275 nu downstream and 15 nu upstream, respectively) – a

part of the lowest percent identity coding DNA among the *NTNG* gene paralogs (52).

Presence of the evolutionary accelerated regions within the *NTNG* genes non-coding areas

underscores them as contributors to the human-specific traits along with other genes (53).

However, not only ROIs of the IQ-affecting alleles but the alleles themselves demonstrate

several unique evolutionary features (see SM: Discussion). To understand evolutionary forces

driving the emergence of cognitive endophenotype-associated alleles we have deduced a set

of rules outlined as follows. 1. An alternative (mutated) allele evolutionary appearance

coincides with the lack of any other mutations within ROI (a conserved island rule); 2.

positioning of the future mutation often represents a turning point of dramatic changes of an

allele ROI (e.g. as seen in marmoset: rs2218404 (50-90%), rs628117 (30-80%), rs96501

(100-40%)); 3. an AE of ROI often precedes the emergence of a mutation allele (e.g.

rs2218404: chimpanzee to human ($k = 1.59$), appearance of "G" in Loschbour; rs628117:

marmoset to chimpanzee ($k = 1.10$), appearance of "T" in AltaiNea; rs96501: marmoset to

chimpanzee ($k = 0.83$), appearance of "C" in AltaiNea; rs2274855: chimpanzee to human (k

$= 0.79$), appearance of "A" in Motala12; 4. low identity percent (equivalent to subsequent

substantial evolutionary changes) among the evolutionary species within the allele

surrounding proximity of as long as $\pm$50 nu is not sufficient for the future mutated allele

significance as a cognitive endophenotype determinant (as deduced by the IQ score) as seen

for the rs1373336, rs1444042, rs4915045 and rs7851893 (none of them are IQ-affecting,

though associated with SCZ, despite showing (very) low identity in mice). Rather some

dramatic changes within the allele's immediate proximity of $\pm$10 nu (defined as a ROI)

preceded or followed by more stringently conserved DNA are necessary (the conserved

island perturbation rule). Currently we are unable to state that the IQ-associated alleles ROIs

represent regulatory loci and an important source of evolutionary innovation (54) but they

may be the smallest functional blocks of a strong positive selection that exerts its action upon,

similarly to the 20–30 nu clusters of strongly conserved non-coding elements (CNEs),

transcription factor binding sites (TFBS), RNA splicing and editing motifs (55).

Extinct hominins and IQ-associated mutation alleles. Availability of archaic genomes allows

excavation for the advantageous alleles that modern humans acquired from archaic extinct

hominins such as Neandertals and Denisovans who used to live 230,000-30,000 years ago

(Middle/Upper Paleolithic, Old Stone Age) defined by distinct morphological features (56),

and from modern extinct humans (hunters, farmers) from Mesolithic (Middle Stone Age,

~10,000 yrs BC, (57)) and Neolithic (New Stone Age, ~5,000 BC, (58, 59) periods. Though

exhibiting several anatomical features, making archaic hominins different from the modern

human, there are studies challenging the idea that reserve symbolism and abstract thinking

was an exclusive prerogative of modern human (60, 61). The time Neanderthals used to live

in is thought to be associated with the onset of cognitive fluidity involving the capacity to

draw analogies (early paintings), to combine concepts (making tools) and to adapt ideas for

new contexts (62). Wynn and Coolidge believe that evolution of WM was central to the

evolution of human cognitive traits consisted from few genetic mutations that led to

"enhanced WM" 200,000-40,000 BC (63). Our work partially supports this idea showing the

perseverance of higher WM score-associated alleles across the hominins such as T of

rs1105684 and G of rs2274855 (Figure 3D) but a Neolithic appearance of rs2149171 T in

Iceman previously found only in mice genome supporting the conclusion made by Crabtree

(64) that modern humans as species "are surprisingly intellectually fragile and perhaps

reached a peak 2,000–6,000 years ago". Mesolithic period has been always considered as a

key gate for the evolution of human languages (65) with our data showing that rs2218404 G-

allele associated with a higher VC score (Figure 1B-1) emerges for the first time in the

Loschbour hunter *NTNG1* gene (Figure 3D). VC, as an example of abstract symbol usage, is

associated with the global network efficiency (as a part of fronto-parietal network (19,66),

and global communication and intellectual performance (67). From this point of view it is not

surprising that appearance of the G allele in Loschbour and Iceman genomes coincides with

the presence of PS enhancing rs96501 T allele (Figure 3D). A wealth of data has been

collected characterising possible look and health status of archaic hominins and modern but

extinct humans (for ref. see (68) and SM: Discussion). Based on our own data we may also

speculate that the extinct hominins may have had a lower VC comparing to us, and

consequently, Neanderthals were unlikely able for a semantic communication due to a global

network inefficiency VC is associated with; they have had similar to us PS (if they had lived

beyond the Mesolithic period), and were likely to have had identical to modern human WM,

corroborating the advanced evolutionary nature of this important human cognitive domain of

a limited capacity and associated with intelligence.

CONCLUSION

Evolution of a novel function relies on enhanced genetic robustness via functional

redundancy potentially provided by a gene duplication event. Further evolutionary outcome

depends on the substrate availability (undergoing its own evolution) upon which the novel

function(s) exerts its action. Nature does not create but tinkers to perfection provided to it

material exploring available evolutionary tools. Half a billion years ago a gene duplication

event had provided a plethora of such substrate thus converting the evolution itself into a

"Creator" of new functions. Here we have described how a pair of twin genes got themselves

involved into the human cognitive functioning believed to be emerged in a primordial state in

primitive vertebrates prior to the first recorded gene duplication. Subsequent process of the

function specialisation made *NTNG* paralogs to subfunctionalise into distinct cognitive

domains in a complementary manner (52).

MATERIALS AND METHODS

Ethics statement. This study was performed in accordance with the World Medical

Association's Declaration of Helsinki and approved by the Osaka University Research Ethics

Committee. A written informed consent was obtained from all subjects after the procedures

had been fully explained.

Subjects. The procedures were performed as per established protocols at Osaka University as

described previously (69). The subjects consisted from 339 patients with SCZ and 716

healthy controls. The sex ratio did not differ significantly between the groups, but the mean

age was significantly different. The subjects were all biologically unrelated Japanese and

recruited from both outpatient and inpatient units at Osaka University Hospital and other

psychiatric hospitals. Each patient with SCZ had been diagnosed by at least two trained

psychiatrists based on unstructured clinical interviews, according to the criteria of the DSM-

IV (36). In case if the diagnosis of the two trained psychiatrists was discordant, it was

resolved through the further negotiations on both specialist opinions. In case of unresolved

diagnostic disputes, the patient was omitted from the study. Psychiatrically healthy controls

were recruited through local advertisements and were evaluated by means of unstructured

interviews to exclude individuals with current or past contact with psychiatric services, those

who experienced psychiatric medications, or who were not Japanese. Controls for family

history of a CD, such as SCZ, BD, or major depressive disorder were not included. Ethnicity

was determined by self-report and was not confirmed by genetic analyses. Additionally,

subjects were excluded from this study if they had neurologic or medical conditions that

could have potentially affect their central nervous system, such as atypical headaches, head

trauma with loss of consciousness, chronic lung disease, kidney disease, chronic hepatic

disease, thyroid disease, active cancer, cerebrovascular disease, epilepsy, seizures, substance

abuse related disorders, or mental retardation.

SNPs selection, genotyping, and genomic sequencing. This study was designed to examine

the association of SCZ patients cognitive performance (through WAIS-III implementation

(17)) with *NTNG* genes. Venous blood was collected from the subjects. Genomic DNA was

extracted from the whole blood using standard procedures. The SNPs (26-28, 30) were

genotyped using the TaqMan allelic discrimination assay (Applied Biosystems, Foster City,

CA). No deviations from the Hardy-Weinberg equilibrium in the examined SNPs were

detected ($p > 0.05$).

Statistical analysis. The effects of the diagnosis, genotype and their interaction on cognitive

performances in the WAIS were analyzed by two-way analyses of covariance (ANCOVA).

Diagnosis and genotype statuses were included in the model as independent variables (ST1).

FIQ and each WAIS subscale score (VIQ, PIQ, VC, PO, WM, PS, Vc, SiM, IF, CH, AM, DS,

LNS, PC, BD, and MR) were included as dependent variables. Sex, age and years of

education were treated as covariates, as they were possible confounding factors. All p values

are two tailed, and statistical significance was defined as $*p < 0.05$ and $**p < 0.01$.

Identity percent calculations and the definition of ROIs. The complete procedure is described

in the Figures 2-3 legend. Stretcher (70) was used for the alignments (the default values were:

gap penalty – 16 (DNA) and 12 (protein), and the extend penalty – 4 (DNA) and 2 (protein)),

for the percent identity calculations and evolutionary rates. A ROI's selection was initially

defined as a minimal area surrounding a SNP mutation allele incorporating the outmost

evolutionary dramatic changes, turns out to be ±10 nu.

Mice, primates and hominins *NTNG* paralogs DNA and encoded protein sequences

reconstruction. Genomes for mouse (GRC38.p3) and marmoset (C_jacchus3.2.1) were from

Ensemble. Since chimpanzee's genome is based only on a single individual (CHIMP2.1.4,

Clint) and contains several questionable information we have reconstructed a consensus

genome sequence for both *NTNG* genes based on 25 primate sequences of *Pan troglodytes*

(71). All datasets used for the *NTNG* paralogs DNA and encoded by them proteins

reconstruction are listed in ST2. For details refer to SM.

SUPPLEMENTARY MATERIALS (SM)

Contain additional Results and Discussion, Supplementary Methods (ancient and primate

genomes reconstructions), and Supplementary Tables (ST1 and ST2) as a single compiled pdf

file. Also included are human Netrin-G1 and Netrin-G2 alignments, as well as 101 nu

alignments for all 11 ROIs across the all analysed species.

ACKNOWLEDGEMENTS

Authors would like to acknowledge the financial support provided by Funding Program for

World-Leading Innovative R&D on Science and Technology (FIRST Programme) and

KAKENHI 15H04290 by the Japan Society for the Promotion of Science (JSPS).

COMPETING INTERESTS

Authors would like to express a lack of any competing interests associated with the work.

REFERENCES

1. Ohno S. 1970. Evolution by Gene Duplication. Springer, New York.

2. Nakashiba T, Nishimura S, Ikeda T, Itohara S. 2002. Complementary expression and

neurite outgrowth activity of netrin-G subfamily members. *Mechanisms of Development*

111:47-60. doi: 10.1016/S0925-4773(01)00600-1.

3. Nishimura-Akiyoshi S, Niimi K, Nakashiba T, and Itohara S. 2007. Axonal netrin-Gs

transneuronally determine lamina-specific subdendritic segments. *Proceedings of the*

National Academy of Sciences of the United States of America 104:14801-14806. doi:

10.1073/pnas.0706919104.

4. Matsukawa H, Akiyoshi-Nishimura S, Zhang Q, Lujan R, Yamaguchi K, Goto H, Yaguchi

K, Hashikawa T, Sano C, Shigemoto R, Nakashiba T, Itohara S. 2014. Netrin-G/NGL

Complexes Encode Functional Synaptic Diversification. *Journal of Neuroscience*

34:15779-15792. doi: 10.1523/jneurosci.1141-14.2014.

5. Sun KLW, Correia JP, Kennedy TE. 2011. Netrins: versatile extracellular cues with

diverse functions. *Development* 138:2153-2169. doi: 10.1242/dev.044529.

6. Yu S, Guo Z, Johnson C, Gu G, Wu Q. 2013. Recent progress in synthetic and biological

studies of GPI anchors and GPI-anchored proteins. *Current Opinion in Chemical Biology*

17: 1006-1013. doi: 10.1016/j.cbpa.2013.09.016.

7. Sevcsik E, Brameshuber M, Foelser M, Weghuber J, Honigmann A, Schütz GJ. 2015. GPI-

anchored proteins do not reside in ordered domains in the live cell plasma membrane.

Nature Communications 6:A6969. doi: 10.1038/ncomms7969.

8. Klotzsch E and Schutz GJ. 2013. A critical survey of methods to detect plasma membrane

rafts. *Philosophical Transactions of the Royal Society B-Biological Sciences*

368:20120033. doi: 10.1098/rstb.2012.0033.

9. Abitua PB, Wagner E, Navarrete IA, Levine M. 2012. Identification of a rudimentary

neural crest in a non-vertebrate chordate. *Nature* 492:104-107. doi: 10.1038/nature11589.

10. Stolfi A, Ryan K, Meinertzhagen IA, Christiaen L. 2015. Migratory neuronal progenitors

arise from the neural plate borders in tunicates. *Nature* 527:371-374. doi:

10.1038/nature15758.

11. Abitua PB, Gainous TB, Kaczmarczyk AN, Winchell CJ, Hudson C, Kamata K,

Nakagawa M, Tsuda M, Kusakabe, TG, Levine M. 2015. The pre-vertebrate origins of

neurogenic placodes. *Nature* 524:462-465. doi: 10.1038/nature14657.

12. Wise SP. 2008. Forward frontal fields: phylogeny and fundamental function. *Trends in*

Neurosciences 31:599-608. doi: 10.1016/j.tins.2008.08.008.

13. Preuss TM. 2012. Human brain evolution: From gene discovery to phenotype discovery.

Proceedings of the National Academy of Sciences of the United States of America

109:10709-10716. doi: 10.1073/pnas.1201894109.

14. Geschwind DH and Rakic P. 2013. Cortical Evolution: Judge the Brain by Its Cover.

Neuron 80:633-647. doi: 10.1016/j.neuron.2013.10.045.

15. Belmonte JCI, Callaway EM, Churchland P, Caddick SJ, Feng G, Homanics GE, Lee K-F,

Leopold DA, Miller CT, Mitchell JF, Mitalipov S, Moutri AR, Movshon JA, Okano H,

Reynolds JH, Ringach D, Sejnowski TJ, Silva AC, Strick PL, Wu J, Zhang F. 2015.

Brains, Genes, and Primates. *Neuron* 86:617-631. doi: 10.1016/j.neuron.2015.03.021.

16. Joshi PK, Esko T, Mattsson H, Eklund N, Gandin I, Nutile T, Jackson AU, Schurmann C,

Smith AV, Zhang W, Okada Y, Stančáková A, Faul JD, Zhao W, Bartz TM, Concas MP,

Franceschini N, Enroth S, Vitart V, Trompet S, Guo X, Chasman DI, O'Connel JR, Corre

T, Nongmaithem SS, Chen Y, Mangino M, Ruggiero D, Traglia M, Farmaki A-E,

Kacprowski T, Bjonnes A, van der Spek A, Wu Y, Giri AK, Yanek LR, Wang L, Hofer E,

Rietveld CA, McLeod O, Cornelis MC, Pattaro C, Verweij N, Baumbach C, Abdellaoui A,

Warren HR, Vuckovic D, Mei H, Bouchard C, Perry JRB, Cappellani S, Mirza SS, Benton

MC, Broeckel U, Medland SE, Lind PA, Malerba G, Drong A, Yengo L, Bielak LF, Zhi D,

van der Most PJ, Shriner D, Maegi R, Hemani G, Karaderi T, Wang Z, Liu T, Demuth I,

Zhao JH, Meng W, Lataniotis L, van der Laan SW, Bradfield JP, Wood AR, Bonnefond

A, Ahluwalia TS, Hall LM, Salvi E, Yazar S, Carstensen L, de Haan HG, Abney M, Afzal

U, Allison MA, Amin N, Asselbergs FW, Bakker SJL, Barr RG, Baumeister SE, Benjamin

DJ, Bergmann S, Boerwinkle E, Bottinger EP, Campbell A, Chakravarti A, Chan Y,

Chanock SJ, Chen C, Chen Y-DI, Collins FS, Connell J, Correa A, Cupples LA, Smith

GD, Davies G, Dörr M, Ehret G, Ellis SB, Feenstra B, Feitosa MF, Ford I, Fox CS,

Frayling TM, Friedrich N, Geller F, Generation Scotland, Gillham-Nasenya I, Gottesman

O, Graff M, Grodstein F, Gu C, Haley C, Hammond CJ, Harris SE, Harris TB, Hastie ND,

Heard-Costa NL, Heikkilä K, Hocking LJ, Homuth G, Hottenga J-J, Huang J, Huffman JE,

Hysi PG, Ikram MA, Ingelsson E, Joensuu A, Johansson Å, Jousilahti P, Jukema JW,

Kähönen M, Kamatani Y, Kanoni S, Kerr SM, Khan NM, Koellinger P, Koistinen HA,

Kooner MK, Kubo M, Kuusisto J, Lahti J, Launer LJ, Lea RA, Lehne B, Lehtimäki T,

Liewald DCM, Lind L, Loh M, Lokki M-L, London SJ, Loomis SJ, Loukola A, Lu Y,

Lumley T, Lundqvist A, Männistö S, Marques-Vidal P, Masciullo C, Matchan A, Mathias

RA, Matsuda K, Meigs JB, Meisinger C, Meitinger T, Menni C, Mentch FD, Mihailov E,

Milani L, Montasser ME, Montgomery GW, Morrison A, Myers RH, Nadukuru R,

Navarro P, Nelis M, Nieminen MS, Nolte IM, O'Connor GT, Ogunniyi A, Padmanabhan S,

Palmas WR, Pankow JS, Patarcic I, Pavani F, Peyser PA, Pietilainen K, Poulter N,

Prokopenko I, Ralhan S, Redmond P, Rich SS, Rissanen H, Robino A, Rose LM, Rose R,

Sala C, Salako B, Salomaa V, Sarin A-P, Saxena R, Schmidt H, Scott LJ, Scott WR,

Sennblad B, Seshadri S, Sever P, Shrestha S, Smith BH, Smith JA, Soranzo N,

Sotoodehnia N, Southam L, Stanton AV, Stathopoulou MG, Strauch K, Strawbridge RJ,

Suderman MJ, Tandon N, Tang S-T, Taylor KD, Tayo BO, Töglhofer AM, Tomaszewski

M, Tšernikova N, Tuomilehto J, Uitterlinden AG, Vaidya D, van Hylckama Vlieg A, van

Setten J, Vasankari T, Vedantam S, Vlachopoulou E, Vozzi D, Vuoksimaa E,

Waldenberger M, Ware EB, Wentworth-Shields W, Whitfield JB, Wild S, Willemsen G,

Yajnik CS, Yao J, Zaza G, Zhu X, The BioBank Japan Project, Salem RM, Melbye M,

Bisgaard H, Samani NJ, Cusi D, Mackey DA, Cooper RS, Froguel P, Pasterkamp G, Grant

SFA, Hakonarson H, Ferrucci L, Scott RA, Morris AD, Palmer CAN, Dedoussis G,

Deloukas P, Bertram L, Lindenberger U, Berndt SI, Lindgren CM, Timpson NJ, Toenjes

A. Munroe PB, Sørensen TIA, Rotimi CN, Arnett DK, Oldehinkel AJ, Kardia SLR,

Balkau B, Gambaro G, Morris AP, Eriksson JG, Wright MJ, Martin NG, Hunt SC, Starr

JM, Deary IJ, Griffiths LR, Tiemeier H, Pirastu N, Kaprio J, Wareham NJ, Pérusse L,

Wilson JG, Girotto G, Caulfield MJ, Raitakari O, Boomsma DI, Gieger C, van der Harst

Pim, Hicks AA, Kraft P, Sinisalo J, Knekt P, Johannesson M, Magnusson PKE, Hamsten

A, Schmidt R, Borecki IB, Vartiainen E, Becker DM, Bharadwaj D, Mohlke KL, Boehnke

M, van Duijn CM, Sanghera DK, Teumer A, Zeggini E, Metspalu A, Gasparini P, Ulivi S,

Ober C, Toniolo D, Rudan I, Porteous DJ, Ciullo M, Spector TD, Hayward C, Dupuis J,

Loos RJF, Wright AF, Chandak GR, Vollenweider P, Shuldiner AR, Ridker PM, Rotter JI,

Sattar N, Gyllensten U, North KE, Pirastu M, Psaty BM, Weir DR, Laakso M, Gudnason

V, Takahashi A, Chambers JC, Kooner JS, Strachan DP, Campbell H, Hirschhorn JN,

Perola M, Polašek O, Wilson JF. 2015. Directional dominance on stature and cognition in

diverse human populations. *Nature* 523:459-462. doi: 10.1038/nature14618.

17. Wechsler D. 1958. *The Measurement And Appraisal Of Adult Intelligence*, 4th edition.

The Williams & Wilkins Company. 324 pp.

18. Deary IJ, Spinath FM, Bates TC. 2006. Genetics of intelligence. *European Journal of*

Human Genetics 14:690-700. doi: 10.1038/sj.ejhg.5201588.

19. Glascher J, Tranel D, Paul LK, Rudrauf D, Rorden C, Hornaday A, Grabowski T,

Damasio H, Adolphs R. 2009. Lesion Mapping of Cognitive Abilities Linked to

Intelligence. *Neuron* 61:681-691. doi: 10.1016/j.neuron.2009.01.026.

20. Spearman C. 1904. "General intelligence" objectively determined and measured.

American Journal of Psychology 15:201-292. doi: 10.2307/1412107.

21. Hampshire A, Highfield RR, Parkin BL, Owen AM. 2012. Fractionating Human

Intelligence. *Neuron* 76:1225-1237. doi: 10.1016/j.neuron.2012.06.022.

22. Lett TA, Voineskos AN, Kennedy JL, Levine B, Daskalakis ZJ. 2014. Treating working

memory deficits in schizophrenia: a review of the neurobiology. Biological Psychiatry

75:361-170. doi: 10.1016/j.biopsych.2013.07.026.

23. Forbes NF, Carrick LA, McIntosh AM, Lawrie SM. 2009. Working memory in

schizophrenia: a meta-analysis. *Psychological Medicine* 39:889-905. doi:

10.1017/s0033291708004558.

24. Leeson VC, Robbins TW, Matheson E, Hutton SB, Ron MA, Barnes TRE, Joyce EM.

2009. Discrimination Learning, Reversal, and Set-Shifting in First-Episode Schizophrenia:

Stability Over Six Years and Specific Associations with Medication Type and

Disorganization Syndrome. *Biological Psychiatry* 66:586-593. doi:

10.1016/j.biopsych.2009.05.016.

25. Barnett JH, Robbins TW, Leeson VC, Sahakian BJ, Joyce EM, Blackwell AD. 2010.

Assessing cognitive function in clinical trials of schizophrenia. *Neuroscience and*

Biobehavioral Reviews 34:1161-1177. doi: 10.1016/j.neubiorev.2010.01.012.

26. Fukasawa M, Aoki M, Yamada K, Iwayama-Shigeno Y, Takao H, Meerabux J, Toyota T,

Nishikawa T, Yoshikawa T. 2004. Case-control association study of human netrin G1

gene in Japanese schizophrenia. *Journal of Medical and Dental Sciences* 51:121-128.

27. Aoki-Suzuki M, Yamada K, Meerabux J, Iwayama-Shigeno Y, Ohba H, Iwamoto K,

Takao H, Toyota T, Suto Y, Nakatani N, Dean B, Nishimura S, Seki K, Kato T, Itohara S,

Nishikawa T, Yoshikawa, T. 2005. A Family-Based Association Study and Gene

Expression Analyses of Netrin-G1 and -G2 Genes in Schizophrenia. *Biological Psychiatry*

57:382–393. doi: 10.1016/j.biopsych.2004.11.022.

28. The Japanese Schizophrenia Sib-Pair Linkage Group (JSSLG), Arinami T, Ohtsuki T,

Ishiguro H, Ujike H, Tanaka Y, Morita Y, Mineta M, Takeichi M, Yamada S, Imamura A,

Ohara K, Shibuya H, Ohara, K, Suzuki Y, Muratake T, Kaneko N, Someya T, Inada T,

Yoshikawa T, Toyota T, Yamada K, Kojima T, Takahashi S, Osamu O, Shinkai T,

Nakamura M, Fukuzako H, Hashiguchi T, Niwa S, Ueno T, Tachikawa H, Hori T, Asada T,

Nanko S, Kunugi H, Hashimoto R, Ozaki N, Iwata N, Harano M, Arai H, Ohnuma T,

Kusumi I, Koyama T, Yoneda H, Fukumaki Y, Shibata H, Kaneko S, Higuchi H, Yasui-

Furukori N, Numachi Y, Itokawa M, Okazaki Y. 2005. Genomewide high-density SNP

linkage analysis of 236 Japanese families supports the existence of schizophrenia

susceptibility loci on chromosomes 1p, 14q, and 20p. *American Journal of Human*

Genetics 77:937-944. doi: 10.1086/498122.

29. Eastwood SL, Harrison PJ. 2008. Decreased mRNA expression of netrin-G1 and netrin-

G2 in the temporal lobe in schizophrenia and bipolar disorder. *Neuropsychopharmacology*

33:933-945. doi: 10.1038/sj.npp.1301457.

30. Ohtsuki T, Horiuchi Y, Koga M, Ishiguro H, Inada T, Iwata N, Ozaki N, Ujike H,

Watanabe Y, Someya T, Arinami T. 2008. Association of polymorphisms in the haplotype

block spanning the alternatively spliced exons of the *NTNG1* gene at 1p13.3 with

schizophrenia in Japanese populations. *Neuroscience Letters* 435:194-197. doi:

10.1016/j.neulet.2008.02.053.

31. Zakharyan R, Boyajyan A, Arakelyan A, Gevorgyan A, Mrazek F, Petrek M. 2011.

Functional variants of the genes involved in neurodevelopment and susceptibility to

schizophrenia in an Armenian population. *Human Immunology* 72:746-748. doi:

10.1016/j.humimm.2011.05.018.

32. Zhu Y, Yang H, Bi Y, Zhang Y, Zhen C, Xie S, Qin H, He J, Liu L, Liu Y. 2011. Positive

association between *NTNG1* and schizophrenia in Chinese Han population. *Journal of*

Genetics 90:499-502.

33. Ayalew M, Le-Niculescu H, Levey DF, Jain N, Changala B, Patel SD, Winiger E, Breier

A, Shekhar A, Amdur R, Koller D, Nurnberger JI, Corvin A, Geyer M, Tsuang MT,

Salomon D, Schork NJ, Fanous AH, O'Donovan MC, Niculescu AB. 2012. Convergent

functional genomics of schizophrenia: from comprehensive understanding to genetic risk

prediction. *Molecular Psychiatry* 17:887-905. doi: 10.1038/mp.2012.37.

34. Wilcox JA and Quadri S. 2014. Replication of NTNG1 association in schizophrenia.

Psychiatric Genetics 24:266-268. doi: 10.1097/ypg.0000000000000061.

35. Nakashiba T, Ikeda T, Nishimura S, Tashiro K, Honjo T, Culotti JG, Itohara, S. 2000.

Netrin-G1: a Novel Glycosyl Phosphatidylinositol-Linked Mammalian Netrin That Is

Functionally Divergent from Classical Netrins. *The Journal of Neuroscience* 20:6540-

6550.

36. APA (American Psychiatric Association). 2013. Diagnostic and Statistical Manual of Mental

Disorders (DSM-V®), Fifth Edition. 991 pp. doi: 10.1176/appi.books.9780890425596.

37. Yoon JH, Sheremata SL, Rokem A, Silver MA. 2014. Windows to the soul: vision

science as a tool for studying biological mechanisms of information processing deficits in

schizophrenia. *Frontiers in Psychology* 4:A681. doi: 10.3389/fpsyg.2014.00681.

38. John B and Lewis KR. 1966. Chromosome variability and geographic distribution in

insects. *Science* 152:711–721. doi: 10.1126/science.152.3723.711.

39. Gottesman II and McGue M. 2015. Endophenotype. In *The Encyclopedia of Clinical*

Psychology, First Edition. Edited by Cautin RL and Lilienfeld SO. JohnWiley & Sons, Inc.

doi: 10.1002/9781118625392.wbecp423.

40. Glahn DC, Knowles EE, McKay DR, Sprooten E, Raventós H, Blangero J, Gottesman I,

Almasy L. 2014. Arguments for the Sake of Endophenotypes: Examining Common

Misconceptions About the Use of Endophenotypes in Psychiatric Genetics. *American*

Journal of Medical Genetics Part B 165B:122–130. doi: 10.1002/ajmg.b.32221.

41. Frydecka D, Eissa AM, Hewedi DH, Ali M, Drapała J, Misiak B, Kłosińska E, Phillips

JR, Moustafa AA. 2014. Impairments of working memory in schizophrenia and bipolar

disorder: the effect of history of psychotic symptoms and different aspects of cognitive

task   demands.   *Frontiers   in   Behavioral   Neuroscience*   8:A416.   doi:

10.3389/fnbeh.2014.00416.

42. Baddeley A. 1992. Working memory. *Science* 255:556-559. doi:10.1126/science.1736359.

43. Mackintosh NJ. 2011. "History of theories and measurement of Intelligence". *In* "The

Cambridge Handbook of Intelligence". Edited by Sternberg RJ, Kaufman SB. p. 3-19. doi:

10.1017/CBO9780511977244.

44. del Rosario RCH, Rayan NA, Prabhakar S. 2014. Noncoding Origins of Anthropoid

Traits and a New Null Model of Transposon Functionalization. *Genome Research*

24:1469-1484. doi: 10.1101/gr.168963.113.

45. Kellis M, Wold B, Snyder MP, Bernstein BE, Kundaje A, Marinov GK, Ward LD, Birney

E, Crawford GE, Dekker J, Dunham I, Elnitski LL, Farnham PJ, Feingold EA, Gerstein M,

Giddings MC, Gilbert DM, Gingeras TR, Green ED, Guigo R, Hubbard T, Kent J, Lieb JD,

Myers RM, Pazin MJ, Ren B, Stamatoyannopoulos JA, Weng Z, White KP, Hardison RC.

2014. Defining functional DNA elements in the human genome. *Proceedings of the*

National Academy of Sciences of the United States of America 111:6131-6138. doi:

10.1073/pnas.1318948111.

46. Hemani G, Shakhbazov K, Westra H-J, Esko T, Henders AK, McRae AF, Yang J, Gibson

G, Martin NG, Metspalu A, Franke L, Montgomery GW, Visscher PM, Powell JE. 2014.

Detection and replication of epistasis influencing transcription in humans. *Nature*

508:249–253. doi: 10.1038/nature13005.

47. Plomin R and Deary IJ. 2015. Genetics and intelligence differences: five special findings.

Molecular Psychiatry 20:98-108. doi: 10.1038/mp.2014.105.

48. Benyamin B, Pourcain BSt, Davis OS, Davies G, Hansell NK, Brion M-JA, Kirkpatrick

RM, Cents RAM, Franic S, Miller MB, Haworth CMA, Meaburn E, Price TS, Evans DM,

Timpson N, Kemp J, Ring S, McArdle W, Medland SE, Yang J, Harris SE, Liewald DC,

Scheet P, Xiao X, Hudziak JJ, de Geus EJC, Wellcome Trust Case Control Consortium 2

(WTCCC2), Jaddoe VWV, Starr JM, Verhulst FC, Pennell C, Tiemeier H, Iacono WG,

Palmer LJ, Montgomery GW, Martin NG, Boomsma DI, Posthuma D, McGue M, Wright

MJ, Davey Smith G, Deary IJ, Plomin R, Visscher PM. 2014. Childhood intelligence is

heritable, highly polygenic and associated with *FNBP1L. Molecular Psychiatry* 19:253-

258. doi: 10.1038/mp.2012.184.

49. Desrivières S, Lourdusamy A, Tao C, Toro R, Jia T, Loth E, Medina LM, Kepa A,

Fernandes A, Ruggeri B, Carvalho FM, Cocks G, Banaschewski T, Barker GJ, Bokde

ALW, Büchel C, Conrod PJ, Flor H, Heinz A, Gallinat J, Garavan H, Gowland P, Brühl R,

Lawrence C, Mann K, Martinot MLP, Nees F, Lathrop M, Poline J-B, Rietschel M,

Thompson P, Fauth-Bühler M, Smolka MN, Pausova Z, Paus T, Feng J, Schumann G and

the IMAGEN Consortium. 2015. Single nucleotide polymorphism in the neuroplastin

locus associates with cortical thickness and intellectual ability in adolescents. *Molecular*

Psychiatry 20: 263-274. doi: 10.1038/mp.2013.197.

50. Davies G, Tenesa A, Payton A, Yang J, Harris SE, Liewald D, Ke X, Hellard SLe,

Christoforou A, Luciano M, McGhee K, Lopez L, Gow AJ, Corley J, Redmond P, Fox HC,

Haggarty P, Whalley LJ, McNeill G, Goddard ME, Espeseth T, Lundervold AJ, Reinvang

I, Pickles A, Steen VM, Ollier W, Porteous DJ, Horan M, Starr JM, Pendleton N, Visscher

PM, Deary IJ. 2011. Genome-wide association studies establish that human intelligence is

highly   heritable   and   polygenic.   *Molecular   Psychiatry*   16:996-1005.   doi:

10.1038/mp.2011.85.

51. Rietveld CA, Medland SE, Derringer J, Yang J, Esko T, Martin NW, Westra HJ,

Shakhbazov K, Abdellaoui A, Agrawal A, Albrecht E, Alizadeh BZ, Amin N, Bamard J,

Baumeister SE, Benke KS, Bielak LF, Boatman JA, Boyle PA, Davies G, de Leeuw C,

Eklund N, Evans DS, Ferhmann R, Fischer K, Gieger C, Gjessing HK, Hägg S, Harris JR,

Hayward C, Holzapfel C, Ibrahim-Verbaas CA, Ingelsson E, Jacobsson B, Joshi PK,

Jugessur A, Kaakinen M, Kanoni S, Karjalainen J, Kolcic I, Kristiansson K, Kutalik Z,

Lahti J, Lee SH, Lin P, Lind PA, Liu YM, Lohman K, Loitfelder M, McMahon G, Vidal

PM, Meirelles O, Milani L, Myhre R, Nuotio ML, Oldmeadow CJ, Petrovic KE, Peyrot

WJ, Polaśek O, Quaye L, Reinmaa E, Rice JP, Rizzi TS, Schmidt H, Schmidt R, Smith

AV, Smith JA, Tanaka T, Terracciano A, van der Loos MJHM, Vitart V, Völzke H,

Wellmann J, Yu L, Zhao W, Allik J, Attia JR, Bandinelli S, Bastardot F, Beauchamp J,

Bennett DA, Berger K, Bierut LJ, Boomsma DI, Bültmann U, Campbell H, Chabris CF,

Cherkas L, Chung MK, Cucca F, de Andrade M, De Jager PL, De Neve J-E, Deary IJ,

Dedoussis GV, Deloukas P, Dimitriou M, Eiríksdóttir G, Elderson MF, Eriksson JG,

Evans DM, Faul JD, Ferrucci L, Garcia ME, Grönberg H, Guonason V, Hall P, Harris JM,

Harris TB, Hastie ND, Heath AC, Hernandez DG, Hoffmann W, Hofman A, Holle R,

Holliday EG, Hottenga JJ, Iacono WG, Illig T, Järvelin MR, Kähönen M, Kaprio J,

Kirkpatrick RM, Kowgier M, Latvala A, Launer LJ, Lawlor DA, Lehtimäki T, Li JM,

Lichtenstein P, Lichtner P, Liewald DC, Madden PA, Magnusson PKE, Mäkinen TE,

Masala M, McGue M, Metspalu A, Mielck A, Miller MB, Montgomery GW, Mukherjee S,

Nyholt DR, Oostra BA, Palmer LJ, Palotie A, Penninx B, Perola M, Peyser PA, Preisig M,

Räikkönen K, Raitakari OT, Realo A, Ring SM, Ripatti S, Rivadeneira F, Rudan I,

Rustichini A, Salomaa V, Sarin AP, Schlessinger D, Scott RJ, Snieder H, St Pourcain B,

Starr JM, Sul JH, Surakka I, Svento R, Teumer A, The LifeLines Cohort Study, Tiemeier

H, van Rooij FJA, Van Wagoner DR, Vartiainen E, Viikari J, Vollenweider P, Vonk JM,

Waeber G, Weir DR, Wichmann H-E, Widen E, Willemsen G, Wilson JF, Wright AF,

Conley D, Davey-Smith G, Franke L, Groenen PJF, Johannesson M, Kardia SLR, Krueger

RF, Laibson D, Martin NG, Meyer MN, Posthuma D, Thurik AR, Timpson NJ,

Uitterlinden AG, van Duijn CM, Visscher PM, Benjamin DJ, Cesarini D, Koellinger PD.

2013. GWAS of 126,559 Individuals Identifies Genetic Variants Associated with

Educational Attainment. *Science* 340:1467-1471. doi: 10.1126/science.1235488.

52. Prosselkov P, Polygalov D, Zhang Q, McHugh TJ, Itohara S. 2015. Cognitive Domains

function complementation by *NTNG* gene paralogs. *bioRxiv* doi:

http://dx.doi.org/10.1101/034413.

53. Prabhakar S, Noonan JP, Pääbo S, Rubin EM. 2006. Accelerated evolution of conserved

noncoding sequences in humans. *Science* 314:786-786. doi: 10.1126/science.1130738.

54. Rubinstein M and de Souza FSJ. 2013. Evolution of transcriptional enhancers and animal

diversity. *Philosophical Transactions of the Royal Society B-Biological Sciences*

368:A20130017. doi: 10.1098/rstb.2013.0017.

55. Harmston N, Baresic A, Lenhard B. 2013. The mystery of extreme non-coding

conservation. *Philosophical Transactions of the Royal Society B-Biological Sciences*

368:A20130021. doi: 10.1098/rstb.2013.0021.

56. Meyer M, Kircher M, Gansauge MT, Li H, Racimo F, Mallick S, Schraiber JG, Jay F,

Prüfer K, de Filippo C, Sudmant PH, Alkan C, Fu QM, Do R, Rohland N, Tandon A,

Siebauer M, Green RE, Bryc K, Briggs AW, Stenzel U, Dabney J, Shendure J, Kitzman J,

Hammer MF, Shunkov MV, Derevianko AP, Patterson N, Andrés AM, Eichler EE,

Slatkin M, Reich D, Kelso J, Pääbo S. 2012. A High-Coverage Genome Sequence from an

Archaic Denisovan Individual. *Science* 338:222-226. doi: 10.1126/science.1224344.

57. Lazaridis I, Patterson N, Mittnik A, Renaud G, Mallick S, Kirsanow K, Sudmant PH,

Schraiber JG, Castellano S, Lipson M, Berger B, Economou C, Bollongino R, Fu Q, Bos

KI, Nordenfelt S, Li H, de Filippo C, Prüfer K, Sawyer S, Posth C, Haak W, Hallgren F,

Fornander E, Rohland N, Delsate D, Francken M, Guinet J-M, Wahl J, Ayodo G, Babiker

HA, Bailliet G, Balanovska E, Balanovsky O, Barrantes R, Bedoya G, Ben-Ami H, Bene J,

Berrada F, Bravi CM, Brisighelli F, Busby GBJ, Cali F, Churnosov M, Cole David EC,

Corach D, Damba L, van Driem G, Dryomov S, Dugoujon J-M, Fedorova SA, Romero IG,

Gubina M, Hammer M, Henn BM, Hervig T, Hodoglugil U, Jha AR, Karachanak-

Yankova S, Khusainova R, Khusnutdinova E, Kittles R, Kivisild T, Klitz W, Kučinskas V,

Kushniarevich A, Laredj L, Litvinov S, Loukidis T, Mahley RW, Melegh B, Metspalu E,

Molina J, Mountain J, Näkkäläjärvi K, Nesheva D, Nyambo T, Osipova L, Parik J,

Platonov F, Posukh O, Romano V, Rothhammer F, Rudan I, Ruizbakiev R, Sahakyan H,

Sajantila A, Salas A, Starikovskaya EB, Tarekegn A, Toncheva D, Turdikulova S,

Uktveryte I, Utevska O, Vasquez R, Villena M, Voevoda M, Winkler CA, Yepiskoposyan

L, Zalloua P, Zemunik T, Cooper A, Capelli C, Thomas MG, Ruiz-Linares A, Tishkoff

SA, Singh L, Thangaraj K, Villems R, Comas D, Sukernik R, Metspalu M, Meyer M,

Eichler EE, Burger J, Slatkin M, Pääbo S, Kelso J, Reich D, Krause J. 2014. Ancient

human genomes suggest three ancestral populations for present-day Europeans. *Nature*

513:409-413. doi: 10.1038/nature13673.

58. Keller A, Graefen A, Ball M, Matzas M, Boisguerin V, Maixner F, Leidinger P, Backes C,

Khairat R, Forster M, Stade B, Franke A, Mayer J, Spangler J, McLaughlin S, Shah M,

Lee C, Harkins TT, Sartori A, Moreno-Estrada A, Henn B, Sikora M, Semino O, Chiaroni

J, Rootsi S, Myres NM, Cabrera VM, Underhill PA, Bustamante CD, Vigl EE, Samadelli

M, Cipollini G, Haas J, Katus H, O'Connor BD, Carlson MRJ, Meder B, Blin N, Meese E,

Pusch CM, Zink A. 2012. New insights into the Tyrolean Iceman's origin and phenotype

as inferred by whole-genome sequencing. *Nature Communications* 3:A698. doi:

10.1038/ncomms1701.

59. Rasmussen M, Li YR, Lindgreen S, Pedersen JS, Albrechtsen A, Moltke I, Metspalu M,

Metspalu E, Kivisild T, Gupta R, Bertalan M, Nielsen K, Gilbert MTP, Wang Y,

Raghavan M, Campos PF, Kamp HM, Wilson AS, Gledhill A, Tridico S, Bunce M,

Lorenzen ED, Binladen J, Guo XS, Zhao J, Zhang XQ, Zhang H, Li Z, Chen MF, Orlando

L, Kristiansen K, Bak M, Tommerup N, Bendixen C, Pierre TL, Grønnow B, Meldgaard

M, Andreasen C, Fedorova SA, Osipova LP, Higham TFG, Ramsey CB, Hansen TVO,

Nielsen FC, Crawford MH, Brunak S, Sicheritz-Pontén T, Villems R, Nielsen R, Krogh A,

Wang J, Willerslev E. 2010. Ancient human genome sequence of an extinct Palaeo-

Eskimo. *Nature* 463:757-762. doi: 10.1038/nature08835.

60. Appenzeller T. 2013. Old masters. *Nature* 497: 302-304.

61. Wong K. 2015. Neandertal Minds. *Scientific American* 312:36-43. doi:

10.1038/scientificamerican0215-36.

62. Gabora L and Russon A. 2011. "The evolution of Intelligence". *In* "The Cambridge

Handbook of Intelligence". Edited by Sternberg RJ, Kaufman SB. p. 328-350.

63. Balter M. 2010. Did Working Memory Spark Creative Culture? *Science* 328:160-163.

doi: 10.1126/science.328.5975.160.

64. Crabtree GR. 2013. Our fragile intellect. Part I. *Trends in Genetics* 29:1-3. doi:

10.1016/j.tig.2012.10.002.

65. Haak W, Lazardis I, Patterson N, Rohland N, Mallick S, Llamas B, Brandt G, Nordenfelt

S, Harney E, Stewardson K, Fu Q, Mittnik A, Bánffy E, Economou C, Francken M,

Friederich S, Pena RG, Hallgren F, Khartanovich V, Khokhlov A, Kunst M, Kuznetsov P,

Meller H, Mochalov O, Moiseyev V, Nicklisch N, Pichler SL, Risch R, Guerra MAR,

Roth C, Szécsényi-Nagy A, Wahl J, Meyer M, Krause J, Brown D, Anthony D, Cooper A,

Alt KW, Reich D. 2015. Massive migration from the steppe was a source for Indo-

European languages in Europe. *Nature* 522:207-211. doi: 10.1038/nature14317.

66. Song M, Zhou Y, Li J, Liu Y, Tian L, Yu C, Tianzi, J. 2008. Brain spontaneous

functional connectivity and intelligence. *Neuroimage* 41:1168-1176. doi:

10.1016/j.neuroimage.2008.02.036.

67. Pamplona GSP, Neto GSS, Rosset SRE, Rogers BP, Salmon CEG. 2015. Analyzing the

association between functional connectivity of the brain and intellectual performance.

Frontiers in Human Neuroscience 9:A61. doi: 10.3389/fnhum.2015.00061.

68. Sarkissian CD, Allentoft ME, Ávila-Arcos MC, Barnett R, Campos PF, Cappellini E,

Ermini L, Fernández R, da Fonseca R, Ginolhac A, Hansen AJ, Jónsson H, Korneliussen T,

Margaryan A, Martin MD, Moreno-Mayar JV, Raghavan M, Rasmussen M, Velasco MS,

Schroeder H, Schubert M, Seguin-Orlando A, Wales N, Gilbert MTP, Willerslev E,

Orlando L. 2015. Ancient genomics. *Philosophical Transactions of the Royal Society B-*

Biological Sciences 370:20130387. doi: 10.1098/rstb.2013.0387.

69. Ohi K, Hashimoto R, Nakazawa T, Okada T, Yasuda Y, Yamamori H, Fukumoto M,

Umeda-Yano S, Iwase M, Kazui H, Yamamoto T, Kano M, Takeda M. 2012. The

p250GAP Gene Is Associated with Risk for Schizophrenia and Schizotypal Personality

Traits. PLoS One:e35696. doi: 10.1371/journal.pone.0035696.

70. McWilliam H, Li W, Uludag M, Squizzato S, Park YM, Buso N, Cowley AP, Lopez R.

2013. Analysis Tool Web Services from the EMBL-EBI. *Nucleic Acids Research* 41:597-

600. doi: 10.1093/nar/gkt376.

71. Prado-Martinez J, Sudmant PH, Kidd JM, Li H, Kelley JL, Lorente-Galdos B, Veeramah

KR, Woerner AE, O'Connor TD, Santpere G, Cagan A, Theunert C, Casals F, Laayouni H,

Munch K, Hobolth A, Halager AE, Malig M, Hernandez-Rodriguez J, Hernando-Herraez I,

Prüfer K, Pybus M, Johnstone L, Lachmann, M, Alkan C, Twigg D, Petit N, Baker C,

Hormozdiari F, Fernandez-Callejo M, Dabad M, Wilson ML, Stevison L, Camprubí C,

Carvalho T, Ruiz-Herrera A, Vives L, Mele M, Abello T, Kondova I, Bontrop RE, Pusey

A, Lankester F, Kiyang JA, Bergl RA, Lonsdorf E, Myers S, Ventura M, Gagneux P,

Comas D, Siegismund H, Blanc J, Agueda-Calpena L, Gut M, Fulton L, Tishkoff SA,

Mullikin JC, Wilson RK, Gut IG, Gonder MK, Ryder OA, Hahn BH, Navarro A, Akey JM,

Bertranpetit J, Reich D, Mailund T, Schierup MH, Hvilsom C, Andrés AM, Wall JD,

Bustamante CD, Hammer MF, Eichler EE, Marques-Bonet T. 2013. Great ape genetic

diversity and population history. *Nature* 499:471-475. doi: 10.1038/nature12228.

72. Green RE, Krause J, Briggs AW, Maricic T, Stenzel U, Kircher M, Patterson N, Li H,

Zhai W, Fritz MH-Y, Hansen NF, Durand EY, Malaspinas A-S, Jensen JD, Marques-

Bonet T, Alkan C, Prüfer K, Meyer M, Burbano HA, Good JM, Schultz R, Aximu-Petri A,

Butthof A, Höber B, Höffner B, Siegemund M, Weihmann A, Nusbaum C, Lander ES,

Russ C, Novod N, Affourtit J, Egholm M, Verna C, Rudan P, Brajkovic D, Kucan Ž,

Gušic I, Doronichev VB, Golovanova LV, Lalueza-Fox C, de la Rasilla M, Fortea J, Rosas

A, Schmitz RW, Johnson PLF, Eichler EE, Falush D, Birney E, Mullikin JC, Slatkin M,

Nielsen R, Kelso J, Lachmann M, Reich D, Pääbo S. 2010. A Draft Sequence of the

Neandertal Genome. *Science* 328:710-722. doi: 10.1126/science.1188021.

73. Prüfer K, Racimo F, Patterson N, Jay F, Sankararaman S, Sawyer S, Heinze A, Renaud G,

Sudmant PH, de Filippo C, Li H, Mallick S, Dannemann M, Fu Q, Kircher M, Kuhlwilm

M, Lachmann M, Meyer M, Ongyerth M, Siebauer M, Theunert C, Tandon A, Moorjani P,

Pickrell J, Mullikin JC, Vohr SH, Green RE, Hellmann I, Johnson PLF, Blanche H, Cann

H, Kitzman JO, Shendure J, Eichler EE, Lein ES, Bakken TE, Golovanova LV,

Doronichev VB, Shunkov MV, Derevianko AP, Viola B, Slatkin M, Reich D, Kelso J,

Pääbo S. 2013. The complete genome sequence of a Neanderthal from the Altai

Mountains. *Nature* 505:43–49. doi: 10.1038/nature12886.

Figure 1. Complementary effect of *NTNG* paralog SNPs on IQ cognitive domains of human subjects as measured by WAIS-III. (A-1, A-2) *NTNG1* and *NTNG2* gene structures with the SNPs location indicated. (B-1, B-2, C) Affected cognitive domains and scores. Red highlights *NTNG1* and green (or blue, in case of heterozygosity) – *NTNG2* located alleles associated with the attenuated IQ scores. The data are presented as a mean±SEM. *$p<0.05$,**$p<0.01$ (two ways ANCOVA (sex, education, and age as co-variates)). The number of human subjects is indicated as N. For statistical and diagnosis details refer to **ST1.**

Figure 2.

Figure 3.

Figures 2 and 3. Accelerated evolution (AE) and definition of a ROI of *NTNG* loci embedding associated with the cognitive endophenotype alleles. Hominins and primate-specific protein amino acid substitutions. (**A**) *NTNG1* and *NTNG2* SNPs' gene locations. Alleles associated with the lower IQ scores are shown either in red (*NTNG1*) or green (*NTNG2*). (**B**) Calculated identity percent of primates and mice to human gene loci as a function of distance from the position of a mutated allele (denoted as a star). Comparison was done in a stepwise manner as ±10 nu to the maximum of ±50 nu using Stretcher (http://www.ebi.ac.uk/Tools/psa (**70**)). The areas were compared based on the positioning relative to the point mutation without any manual curation for "the best-fit" alignment, but as per the algorithm output only. Due to low level of identity for mouse and marmoset the initial search of the mutation allele positioning was done aligning against the human corresponding full-length intron, and then second time against the obtained 101 nu query ("-50 nu-SNP +50 nu"). Two dashed lines define an area of -10 nu to +10 nu from the mutation allele position. This area (21 nu in total) is defined as a ROI of the given mutation allele of a representing SNP. An arrow indicates a position of extinct hominins-specific mutations (see below). (**C**) Evolutionary rates for the ROIs calculated as a percent identity change relative to the hg19 over the mln of years of evolution. The spectrum color reflects the mutations' positioning order on a gene, as purple-yellow for *NTNG1*, and yellow-blue for *NTNG2*. (**D**) ROIs DNA sequences across hominins, primates and mice loci. The extinct and ancient hominin's *NTNG* paralogs were reconstructed from the available datasets (see **SM**): Saqqaq-Eskimos (Es: **59**); Iceman (Ice: **58**); Hunters (Loschbour (Lo), Motala12 (Mo12), Motala3 (Mo3), Motala1: **57**); VindijaNea (Vi: **72**); Denisovan (De), AltaiNea (Al) and MezmaiskayaNea (Me): **73**; chimpanzee (Chi: reconstructed from (**71**), n=25 animals). Non-available sequences due to poor reads quality are denoted as X. Ensemble was used for the initial *NTNG* paralogs retrieval in marmoset (Ma) and mouse (Mo). Yellow (vertical strip) denotes the position of the SNP-related mutation allele and non-matched to human substitutions. (**E**) Evolution rates for the proteins encoded by the *NTNG1* (Netrin-G1) and *NTNG2* (Netrin-G2). (**F**) Amino acid changes for Netrin-Gs across hominins, primates and mice. For Netrin-G1, there are no common mutations among primates, hominins and modern human. Es contains 5 point mutations absent in other hominins; Chi contains 1 mutation outside the depicted area (A81S); all 4 mutations for marmoset are shown, and 10 more extra mutations for mouse are not shown. For Netrin-G2, all hominins except Ice (relative to hg19) carry a T346A mutation (exon 5-located and known as rs4962173), also detectable in primates and mouse. Chi's Netrin-G2 differs from all hominins by S371A mutation (exon 6 - nested) and present in marmoset as S371V. Non-matched to hg19 amino acids are highlighted as pink. Refer to **SM** for the full alignments.