MeSH annotation of the chicken genome: MeSH-informed enrichment analysis and MeSH-guided semantic similarity among functional terms and gene products

Gota Morota[*], Timothy M Beissinger[† ‡], and Francisco Peñagaricano[§ **]

[*]Department of Animal Science, University of Nebraska-Lincoln, Lincoln, Nebraska

[†]United States Department of Agriculture, Agricultural Research Service, Columbia, Missouri

[‡]Division of Plant Sciences, University of Missouri, Columbia, Missouri

[§]Department of Animal Sciences, University of Florida, Gainesville, Florida

[**]University of Florida Genetics Institute, University of Florida, Gainesville, Florida

Keywords: annotation, chicken, enrichment analysis, MeSH, semantic similarity

Running title: MeSH annotation of the chicken genome

Corresponding author:
Gota Morota
Department of Animal Science
University of Nebraska-Lincoln
PO Box 830908
Lincoln, NE 68583-0908, USA.
E-mail: morota@unl.edu

Abstract

Biomedical vocabularies and ontologies aid in recapitulating biological knowledge. The annotation of gene products is mainly accelerated by Gene Ontology (GO) and more recently by Medical Subject Headings (MeSH). Here we report the MeSH annotation of the chicken genome and illustrate some features of different MeSH-based analyses, including MeSH-informed enrichment analysis and MeSH-guided semantic similarity among terms and gene products, using two lists of chicken genes available in public repositories. The two published datasets that were employed represent (i) differentially expressed genes and (ii) candidate genes under selective sweep or epistatic selection. The comparison of MeSH with GO overrepresentation analyses suggested not only that MeSH supports the findings obtained from GO analysis but also that MeSH is able to further enrich the representation of biological knowledge and often provide more interpretable results. Based on the hierarchical structures of MeSH and GO, we computed semantic similarities among vocabularies as well as semantic similarities among selected genes. These yielded the similarity levels between significant functional terms, and the annotation of each gene yielded the measures of gene similarity. Our findings show the benefits of using MeSH as an alternative choice of annotation in order to draw biological inferences from a list of genes of interest, and we demonstrate that since it is based on keywords from published studies, it has the potential provide easily interpretable functional implications. We argue that the use of MeSH in conjunction with GO will be instrumental in facilitating the understanding of the genetic basis of complex traits.

Introduction

Understanding the genetic basis of variation for complex traits remains a fundamental goal of biology. Different approaches, including whole-genome scans and genome-wide expression studies, have been used in order to identify individual genes underlying economically relevant traits in a wide spectrum of agricultural species. These studies usually generate lists of genes potentially involved in the phenotypes under study. The challenge is to translate these lists of candidates genes into a better understanding of the biological phenomena involved. It is increasingly accepted that overrepresentation or enrichment analysis (Drăghici et al., 2003) can provide further insights into the biological pathways and processes affecting complex traits.

Recently, the Medical Subject Headings (MeSH) vocabulary (Nelson et al., 2004) has been proposed for defining functional sets of genes in the context of enrichment analysis. MeSH is a controlled life sciences vocabulary maintained by the National Library of Medicine to index documents in the MEDLINE database. Each bibliographic reference in the MEDLINE database is associated with a set of MeSH terms that describe the content of the publication. Importantly, MeSH contains a substantially more diverse and extensive range of categories than that of Gene Ontology (GO) (Ashburner et al., 2000), which is probably the most popular among the initiatives for defining functional classes of genes (Nakazato et al., 2008). Therein, GO terms are classified into three domains: biological processes, molecular functions, and cellular components. This ontology has been successfully used for dissecting relevant traits in livestock species (e.g, Peñagaricano et al., 2013; Gambra et al., 2013). Similarly, each MeSH term is clustered into 19 different categories; some MeSH categories, such as Diseases, are not included in GO, whereas other functional categories, such as Phenomena and Processes or Chemicals and Drugs, share similar concepts with those of GO. The recent availability of MeSH software packages has rendered agricultural species amenable to MeSH-based analysis (Tsuyuzaki et al., 2015). For instance, MeSH enrichment analysis has been successfully applied to dairy cattle, swine, and horse datasets (Morota et al., 2015). This study showed the potential of MeSH for enhancing the biological interpretation of sets of genes in these three domestic animals.

The main objective of the current study was to report for the first time the MeSH annotation of the chicken genome, and to illustrate the features of different MeSH-based analyses, including MeSH-informed enrichment analysis and MeSH-guided semantic similarity among terms and gene products. For this purpose, we used two lists of selected genes available in public repositories: (i) differentially expressed genes reported in a RNA-seq study (Zhuo et al., 2015) and (ii) candidate genes historically impacted by selection detected in a whole-genome scan using a broad spectrum of populations (Beissinger et al., 2015). The results of the MeSH-based enrichment analysis were contrasted with GO terms. The use of MeSH and GO terms in functional genomics studies was further explored through computing the similarity between significant functional terms as well as the similarity between significant genes by leveraging the hierarchies of these two controlled vocabularies.

Materials and Methods

We used two datasets from previously published studies with the objective of demonstrate some capabilities of different MeSH-based analyses in chicken. The first dataset includes 263 genes that showed differential expression in abdominal fat tissue between high and low feed efficiency broiler chickens (Zhuo et al., 2015). The second dataset contains 352 genes identified by a whole-genome scan using Ohta's between-population linkage disequilibrium measure, D_{IS}^2, in a panel that included 72 different chicken breeds (Beissinger et al., 2015). In both datasets, the list of background genes was defined as all annotated genes in the chicken genome available in NCBI.

The suite of MeSH (Tsuyuzaki et al., 2015) and the GOstats (Falcon and Gentleman, 2007) packages in Bioconductor were used for performing a hypergeometric test in the enrichment analysis. This test evaluates whether a given functional term or vocabulary is enriched or overrepresented with selected genes. In particular, the P-value of observing g significant genes in a functional term (i.e. MeSH or GO term) was calculated by

$$Pvalue = 1 - \sum_{i=0}^{g-1} \frac{\binom{S}{i}\binom{N-S}{k-i}}{\binom{N}{k}}$$

where S is the total number of selected genes, N is the total number of analyzed genes, and k is the total number of genes in the functional term under study. The hierarchical structures of MeSH and GO permitted us to compute semantic similarities between functional terms (Lord et al., 2003; Pesquita et al., 2009). This is a metric between two terms on the basis of their biological meanings of annotation: the closer two terms are in the hierarchy, the higher the similarity measure is between these terms. We employed the information content-based Jiang and Conrath's measure (Jiang and Conrath, 1998) to compute the pairwise similarities within GO ontologies and MeSH headings. The semantic similarity measure between two terms t_1 and t_2 is given by the information content $IC(t) = -\log p(t)$, where $p(t)$ is the probability of occurrence of the term t and its children terms in MeSH or GO hierarchy. The semantic distance metric is a function of

$$Dist = IC(t_1) + IC(t_2) - 2IC(MICA),$$

where MICA is the most informative common ancestor.

We further computed semantic similarity between selected genes by aggregating their MeSH or GO terms assigned. This is a similarity measure at the level of genes which is analogous to a similarity matrix among SNPs (Morota and Gianola, 2013). We calculated similarity scores over all pairs of terms between the two vocabulary sets of genes under consideration. All these GO and MeSH-guided semantic similarity analyses were carried out using the GOSemSim (Yu et al., 2010) and the MeSHSim (Zhou et al., 2015) Bioconductor packages, respectively. GO-based gene semantic similarity yielded category specific measures, whereas the MeSH counterpart produced a single measure by setting similarity to zero if two terms belong to different MeSH categories. For this reason, we selected exactly the same genes as were identified in GO categories when computing MeSH-based gene similarity to allow direct comparisons between these two functional vocabularies. Source code and reproducible output reports generated by R Markdown are available as Supporting Files.

Data Availability

The two datasets used in the current study have already been published. The gene expression data can be downloaded from http://journals.plos.org/plosone/article?id=10.1371/journal.pone.0135810#sec025. Raw data for the selective sweep data are available from http://dx.doi.org/10.6084/m9.figshare.1497961, and selected genes can be found in Beissinger et al. (2015).

Results

Summary of MeSH and GO annotations

The organism and the biomaRt Bioconductor packages were queried to annotate genes by MeSH and GO terms. Table 1 shows the total number of genes (background and selected genes) annotated by MeSH and GO in each of the datasets under study. Both MeSH and GO terms had a similar number of annotated known genes, whereas the number of selected genes with MeSH terms assigned was about one-half of that of GO. It is important to note that this difference could be because the majority of chicken genes are annotated by Inferred from Electronic Annotation (evidence code: IEA) in GO, whereas all MeSH terms are assigned by manual curation at NCBI. We expect that over time, MeSH will improve as new knowledge is created and published in the scientific literature.

Enrichment analysis

Gene Expression Data: A subset of significant MeSH terms (P-value ≤ 0.05) enriched with differentially expressed genes detected in fat tissue between high and low feed efficiency chickens are highlighted in Table 2. The majority of the MeSH terms in the Chemicals and Drugs category are related to lipid deposition and lipid metabolism. For instance, *Lipoproteins* (MeSH:D008074), and *Apolipoproteins* (MeSH:D001053) are closely related to lipid transportation. Additionally, *Fatty Acid-Binding Proteins* (MeSH:D050556) regulates diverse lipid signals, while *PPAR alpha* (MeSH:D047493) controls lipid and lipoprotein metabolism. Interestingly, many GO terms re-

lated to lipid deposition and metabolism, such as *cholesterol metabolic process* (GO:0008203), *high-density lipoprotein particle assembly* (GO:0034380), *spherical high-density lipoprotein particle* (GO:0034366), and *high-density lipoprotein particle binding* (GO:0008035), were also significantly enriched with differentially expressed genes (File S1). Similarly, MeSH terms related to Wnt proteins and signalling pathways, such as *Wnt Proteins* (MeSH:D051153), *Wnt4 Protein* (MeSH: D060528), *Wnt1 Protein* (MeSH:D051155), and their counterparts in GO, such as *regulation of Wnt signaling pathway* (GO:0030111) and *Wnt signaling pathway* (GO:0016055), were found as significant. The Wnt proteins are known to interact with lipids. We also found *Steroid 17-alpha-Hydroxylase* (MeSH:D013254) and *steroid 17-alpha-monooxygenase activity* (GO:0004508) as significant terms; these two categories are enriched in genes involved in the synthesis of lipids. Moreover, we detected some MeSH terms related to the immune system regulation (e.g., *Interleukin-6* (MeSH:D015850) and *Chemokines* (MeSH:D018925)). Lastly, *Glycoproteins* (MeSH:D006023), is produced from the gene *AHSG* and plays a role in glucose metabolism and the regulation of insulin signaling. Taken together, our findings confirm that MeSH enrichment analysis can either reinforce findings from GO or even bring an additional biological insight. Figure 1 depicts the semantic similarity between significant MeSH terms in the Chemicals and Drugs category. In general, this subset of MeSH terms showed low to high levels of semantic similarity.

For the Diseases category, which is unique to MeSH-based analysis, a subset of significant MeSH terms that deserves particular attention in the area of feed efficiency and lipid metabolism in poultry is highlighted in Table 2. For instance, *Hyperplasia* (MeSH:D006965) is a potential contributor to abdominal fat mass in broiler chickens; its relationship with *Diabetes Mellitus, Type 2* (MeSH:D003924) is well-documented in humans. Some MeSH terms directly related to the immune function, such as *Newcastle Disease* (MeSH:D009521) and *Inflammation* (MeSH:D007249), also showed a significant enrichment with differentially expressed genes. Interestingly, *Hyperplasia* and *Inflammation* showed a moderate semantic similarity according to the MeSH hierarchy (File S1).

Selective Sweep Data: Table 2 shows the results of the MeSH-informed enrichment analysis using genes putatively swept or under epistatic selection derived from a chicken diversity panel. Most of these terms are related to insulin metabolism. For instance, resistance to insulin occurs in birds due to high plasma glucose and fatty acid levels; this is supported by *Insulin Resistance*

(MeSH:D007333) in both the Diseases and Phenomena and Processes categories, as well as *Receptor, Insulin* (MeSH:D011972) and *Insulin* (MeSH:D007328) in the Chemicals and Drugs category. Moreover, we identified MeSH terms involved in the circadian clock of chicken. These are *Period Circadian Proteins* (MeSH:D056950), *CLOCK Proteins* (MeSH:D056926) and *ARNTL Transcription Factors* (MeSH:D056930) in Chemicals and Drugs, as well as *E-Box Elements* (MeSH:D024721), *Biological Clocks* (MeSH:D001683), and *Light* (MeSH:D008027) in Phenomena and Processes. Figure 2 shows the semantic similarities among MeSH terms in the Chemicals and Drugs category. Biological clock-related annotations, such as *Period Circadian Proteins* and *CLOCK Proteins*, exhibited moderate to high similarity. The results obtained from the other MeSH and GO categories were shown in File S2.

Gene semantic similarity

Gene Expression Data: Comparison of gene semantic similarity between MeSH and GO Biological Process for a subset of significant genes from the RNA-seq dataset is depicted in Figure 3. MeSH-based gene semantic similarity analysis showed that genes related to energy reserve metabolic process are highly related. For instance, genes that are involved in triacylglycerol and cholesterol biosynthesis, such as methylsterol monooxygenase 1 (*MSMO1*), insulin induced gene 1 (*INSIG1*), 1-acylglycerol-3-phosphate O-acyltransferase 9 (*AGPAT9*), and ADP ribosylation factor like GTPase 2 binding protein (*ARL2BP*), were highly similar to each other based on the MeSH hierarchy. Interestingly, GO-based analysis produced slightly different results; for instance, the gene *MSMO1* was highly similar to *INSIG1* but moderately similar to *AGPAT9* and *ARL2BP*. Additionally, genes *MSMO1* and *INSIG1* were moderately or highly related to lecithin-cholesterol acyltransferase (*LCAT*) and cytochrome b5 type A (microsomal) (*CYB5A*) based on the GO structure. These two genes, involved in lipid metabolism, also showed high similarity to apolipoprotein A-I (*APOA1*) and cytochrome P450, family 17, subfamily A, polypeptide 1 (*CYP17A1*). The relationship among these genes were low to moderate based on the MeSH hierarchy. The results based on the GO Molecular Function and Cellular Component categories were presented in File S3.

Selective Sweep Data: Gene semantic similarity based on both MeSH and GO Biological Process

among a subset of genes under selection is shown in Figure 4. Notably, a large group of genes, including strawberry notch homolog 1 (Drosophila) (*SBNO1*), ARP5 actin-related protein 5 (*ACTR5*), SET domain containing 1B (*SETD1B*), Obg-like ATPase 1 (*OLA1*), and histone deacetylase 9 (*HDAC9*) were highly related based on both MeSH and GO-guided semantic similarity analyses. All these genes are involved in chromatin organization and regulation of gene expression. Moreover, particular attention was paid to the top five candidates under epistatic selection reported by Beissinger et al. (2015). These genes are adenylate cyclase 5 (*ADCY5*), myosin light chain kinase (*MYLK*), phosphatidylinositol-4,5-bisphosphate 3-kinase, catalytic subunit beta (*PIK3CB*), calcium binding protein 39 (*CAG39*), and interleukin 1 receptor accessory protein (*IL1RAP*). Although none of these pair of genes appeared in a GO-based similarity matrix, *ADCY5* and *MYLK* presented a low to moderate gene semantic similarity based on the MeSH hierarchy (File S4).

Discussion

This article reports the MeSH annotation of the chicken genome. This new set of information enabled us to carry out different MeSH-based analyses, including enrichment analysis and MeSH-guided semantic similarity among functional terms and gene products. We exemplified the potential usefulness of these MeSH-based approaches by using two different publicly available chicken data.

The adipose tissue is the major site for lipid deposition and lipid metabolism, and it plays a central role in energy homeostasis. Unsurprisingly, several MeSH terms closely related to fat metabolism, such as Lipoproteins, Apolipoproteins, Fatty Acid-Binding Proteins, and PPAR alpha, were significantly enriched with genes that showed differential expression in fat tissue between high and low feed efficiency broiler chickens. Moreover, adipose tissue is now recognized as a metabolically active tissue that has important endocrine and immune regulatory functions (Kershaw and Flier, 2004). Interestingly, we found many significant MeSH terms, such as Interleukin-6, Chemokines, and Immunoglobulins, that are closely associated with the regulation of the immune function. Overall, our MeSH-based findings provide further insights into the biological mechanisms underlying differences in adiposity between high and low feed efficiency broiler chickens.

Included in our exemplary applications of MeSH annotations is a set of 352 genes previously iden-

tified as putatively affected by selection. Genes identified through population-genetic approaches such as this can be elusive, because their identification does not rely on phenotypes. Therefore associating selection with any specific trait is often very difficult (Akey, 2009). As we demonstrate in this study, tools such as GO and now MeSH are useful for suggesting biological interpretations that can later be followed up on or drive future biological hypotheses. For instance, our results showed that insulin-related MeSH terms appeared unusually often in the set of genes impacted by selection. This implies that selection for insulin-related traits may have played an important role in differentiating chicken breeds. Furthermore, our analysis involved testing for semantic similarity between pairs of genes, which was particularly useful for evaluating the most promising gene-pairs highlighted by Beissinger et al. (2015) as candidates for epistatic selection. Our expectation was that these pairs of genes are likely to be related to each other, as they have been predicted to be involved in the same selected phenotype. Our finding that one pair showed at least a weak semantic similarity may be interpreted as evidence that these two genes, *ADCY5* and *MYLK* are the most likely among the set to truly be epistatic.

The recent advancement in cataloguing genes with MeSH and GO has made it possible to assess the role of selected genes and has opened new opportunities for genetic research. Enrichment analysis recapitulates a set of genes into higher-level biological features. We argue that obtaining a complete picture of genes of interest using MeSH and GO is an important initial step toward functional genomics studies in poultry as well as other agricultural species as it facilitates efforts to illuminate the genetic basis of phenotypic variation.

Acknowledgements

This work was supported in part by the University of Nebraska Layman Fund to G.M. Support for T.M.B. was provided by USDA-ARS. F.P. acknowledges financial support from the Florida Agricultural Experiment Station and the Department of Animal Sciences, University of Florida.

References

Akey, J. M. (2009). Constructing genomic maps of positive selection in humans: Where do we go from here? *Genome Res.*, 19:711–722.

Ashburner, M., Ball, C. A., Blake, J. A., Botstein, D., Butler, H., Cherry, J. M., Davis, A. P., Dolinski, K., Dwight, S. S., Eppig, J. T., Harris, M. A., Hill, D. P., Issel-Tarver, L., Kasarskis, A., Lewis, S., Matese, J. C., Richardson, J. E., Ringwald, M., Rubin, G. M., and Sherlock, G. (2000). Gene Ontology: tool for the unification of biology. *Nat Genet.*, 25:25–29.

Beissinger, T. M., Gholami, M., Erbe, M., Weigend, S., Weigend, A., de Leon, N., Gianola, D., and Simianer, H. (2015). Using the variability of linkage disequilibrium between subpopulations to infer sweeps and epistatic selection in a diverse panel of chickens. *Heredity*, Advance online.

Drăghici, S., Khatria, P., Martinsb, R. P., Ostermeier, G. C., and Krawetz, S. A. (2003). Global functional profiling of gene expression. *Genomics*, 81:98–104.

Falcon, S. and Gentleman, R. (2007). Using GOstats to test gene lists for GO term association. *Bioinformatics*, 23:257–258.

Gambra, R., Peñagaricano, F., Kropp, J., Khateeb, K., Weigel, K. A., Lucey, J., and Khatib, H. (2013). Genomic architecture of bovine κ-casein and β-lactoglobulin. *J Dairy Sci.*, 96:5333–5343.

Jiang, J. and Conrath, D. (1998). Semantic similarity based on corpus statistics and lexical taxonomy. In *Proceedings of the 10th International Conference on Research in Computational Linguistics*, Taiwan.

Kershaw, E. E. and Flier, J. S. (2004). Adipose tissue as an endocrine organ. *J Clin Endocrinol Metab.*, 89:2548–2556.

Lord, P. W., Stevens, R. D., Brass, A., and Goble, C. A. (2003). Investigating semantic similarity measures across the Gene Ontology: the relationship between sequence and annotation. *Bioinformatics*, 19:1275–1283.

Morota, G. and Gianola, D. (2013). Evaluation of linkage disequilibrium in wheat with an L1-regularized sparse Markov network. *Theor Appl Genet.*, 126:1991–2002.

Morota, G., Peñagaricano, F., Petersen, J. L., Ciobanu, D. C., Tsuyuzaki, K., and Nikaido, I. (2015). An application of MeSH enrichment analysis in livestock. *Anim Genet.*, 46:381–387.

Nakazato, T., Takinaka, T., Mizuguchi, H., Matsuda, H., Bono, H., and Asogawa, M. (2008). Biocompass: a novel functional inference tool that utilizes MeSH hierarchy to analyze groups of genes. *In Silico Biol.*, 8:53–61.

Nelson, S. J., Schopen, M., Savage, A. G., Schulman, J. L., and Arluk, N. (2004). The MeSH translation maintenance system: structure, interface design, and implementation. *Stud. Health Technol. Inform*, 107:67–69.

Peñagaricano, F., Weigel, K. A., Rosa, G. J. M., and Khatib, H. (2013). Inferring quantitative trait pathways associated with bull fertility from a genome-wide association study. *Front Genet.*, 3:307.

Pesquita, C., Faria, D., Falcão, A. O., Lord, P., and Couto, F. M. (2009). Semantic similarity in biomedical ontologies. *PLoS Comput. Biol.*, 5:e1000443.

Tsuyuzaki, K., Morota, G., Ishii, M., Nakazato, T., Miyazaki, S., and Nikaido, I. (2015). MeSH ORA framework: R/Bioconductor packages to support MeSH over-representation analysis. *BMC Bioinformatics*, 16:45.

Yu, G., Li, F., Qin, Y., Bo, X., Wu, Y., and Wang, S. (2010). GOSemSim: an R package for measuring semantic similarity among GO terms and gene products. *Bioinformatics*, 26:976–978.

Zhou, J., Shui, Y., Peng, S., Li, X., Mamitsuka, H., and Zhu, S. (2015). MeSHSim: An R/Bioconductor package for measuring semantic similarity over MeSH headings and MEDLINE documents. *J. Bioinform. Comput. Biol*, Online Ready.

Zhuo, Z., Lamont, S. J., Lee, W. R., and Abasht, B. (2015). RNA-seq analysis of abdominal fat reveals differences between modern commercial broiler chickens with high and low feed efficiencies. *PLoS ONE*, 10:e0135810.

Supporting Information

- File S1: MeSH over-representation analysis (RNA-seq data)

- File S2: MeSH over-representation analysis (Selective sweep data)

- File S3: Gene Semantic Similarity (RNA-seq data)

- File S4: Gene Semantic Similarity (Selective sweep data)

 # Tables

Table 1: Number of known and selected genes annotated by MeSH (Medical SubjectHeadings) and GO (Gene Ontology).

	Annotated Genes		Selected Genes		
Data	MeSH	GO	Total	MeSH	GO
RNA-seq	10227	12460	263	110	245
Selective Sweep			352	145	333

Table 2: A subset of statistically signicant MeSH (Medical Subject Headings) terms. Background and Selected denote the number of background genes and selected genes annotated by the MeSH term, respectively.

Data	Category	MeSH ID	Background	Selected	MeSH Term
RNA-seq	Chemicals and Drugs	D008074	14	4	*Lipoproteins*
		D001054	7	2	*Apolipoproteins A*
		D001053	5	2	*Apolipoproteins*
		D050556	17	3	*Fatty Acid-Binding Proteins*
		D047493	7	2	*PPAR alpha*
		D012177	6	2	*Retinol-Binding Proteins*
		D051153	91	8	*Wnt Proteins*
		D060528	8	3	*Wnt4 Proteins*
		D051155	19	2	*Wnt1 Proteins*
		D015850	25	4	*Interleukin-6*
		D018925	14	2	*Chemokines*
		D007136	76	5	*Immunoglobulins*
		D013254	1	1	*Steroid 17-alpha-Hydroxylase*
		D006023	120	15	*Glycoproteins*
	Diseases	D006965	1	1	*Hyperplasia*
		D003924	2	1	*Diabetes Mellitus, Type 2*
		D009521	9	3	*Newcastle Disease*
		D014802	5	2	*Vitamin A Deficiency*
		D007249	12	2	*Inflammation*
Sweeps	Chemicals and Drugs	D011972	2	8	*Receptor, Insulin*
		D007328	26	3	*Insulin*
		D056950	5	2	*Period Circadian Proteins*
		D056926	8	2	*CLOCK Proteins*
		D056930	6	2	*ARNTL Transcription Factors*
	Diseases	D007333	1	1	*Insulin Resistance*
	Phenomena and Processes	D007333	1	1	*Insulin Resistance*
		D024721	8	2	*E-Box Elements*
		D001683	13	2	*Biological Clocks*
		D008027	28	3	*Light*

 # Figures

Figure 1: MeSH semantic similarity for the RNA-seq dataset.

Figure 2: MeSH semantic similarity for the selective sweep dataset.

Figure 3: Gene semantic similarity for the RNA-seq dataset. Top:MeSH, Bottom:GO

Figure 4: Gene semantic similarity for the selective sweep dataset. Top:MeSH, Bottom:GO