An Ancestry Based Approach for Detecting Interactions

Danny S. Park[1*], Itamar Eskin[2], Eun Yong Kang[3], Eric R. Gamazon[4,5], Celeste Eng[6], Christopher R. Gignoux[1,7], Joshua M. Galanter[6], Esteban Burchard[1,6], Chun J. Ye[8], Hugues Aschard[9], Eleazar Eskin[3], Eran Halperin[2], Noah Zaitlen[1,6*]

Affiliations:
1. Department of Bioengineering and Therapeutic Sciences. University of California San Francisco. San Francisco, CA.
2. The Blavatnik School of Computer Science. Tel-Aviv University. Tel Aviv, Israel.
3. Department of Computer Science. University of California Los Angeles. Los Angeles, CA.
4. Division of Genetic Medicine, Department of Medicine. Vanderbilt University. Nashville, TN.
5. Academic Medical Center, University of Amsterdam, Amsterdam, The Netherlands
6. Department of Medicine. University of California San Francisco. San Francisco, CA.
7. Department of Genetics. Stanford University. Palo Alto, CA.
8. Institute of Human Genetics. University of California San Francisco. San Francisco, CA.
9. Department of Epidemiology. Harvard School of Public Health. Boston, MA.

* Corresponding Author

Email: danny.park@ucsf.edu, noah.zaitlen@ucsf.edu

I. Abstract

Background: Epistasis and gene-environment interactions are known to contribute significantly to

variation of complex phenotypes in model organisms. However, their identification in human

association studies remains challenging for myriad reasons. In the case of epistatic interactions, the

large number of potential interacting sets of genes presents computational, multiple hypothesis

correction, and other statistical power issues. In the case of gene-environment interactions, the lack

of consistently measured environmental covariates in most disease studies precludes searching for

interactions and creates difficulties for replicating studies.

Results: In this work, we develop a new statistical approach to address these issues that leverages

genetic ancestry in admixed populations. We applied our method to gene expression and methylation

data from African American and Latino admixed individuals respectively, identifying nine

interactions that were significant at $p < 5 \times 10^{-8}$, we show that two of the interactions in methylation

data replicate, and the remaining six are significantly enriched for low p-values ($p < 1.8 \times 10^{-6}$).

Conclusion: We show that genetic ancestry can be a useful proxy for unknown and unmeasured

covariates in the search for interaction effects. These results have important implications for our

understanding of the genetic architecture of complex traits.

Keywords: Gene-environment interaction, gene-gene interactions, admixture

II. Background

Genetic association studies in humans have focused primarily on the identification of

additive SNP effects through marginal tests of association. There is growing evidence that both

epistatic and gene-environment $(G \times E)$ interactions contribute significantly to phenotypic variation

in humans and model organisms[1-5]. In addition to explaining additional components of missing

heritability, interactions lend insights into biological pathways that regulate phenotypes and improve

our understanding of their genetic architectures. However, identification of interactions in human

studies has been complicated by the computational and multiple testing burden in the case

of epistatic interactions, and the lack of consistently measured environmental covariates in the case

of $G \times E$ interactions[6,7].

To overcome these challenges, we leverage the unique nature of genomes from recently

admixed populations such as African Americans, Latinos, and Pacific Islanders. Admixed genomes are

mosaics of different ancestral segments[8] and for each admixed individual it is possible to

accurately estimate θ, the proportion of ancestry derived from each ancestral population (e.g. the

fraction of European/African ancestry in African Americans)[9]. Ancestry has been previously

leveraged to demonstrate that an array of environmental and biomedical covariates are correlated

with θ [10-20] and we therefore consider its use as a surrogate for unmeasured and unknown

environmental exposures. θ is also correlated with the genotypes of SNPs that are differentiated

between the ancestral populations, suggesting that θ may be effectively used as a proxy for detecting

multi-way epistatic interactions. Therefore, we propose a new SNP by θ test of interaction in order to

detect evidence of interaction in admixed populations.

We first investigate the properties of our method through simulated genotypes and

phenotypes of admixed populations. In our simulations we demonstrate that differential linkage-

disequilibrium (LD) between ancestral populations can produce false positive SNP by θ interactions

when local ancestry is ignored. To accommodate differential LD, we include local ancestry in our

statistical model and demonstrate that this properly controls this confounding factor. We also show

that our approach, the Ancestry Test of Interaction with Local Ancestry (AITL), is well-powered to

detect $G \times E$ interactions when θ is correlated with the environmental covariates of interest and

multi-way epistatic interactions. The power for detecting pairwise $G \times G$ interactions at highly

differentiated SNPs is lower than direct interaction tests even after accounting for the additional

multiple testing burden. However, the results of our simulations show that AITL is well powered to

detect multi-way epistasis involving tens or hundreds of SNPs of small effects, not detectable by

pairwise tests.

We first examined molecular phenotypes by applying our method to gene expression data

from African Americans, as well as DNA methylation data from Latinos. Gene expression traits have

previously been shown to have large-scale differences as a function of genetic ancestry[13]. Other

molecular phenotypes, such as LDL levels, have also been shown to be associated with genetic

ancestry [13,16,21-24]. For gene expression in particular, Price *et al.* showed that the effects of

ancestry on expression are widespread and not restricted to a handful of genes. Additionally,

molecular phenotypes are often used in deep phenotyping and Mendelian randomization studies and

are thus directly relevant to elucidating disease biology[25,26].

We identified one genome-wide significant interaction ($p < 5 \times 10^{-8}$) associated with gene

expression in the African Americans and eight significant interactions ($p < 5 \times 10^{-8}$) associated with

methylation in the Latinos. Two of the eight interactions associated with DNA methylation in the

Latinos also replicated and the remaining six were enriched for low p-values ($p < 1.8 \times 10^{-6}$). To

demonstrate that our approach works in larger data sets we also applied AITL to asthma case-control

data from Latinos and observed well-calibrated test statistics. Together, these results provide

evidence for the existence of interactions regulating expression and methylation and show that our

approach is statistically sound.

III.   Results

Simulated Data

To determine the utility of using θ as a proxy for unmeasured and unknown environmental

covariates, we applied the AITL to simulated 2-way admixed individuals. We tested θ_1, the

proportion of ancestry from ancestral population 1, for interaction with simulated SNPs (see

Simulation Framework). Power was computed over 1,000 simulations, assuming 10,000 SNPS being

tested, and using a Bonferroni correction p-value cutoff of 5×10^{-6}. We calculated the power using

assumed interaction effect sizes (either $\beta_{G\times G}$ or $\beta_{G\times E}$) of 0.1, 0.2, 0.3, and 0.4 (see Simulation

Framework). Although the few interactions reported for human traits and diseases have smaller

effects in terms of the phenotypic variance they explain, we simulated large effects because genetic

and environmental effect sizes in omics data, such as the expression and methylation data considered

here, are known to be of larger magnitude. For example, some cis-eQTL SNPs explain up to 50% of

the variance of gene expression[27]. However for most phenotypes, known interactions will explain a

very small proportion of the phenotypic variance, mainly due to the fact that so few interactions have

been identified and replicated[28].

Power When Using θ as a Proxy for Highly Differentiated SNPs

To determine whether using θ as a proxy for highly differentiated SNPs is more powerful

than testing all pairs of potentially interacting SNPs directly, we simulated two interacting SNPS in

1000 admixed individuals (see Simulation Framework). We then tested for an interaction using AITL

by replacing the genotypes at the highly differentiated SNP with $\vec{\theta}_1$. We observed that even with

moderate effect sizes, using θ in place of the actual genotypes does not provide any increase in power

even after accounting for multiple corrections (see Figure 1a). This is in agreement with recent work

showing the limited utility of local ancestry by local ancestry interaction test to identify underlying

SNP by SNP interaction when genotype data are available[29]. For the larger effect sizes we

simulated, we do see power increasing as the delta between ancestral frequencies increases. The

plots show that AITL has little power unless the effect was very strong. Figure 1b reveals that even

with the multiple correction penalty, testing all pairwise SNPS directly is always more powerful. We

note that when testing the interacting SNPs directly, we used a cutoff p-value of 1×10^{-9} since in

theory we were testing all unique pairs of 10,000 SNPs. Based on these results, we would

recommend testing for pairs of interacting SNPs directly if pairwise $G{\times}G$ interactions are a subject of

interest in the study.

127   However, when multi-way interactions are considered, AITL may become more powerful

since differentiated SNPs across the genome will be correlated with genetic ancestry. These

simulations are important as other studies have suggested that higher order interactions may be

important for some traits[1,30,31]. To evaluate the ability of θ to serve as a proxy for multiple

(independent) differentiated SNPs, we simulated a scenario where a candidate SNP z had

interactions with m SNPs (see Simulation Framework). For each interaction, we assumed a small

interaction effect size ($\beta_{G{\times}G} = 0.025$), which would not be detectable using a pairwise approach, as

we demonstrated in the pairwise simulation. Figure 2 shows that AITL is better powered to detect

the existence of interactions than a pairwise approach in the presence of multiple interacting SNPs

with a candidate SNP.

Power When Using θ as a Proxy Environmental Covariate

139   When assessing the utility of θ as a proxy for an environmental covariate E, we simulated

3000 individuals. E was simulated such that it was correlated with the global ancestries in varying

degrees (see Simulation Framework). Figure 3 shows the power of the AITL as a function of the

Pearson correlation between $\vec{\theta}_1$ and E. The power of testing E directly is exactly the power of the

AITL when the correlation is equal to 1. As expected, as the correlation increases, the power

increases as well. When the effect size is 0.1, the power to detect a $G{\times}E$ interaction is low whether

one uses θ_1 or E. However, both tests are much better powered for effect sizes greater or equal to 0.2,

with the AITL's power being dependent on the level of correlation. Note that using θ as a proxy for E

is equivalent to testing GxE in the presence of measurement error. Under the assumption of non-

differential error with regard to the outcome (e.g. the correlation between θ and E is equal among

cases and control) such a test is underpowered but has a controlled type I error rate under the

null[32].

Differential LD

To demonstrate that differential LD has the potential to cause inflated test-statistics, we ran

10,000 simulations of 1000 admixed individuals. For each individual we simulated 2 SNPs, a causal

SNP and a tag SNP. The LD between the tag SNP and causal SNP was different based on the ancestral

background the SNPs were on (see Simulation Framework). Over 10,000 simulations, we computed

the mean χ_1^2 test-statistic for the AIT and the AITL. We note that the phenotypes for these

simulations were generated under a model that assumed no interaction. We observed a mean $\chi_1^2 =$

0.996 with a standard deviation of 1.53 for AITL. AIT, which does not condition on local ancestry, had

a mean $\chi_1^2 = 3.59$ with a standard deviation of 3.60. We also looked at genomic control λ_{GC}, the ratio

of the observed median χ^2 over the expected median χ^2 under the null[33]. λ_{GC} compares the

median observed χ^2 test-statistic versus the true median under the null. In our simulations, we

observed $\lambda_{GC} = 5.81$ for AIT and $\lambda_{GC} = 0.980$ for AITL (see Supplementary Figure S1). Last, we

computed the proportion of test-statistics that passed a p-value threshold of .05 and .01 in our

simulations. The AIT had 3687 statistics passing a p-value of .05 and 1687 at a threshold of .01,

whereas AITL had 464 and 96 at the same p-value thresholds. The results for AITL are as expected

under a true null. The results from our simulations show that not accounting for local ancestry can

result in inflated test-statistics and can potentially lead to false positive findings.

Real Data

Coriell Gene Expression Results

We first applied our method to the Coriell gene expression dataset[34]. The Coriell cohort is

composed of 94 African-American individuals and the gene expression values of ~8800 genes in

lymphoblastoid cell lines (LCLs). Since African Americans derive their genomes from African and

European ancestral backgrounds, we tested for interaction between a given SNP and the proportion

of European ancestry, θ_{EUR}. Each SNP by θ_{EUR} term was tested once for association with the

expression of the gene closest to the SNP. We observed well-calibrated statistics with a λ_{GC} equal to

1.04 (see Supplementary Figure S2). In the LCLs, we found that interaction of rs7585465 with θ_{EUR}

was associated with ERBB4 expression (AITL $p = 2.95\times10^{-8}$, marginal $p = 0.404$) at a genome-wide

significant threshold ($p \leq 5\times10^{-8}$). rs7585465 has a 'C' allele frequency of 0.218 in the Corriell data

and appears to be differentiated between CEU and YRI with allele frequencies of 0.619 and 0.097 in

the respective populations.

Given that the gene expression values come from LCLs (all cultured according to the same

standards), the SNPs may be interacting with epigenetic alterations due to environmental exposures

that have persisted since transformation into LCLs. This scenario is unlikely, and we believe that

signals are driven by multi-way epistatic interactions. In our simulations, we showed that using θ as

a proxy for a single highly differentiated SNP is underpowered compared to testing all pairs of

potentially interacting SNPs directly. However, there are many SNPs that are highly differentiated

across the genome with which θ will be correlated. It is therefore possible that θ is capturing the

interaction between the aggregate of many differentiated trans-SNPs (i.e. global genetic background)

and the candidate SNP. This is consistent with a recently reported finding, conducted in human iPS

cell lines, that genetic background accounts for much of the transcriptional variation[2,35].

Although we believe the ERBB4 result to be representative of multi-way epistasis, we

performed a standard pairwise interaction test (see Methods) to check for interaction between

rs7585465 and other SNPs genome-wide. Interestingly, we found that the standard interaction test

(see Methods) showed substantial departure from the null with a λ_{GC} equal to 1.8 (see

Supplementary Figure S3). Since the interaction of rs7585465 by θ was significant, the pairwise

interaction test-statistics of rs7585465 by any SNP j can be inflated if j is correlated with θ. We found

that including the original significant SNP by θ term in the null (see Methods) brought the λ_{GC} down

to 1.05, and controlled for such scenarios in this dataset (See Supplementary Figure S3). As we had

previously anticipated, identifying the exact interactions driving the SNP by θ interaction proved to

be difficult. We found one borderline significant SNP (rs4839709, $p = 3.08 \times 10^{-7}$) but no

interactions that passed genome-wide significance. These results are consistent with what we have

observed in simulations, in which even though a standard pairwise interaction test is underpowered

to detect interactions, AITL is able to identify the main locus involved in a multi-way interaction.

206
207

GALA II Case-Control

To determine if our method is biased in large structured GWAS data, we applied AITL to

case-control data from a study of asthmatic Latino individuals called the Genes-environments and

Admixture in Latino Americans (GALA II)[36]. The dataset includes 1158 Mexicans and 1605 Puerto

Ricans, which were analyzed separately. Case status was assigned to individuals if they were

between the ages of 8 and 40 years with a physician-diagnosed mild to moderate-to-severe asthma.

Additionally, they had to have experienced 2 or more asthma related symptoms in the previous 2

years at the time of recruitment[37]. In the Mexicans and Puerto Ricans there were 548 and 797

cases, respectively. In our analysis, we also included BMI, age, and sex as additional covariates. We

observed well-calibrated statistics with a λ_{GC} equal to 1.00 and 0.98 in the Mexicans and Puerto

Ricans, respectively (see Supplementary Figure S5). In contrast to the molecular phenotype data,

searches for interactions in these phenotypes did not yield any findings passing genome-wide

significance. This is consistent with previous disease studies that have failed to find many replicable

interactions in disease studies[28]. In the data here, the lack of any findings may be due to the

relatively small sample size or because the effects of the interactions are extremely small (if they

exist for covariates correlated with θ_{EUR}).

GALA II Methylation Results

We searched for interactions in methylation data derived from a study of GALA II asthmatic

Latino individuals[36]. The methylation data is composed of 141 Mexicans and 184 Puerto Ricans. As

the phenotype, we used DNA methylation measurements on ~300,000 markers from peripheral

blood. As we had done with gene expression, we tested for interaction between a given SNP and θ_{EUR}

using AITL. All SNPs within a 1 MB window centered around the methylation probe were tested. We

used the European component of ancestry because it is the component shared most between

Mexicans and Puerto Ricans (see Table 1). We observed well-calibrated test-statistics with λ_{GC} equal

to 1.06 in the Mexicans and 0.96 in the Puerto Ricans (see Supplementary Figure S6). We tested

128,794,325 methylation-SNP pairs, which result in a Bonferroni corrected p-value cutoff of

3.88×10^{-10}. However, this cutoff is extremely conservative given the tests are not independent. We

therefore report all results that are significant at 5×10^{-8} in either set as an initial filter. We found 5

interactions in the Mexicans and 3 in the Puerto Ricans that are significant at this threshold (see

Table 2).

Unlike the Coriell individuals, who are 2-way admixed, the GALA II Latinos are 3-way

admixed and derive their ancestries from European, African, and Native American ancestral groups.

Consequently, to confirm that incomplete modeling or better tagging on one of the non-European

ancestries was not driving the results, we retested all significant interactions including a second

component of ancestry for AITL. In the case of the Mexicans, we included African and European

ancestry, and in the case of the Puerto Ricans, we included European and Native American ancestry.

Even after adjusting for the second ancestry the interactions between SNP and θ_{EUR} remained highly

significant (see Supplementary Table 1).

As we did for the gene expression data, we attempted to identify pairwise interactions

involved in the methylation data results. For each genome-wide significant result, we performed a

standard pairwise interaction test of all SNPs with the original SNP found to be significant with AITL.

We were unable to identify any significant interactions after applying genomic control to the results.

For all tests, we included the significant SNP by θ term (see Methods) in the null. For this dataset,

unlike the gene expression data, we observed substantial remaining departure from the null (see

Supplementary Table S2) even after including the original significant SNP by θ term, suggesting there

may be other factors that need to be accounted for when testing for interactions in admixed

populations. The results from our pairwise scan are what we would anticipate, given that in

simulations only AITL (not the standard pairwise interaction test) was able to identify the main locus

involved in the multi-way interaction.

We then performed a replication study of the significant Puerto Rican associations in the

Mexican cohort and vice versa. To account for the fact that we are replicating eight total results

across both populations, we used a Bonferroni corrected p-value threshold equal to $.05/8 =$

6.25×10^{-3}. The interaction of rs4312379 and rs4312379 with ancestry in the Puerto Ricans

replicated in the Mexicans. Furthermore, there was a highly significant enrichment of low p-values in

the replication study among the discovery results (permutation $p < 1\times10^{-4}$). Furthermore, 5 out of

the 6 non-replicating results have a p-value less than 0.05 (binomial test $p < 1.8\times10^{-6}$). The results

of the permutation and binomial test suggests that the interactions that did not replicate are likely to

do so with bigger sample sizes. It is important to note that replicated interactions and the enrichment

for low p-values do not necessarily indicate that the same genetic or environmental covariates are

interacting with the genetic locus in both populations. The covariates correlated with θ_{EUR} in one

population are not necessarily those correlated with θ_{EUR} in the other population. There may be

correlations which exist in both populations but θ_{EUR} serves as a proxy for all such correlated

covariates and therefore should not be necessarily viewed as a proxy for any specific one. Overall,

our results from the GALA II (methylation) cohort suggest there are both genetic and environmental

variables contributing to epistasis that have yet to be discovered in admixed individuals.

IV.   Discussion and Conclusions

For many disease architectures, interactions are believed to be a major component of

missing heritability[38]. Finding new interactions has proven to be difficult for logistical, statistical,

biological, and computational reasons. In this study, we have demonstrated that in admixed

populations, testing for $G \times \theta$ interactions can be leveraged to overcome some of the difficulties

typically encountered when searching for interactions. The computational cost is minimal and has

the same order as running a standard GWAS.

One drawback of our method is that it does not identify which covariate is interacting with a

genetic locus. Nevertheless, the approach can show whether an interaction effect exists in a given

dataset and if it does exist, our method ensures that an underlying genetic or environmental

covariate(s) is correlated with ancestry. Additionally, in the case where there is no marginal effect,

our approach identifies new loci and shows that the genetic locus influences the phenotype and

exerts its effects through interactions, which has important implications for the genetic architecture

of the phenotype. The relative contribution of additive and non-additive genetic effects to variability

in molecular phenotypes and disease risk is an important area of investigation, and our approach

provides a direct test for detecting non-additive contributions[39].

Environmental covariates are often not consistently measured across cohorts whereas

genetic ancestry is nearly perfectly replicable. Testing for the presence of interaction using a nearly

perfectly reproducible covariate may enhance our understanding of the genetic basis of disease and

other traits. Our method also provides the additional benefit of not being confounded by interactions

between unaccounted-for covariates[40].

Association testing for interaction effects involving continuous environmental exposures in

the context of mixed-models remains an open problem. For binary environmental exposures, it has

been shown that mixed-models control for population structure nominally better than including

genetic ancestry (or principal components) as a covariate[41]. Because it is unclear how mixed-

models perform with continuous environmental exposures, especially those correlated with

ancestry, in our analyses we took the standard approach of filtering related individuals and including

ancestry as a covariate.

It has been shown that 2-step analyses may be more powerful for detecting interactions

when exposures are binary [42-44]. However, these studies have primarily been done in a single

homogeneous population, and the correct null distribution for the interaction effect must assume

that the 2nd stage procedure is independent of the marginal effect test-statistic. In real data, using a 2-

step approach in conjunction with AITL to test for interactions may be problematic because the

interaction effect size will not necessarily be independent of the marginal effect size, as the allele

frequency at any SNP will be a function of ancestry in an admixed population. Additionally, only 1 of

the interaction results that we report here had a marginal effect (p< 0.05) and thus would have been

missed by a 2-step approach. Thus, our approach can serve to complement or extend the frequently

used 2-step procedure for detecting interaction effects.

Results from our multi-way epistasis simulation analyses and empirical data in cell lines

suggest that genetic ancestry is a good proxy for genetic background, since all highly differentiated

SNPs across the genome will be correlated with genetic ancestry. Our simulations also demonstrated

that genetic ancestry can be a good proxy for an environmental covariate depending on the

correlation between the two. However, it may be the case that there are multiple environmental

factors interacting with a genetic locus, all of which are correlated with θ in differing degrees and

effect sizes. Such a situation would mirror what we saw in our multi-way $G \times G$ simulations where a

single interaction may not be detectable by using a traditional $G \times E$ test, but because θ aggregates the

effects of all interacting covariates, AITL would be able to detect it. There are also other contexts in

which modeling SNP by θ may be useful, such as using variance components. For example, SNP by θ

interaction terms can be used in a mixed-model framework to test for interaction effects because

genetic ancestry is correlated with many genetic markers and environmental covariates[45].

For some traits, there may be systematic differences between ancestral populations in the

genetic effects on the trait. In admixed individuals with these ancestral populations, the effect of

genetic variation on phenotype will be reflected in the correlation between phenotype and θ, thereby

affecting epistatic and $G{\times}E$ interactions. It will be interesting to see how much of the phenotype-

ancestry correlations are due to epistatic and $G{\times}E$ interactions.

In our analysis of real data, we discovered gene by θ interactions associated with genes that

have known interactions. In the GALA II Mexicans, the interaction of rs925736 with ancestry was

associated with the methylation of HDAC4, a known histone deaceytlase (HDAC). In concert with

DNA methylases, HDACs function to regulate gene expression by altering chromatin state[46]. In

Europeans, HDACs have been shown to be associated with lung function through direct genetic

effects and through environmental interactions[47,48]. For the GALA II Puerto Ricans, rs17091085

showed an interaction associated with the methylation state of SERPINA6. Of note, interaction

between birth weight and SERPINA6 has been previously associated with Hypothalamic-Pituitary-

Adrenal axis function[49]. Further investigations of our interaction findings are thus warranted.

In the GALA II (methylation) dataset, two of the eight significant associations replicated and,

in general, the results had an enrichment of low p-values in the replication dataset. However, we note

that if the interactions detected by AITL are multi-way epistasis it is more likely that the results will

replicate. This is because most SNPs differentiated in the Mexicans will still be differentiated in the

Puerto Ricans, and thus still be correlated with θ. If the interactions detected by AITL are $G{\times}E$

interactions, then the interactions are less likely to replicate because the same environmental

covariate(s) will need to be correlated with ancestry in both groups.

Another caveat is that the Mexicans and Puerto Ricans, though independent, are part of the

same study and occasionally technical artifacts, such as issues with genotyping or measuring

methylation, can affect downstream analyses of both populations. For our analyses, we have taken

careful quality-control steps to ensure that this is not the case and there is no apparent inflation of

test-statistics as demonstrated by our values for genomic control. Future research of interactions

using AITL should keep such caveats in mind.

We investigated in detail the potential of single SNP-SNP interactions driving the results that

were found both in the gene expression and methylation datasets. As demonstrated by the wide

range of λ_{GC} values, we observed that non-linear effects can cause substantial departure from the null

when testing for pairwise SNP-SNP interactions. This is especially true when testing for interaction

between SNPs s and j, where s has a significant interaction with θ and j is correlated covariates that

are also correlated with θ. As we saw in the gene expression data, including the significant SNP by θ

term can properly control for such situations, but its use in standard pairwise interaction tests

warrants further investigation.

Our analysis revealed the existence of interactions but does not provide a direct way to

determine the covariate that is interacting with a SNP. Further methodological work is required to

uncover the exact environmental exposures or genetic loci with which SNPs are interacting. The

existence of gene by θ interactions in GALA II underscores why modeling interactions should be

considered for future association studies and for heritability estimation in admixed populations.

V.   Materials and Methods

Our approach is best illustrated with an example. First consider testing a SNP s for

interaction with an environmental covariate E. θ can serve as a proxy for E if the two are correlated,

even if E is unknown or unmeasured (see Figure 4a). Now consider testing s for interaction with a

SNP $j \neq s$ that is highly differentiated in terms of ancestral allele frequencies. For example, a SNP that

has a high allele frequency in one ancestral population and a low allele frequency in the other

ancestral population. θ can be used as a proxy for j because θ and the genotypes of SNP j will be

correlated. Consider the case where j has a frequency of 0.9 in population 1 and frequency of 0.1 in

population 2. Individuals with large values of θ_1 are more likely to have derived j from population 1

and on average have greater genotype values at j. Similarly, individuals with small values of θ_1 are

more likely to have derived *j* from population 2 and on average have smaller genotype values. Thus, θ

will be correlated with the genotypes of the individuals for highly differentiated SNPs and can serve

as a proxy for detecting interactions (see Figure 4b).

379   Consider an admixed individual *i* who derives his or her genome from *k* ancestral

populations. We denote individual *i*'s global ancestry proportion as $\theta_i =$

$\langle \theta_{i1}, \theta_{i2}, \quad , \theta_{ik} \rangle$, where $\sum_k \theta_{ik} = 1$. The local ancestry of individual *i* at a SNP *s* is denoted as $\gamma_{ais} \in$

$\{0, 1, 2\}$ and is equal to the number of alleles from ancestry $a \in \{1 \ldots k\}$ inherited at SNP *s.* Current

methods allow us to estimate ancestry directly from genotype data both globally and at specific

SNPs[9,50,51]. We denote the genotype of an individual *i* at SNP *s* as $g_{is} \in \{0, 1, 2\}$ and the

corresponding phenotype as y_i.

386   In this work, we model continuous phenotypes in an additive linear regression framework.

Assuming *n* (unrelated) individuals, define $\vec{y}$ to be the vector of all individuals' phenotypes. The

model for the phenotype is then

$$\vec{y} = \boldsymbol{X}\vec{\beta} + \vec{\varepsilon}$$

where $\vec{\varepsilon} \sim \mathcal{N}(0, \sigma)$ is a *n*×1 vector of error terms, $\boldsymbol{X}$ is a *n*×*v* matrix of *v* covariates, and $\vec{\beta}$ is a *v*×*1*

vector of the covariate effect sizes. We note that in our notation $\vec{v}^2 = \vec{v}^T \vec{v}$ for a vector $\vec{v}$. Assuming

independence, the likelihood under this model is:

$$L = \left(\frac{1}{\sigma\sqrt{2\pi}}\right)^n \exp\left(-\frac{1}{2\sigma^2}(\vec{y} - \boldsymbol{X}\beta)^2\right)$$

We can compute the log-likelihood ratio statistic (*D*) using a maximum likelihood approach:

$$D = -2(\log L_1 - \log L_0) = -2\left(n\log(\hat{\sigma}_{L_1}) + \frac{(\vec{y} - \boldsymbol{X}\hat{\beta}_{L_1})^2}{2\hat{\sigma}_{L_1}^2}\right) + 2\left(n\log(\hat{\sigma}_{L_0}) + \frac{(\vec{y} - \boldsymbol{X}\hat{\beta}_{L_0})^2}{2\hat{\sigma}_{L_0}^2}\right)$$

We note that for a case-control phenotype we would use the following likelihood and log-likelihood

ratio statistic:

398
$$L = \prod_{i=1}^{n} \left[\frac{1}{1 + e^{-X_i\beta}}\right]^{y_i} \left[1 - \frac{1}{1 + e^{-X_i\beta}}\right]^{1-y_i}$$

399
$$D = -2\left(\log L_1 - \log L_0\right)$$

400
$$= -2\left(\sum_{i=1}^{n} -\log(1) + e^{-X_i\hat{\beta}_{L1}} + \sum_{i=1}^{n} y_i\left(X_i\hat{\beta}_{L_1}\right)\right)$$

401
$$+ 2\left(\sum_{i=1}^{n} -\log(1) + e^{-X_i\hat{\beta}_{L0}} + \sum_{i=1}^{n} y_i\left(X_i\hat{\beta}_{L_0}\right)\right)$$

where X_i is the i-th row of the matrix $\boldsymbol{X}$, which correspond to the covariates of individual i.

For linear regression, the maximum likelihood estimator (MLE) of the effect sizes is $\hat{\beta} =$

$(\boldsymbol{X}^T\boldsymbol{X})^{-1}\boldsymbol{X}^T\vec{y}$, and the MLE of the error variance is $\hat{\sigma}^2 = \frac{1}{n}\left(\vec{y} - \boldsymbol{X}\hat{\beta}\right)^2$. Here, L_1 is the likelihood under

the alternative and L_0 is the likelihood under the null. $(\hat{\beta}_{L_1}, \hat{\sigma}^2_{L_1})$ and $(\hat{\beta}_{L_0}, \hat{\sigma}^2_{L_0})$ are the effect sizes and

error variance estimates that maximize the respective likelihoods. D is distributed as χ^2 with k

degrees of freedom (df), where k is the number of parameters constrained under the null.

1-df Ancestry Interaction Test (AIT)

The first test we present is the standard direct test of interaction. We test for a SNP's

interaction with θ instead of an environmental covariate or another genotype. Let $\vec{g}_s = \langle g_{1s} \dots g_{ns}\rangle$ be

the vector of the individuals' genotypes at SNP s, $\vec{\theta}_a = \langle\theta_{1a} \dots \theta_{na}\rangle$ be the vector of their global

ancestries for ancestry a, and $\vec{g}_s \times \vec{\theta}_a$ be the vector of interaction terms which result from the

component-wise multiplication of the genotype and global ancestry vectors. We test the alternative

hypothesis $\left(\hat{\beta}_{G\times\theta} \neq 0\right)$ against the null hypothesis $\left(\hat{\beta}_{G\times\theta} = 0\right)$.

418
$$H_1: \vec{y} = \vec{g}_s + \vec{g}_s \times \vec{\theta}_a + \vec{\theta}_a$$

419
$$H_0: \vec{y} = \vec{g}_s + \vec{\theta}_a$$

In this test of interaction, we test a single ancestry versus the other ancestries that may be present in

the population of interest. One parameter is constrained under the null which results in a statistic

with $k=1$ df. Let $\hat{\beta}_{L_{\{0,1\}}(s)}$, $\hat{\beta}_{L_{\{0,1\}}(G \times \theta)}$, and $\hat{\beta}_{L_{\{0,1\}}(\theta)}$ denote the effect sizes of genotype, interaction, and

global ancestry under a given hypothesis respectively. The statistic is given below.

$$D = -2\left(n\log(\hat{\sigma}_{L_1}) + \frac{[\vec{y} - \boldsymbol{X}\langle\hat{\beta}_{L_1(s)}, \hat{\beta}_{L_1(G \times \theta)}, \hat{\beta}_{L_1(\theta)}\rangle]^2}{2\hat{\sigma}_{L_1}^2}\right) + 2\left(n\log(\hat{\sigma}_{L_0}) + \frac{[\vec{y} - \boldsymbol{X}\langle\hat{\beta}_{L_0(s)}, 0, \hat{\beta}_{L_0(\theta)}\rangle]^2}{2\hat{\sigma}_{L_0}^2}\right)$$

where $\boldsymbol{X}$ is an $n \times 3$ matrix composed of $\vec{g}_s$, $\vec{\theta}_a$, and $\vec{g}_s \times \vec{\theta}_a$ as columns.

1-df Ancestry Interaction Test with Local Ancestry (AITL)

Given that the individuals we analyze in this work are assumed to be admixed, there is

potential for confounding due to differential LD. An interaction that is not driven by biology could

occur due to the possibility that a causal variant may be better tagged by a SNP being tested on one

ancestral background versus another (See Figure 4c). We account for the different LD patterns on

varying ancestral backgrounds by including local ancestry as an additional covariate in AITL. By

including local ancestry, we assume that the SNP being tested is on the same local ancestry block as

the causal SNP that it may be tagging. Such an assumption is reasonable because admixture in

populations such as Latinos and African Americans are relatively recent events and their genomes

have not undergone many recombination events. As a result, local ancestry blocks on average stretch

for several hundred kilobases[52,53].

Let $\vec{\gamma}_{as} = \langle\gamma_{a1s} \ldots \gamma_{a1s}\rangle$ be the vector of local ancestry calls for all individuals for ancestry a

and let $\vec{g}_s \times \vec{\gamma}_{as}$ be the interaction terms from piecewise multiplication of the two vectors. We use the

following alternative and null hypotheses:

$$H_1: \vec{y} = \vec{g}_s + \vec{g}_s \times \vec{\theta}_a + \vec{\theta}_a + \vec{\gamma}_{as} + \vec{g}_s \times \vec{\gamma}_{as}$$

$$H_0: \vec{y} = \vec{g}_s + \vec{\theta}_a + \vec{\gamma}_{as} + \vec{g}_s \times \vec{\gamma}_{as}$$

Here we are testing for an interaction effect, i.e. $\hat{\beta}_{G \times \theta} \neq 0$, and constrain one parameter under the

null resulting in a statistic with $k=1$ *df*. Let $\hat{\beta}_{L_{\{0,1\}}(G \times \gamma)}$ and $\hat{\beta}_{L_{\{0,1\}}(\gamma)}$ denote the effect sizes of the

interaction between genotype and local ancestry and just local ancestry, respectively. The log

likelihood ratio statistic is given by

452
$$D = -2 \left(n \log(\hat{\sigma}_{L_1}) + \frac{\left[\vec{y} - X \langle \hat{\beta}_{L_1(s)}, \hat{\beta}_{L_1(G \times \theta)}, \hat{\beta}_{L_1(\theta)}, \hat{\beta}_{L_1(\gamma)}, \hat{\beta}_{L_1(G \times \gamma)} \rangle \right]^2}{2 \hat{\sigma}_{L_1}^2} \right)$$

453
$$+ 2 \left(n \log(\hat{\sigma}_{L_0}) + \frac{\left[\vec{y} - X \langle \hat{\beta}_{L_0(s)}, 0, \hat{\beta}_{L_0(\theta)}, \hat{\beta}_{L_0(\gamma)}, \hat{\beta}_{L_0(G \times \gamma)} \rangle \right]^2}{2 \hat{\sigma}_{L_0}^2} \right)$$

where X is an $n \times 5$ matrix composed of $\vec{g}_s$, $\vec{\theta}_a$, $\vec{g}_s \times \vec{\theta}_a$, $\vec{\gamma}_{as}$, and $\vec{g}_s \times \vec{\gamma}_{as}$ as columns. All of these test-

statistics are straightforwardly modified to jointly incorporate several ancestries in the case of multi-

way admixed populations.

Standard Pairwise Test of Interaction and Controlling Confounding in Admixed Populations

Here we present the standard approach for testing for interaction between two SNPs. We

use the following alternative and null hypotheses.

461
$$H_1: \vec{y} = \vec{g}_1 + \vec{g}_2 + \vec{g}_1 \times \vec{g}_2 + \vec{\theta}_a$$

462
$$H_0: \vec{y} = \vec{g}_1 + \vec{g}_2 + \vec{\theta}_a$$

463 If AITL is significant for a given SNP *s*, then any SNP *j* tested for interaction with *s* may be biased if *j* is

464 correlated with covariates that are also correlated with θ. Furthermore, if the effects of the covariates

465 correlated with θ are non-linear then controlling for the main effects of the SNPs and ancestry will

466 account for the non-linear effects. We thus, propose the use of the following alternative and null

467 hypotheses.

468
$$H_1: \vec{y} = \vec{g}_s + \vec{g}_j + \vec{g}_s \times \vec{g}_j + \vec{\theta}_a + \vec{g}_s \times \vec{\theta}_a$$

469
$$H_0: \vec{y} = \vec{g}_s + \vec{g}_j + \vec{\theta}_a + \vec{g}_s \times \vec{\theta}_a$$

We note that the utility of this test will require further investigation (see Discussion).

Simulation Framework

For all our simulations, we simulated 2-way admixed individuals. Global ancestry for

ancestral population 1 (θ_1) was drawn from a normal distribution with $\mu = 0.7$ and $\sigma = 0.2$.

Individuals with $\theta_1 > 1$ or $\theta_1 < 0$ were assigned a value of 1 or 0, respectively. We simulated

phenotypes of individuals to investigate our method in four different scenarios: $G{\times}E$ interactions,

pairwise epistatic interactions, multi-way epistatic interactions, and false positive interactions due to

local differential tagging.

To simulate phenotypes under the situation of a $G{\times}E$ interaction, we simulated a single SNP.

For each individual i, we assigned the local ancestry or the number of alleles derived from population

1 (γ_{ai}) for each haplotype by performing two binomial trials with the probability of success equal to

θ_{i1}. We then drew ancestry specific allele frequencies following the Balding-Nichols model by

assuming a $F_{ST} = 0.16$ and drawing two ancestral frequencies, p_1 and p_2, from the following beta

distribution[54].

488
$$p_1, p_2 \sim Beta\left(\frac{p(1 - F_{ST})}{F_{ST}}, \frac{(1 - p)(1 - F_{ST})}{F_{ST}}\right)$$

where p is the underlying MAF in the entire population and is set to 0.2. Genotypes were drawn using

a binomial trial for each local ancestry haplotype with the probability of success equal to p_1 or p_2 for

values of $\gamma_{ai} = 0$ or 1, respectively. Environmental covariates correlated with θ_1, E_i, were generated

for each individual i by drawing from a normal distribution $\mathcal{N}(\mu = \theta_{i1}, \sigma_E)$, where σ_E is the standard

deviation of the environmental covariates. σ_E was varied from 0 to 5 in increments of 0.005 to create

E_i's that were correlated with individuals' global ancestries in varying degrees. We generated

phenotypes for individuals assuming only an interaction effect by drawing from a normal

distribution, $\mathcal{N}(\mu = \beta_{G{\times}E} \times g_{i1} \times E_i, \sigma = 1)$ for a given interaction effect size ($\beta_{G{\times}E}$).

To simulate phenotypes based on pairwise epistatic interactions, we simulated two SNPs. At

both SNPs, we assigned the local ancestry values as described for the $G{\times}E$ case. We assigned

genotypes for individuals at the first SNP assuming an allele frequency of 0.5 for both populations

and drawing from two binomial trials. We assigned genotypes at the second SNP over a wide range of

ancestry specific allele frequencies to simulate different levels of SNP differentiation. Ancestry

specific allele frequencies were initially $p_1 = p_2 = 0.5$ and iteratively increasing p_1 by 0.005 while

simultaneously decreasing p_2 by 0.005 until $p_1 = 0.05$ and $p_2 = 0.95$. Genotypes at the second SNP

were drawn using the same approach described for $G{\times}E$. Using the simulated genotypes, phenotypes

were drawn from a normal distribution, $\mathcal{N}(\mu = \beta_{G\times G}\times g_{i1}\times g_{i2}, \sigma = 1)$, where g_{is} is the genotype for

individual i at the simulated SNP s.

To simulate phenotypes based on multi-way epistatic interactions, we simulated a SNP z and

m (independent) SNPs with pairwise interactions with z. Genotypes for individuals at SNP z were

assigned assuming an allele frequency of 0.5 for both populations and drawing from two binomial

trials. Genotypes at the m interacting SNPs were assigned in the same manner as the 2nd SNP in the

pairwise interaction simulations. Using the simulated genotypes, phenotypes were drawn from a

normal distribution, $\mathcal{N}(\mu = \sum_{x=1}^{m}\beta_{G\times G}\times g_{iz}\times g_x, \sigma = 1)$ where g_{is} is the genotype for individual i at

the simulated SNP s.

To simulate the scenario of differential LD on different ancestral backgrounds leading to

false positives, we simulated phenotypes based on a single causal SNP that was tagged by another

SNP. At both SNPs, local ancestries were assigned as described previously and genotypes were drawn

using ancestry specific allele frequencies. Ancestral allele frequencies were assigned such that the

average r^2 between the causal and tag SNP was 0.272 on the background of ancestral population 1

and 0.024 on the background of ancestral population 2. Thus, the tag SNP was only a tag on the

population1 background and not on the population 2 background. Phenotypes were drawn from a

normal distribution, $\mathcal{N}(\mu = \beta_{Causal}\times g_{ic}, \sigma = 1)$, assuming no interaction and $\beta_{Causal} = 0.7$, where

g_{ic} is the genotype of individual i at the causal variant.

We implemented our approach in an R package (GxTheta), which is available for download

at http://www.scandb.org/newinterface/GxTheta.html

Ancestry Inference

Global ancestry inference was done using ADMIXTURE [9] and local ancestry inference was

done using LAMP-LD [55]. CEU and YRI from 1000 Genomes Phase 3 [56] were used as the European

and African reference panels. For the Native American reference panels, 95 Native Americans

genotyped on the Axiom LAT1 array were used[57].

Filtering for Related Individuals

All analyses in real data were filtered for related individuals due to the possibility of cryptic

relatedness causing false positives. To filter for related individuals, we estimated kinship coefficients

between all pairs of individuals using REAP [58]. We defined two individuals as related if they had a

kinship coefficient greater than 0.025. For a pair of related individuals, we removed the one with a

greater number of other individuals to whom he or she was related. In the case of a tie, we removed

one of the pair at random.

Data Normalization

Gene Expression Normalization

Gene expression data (see Results) were first standardized for each gene such that mean

expression was 0 and variance was 1. We then computed a covariance matrix of individual's

expression values and performed PCA on the covariance matrix. Residuals were computed for all

expression values by adjusting for the top 10 principal components and the mean for each gene was

added back to the residuals. Due to the high dynamic range of gene expression compared to

methylation we conservatively chose to additionally perform quantile normalization. We then sorted

the gene expression residuals and used the quantiles of their rank order to draw new expression

values from a normal distribution, $\mathcal{N}(\mu = 0, \sigma = 1)$, by using the inverse cumulative density

function[24,25].

Methylation Data Normalization

Raw methylation values (see Results) were first normalized using Illumina's control probe

scaling procedures. All probes with median methylation less than 1% or greater than 99% were

removed and the remaining probes were logit-transformed as previously described[59]. To control

for extreme outliers, we truncated the distribution of methylation values. For a given probe, we first

computed the mean and standard deviation of the methylation values. We then set any methylation

values deviating more than 2.58 standard deviations from the mean to the methylation value

corresponding to the 99.5th quantile.

Availability of Supporting Data

The Coriell data is available from dbGAP under accession number phs000211.v1.p1. The GALA

and SAGE data is available by emailing the study organizers at https://pharm.ucsf.edu/gala/contact.

Competing Interests

The authors declare that they have no competing interests.

Authors' Contributions

DSP, IE, EK, EE, EH and NZ designed research. DSP, IE, EK, ERG, and NZ performed research. DSP, IE,

EK, EE, CE, CRG, JMG, EG, HA, CJY, EE, EH, and NZ contributed new reagents/analytic tools. DSP, ERG,

and NZ wrote the manuscript. All authors read and approved the final manuscript.

Description of Additional Data Files

The following data are available with the online version of this paper. The Supplemental contains QQ-

plots for the simulations and real analyses performed as well as a table containing p-values for the 2-

component ancestry analysis of the GALA methylation data.

Acknowledgements
We would like to thank Lancelote Leong for his helpful manuscript comments.

583
References
1. Hemani G, Shakhbazov K, Westra H-J, Esko T, Henders AK, Mcrae AF, et al. Detection and
replication of epistasis influencing transcription in humans. Nature. Nature Publishing Group;
2014 Apr 10;508(7495):249–53.

2. Rouhani F, Kumasaka N, de Brito MC, Bradley A, Vallier L, Gaffney D. Genetic Background
Drives Transcriptional Variation in Human Induced Pluripotent Stem Cells. Gibson G, editor.
PLoS Genet. 2014;10(6):e1004432.

3. Kang EY, Han B, Furlotte N, Joo JWJ, Shih D, Davis RC, et al. Meta-Analysis Identifies Gene-by-
Environment Interactions as Demonstrated in a Study of 4,965 Mice. Gibson G, editor. PLoS
Genet. Public Library of Science; 2014 Jan 9;10(1):e1004022.

4. Jemal A, Bray F, Center MM, Ferlay J, Ward E, Forman D. Global cancer statistics. CA Cancer J
Clin. 2011 Mar;61(2):69–90.

5. Lee M, Raj T, Castillo IW. ImmVar Project: Genetic architecture of leukocyte gene expression
in healthy humans. JOURNAL OF …; 2012.

6. Manolio TA, Collins FS, Cox NJ, Goldstein DB, Hindorff LA, Hunter DJ, et al. Finding the missing
heritability of complex diseases. Nature. Nature Publishing Group; 2009 Oct
8;461(7265):747–53.

7. Eichler EE, Flint J, Gibson G, Kong A, Leal SM. Missing heritability and strategies for finding the
underlying causes of complex disease. Nature Reviews …. 2010.

8. Seldin MF, Pasaniuc B, Price AL. New approaches to disease mapping in admixed populations.
Nature Reviews Genetics. Nature Publishing Group; 2011 Aug 1;12(8):523–8.

9. Alexander DH, Novembre J, Lange K. Fast model-based estimation of ancestry in unrelated
individuals. Genome Res. Cold Spring Harbor Lab; 2009 Sep 1;19(9):1655–64.

10. Burchard EG, Ziv E, Coyle N, Gomez SL. The importance of race and ethnic background in
biomedical research and clinical practice. New England Journal …. 2003.

11. Kumar R, Seibold MA, Aldrich MC, Williams LK, Reiner AP, Colangelo L, et al. Genetic Ancestry
in Lung-Function Predictions. N Engl J Med. 2010 Jul 22;363(4):321–30.

12. Kumar R, Nguyen EA, Roth LA, Oh SS, Gignoux CR, Huntsman S, et al. Factors associated with
degree of atopy in Latino children in a nationwide pediatric sample: The Genes-environments
and Admixture in Latino Asthmatics (GALA II) study. J Allergy Clin Immunol. Elsevier; 2013
May 16;132(4):896–905.e1.

13. Price AL, Patterson N, Hancks DC, Myers S, Reich D, Cheung VG, et al. Effects of cis and trans
Genetic Ancestry on Gene Expression in African Americans. Gibson G, editor. PLoS Genet.
Public Library of Science; 2008 Dec 5;4(12):e1000294.

14. Shaffer JR, Kammerer CM, Reich D, McDonald G, Patterson N, Goodpaster B, et al. Genetic
markers for ancestry are correlated with body composition traits in older African Americans.
Osteoporos Int. Springer-Verlag; 2007;18(6):733–41.

15. Florez JC, Price AL, Campbell D, Riba L, Parra MV, Yu F, et al. Strong Association of
Socioeconomic Status and Genetic Ancestry in Latinos: Implications for Admixture Studies of

Type 2 Diabetes. In: Racial Identities, Genetic Ancestry, and Health in South America. Palgrave
Macmillan US; 2011. pp. 137–53.

16.   Reiner AP, Carlson CS, Ziv E, Iribarren C, Jaquish CE, Nickerson DA. Genetic ancestry,
population sub-structure, and cardiovascular disease-related traits among African-American
participants in the CARDIA Study. Hum Genet. Springer-Verlag; 2007;121(5):565–75.

17.   Sanchez E, Webb RD, Rasmussen A, Kelly JA, Riba L, Kaufman KM, et al. Genetically
determined Amerindian ancestry correlates with increased frequency of risk alleles for
systemic lupus erythematosus. Arthritis & Rheumatism. Wiley Subscription Services, Inc., A
Wiley Company; 2010 Dec 1;62(12):3722–9.

18.   Ziv E, John EM, Choudhry S, Kho J, Lorizio W, Pérez-Stable EJ, et al. Genetic Ancestry and Risk
Factors for Breast Cancer among Latinas in the San Francisco Bay Area. Cancer Epidemiol
Biomarkers Prev. American Association for Cancer Research; 2006 Oct 1;15(10):1878–85.

19.   Cheng C-Y, Reich D, Haiman CA, Tandon A, Patterson N, Elizabeth S, et al. African Ancestry and
Its Correlation to Type 2 Diabetes in African Americans: A Genetic Admixture Analysis in
Three U.S. Population Cohorts. Atkin SL, editor. PLoS ONE. Public Library of Science; 2012
Mar 16;7(3):e32840.

20.   Choudhry S, Burchard EG, Borrell LN, Tang H, Gomez I, Naqvi M, et al. Ancestry–Environment
Interactions and Asthma Risk among Puerto Ricans. Am J Respir Crit Care Med [Internet].
American Thoracic Society; 2006 Nov 15;174(10):1088–93. Available from:
http://www.atsjournals.org/doi/abs/10.1164/rccm.200605-596OC

21.   Spielman RS, Bastone LA, Burdick JT, Morley M, Ewens WJ, Cheung VG. Common genetic
variants account for differences in gene expression among ethnic groups. Nat Genet. Nature
Publishing Group; 2007 Feb 1;39(2):226–31.

22.   Peralta CA, Risch N, Lin F, Shlipak MG, Reiner A, Ziv E, et al. The Association of African
Ancestry and Elevated Creatinine in the Coronary Artery Risk Development in Young Adults
(CARDIA) Study. Am J Nephrol. Karger Publishers; 2009 Dec 21;31(3):202–8.

23.   Galanter JM, Gignoux CR, Oh SS, Torgerson D, Pino-Yanes M, Thakur N, et al. Methylation
Analysis Reveals Fundamental Differences Between Ethnicity and Genetic Ancestry. bioRxiv.
Cold Spring Harbor Labs Journals; 2016 Jan 15;:036822.

24.   Population-specificity of human DNA methylation. 2012.

25.   Delude CM. Deep phenotyping: The details of disease. Nature. Nature Publishing Group; 2015
Nov 5;527(7576):S14–5.

26.   Vimaleswaran KS, Berry DJ, Lu C, Tikkanen E, Pilz S, Hiraki LT, et al. Causal Relationship
between Obesity and Vitamin D Status: Bi-Directional Mendelian Randomization Analysis of
Multiple Cohorts. Minelli C, editor. PLOS Med. Public Library of Science; 2013 Feb
5;10(2):e1001383.

27.   Grundberg E, Small KS, Hedman ÅK, Nica AC, Buil A, Keildson S, et al. Mapping cis- and trans-
regulatory effects across multiple tissues in twins. Nat Genet. Nature Publishing Group; 2012
Oct 1;44(10):1084–9.

28.   Aschard H, Lutz S, Maus B, Duell EJ, Fingerlin TE, Chatterjee N, et al. Challenges and
opportunities in genome-wide environmental interaction (GWEI) studies. Hum Genet.

664   Springer-Verlag; 2012;131(10):1591–613.

665   29.   Aschard H, Gusev A, Brown R, Pasaniuc B. Leveraging local ancestry to detect gene-gene
666   interactions in genome-wide data. BMC Genetics. BioMed Central Ltd; 2015 Oct 24;16(1):124.

667   30.   De R, Hu T, Moore JH, Gilbert-Diamond D. Characterizing gene-gene interactions in a
668   statistical epistasis network of twelve candidate genes for obesity. BioData Mining. BioMed
669   Central; 2015;8(1):1–16.

670   31.   Ritchie MD, Hahn LW, Roodi N, Bailey LR, Dupont WD, Parl FF, et al. Multifactor-
671   Dimensionality Reduction Reveals High-Order Interactions among Estrogen-Metabolism
672   Genes in Sporadic Breast Cancer. The American Journal of Human Genetics. 2001
673   Jul;69(1):138–47.

674   32.   Wong MY, Day NE, Luan JA, Chan KP, Wareham NJ. The detection of gene-environment
675   interaction for continuous traits: should we deal with measurement error by bigger studies or
676   better measurement? Int J Epidemiol. 2003 Feb;32(1):51–7.

677   33.   Devlin B, Roeder K. Genomic Control for Association Studies. Biometrics [Internet]. Blackwell
678   Publishing Ltd; 2004 May 25;55(4):997–1004. Available from:
679   http://doi.wiley.com/10.1111/j.0006-341X.1999.00997.x

680   34.   Simon-Sanchez J, Scholz S, Fung H-C, Matarin M, Hernandez D, Gibbs JR, et al. Genome-wide
681   SNP assay reveals structural genomic variation, extended homozygosity and cell-line induced
682   alterations in normal individuals. Hum Mol Genet. Oxford University Press; 2007 Jan
683   1;16(1):1–14.

684   35.   Martin AR, Costa HA, Lappalainen T, Henn BM, Kidd JM, Yee M-C, et al. Transcriptome
685   Sequencing from Diverse Human Populations Reveals Differentiated Regulatory Architecture.
686   Gibson G, editor. PLoS Genet. Public Library of Science; 2014 Aug 14;10(8):e1004549.

687   36.   Borrell LN, Nguyen EA, Roth LA, Oh SS, Tcheurekdjian H, Sen S, et al. Childhood Obesity and
688   Asthma Control in the GALA II and SAGE II Studies. dx.doi.org. American Thoracic Society;
689   2013. 6 p.

690   37.   Torgerson DG, Gignoux CR, Galanter JM, Drake KA, Roth LA, Eng C, et al. Case-control
691   admixture mapping in Latino populations enriches for known asthma-associated genes. J
692   Allergy Clin Immunol. 2012 Jul;130(1):76–82.e12.

693   38.   Eichler EE, Flint J, Gibson G, Kong A, Leal SM, Moore JH, et al. Missing heritability and
694   strategies for finding the underlying causes of complex disease. Nature Reviews Genetics.
695   Nature Publishing Group; 2010 Jun 1;11(6):446–50.

696   39.   Powell JE, Henders AK, Mcrae AF, Kim J, Hemani G, Martin NG, et al. Congruence of Additive
697   and Non-Additive Effects on Gene Expression Estimated from Pedigree and SNP Data. Spector
698   TD, editor. PLoS Genet. Public Library of Science; 2013 May 16;9(5):e1003502.

699   40.   Keller MC. Gene × Environment Interaction Studies Have Not Properly Controlled for
700   Potential Confounders: The Problem and the (Simple) Solution. Biological Psychiatry. 2014
701   Jan;75(1):18–24.

702   41.   Sul JH, Bilow M, Yang W-Y, Kostem E, Furlotte N, He D, et al. Accounting for Population
703   Structure in Gene-by-Environment Interactions in Genome-Wide Association Studies Using
704   Mixed Models. Schork NJ, editor. PLoS Genet. Public Library of Science; 2016 Mar

4;12(3):e1005849.

42.   Kooperberg C, LeBlanc M. Increasing the power of identifying gene × gene interactions in
genome-wide association studies. Genet Epidemiol. Wiley Subscription Services, Inc., A Wiley
Company; 2008 Apr 1;32(3):255–63.

43.   Murcray CE, Lewinger JP, Gauderman WJ. Gene-Environment Interaction in Genome-Wide
Association Studies. Am J Epidemiol. Oxford University Press; 2009 Jan 15;169(2):219–26.

44.   Hsu L, Jiao S, Dai JY, Hutter C, Peters U, Kooperberg C. Powerful Cocktail Methods for
Detecting Genome-Wide Gene-Environment Interaction. Genet Epidemiol. 2012 Apr
1;36(3):183–94.

45.   Yang J, Benyamin B, McEvoy BP, Gordon S, Henders AK, Nyholt DR, et al. Common SNPs
explain a large proportion of the heritability for human height. Nat Genet. 2010 Jun
20;42(7):565–9.

46.   Smith ZD, Meissner A. DNA methylation: roles in mammalian development. Nature Reviews
Genetics. Nature Publishing Group; 2013 Mar 1;14(3):204–20.

47.   Artigas MS, Loth DW, Wain LV, Gharib SA, Obeidat M, Tang W, et al. Genome-wide association
and large-scale follow up identifies 16 new loci influencing lung function. Nat Genet. Nature
Publishing Group; 2011 Nov 1;43(11):1082–90.

48.   Liao SY, Lin X, Christiani DC. Gene-environment interaction effects on lung function-a
genome-wide association study within the Framingham heart study. Environ Health. 2013.

49.   Anderson LN, Briollais L, Atkinson HC, Marsh JA, Xu J, Connor KL, et al. Investigation of
Genetic Variants, Birthweight and Hypothalamic-Pituitary-Adrenal Axis Function Suggests a
Genetic Variant in the SERPINA6 Gene Is Associated with Corticosteroid Binding Globulin in
the Western Australia Pregnancy Cohort (Raine) Study. Hsu Y-H, editor. PLoS ONE. Public
Library of Science; 2014 Apr 1;9(4):e92957.

50.   Baran Y, Pasaniuc B, Sankararaman S, Torgerson DG, Gignoux C, Eng C, et al. Fast and accurate
inference of local ancestry in Latino populations. Bioinformatics. Oxford University Press;
2012 May 15;28(10):1359–67.

51.   Sankararaman S, Sridhar S, Kimmel G. Estimating local ancestry in admixed populations. The
American Journal of …. 2008.

52.   Price AL, Patterson N, Yu F, Cox DR, Waliszewska A, McDonald GJ, et al. A Genomewide
Admixture Map for Latino Populations. The American Journal of Human Genetics. 2007
Jun;80(6):1024–36.

53.   Smith MW, Patterson N, Lautenberger JA, Truelove AL, McDonald GJ, Waliszewska A, et al. A
High-Density Admixture Map for Disease Gene Discovery in African Americans. The American
Journal of Human Genetics. 2004 May;74(5):1001–13.

54.   Balding DJ, Nichols RA. A method for quantifying differentiation between populations at
multi-allelic loci and its implications for investigating identity and paternity. Human
Identification: The Use of DNA Markers. 1995.

55.   Baran Y, Pasaniuc B, Sankararaman S, Torgerson DG, Gignoux C, Eng C, et al. Fast and accurate
inference of local ancestry in Latino populations. Bioinformatics. 2012 May 15;28(10):1359–

67.

56.   Consortium T1GP. An integrated map of genetic variation from 1,092 human genomes.
Nature. Nature Publishing Group; 2012 Nov 1;491(7422):56–65.

57.   Drake KA, Torgerson DG, Gignoux CR, Galanter JM, Roth LA, Huntsman S, et al. A genome-wide
association study of bronchodilator response in Latinos implicates rare variants. Journal of
Allergy and Clinical Immunology. 2014 Feb;133(2):370–378.e15.

58.   Thornton T, Tang H, Hoffmann TJ, Ochs-Balcom HM, Caan BJ, Risch N. Estimating kinship in
admixed populations. Am J Hum Genet. 2012 Jul 13;91(1):122–38.

59.   Du P, Zhang X, Huang C-C, Jafari N, Kibbe WA, Hou L, et al. Comparison of Beta-value and M-
value methods for quantifying methylation levels by microarray analysis. BMC
Bioinformatics. BioMed Central Ltd; 2010 Nov 30;11(1):587.

Figure Legends
Figure 1. Power Plots for Pairwise Interaction Simulations.
Power of testing $G \times \theta$ (a) versus testing pairwise SNPs directly (b) as a function of the difference in
the ancestral allele frequencies at a differentiated SNP.
762
763 Figure 2. Power Plots for Multi-way Pairwise Interaction Simulations
764 Power of testing $G \times \theta$ as a function of the difference in the ancestral allele frequencies for multiple
765 interacting SNPs.
766
767 Figure 3. Power Plots for $G \times E$ Interaction Simulations.
768 Power of testing $G \times \theta$ as a function of the correlation between an environmental covariate and
769 genetic ancestry.
770
Figure 4. Examples of How Genetic Ancestry Can Be A Proxy for Interacting Covariates.
(a) Model of how genetic ancestry θ can be correlated with various environmental exposures, some
of which affect a phenotype. (b) Example of how the correlation between the probability of an AA
genotype (bars 2-4) and values of θ (bar 1) increase with higher levels of SNP allele frequency
differentiation. In this plot p_1 and p_2 denote the allele frequency of allele A in ancestral populations 1
and 2 respectively. (c) Example of how effect sizes at a tag-SNP may differ due to differential LD on
distinct ancestral backgrounds (here, EUR and AFR).
778
779

Tables

782 **Table 1. Distribution of Ancestry in Coriell and GALA II.**

Dataset	θ_{EUR}	θ_{AFR}	θ_{NAM}
Coriell	μ=0.212, σ=0.021	μ=0.788, σ=0.021	NA
GALA II MX	μ=0.396, σ=0.149	μ=0.043, σ=0.025	μ=0.561, σ=0.159
GALA II PR	μ=0.641, σ=0.094	μ=0.246, σ=0.101	μ=0.113, σ=0.024

783 Mean and variance of the global ancestry distributions for each dataset.

785 **Table 2. GALA II DNA Methylation Analysis Results.**

GALA II Population	Probe Gene	Probe ID	rsid	Distance of SNP to Probe	Marginal p-value	AITL p-value	AITL Replication p-value
MX	CNFN	cg14327995	rs16975986	280795	2.49E-09	5.69E-09	9.27E-03
MX	C11orf95	cg16678159	rs7106153	249768	2.58E-01	2.52E-08	9.39E-02
MX	NA	cg05697734	rs1560919	13711	1.14E-01	2.21E-08	8.18E-03
MX	TNK2	cg01792640	rs67217828	278866	4.49E-01	6.38E-09	1.43E-02
MX	HDAC4	cg06533788	rs925736	9548	4.51E-01	3.09E-09	2.80E-02
PR	NA	cg07436864*	rs8117083	31813	7.46E-02	1.34E-09	5.34E-03
PR	NA	cg16803083*	rs4312379	63847	3.69E-01	2.29E-08	2.31E-04
PR	SERPINA6	cg10025865	rs17091085	247796	6.83E-01	2.97E-08	8.05E-03

786 P-values for AITL applied to the methylation data in the GALA II Latinos. MX and PR denote Mexicans
787 and Puerto Ricans respectively in the GALA II population columns. The probe gene column shows the
788 gene that the methylation probe lies in. The marginal column is the p-value for standard linear
789 regression of methylation on genotype while controlling for population structure. * indicates results
790 that replicated between the Mexicans and Puerto Ricans.