# Evolution of corticosteroid specificity for human, chicken, alligator and frog glucocorticoid receptors

Yoshinao Katsu[1, 2, *], Satomi Kohno[3], Kaori Oka[1], Michael E. Baker[4, *]

[1]Graduate School of Life Science, Hokkaido University, Sapporo, Japan; [2]Department of Biological Sciences, Hokkaido University, Sapporo, Japan; [3]Department of Obstetrics and Gynecology, Medical University of South Carolina, and Marine Biomedicine and Environmental Science Center, Hollings Marine Laboratory, Charleston, SC, USA; [4]Department of Medicine, University of California, San Diego, CA, USA

Correspondence:

ykatsu@sci.hokudai.ac.jp

mbaker@ucsd.edu

**Abstract**. We investigated the evolution of the response of human, chicken, alligator and frog glucocorticoid receptors (GRs) to dexamethasone, cortisol, corticosterone, 11-deoxycorticosterone, 11-deoxycortisol and aldosterone.   We find significant differences among these vertebrates in the transcriptional activation of their full length GRs by these steroids, indicating that there were changes in the specificity of the GR for steroids during the evolution of terrestrial vertebrates.   To begin to study the role of interactions between different domains on the GR in steroid sensitivity and specificity for terrestrial GRs, we investigated transcriptional activation of truncated GRs containing their hinge domain and ligand binding domain (LBD) fused to a GAL4 DNA binding domain (GAL4 DBD).   Compared to corresponding full length GRs, transcriptional activation of GAL4 DBD-GR hinge/LBD constructs required higher steroid concentrations and displayed altered steroid specificity, indicating that interactions between the hinge/LBD and other domains are important in glucocorticoid activation of these terrestrial GRs.

**Short Title:** Evolution of steroid specificity for terrestrial GRs

**Key Words:** Glucocorticoid Receptor, Evolution, Allosteric Regulation, terrestrial vertebrates

## 1. Introduction

Glucocorticoids (Figure 1) regulate a variety of physiological functions

including carbohydrate and protein metabolism, blood pressure, immune function and

the body's anti-inflammatory processes via transcriptional activation of the

glucocorticoid receptor (GR) [1-5].   The GR and other steroid receptors belong to the

nuclear receptor family, a large family of transcription factors, which includes receptors

for thyroid hormone, retinoids and other small lipophilic molecules [6-10].   The GR

and other steroid receptors have a characteristic modular structure consisting of an

N-terminal domain (NTD) (domains A and B), a central DNA-binding domain (DBD)

(domain C), a hinge domain (D) and a C-terminal ligand-binding domain (LBD)

(domain E) [9, 11-14] (Figure 2).   The E domain alone is competent to bind steroids

[11, 12, 15-18].

The NTD contains an activation function 1 [AF1] domain, which is a strong

transcriptional activator of the GR [19-21].   Interestingly, AF1 is intrinsically

disordered, unlike the DBD and LBD [21-23].   Allosteric interactions between AF1

and other domains on the GR and coactivators lead to a conformational rearrangement

of AF1 that is important in transcriptional activation of the GR [23-26].   In rat GR,

there is evidence that allosteric interactions between DBD and other domains regulate

gene transcription [27, 28].   Recent crystal structures of the DBD-Hinge-LBD domains

of other nuclear receptors [13, 22] identified allosteric signaling between the DBD and

LBD domains.   [27, 28].

Although dexamethasone (DEX) and cortisol (F) activation of rodent [29] and

human [20, 30-33] GRs has been investigated,    there has been no systematic

assessment of ligand specificity among phylogenetically diverse terrestrial vertebrate

GRs, such as amphibians, reptiles, birds and mammals.   It is possible that more than

one corticosteroid may act as a physiological glucocorticoid in terrestrial vertebrates.

Reports of transcriptional activation by corticosteroids of the GR for other terrestrial

vertebrates: amphibians, reptiles and birds, are limited [34, 35].   Oka et al. [34]

reported half-maximal response (EC50) values for transcriptional activation of full

length alligator GR by F, corticosterone (B), 11-deoxycorticosterone (DOC) and

aldosterone (Aldo).   The EC50s for F and B were 0.29 nM and 0.16 nM, respectively,

which is consistent with the known role of these two steroids as glucocorticoids in

mammals.   However, the EC50s for Aldo and DOC were 2.9 nM and 2.8 nM, which is

unexpected because both steroids have a lower binding affinity for human GR [30,

32]and are weak transcriptional activators of human GR [30, 36].   Similar intriguing

findings for Aldo were reported for chicken GR by Proszkowiec-Weglarz and Porter

[35], who found that the EC50s for transcriptional activation of chicken GR by Aldo

and B were 0.8 nM and 1.8 nM, respectively, with the level of transcription due to B

being about 30% higher than to Aldo.   The EC50s of DOC and other corticosteroids

for chicken GR and of DEX for alligator GR were not determined.

**Figure 1. Structures of various corticosteroids.**

Cortisol and corticosterone are physiological glucocorticoids in terrestrial vertebrates and

ray-finned fish [12, 37, 38].   Aldosterone, 11-deoxycorticosterone and 11-deoxycortisol are

physiological mineralocorticoids [12, 39-41].   Aldo and DOC are weak transcriptional

activators of human GR [30, 32, 36].   11-deoxycortisol is both a mineralocorticoid and a

glucocorticoid in lamprey [42].

**Figure 2. Comparison of domains in some terrestrial vertebrate GRs.**

GRs from human, chicken, alligator and *X. laevis* are compared.    The functional A/B domain

to E domains are schematically represented with the numbers of amino acid residues and the

percentage of amino acid identity between the domain in the human GR and the corresponding

domain in the other vertebrate GRs.    For example, the entire human GR sequence is 75%

identical to that of chicken GR, while domain E (LBD) on human GR is 90% identical to that of

chicken GR.

GenBank accession numbers: human GR (NM_000176), chicken GR (NM_001037826),

alligator GR (AB701407), *X. laevis* GR (NM_001088062).

These unexpected responses of alligator and chicken GRs to Aldo and our

interest in the evolution of specificity for corticosteroids in the GR in vertebrates [12, 34,

41, 43, 44] motivated us to investigate the response to a panel corticosteroids of the GR

from chicken and the amphibian [*Xenopus laevis*] for comparison to human and

alligator GR with the goal of clarifying the evolution of corticosteroid specificity in

terrestrial vertebrates.    In addition, we were interested in investigating the role of

domains A-C and domains D-E [13, 21-23, 44-47] in the response of GRs to steroids.

The influence of domains A-C on steroid responses for the GR has not been studied

previously in non-mammalian terrestrial vertebrates.    For these studies we constructed

a plasmid containing the GAL4 DBD fused to the D domain and E domain of the GR

(GR-LBD).

Interestingly, we found significant differences in the EC50s of these full length

GRs to corticosteroids indicating that during the evolution of these terrestrial vertebrates

there were changes in their response to various corticosteroids.    Moreover, in the

presence of corticosteroids, truncated GRs containing a GR LBD fused to a GAL4 DBD

had a higher EC50 value (weaker activation) than their corresponding full length GRs,

indicating altered steroid specificity among these terrestrial vertebrate GRs and that the

evolution of the response of terrestrial vertebrate GRs to different steroids was complex.

These differences may involve allosteric signaling between the domains D-E and

other GR domains [22, 44, 45] or alterations in the binding of co-regulator proteins [24,

48, 49] or the absence of post-translational modification of domains A, B or C [49-51],

or combinations of these mechanisms.

**2. Materials and Methods**

**2.1 Chemical reagents**

DEX, F, B, aldosterone (Aldo), DOC and 11-deoxycortisol (S) were purchased

from Sigma-Aldrich. For the reporter gene assays, all hormones were dissolved in

dimethylsulfoxide (DMSO) and the final concentration of DMSO in the culture medium

did not exceed 0.1%.

**2.2 Construction of plasmid vectors**

The full-coding regions and D/E domains of the GR from *X. laevis*, alligator,

chicken and human were amplified by PCR with KOD DNA polymerase (TOYOBO

Biochemicals, Osaka, Japan). The PCR products were gel-purified and ligated into

pcDNA3.1 vector *(KpnI-NotI* site for human, chicken and alligator GRs, and

*HindIII-NotI* site for *X. laevis* GR) (Invitrogen) for the full-coding region or pBIND

vector (*MluI-NotI* site) (Promega) for D-E domains. As shown in Figure 2, the D

domain begins at human GR (N487), chicken GR (N482), alligator GR (N490) and *X.*

*laevis* GR (N486) [34].

**2.3 Transactivation Assay and Statistical Methods**

CHO-K1 cells (Chinese hamster ovary cell) were used in the reporter gene assay.

Transfection and reporter assays were carried out as described previously [34, 52].

The use of CHO-K1 cells and an assay temperature of 37C does not replicate the

physiological environment of *X. laevis*, alligator and chicken. Nevertheless, studies

with mammalian cell lines at 37C have proven useful for other studies of transcriptional

activation by corticosteroids of teleost fish GRs [53-56] and other non-mammalian GRs

[35, 57, 58]. Levels of expression of the different non-mammalian GRs and their

truncated counterparts may differ in CHO-K1 cells. However, comparisons of the

EC50 of different corticosteroids for each GR would be valid, which is the goal of our

study. All transfections were performed at least three times, employing triplicate

sample points in each experiment. The values shown are mean ± SEM from three

separate experiments, and dose-response data and EC50 were analyzed using GraphPad

Prism.   Comparisons between two groups were performed using *t*-test, and all

multi-group comparisons were performed using one-way ANOVA followed by

Bonferroni test.   $P < 0.05$ was considered statistically significant.

## 3. Results

### 3.1 Different steroid-response for full length and truncated human, chicken, alligator and *X. laevis* GRs.

### 3.11 Human GR

In Figures 3A and B, we show corticosteroid-inducible transcriptional activation of full length and truncated (GAL4 DBD-GR LBD) human GRs by DEX, F, B, Aldo, DOC and S.   At $10^{-7}$ M, all corticosteroids induced transcription of full length human GR via the MMTV-reporter gene.   In contrast, truncated human GR had a strong response to DEX, a much weaker response to F, a small response to B and no response to Aldo, DOC and S.

### 3.12 Chicken GR

Transcription of full length chicken GR was activated by all corticosteroids at $10^{-7}$ M, with a similar strong response to B, Aldo and DOC and a lesser response to DEX, F and S (Figure 3C).   Truncated chicken GR was strongly activated by B, F, DEX and Aldo, with a weaker response to DOC and S (Figure 3D).

### 3.13 Alligator GR

Transcription of full length alligator GR was activated by all corticosteroids at $10^{-7}$ M, with a similar strong response to B, Aldo and DOC and a lower response to DEX, F and S (Figure 3E).   Truncated alligator GR was strongly activated by DEX, F and B, with lower response to Aldo and S and a very weak response to DOC (Figure 3F).

### 3.14 *X laevis* GR

Transcription of full length *X. laevis* GR was activated by all corticosteroids at $10^{-7}$ M, with a similar strong response to DEX and B and a lower response to F, Aldo and DOC and much lower response to S (Figure 3G).   Truncated *X. laevis* GR was strongly activated by DEX and B, with a much lower response to F and Aldo and a no response to DOC and S (Figure 3H).

Figure 3. Ligand-specificities of human, chicken, alligator and *X. laevis* full length GRs

and LBD GRs.

Full-length human GR (A), chicken GR (C), alligator GR (E), and *X. laevis* GR (G) were

expressed in CHO-K1 cells with an MMTV-luciferase reporter.   Plasmids for corresponding

truncated GRs (human (B), chicken (D), alligator (F) and *X. laevis* (H) containing the D domain

and LBD (E domain) fused to a GAL4-DBD were expressed in CHO-K1 cells with a luciferase

reporter containing GAL4 binding site.   Cells were treated with $10^{-7}$ M DEX, F, B, Aldo, DOC,

S or vehicle alone (DMSO).   Results are expressed as means ± SEM, n=3.   Y-axis indicates

fold-activation compared to the activity of control vector with vehicle (DMSO) alone as 1.

Transcriptional activation of the different GRs in the presence of the DMSO control is at

background level, indicating that these GRs do not have constitutive activity.

**3.15 EC50 values for transcriptional activation of full length human, chicken,**

**alligator and *X. laevis* GRs**

Next we examined the concentration-dependence of transcriptional activation of

full length terrestrial vertebrate GRs by DEX, F, B, Aldo, DOC and S (Figure 4, Table

1).   Compared to the other steroids, DEX has the lowest EC50 for all of the full length

GRs (Table 1).   Interestingly, there are significant differences among the GRs of the

EC50s for other corticosteroids, including F and B, which are the major physiological

glucocorticoids in terrestrial vertebrates.   For example, for full length GRs, B has a

lower EC50 than F for *X. laevis* GR, while F has a lower EC50 than B for human,

chicken and alligator GR.

Aldo, which is a mineralocorticoid, has an EC50 of 2.7 nM and 44 nM

respectively, for alligator GR and *X. laevis* GR and an EC50 of 2 nM and 82 nM,

respectively, for chicken and human GR.   DOC, which also is a mineralocorticoid, has

an EC50 of 2.6 nM and 23 nM, respectively, for alligator GR and *X. laevis* GR, and an

EC50 of 0.63 nM and 110 nM, respectively, for chicken GR and human GR.

Interestingly, S has an EC50 of 0.17 nM and 0.35 nM, respectively, for chicken and

alligator GR, and a much higher EC50 for human GR [50 nM] and *X. laevis* GR [530

nM].

**3.16 EC50 values for transcriptional activation of truncated (GAL4 DBD-GR**

**LBD) terrestrial vertebrate GRs**

The concentration-dependence of transcriptional activation of truncated

terrestrial vertebrate GRs by DEX, F, B, Aldo, DOC and S is shown in Figure 4 and

Table 1.   Transcriptional activation by several steroids was dramatically different

among the terrestrial vertebrate GRs that lacked the A-C domains.   For example,

truncated human GR has a strong response to DEX (EC50 = 8.3 nM) and a very weak

response to F (EC50 = 1.2 µM), and no significant response to B, Aldo, DOC or S.

This contrasts to truncated chicken GR, which has nM EC50s for DEX, F and B, and a

weaker but significant response to Aldo and S.   Only DOC does not activate truncated

chicken GR.   Truncated alligator GR has nM EC50s for DEX and F, a weaker but

significant response to B (EC50 = 49 nM), a weak response to Aldo (EC50 = 0.16 µM)

and S (EC50 = 0.12 µM) and no response to DOC.

These results suggest that allosteric signaling between the hinge/LBD and one or

more of the A, B and C domains influences the response of terrestrial vertebrate GRs to

corticosteroids.

**Table 1. EC50 values for transcriptional activation by corticosteroids of terrestrial**

**vertebrate GRs**

**A. Full length GR (A-E domains)**

|  | DEX | F | B | Aldo | DOC | S |
|---|---|---|---|---|---|---|
| **Human GR-Full** | $1.7 \times 10^{-10}$ | $5.6 \times 10^{-9}$ | $2.0 \times 10^{-8}$ | $8.2 \times 10^{-8}$ | $1.1 \times 10^{-8}$ | $5.0 \times 10^{-8}$ |
| **Chicken GR-Full** | $2.8 \times 10^{-11}$ | $6.0 \times 10^{-11}$ | $2.3 \times 10^{-10}$ | $2.0 \times 10^{-9}$ | $6.3 \times 10^{-10}$ | $1.7 \times 10^{-10}$ |
| **Alligator GR-Full** | $1.4 \times 10^{-10}$ | $2.0 \times 10^{-10}$ | $3.5 \times 10^{-10}$ | $2.7 \times 10^{-9}$ | $2.6 \times 10^{-9}$ | $3.5 \times 10^{-10}$ |
| **Xenopus GR-Full** | $7.3 \times 10^{-9}$ | $5.6 \times 10^{-8}$ | $5.1 \times 10^{-9}$ | $4.4 \times 10^{-8}$ | $2.3 \times 10^{-8}$ | $5.3 \times 10^{-7}$ |

15 **B. Truncated GR (GAL4-DBD+GR-D+E domains)**

|  | DEX | F | B | Aldo | DOC | S |
|---|---|---|---|---|---|---|
| **Human GR-LBD** | $8.3 \times 10^{-9}$ | $1.2 \times 10^{-6}$ | - | - | - | - |
| **Chicken GR-LBD** | $2.5 \times 10^{-10}$ | $1.6 \times 10^{-9}$ | $6.5 \times 10^{-9}$ | $7.7 \times 10^{-8}$ | - | $6.6 \times 10^{-8}$ |
| **Alligator GR-LBD** | $3.1 \times 10^{-9}$ | $7.7 \times 10^{-9}$ | $4.9 \times 10^{-8}$ | $1.6 \times 10^{-7}$ | - | $1.2 \times 10^{-7}$ |
| **Xenopus GR-LBD** | $6.7 \times 10^{-8}$ | - | $4.8 \times 10^{-8}$ | - | - | - |

Figure 4. Concentration-dependent transcriptional activation by corticosteroids of full length and truncated human, chicken, alligator and *X. laevis* GRs.

Plasmids encoding full length GRs (A: human GR, C: chicken GR, E: alligator GR, G: *Xenopus* GR) or the GAL4-DBD fused to the D domain and LBD of GRs (B: human GR, D: chicken GR, F: alligator GR, H: *Xenopus* GR) were expressed in CHO-K1 cells and treated with increasing concentrations of DEX, F, B, Aldo, DOC, S or vehicle alone (DMSO). Y-axis indicates fold-activation compared to the activity of control vector with vehicle (DMSO) alone as 1.

**3.17 Analysis of a 25 residue segment on human GR and MR that influences corticosteroid specificity**

Rogerson et al. [59] identified a combination of 12 amino acids in a 25 residue segment, corresponding to the c-terminus of helix 5, a β-turn and helix 6 on human GR, that could be replaced with corresponding residues from human MR to yield a hybrid GR that had an EC50 of 3 nM for Aldo.   In Figure 5, we compare this segment in chicken, alligator and *X. laevis* GRs with the corresponding segments in human MR and GR.   The alignment does not reveal a pattern of similarity between chicken and alligator GRs and human MR that can explain the lower EC50s that Aldo has for chicken and alligator GRs compared to human and *X. laevis* GRs.

```
Figure 5.  Exchange of Amino Acids in Human MR to Human GR that
increase binding of Aldosterone.

Human GR                    615   620   625   630   635

                             |     |     |     |     |

Human GR            614 RQSSANLLCFAPDLIINEQRMTLPC 638

Chicken GR          609 KQSNGNLLCFAPDLIINEQRMNLPC 633

Alligator GR        617 KQSNGNLLCFAPDLIINEQRMSLPC 641

X.laevis GR         613 KQTNGSILYFAPDLVITEDRMHLPF 637

Human MR            820 KHTNSQFLYFAPDLVFNEEKMHQSA 844

                             |     |     |     |

Human MR               825   830   835   840

Exchange of underlined amino acids from hMR to hGR yields a
mutant GR with high affinity (Kd= 3nM) and low EC50 for Aldo.
```

**Figure 5. Analysis of a region in human GR and MR that is important for mineralocorticoid specificity.**

Rogerson et al. [59] identified a segment in human MR that could be inserted into human GR and increase its response to Aldo.   We underline specific residues in the MR that replaced residues in the GR to yield an increase in the response of the GR to Aldo.   Also shown is the corresponding region in chicken, alligator and *X. laevis* GRs.   Residues that are conserved in all vertebrate GRs are underlined.

**4. Discussion**

Although there are several reports of the response to different corticosteroids of the mammalian GR [20, 29-33, 36, 44], the corticosteroids that activate GRs from other terrestrial vertebrates have not been studied in depth.    In birds and amphibians, B appears to be the physiological glucocorticoid [60, 61], and S has been found to be a physiological glucocorticoid in lamprey [42].    However, as discussed below, our data [Table 1] supports the presence of more than one physiological glucocorticoids in some terrestrial vertebrates.    Also, our data indicate that there were changes in specificity for corticosteroids in the GR at key transitions in the evolution of terrestrial vertebrates.

As shown in Figures 3 and 4 and Table 1, we find significant differences in the response of full length GRs from *X. laevis*, alligator, chicken and humans to a panel of corticosteroids, providing evidence for the evolution of selectivity of terrestrial vertebrate GRs for F, B, Aldo, DOC and S.    We confirm previous studies [34, 35] that Aldo has nM EC50s for full length chicken and alligator GR [34, 35].    This contrasts with the response to Aldo of full length human and *X. laevis* GR, for which the EC50 is 82 nM and 44 nM, respectively.    The low EC50s of B for chicken GR (0.23 nM), alligator GR (0.35 nM) and *X. laevis* GR (5.1 nM) are consistent with a role for B as a physiological glucocorticoid in these vertebrates [60, 61].    We also find that DOC, another mineralocorticoid [40, 41, 62], has a low EC50 for full length chicken GR (0.6 nM) and alligator GR (2.6 nM), in contrast to DOC's higher EC50 for human GR (110 nM) and *X. laevis* GR (23 nM).    S also has a substantially lower EC50 for chicken GR (0.17 nM) and alligator GR (0.35 nM) compared to human GR (50 nM) and *X. laevis* GR (953 nM).    The low EC50s of B, DOC and S for chicken and alligator GRs and of B for *X. laevis* GR leaves open the possibility that these steroids are physiological glucocorticoids in these vertebrates.    There are regulatory implications for DOC and S as glucocorticoids because these steroids lack an 11β-OH group that is present in F and B [Figure 1].    Thus, DOC and S would be inert to 11β-HSD2, and could activate chicken and alligator GRs in tissues containing 11β-HSD2, which would inactivate B and F [1, 63-65].

Our studies with truncated GRs (hinge-LBD) reveal that one or more of the A, B and C domains are important in the response of terrestrial vertebrate GRs to corticosteroids.    We find that compared to full length GRs, all of the truncated GRs (hinge-LBD) have substantially higher EC50s for all corticosteroids.    For example, the EC50s of DEX and F for truncated human GR increased to 8.3 nM and 1.2 μM, respectively.    Moreover, Aldo, B, DOC and S have an EC50 greater than 1 μM for truncated human GR.    Similar changes to higher EC50s were found for the

1   non-mammalian vertebrate GRs.   Thus, F, Aldo, DOC and S have EC50s greater 1 $\mu$M

2   for truncated *X. laevis* GR.   DOC has an EC50 greater 1 $\mu$M for truncated chicken and

3   alligator GR.   Among the corticosteroids that we studied, DEX is least sensitive and

4   DOC is most sensitive to the loss of the A, B and C domains.

5           There are several overlapping mechanisms that could account for stronger

response to corticosteroids of full length GRs compared to that of their truncated GR

counterparts.   Allosteric interactions between the LBD and the NTD [44, 45, 49] or

DBD [27, 47] are known to influence transcriptional activation of human and rat GRs.

These allosteric interactions may be influenced by post-translational modification of the

NTD by phosphorylation [49, 50] or SUMOylation [51], which also may influence

binding of co-activators [24, 45, 49].

Based on Rogerson et al.'s [59] identification of a region in hMR that could

be substituted into hGR and increase its response to Aldo, we analyzed the

corresponding segment on chicken, alligator, *X laevis* and human GRs for clues to

differences in their responses to corticosteroids.   Our analysis of this segment (Figure

5) did not find a pattern that can explain the relatively strong responses to Aldo of

chicken and alligator GRs, suggesting that other mechanisms such as interactions with

the LBD of the NTD on alligator and chicken GRs may contribute to the differences in

their response to corticosteroids compared to human and *X. laevis* GRs.

4.1 **Evolution**

Our data indicate that there were significant changes in the response to

corticosteroids during the evolution of terrestrial vertebrates.   Among the species that

we have studied, chicken and alligator are closest, having diverged about 150 million

years ago (myr) from a common ancestor.   Consistent with this close relationship, full

length chicken and alligator GRs have similar EC50s for B, Aldo, S and E.   In contrast,

full length GR from *X. laevis*, which is the most divergent of the studied

non-mammalian species, has the high EC50s for all tested corticosteroids.

It is interesting that human mineralocorticoid receptor [MR], which descended

with the GR from a common ancestor [37, 66, 67], also has an interaction between

domains A and B and the LBD that regulates transcriptional activation by Aldo [39, 68,

69] as does zebrafish MR [70].   The A/B domains on human and zebrafish MR can

interact with each other's LBD, indicating that this is an ancient property of the MR.

This suggests that the role in transcriptional activation of the interaction between the

A/B and LBD domains arose in the common ancestor of the GR and MR.   Further

studies of the role in transcriptional activation of the A, B and C domains on the GR and

MR should provide insights into the evolution of steroid specificity in these receptors.

**Author Contributions**

Yoshinao Katsu and Michael E. Baker conceived and designed the experiments and wrote the paper.    Satomi Kohno and Kaori Oka carried out the research.

**Declaration of interests:** The authors have no conflict of interest to declare.

**Acknowledgments**

        We thank colleagues in our laboratories.    K.O. was supported by the Japan Society for the Promotion of Science (JSPS) Research Fellowships for Young Scientists. This work was supported in part by Grants-in-Aid for Scientific Research 23570067 and 26440159 (YK) from the Ministry of Education, Culture, Sports, Science and Technology of Japan.

**References**

[1] Chapman K, Holmes M, Seckl J. 11beta-hydroxysteroid dehydrogenases: intracellular gate-keepers of tissue glucocorticoid action. Physiological reviews. 2013;93:1139-206.

[2] Kadmiel M, Cidlowski JA. Glucocorticoid receptor signaling in health and disease. Trends in pharmacological sciences. 2013;34:518-30.

[3] McEwen BS. Physiology and neurobiology of stress and adaptation: central role of the brain. Physiological reviews. 2007;87:873-904.

[4] Rose AJ, Herzig S. Metabolic control through glucocorticoid hormones: an update. Molecular and cellular endocrinology. 2013;380:65-78.

[5] Chrousos GP. Stress and sex versus immunity and inflammation. Sci Signal. 2010;3:pe36.

[6] Baker ME. Origin and diversification of steroids: co-evolution of enzymes and nuclear receptors. Molecular and cellular endocrinology. 2011;334:14-20.

[7] Bridgham JT, Eick GN, Larroux C, Deshpande K, Harms MJ, Gauthier ME, et al. Protein evolution by molecular tinkering: diversification of the nuclear receptor superfamily from a ligand-dependent ancestor. PLoS biology. 2010;8.

[8] Markov GV, Tavares R, Dauphin-Villemant C, Demeneix BA, Baker ME, Laudet V. Independent elaboration of steroid hormone signaling pathways in metazoans. Proc Natl Acad Sci U S A. 2009;106:11913-8.

[9] Sladek FM. What are nuclear receptor ligands? Molecular and cellular

endocrinology. 2011;334:3-13.

[10] Bertrand S, Belgacem MR, Escriva H. Nuclear hormone receptors in chordates. Molecular and cellular endocrinology. 2011;334:67-75.

[11] Huang P, Chandra V, Rastinejad F. Structural overview of the nuclear receptor superfamily: insights into physiology and therapeutics. Annu Rev Physiol. 2010;72:247-72.

[12] Baker ME, Funder JW, Kattoula SR. Evolution of hormone selectivity in glucocorticoid and mineralocorticoid receptors. J Steroid Biochem Mol Biol. 2013;137:57-70.

[13] Rastinejad F, Huang P, Chandra V, Khorasanizadeh S. Understanding nuclear receptor form and function using structural biology. Journal of molecular endocrinology. 2013;51:T1-T21.

[14] Olefsky JM. Nuclear receptor minireview series. The Journal of biological chemistry. 2001;276:36863-4.

[15] Bledsoe RK, Montana VG, Stanley TB, Delves CJ, Apolito CJ, McKee DD, et al. Crystal structure of the glucocorticoid receptor ligand binding domain reveals a novel mode of receptor dimerization and coactivator recognition. Cell. 2002;110:93-105.

[16] Li Y, Suino K, Daugherty J, Xu HE. Structural and biochemical mechanisms for the specificity of hormone binding and coactivator assembly by mineralocorticoid receptor. Mol Cell. 2005;19:367-80.

[17] Williams SP, Sigler PB. Atomic structure of progesterone complexed with its receptor. Nature. 1998;393:392-6.

[18] Brzozowski AM, Pike AC, Dauter Z, Hubbard RE, Bonn T, Engstrom O, et al. Molecular basis of agonism and antagonism in the oestrogen receptor. Nature. 1997;389:753-8.

[19] Godowski PJ, Rusconi S, Miesfeld R, Yamamoto KR. Glucocorticoid receptor mutants that are constitutive activators of transcriptional enhancement. Nature. 1987;325:365-8.

[20] Hollenberg SM, Giguere V, Segui P, Evans RM. Colocalization of DNA-binding and transcriptional activation functions in the human glucocorticoid receptor. Cell. 1987;49:39-46.

[21] Simons SS, Jr., Kumar R. Variable steroid receptor responses: Intrinsically disordered AF1 is the key. Molecular and cellular endocrinology. 2013;376:81-4.

[22] Rastinejad F, Ollendorff V, Polikarpov I. Nuclear receptor full-length architectures: confronting myth and illusion with high resolution. Trends in biochemical sciences. 2015;40:16-24.

[23] Kumar R, Thompson EB. Folding of the glucocorticoid receptor N-terminal

transactivation function: dynamics and regulation. Molecular and cellular endocrinology.

2012;348:450-6.

[24] Khan SH, Awasthi S, Guo C, Goswami D, Ling J, Griffin PR, et al. Binding of the

N-terminal region of coactivator TIF2 to the intrinsically disordered AF1 domain of the

glucocorticoid receptor is accompanied by conformational reorganizations. The Journal

of biological chemistry. 2012;287:44546-60.

[25] Kumar R, Thompson EB. Influence of flanking sequences on signaling between the

activation function AF1 and DNA-binding domain of the glucocorticoid receptor. Arch

Biochem Biophys. 2010;496:140-5.

[26] Li J, Motlagh HN, Chakuroff C, Thompson EB, Hilser VJ. Thermodynamic

dissection of the intrinsically disordered N-terminal domain of human glucocorticoid

receptor. The Journal of biological chemistry. 2012;287:26777-87.

[27] Meijsing SH, Pufall MA, So AY, Bates DL, Chen L, Yamamoto KR. DNA binding

site sequence directs glucocorticoid receptor structure and activity. Science.

2009;324:407-10.

[28] Watson LC, Kuchenbecker KM, Schiller BJ, Gross JD, Pufall MA, Yamamoto KR.

The glucocorticoid receptor dimer interface allosterically transmits sequence-specific

DNA signals. Nature structural & molecular biology. 2013;20:876-83.

[29] Miesfeld R, Godowski PJ, Maler BA, Yamamoto KR. Glucocorticoid receptor

mutants that define a small region sufficient for enhancer activation. Science.

1987;236:423-7.

[30] Hellal-Levy C, Couette B, Fagart J, Souque A, Gomez-Sanchez C, Rafestin-Oblin

M. Specific hydroxylations determine selective corticosteroid recognition by human

glucocorticoid and mineralocorticoid receptors. FEBS Lett. 1999;464:9-13.

[31] Rupprecht R, Arriza JL, Spengler D, Reul JM, Evans RM, Holsboer F, et al.

Transactivation and synergistic properties of the mineralocorticoid receptor: relationship

to the glucocorticoid receptor. Molecular endocrinology (Baltimore, Md ).

1993;7:597-603.

[32] Zhang J, Simisky J, Tsai FT, Geller DS. A critical role of helix 3-helix 5 interaction

in steroid hormone receptor function. Proc Natl Acad Sci U S A. 2005;102:2707-12.

[33] Lim-Tio SS, Keightley MC, Fuller PJ. Determinants of specificity of

transactivation by the mineralocorticoid or glucocorticoid receptor. Endocrinology.

1997;138:2537-43.

[34] Oka K, Kohno S, Urushitani H, Guillette LJ, Jr., Ohta Y, Iguchi T, et al. Molecular

cloning and characterization of the corticoid receptors from the American alligator.

Molecular and cellular endocrinology. 2013;365:153-61.

[35] Proszkowiec-Weglarz M, Porter TE. Functional characterization of chicken

glucocorticoid and mineralocorticoid receptors. American journal of physiology

Regulatory, integrative and comparative physiology. 2010;298:R1257-68.

[36] Arriza JL, Simerly RB, Swanson LW, Evans RM. The neuronal mineralocorticoid

receptor as a mediator of glucocorticoid response. Neuron. 1988;1:887-900.

[37] Bridgham JT, Carroll SM, Thornton JW. Evolution of hormone-receptor

complexity by molecular exploitation. Science. 2006;312:97-101.

[38] Sturm A, Colliar L, Leaver MJ, Bury NR. Molecular determinants of hormone

sensitivity in rainbow trout glucocorticoid receptors 1 and 2. Molecular and cellular

endocrinology. 2011;333:181-9.

[39] Fuller PJ, Yao Y, Yang J, Young MJ. Mechanisms of ligand specificity of the

mineralocorticoid receptor. J Endocrinol. 2012;213:15-24.

[40] Hawkins UA, Gomez-Sanchez EP, Gomez-Sanchez CM, Gomez-Sanchez CE. The

ubiquitous mineralocorticoid receptor: clinical implications. Curr Hypertens Rep.

2012;14:573-80.

[41] Rossier BC, Baker ME, Studer RA. Epithelial sodium transport and its control by

aldosterone: the story of our internal environment revisited. Physiological reviews.

2015;95:297-340.

[42] Close DA, Yun SS, McCormick SD, Wildbill AJ, Li W. 11-deoxycortisol is a

corticosteroid hormone in the lamprey. Proc Natl Acad Sci U S A. 2010;107:13942-7.

[43] Baker ME, Chandsawangbhuwana C, Ollikainen N. Structural analysis of the

evolution of steroid specificity in the mineralocorticoid and glucocorticoid receptors.

BMC Evol Biol. 2007;7:24.

[44] Oka K, Hoang A, Okada D, Iguchi T, Baker ME, Katsu Y. Allosteric role of the

amino-terminal A/B domain on corticosteroid transactivation of gar and human

glucocorticoid receptors. J Steroid Biochem Mol Biol. 2015;154:112-9.

[45] Christopoulos A, Changeux JP, Catterall WA, Fabbro D, Burris TP, Cidlowski JA,

et al. International union of basic and clinical pharmacology. XC. multisite

pharmacology: recommendations for the nomenclature of receptor allosterism and

allosteric ligands. Pharmacological reviews. 2014;66:918-47.

[46] Kumar R, Litwack G. Structural and functional relationships of the steroid hormone

receptors' N-terminal transactivation domain. Steroids. 2009;74:877-83.

[47] Thomas-Chollier M, Watson LC, Cooper SB, Pufall MA, Liu JS, Borzym K, et al.

A naturally occurring insertion of a single amino acid rewires transcriptional regulation

by glucocorticoid receptor isoforms. Proc Natl Acad Sci U S A. 2013;110:17826-31.

[48] Cho S, Blackford JA, Jr., Simons SS, Jr. Role of activation function domain-1,

DNA binding, and coactivator GRIP1 in the expression of partial agonist activity of

glucocorticoid receptor-antagonist complexes. Biochemistry. 2005;44:3547-61.

[49] Lu NZ, Cidlowski JA. Glucocorticoid receptor isoforms generate transcription

specificity. Trends Cell Biol. 2006;16:301-7.

[50] Arango-Lievano M, Lambert WM, Bath KG, Garabedian MJ, Chao MV, Jeanneteau

F. Neurotrophic-priming of glucocorticoid receptor signaling is essential for neuronal

plasticity to stress and antidepressant treatment. Proc Natl Acad Sci U S A.

2015;112:15737-42.

[51] Hua G, Paulen L, Chambon P. GR SUMOylation and formation of an

SUMO-SMRT/NCoR1-HDAC3 repressing complex is mandatory for GC-induced IR

nGRE-mediated transrepression. Proc Natl Acad Sci U S A. 2016;113:E626-34.

[52] Katsu Y, Kubokawa K, Urushitani H, Iguchi T. Estrogen-dependent transactivation

of amphioxus steroid hormone receptor via both estrogen and androgen response

elements. Endocrinology. 2010;151:639-48.

[53] Bury NR, Sturm A, Le Rouzic P, Lethimonier C, Ducouret B, Guiguen Y, et al.

Evidence for two distinct functional glucocorticoid receptors in teleost fish. J Mol

Endocrinol. 2003;31:141-56.

[54] Greenwood AK, Butler PC, White RB, DeMarco U, Pearce D, Fernald RD.

Multiple corticosteroid receptors in a teleost fish: distinct sequences, expression patterns,

and transcriptional activities. Endocrinology. 2003;144:4226-36.

[55] Stolte EH, de Mazon AF, Leon-Koosterziel KM, Jesiak M, Bury NR, Sturm A, et al.

Corticosteroid receptors involved in stress regulation in common carp, Cyprinus carpio.

J Endocrinol. 2008;198:403-17.

[56] Sturm A, Bury N, Dengreville L, Fagart J, Flouriot G, Rafestin-Oblin ME, et al.

11-deoxycorticosterone is a potent agonist of the rainbow trout (Oncorhynchus mykiss)

mineralocorticoid receptor. Endocrinology. 2005;146:47-55.

[57] Carroll SM, Bridgham JT, Thornton JW. Evolution of hormone signaling in

elasmobranchs by exploitation of promiscuous receptors. Molecular biology and

evolution. 2008;25:2643-52.

[58] Carroll SM, Ortlund EA, Thornton JW. Mechanisms for the evolution of a derived

function in the ancestral glucocorticoid receptor. PLoS Genet. 2011;7:e1002117.

[59] Rogerson FM, Yao YZ, Elsass RE, Dimopoulos N, Smith BJ, Fuller PJ. A critical

region in the mineralocorticoid receptor for aldosterone binding and activation by

cortisol: evidence for a common mechanism governing ligand binding specificity in

steroid hormone receptors. Mol Endocrinol. 2007;21:817-28.

[60] Romero LM. Seasonal changes in plasma glucocorticoid concentrations in free-living vertebrates. Gen Comp Endocrinol. 2002;128:1-24.

[61] Cockrem JF. Individual variation in glucocorticoid stress responses in animals. Gen Comp Endocrinol. 2013;181:45-58.

[62] Martinerie L, Munier M, Le Menuet D, Meduri G, Viengchareun S, Lombes M. The mineralocorticoid signaling pathway throughout development: expression, regulation and pathophysiological implications. Biochimie. 2013;95:148-57.

[63] Odermatt A, Atanasov AG. Mineralocorticoid receptors: emerging complexity and functional diversity. Steroids. 2009;74:163-71.

[64] Funder JW. Aldosterone and mineralocorticoid receptors: a personal reflection. Molecular and cellular endocrinology. 2012;350:146-50.

[65] Odermatt A, Kratschmar DV. Tissue-specific modulation of mineralocorticoid receptor function by 11beta-hydroxysteroid dehydrogenases: an overview. Molecular and cellular endocrinology. 2012;350:168-86.

[66] Baker ME, Nelson DR, Studer RA. Origin of the response to adrenal and sex steroids: Roles of promiscuity and co-evolution of enzymes and steroid receptors. J Steroid Biochem Mol Biol. 2015;151:12-24.

[67] Mani O, Nashev LG, Livelo C, Baker ME, Odermatt A. Role of Pro-637 and Gln-642 in human glucocorticoid receptors and Ser-843 and Leu-848 in mineralocorticoid receptors in their differential responses to cortisol and aldosterone. J Steroid Biochem Mol Biol. 2016;159:31-40.

[68] Pippal JB, Yao Y, Rogerson FM, Fuller PJ. Structural and functional characterization of the interdomain interaction in the mineralocorticoid receptor. Mol Endocrinol. 2009;23:1360-70.

[69] Rogerson FM, Fuller PJ. Interdomain interactions in the mineralocorticoid receptor. Molecular and cellular endocrinology. 2003;200:45-55.

[70] Pippal JB, Cheung CM, Yao YZ, Brennan FE, Fuller PJ. Characterization of the zebrafish (Danio rerio) mineralocorticoid receptor. Molecular and cellular endocrinology. 2011;332:58-66.