1   **Title: How clonal are bacteria over time?**

3   **Author: B. Jesse Shapiro[1]**

5   **Affiliation**

6   1. Département de sciences biologiques, Université de Montréal, Montréal, QC H3C 3J7,

7   Canada

* jesse.shapiro@umontreal.ca

Abstract

Bacteria and archaea reproduce clonally (vertically), but exchange genes by recombination

(horizontal transfer). Recombination allows adaptive mutations or genes to spread within (or

between) species. Clonality – the balance between vertical and horizontal inheritance – is

therefore a key microbial trait, determining how quickly a population can adapt. Here, I consider

whether clonality can be considered a stable trait of a given population. In some cases, clonality

changes over time: non-clonal (recombining) populations can give rise to clonal expansions.

However, an analysis of time-course metagenomic data suggests that a bacterial population's past

clonality is indicative of its future clonality. Thus, a population's evolutionary potential – whether

it is likely to retain genetic diversity or not – can in principle be predicted from its past.

Introduction

Here, I revisit the question posed in the title of a classic paper by John Maynard Smith and colleagues [1]: How clonal are bacteria? More specifically how does clonality vary among different microbial populations and over time? First, what do we mean by clonality? Perfectly clonal bacteria replicate by cell division (vertical descent) and evolve by random mutations that occur during DNA replication. There is negligible horizontal transfer of DNA by recombination across the resulting tree of vertical descent. Very few (if any) natural bacterial populations fit this theoretical definition of clonality. Or, as discussed below, they might only fit it for a short amount of time. However, knowing where a bacterial population happens to fall along a spectrum of clonality can help us understand its biology, and even make predictions about its evolution.

The opposite of clonality is panmixis – a situation in which the rate of horizontal transfer is higher than the rate of vertical cell division [1,2]. However, rates of horizontal transfer (recombination) vary widely across the genome, such that a population can be mostly clonal, except for a few loci in the genome. These loci came to be termed genomic islands – a metaphor I will build upon below. Some of the first islands identified were called pathogenicity islands because they contained virulence factors [3]. However, non-pathogenic environmental bacteria also contain islands, conferring adaptation to different ecological niches. For example, genes in *Prochlorococcus* genomic islands confer adaptation to light and nutrient conditions [4,5]. But islands need not confer niche adaptation to their host genome; they can be neutral to host fitness or even detrimental, selfish parasites. Here, I define genomic islands broadly as any piece of DNA that is transferred horizontally (by either homologous or nonhomologous recombination) from cell to cell and therefore evolves independently (*i.e.* is unlinked) from the rest of the genome.

Are some islands really peninsulas?

In the classic analogy, an island is totally disconnected from the mainland, meaning that genes in

the island evolve independently of the genome (**Table 1**). Examples of islands that fit this strict

independence might include integrated phages and other "selfish" elements, or genes that reside

in a particular niche but not in a particular genome (*e.g.* a gene ecology model [6]). Peninsulas

provide an analogy that might better describe how islands are related to microbial genomes. A

peninsula (or "presque-ile," from the French for "almost island") is a geographic term for a very

narrow strip of land connected to (but distinct from) the mainland. An island is evolutionarily

independent of the mainland genome, but their fates may become linked, forming a peninsula. For

example, a bacterium may acquire a gene from a vast microbial gene pool. This gene allows the

bacterium to invade a new ecological niche, triggering a clonal expansion in which the fate of the

gene and its new host genome are linked, at least for the duration of the clonal expansion. One

such example could be *Yersinia pestis,* which acquired a single gene allowing flea-borne

transmission and triggering a clonal expansion in the form of Plague pandemics [7]. Another

peninsula, the prophage-encoded cholera toxin, and its links to the mainland *Vibrio cholerae*

genome [8,9], is discussed below.

Are some genomes archipelagos?

The very concept of one or a few islands implies a contrast with the large, clonal genomic

mainland or continent. But some microbial genomes may contain so many islands that there is no

mainland, only a vast archipelago (**Table 1**). A striking recent example is a population of

hotspring cyanobacteria in which virtually every gene in the genome evolved independently due

to frequent recombination [10], leading the authors to call the population "quasi-sexual"

(panmictic). The Asian ocean population of *Vibrio parahaemolyticus* also form a panmictic gene

pool, with each recombination block of ~1.8 kbp evolving independently [11]. Similarly, almost

every gene in a population of *Vibrio cyclitrophicus* genomes showed signals of recombination

over relatively recent time scales [12]. Such apparently high rates of recombination in natural

populations were mysterious at first, contradicting recombination rates measured in the lab

[13,14] and predicted by theory [15,16]. However, new models (discussed below) suggest

mechanisms capable of explaining such surprisingly panmictic populations [17,18].

Clonal expansions from panmictic pools

Archipelagos are not necessarily static over time; they can sometimes coalesce into continents.

Given the right ecological opportunity, a genome from a panmictic gene pool can escape the

"gravitational pull" of recombination and take off into a clonal expansion (resulting in "epidemic"

population structure [1]). An example mentioned earlier is *V. cholerae*, a genetically diverse

group of coastal marine bacteria, some of which cause cholera. Virulence is mainly determined

by two loci in the genome: the cholera toxin and the toxin-coregulated pilus. Both genes are

frequently gained and lost by recombination [19,20], but are always found in one lineage of *V.*

cholerae – the lineage causing severe disease with pandemic potential, known as the phylocore

genome (PG) group [9]. It remains a mystery why the PG lineage evolved once, and only once. If

PG *V. cholerae* really did evolve just once, this would be surprising because *V. cholerae* draws on

a diverse, global gene pool and can be considered panmictic [21]. Therefore multiple different

lineages would be expected to acquire the two (or perhaps a handful of) genetic elements required

for pandemic disease. This leads to the hypothesis that pandemic cholera emergence is *selection*

limited rather than *diversity limited*. In other words, benign *V. cholerae* strains constantly acquire

virulence genes. However, these strains rarely encounter the right ecological niche to flourish,

e.g. a human population consuming brackish water. "The right niche" has appeared a few times in

human history: for example in India in the 1800s, when the Classical lineage evolved, and again

in Indonesia in the 1950s, when the El Tor lineage evolved [22]. El Tor and Classical lineages

have different cholera toxin alleles, but both evolved from within the PG group, suggesting that

the PG genomic background is adapted to acquire and exploit virulence genes. When the "right"

conditions appear, the PG lineage, along with its virulence factors, takes off in a clonal

expansions which continue to wreak havoc today (*e.g.* cholera pandemics from the 1800s to

today, all caused by the PG clonal group). The virulence factors, previously islands in an

archipelago, became a peninsula connected to the PG mainland. The linkage between virulence

factors and PG remains imperfect because different variants of the cholera toxin continue to flow

in and out of the PG continent [9,19]; hence the toxin remains a peninsula, not firmly part of the

mainland.

V. cholerae is a particularly well-characterized example of a panmictic gene pool giving

rise to a clonal expansion, but similar evolutionary dynamics are seen in other pathogens as well.

For example, enterotoxigenic *Escherichia coli* (ETEC) seems to behave similarly, with deep

branches of the phylogeny obscured by frequent recombination and plasmid exchange, but more

recent branches experiencing mostly clonal descent, with tight linkage between virulence factors

and the genomic mainland [23]. These observations are consistent with an ancient, diverse

panmictic gene pool giving rise to clonal expansions, which can last for decades or centuries.

The balance between recombination and selection

Let us consider the evolutionary forces that determine clonality: natural selection and

recombination. The effect of recombination on clonality is straightforward: more recombination

means less clonality. The effect of natural selection is more complex, but is defined here simply

as a force which favors clonal expansions of adaptive mutants within an ecological niche. When

driven by ecological selection, clonal expansions are called selective sweeps, in which one clone

outcompetes all others, purging genetic diversity in the population [15,16].

Recombination and selection interact to determine the clonality of a population.

Recombination rates depend both on the ability of DNA to enter a cell and be incorporated into

the genome (the baseline rate) and the ability of that DNA to be retained by a balance of genetic

drift and natural selection (the realized rate). Some bacteria, such as *Helicobacter pylori,* have

realized recombination rates that are much higher than point mutation rates, exchanging at least

10% of their genome within a single four-year human infection [24]. Others, such as

Staphylococcus aureus [25,26] and *Mycobacterium tuberculosis* [27-29] are decidedly more

clonal. Recombination rates (both realized and baseline) vary widely across the genome. Of 10

pathogenic bacterial species studied, all had identifiable recombination 'hot' regions, although

their length, genomic location and gene content varied [30]. Genes of different functions had

different realized recombination rates, implying a role for natural selection on gene function in

determining whether newly acquired genes are retained.

Modeling the recombination-selection balance

When rates of recombination are relatively low compared to rates of natural selection (s) on

adaptive genes within niches, entire genomes will sweep to fixation before they can be shuffled

by recombination. The $s \gg r$ regime is well described in the Stable Ecotype Model [15], which

predicts that most of the genome will follow a single, clonal phylogeny. Genome-wide sweeps

thus increase clonality and can be considered a hallmark of clonal populations (**Table 1, Figure

1a**). In the $r \gg s$ regime, individual genes (rather than entire genomes) will sweep to fixation

(*i.e.* reach 100% frequency) in ecological niches to which they are adapted, without affecting

genetic diversity elsewhere in the genome (**Figure 1b**). A recent model showed how these gene

sweeps can occur at moderate (not unrealistically high) rates of recombination [18]. In this model,

a microbial habitat is bombarded with genetically maladapted migrants, allowing gene sweeps to

occur, although the adaptive allele never reaches 100% frequency due to the constant input of

migrants. In another model, Takeuchi et al. [17] show that gene sweeps can occur when r is either

very high or – counter-intuitively – when r is very low, but only when negative frequency-

dependent selection (NFDS) reduces the rate of genome-wide selective sweeps. NFDS might be

commonly imposed on bacteria and archaea by viral (phage) predation, providing a selective

advantage to rare alleles of phage receptor genes [31].

Genome-wide and gene-specific sweeps in nature

To date, empirical evidence for gene-specific and genome-wide sweeps has come mostly from

cross-sectional studies of a single population of genomes at a single point in time, with

recombination and selection inferred backward in time [10-12,32]. Sequencing microbial

genomes or metagenomes sampled over time – already a typical practice in genomic

epidemiology (e.g. [26,33]) – promises to elucidate the rates of gene-specific and genome-wide

sweeps in nature (**Figure 1**).

In a pioneering study, Bendall et al. [34] sampled a lake over 9 years and followed single-

nucleotide polymorphism (SNP) and gene frequencies in 30 bacterial populations by

metagenomic sequencing. They inferred that one of the populations (*Chlorobium*-111) had

undergone a near-complete genome-wide sweep over the 9-year study, with most SNP diversity

purged genome-wide (**Figure 1a**). In 6 other populations, they identified regions of the genome

with unexpectedly low diversity compared to the genome-wide average, throughout the time-

course. Either strong purifying selection on these regions or gene-specific selective sweeps

(completed before the start of the time-course) could explain the low diversity. During the 9-year

study, they also observed examples "where a few adjacent SNPs trended toward fixation while

genome-wide diversity was maintained" (**Figure 1b**). They took this observation as consistent

with gene-specific selective sweeps, but did not attempt to determine whether the sweeps were

due to selection or drift. As a whole, the study showed that both genome-wide and gene-specific

sweeps can occur in different microbial populations from the same environment. Whether

microbial populations behaved differently due to differences in their ecology (*i.e.* regime of

natural selection) or in their baseline recombination rates remains a question for future study. The

fact that a genome-wide sweep was observed over a 9 year period suggests that such events might

be relatively rapid but rare (only observed in 1 of 30 populations). Meanwhile, gene-sweeps

might be more common historically (affecting 6 of 30 populations), but could take longer to

proceed to completion.

Is clonality a stable trait?

As described in the *V. cholerae* example, some pathogenic bacterial populations can switch

between panmictic and clonal lifestyles. Therefore clonality can vary over time, but how much

and how often? To quantify the stability of clonality over time, I re-analyzed the lake time-course

of Bendall et al. [34]. Because estimates of selection and recombination rates were not readily

available for this dataset, I defined clonality based on the frequency of genome-wide selective

sweeps. Frequent genome-wide selective sweeps suggest $s \gg r$, suggesting clonality. I identified

20 "old, diverse" populations as those with a high density of SNPs (>1500 SNPs/Mbp) at the

beginning of the time-course. These populations are likely "old" because they have gone a

relatively long time since the last genome-wide purge of genetic diversity and are relatively non-

clonal. They include the 6 populations inferred to have undergone gene-specific sweeps [34]. The

remaining 10 populations were defined as "young, low-diversity," having more recently

experienced a genome-wide purge of diversity. The "old, diverse" populations have a relatively

low ratio of nonsynonymous (N) to synonymous (S) SNPs, suggesting large effective population

sizes and ample time for purifying selection to remove (mostly deleterious) nonsynonymous

mutations (**Figure 2**). In contrast, the "young, low-diversity" populations are more likely to have

high N:S ratios, suggesting smaller effective population sizes and less time for purifying selection

to have acted.

With young (clonal) and old (less clonal) populations thus defined, I asked whether "old,

diverse" populations tended to maintain their diversity through the 9-year period of the study.

Bendall et al. defined two alternative population types: 1) those that maintained stable SNP

diversity over 9 years, and 2) those that experienced significant fluctuations in diversity due to

clonal expansions – defined when one, but not all timepoints are dominated by a single allele

(≥95% frequency) at >40% of SNP sites in the genome [34]. By definition, the 19 populations of

the first type did not experience genome-wide sweeps during the study, while the 11 populations

of the second type did experience genome-wide purges of diversity, which were transient in 10

cases and apparently permanent in 1 case (*Chlorobium*-111). Strikingly, 17 out of 20 "old,

diverse" populations maintained their diversity over the 9-year study, compared to only 2 out of

10 "young, low-diversity" populations (Fisher test, Odds Ratio = 19.4, $P < 0.001$). This result

suggests that populations with a history of genome-wide sweeps (and/or population bottlenecks)

tend to experience subsequent genome-wide sweeps, and those that have maintained genetic

diversity in the past tend to maintain their diversity into the future. In other words, clonality can

be considered a relatively stable microbial trait.

History repeats itself

It appears that pathogens are more likely than free-living bacteria to undergo clonal expansions,

due in part to their ecology and transmission dynamics [1,35]. Free-living aquatic bacteria, on the

other hand, seem to be more likely to live in large, panmictic populations and behave like

archipelagos [10-12,34]. If clonality is indeed a stable trait, this implies that history will repeat

itself, and that the future behavior of microbial populations can be predicted with some

confidence from their past behavior. Diverse populations tend to stay diverse. Clonal populations

(that experience frequent genome-wide sweeps) tend to stay clonal. But history is not doomed to

repeat itself forever. As we have seen, clonal expansions, such as pandemic *V. cholerae,* may

originate when a panmictic gene pool (an archipelago) coalesces into a clonal continent, with

virulence factors linked as peninsulas. Many such pathogenic clones have been documented, with

life spans of decades to thousands of years [23,26,36,37]. Other pathogens, such as *Streptococcus*

pneumoniae, may retain their panmictic population structure throughout an outbreak [38-40].

Why some pathogens are clonal and others are panmictic is an open question, but surely depends

on the balance between recombination and selection, and on the time scales considered.

Acknowledgments. I am grateful to Yan Boucher and Salvador Almagro-Moreno for their

thoughtful comments which improved the manuscript. I was supported by a Canada Research

Chair.

Table 1. Extended island metaphors of microbial genome evolution.
232

Geographic metaphor	Genetic unit to which the metaphor applies	Type of selective sweep experience by the unit	Dominant mode of genetic transmission	Example
Island	Gene	Gene-specific	horizontal	genes in the *V. cholerae* integron [20,21]
Peninsula	Gene	Genome-wide	vertical (clonal)	the cholera toxin gene, acquired horizontally, then linked to a clonal *V. cholerae* genome [8,19]
Continent	Genome	Genome-wide	vertical (clonal)	clonal expansions of *S. aureus* [26] and *M. tuberculosis* [29,37]
Archipelago	Genome	Gene-specific	horizontal	hotspring cyanobacteria [10], ocean vibrios [11,12], pneumococcus [38,40]

233
234
235   **Figures**
236
Figure 1. Temporal dynamics of genome-wide and gene-specific selective sweeps
inferred from metagenomic data. Genetic diversity can be measured by mapping
metagenomic sequence reads to a reference genome, identifying SNPs, and calculating
the allele frequencies at each SNP position in the genome over time. The lowest possible
genetic diversity occurs when a single allele is present in 100% of metagenomic reads.
Alternatively, diversity could be defined in terms of gene presence/absence, based on
relative coverage of a gene in the reference genome by metagenomic reads. **(a)** In a
hypothetical genome-wide selective sweep, all positions in the genome tend toward low
diversity over time. **(b)** In a hypothetical gene-specific selective sweep, only one or a few
positions in the genome tend toward low diversity, while the rest of the genome maintains
high or intermediate diversity.
248
Figure 2. Past diversity predicts future diversity. Based on data from Table 2 of
Bendall et al. [34], the genome-wide average nonsynonymous to synonymous (N:S) SNP
ratio was plotted against the total SNP density (SNPs per megabasepair) for each of 30
bacterial populations. A pseudocount of 1 was added to both N and S counts. These 30
populations were divided into 20 "old, diverse" populations (>1500 SNPs/Mbp) and 10
"young, low-diversity" populations (<1500 SNPs/Mbp), highlighted in blue and yellow,
respectively. Each point represents one of the 30 populations, colored in black if diversity
was maintained over a 9-year metagenomic time-course, or in red if it was not. Seventeen
out of 20 "old, diverse" populations maintained their diversity over the 9-year study,
compared to only 2 out of 10 "young, low-diversity" populations (Fisher test, Odds Ratio
= 19.4, $P < 0.001$). Consistent with previous observations that N:S depends on the
evolutionary time available for purifying selection to act [41,42], N:S is negatively
correlated with SNPs/Mbp, a proxy for evolutionary time or the time since the last
genome-wide purge of genetic diversity in this dataset (Pearson's correlation of $\log_{10}$
transformed data, r = -0.81, P = 5.6e-8).

References

1. Smith JM, Smith NH, O'Rourke M, Spratt BG: **How clonal are bacteria?** *Proc. Natl. Acad. Sci. USA* 1993, **90**:4384–4388.

2. Guttman DS, Dykhuizen DE: **Clonal Divergence in Escherichia coli as a Result of Recombination, Not Mutation**. *Science* 1994, **266**:1380–1383.

3. Hacker J, Blum-Oehler G, Mühldorfer I, Tschäpe H: **Pathogenicity islands of virulent bacteria: structure, function and impact on microbial evolution.** *Mol Microbiol* 1997, **23**:1089–1097.

4. Coleman ML, Sullivan MB, Martiny AC, Steglich C, Barry K, Delong EF, Chisholm SW: **Genomic islands and the ecology and evolution of *Prochlorococcus*.** *Science* 2006, **311**:1768–1770.

5. Coleman ML, Chisholm SW: **Ecosystem-specific selection pressures revealed through comparative population genomics**. *Proc. Natl. Acad. Sci. USA* 2010, **107**:18634–18639.

6. Shapiro BJ: **Signatures of natural selection and ecological differentiation in microbial genomes.** *Advances in experimental medicine and biology* 2014, **781**:339–359.

7. Rasmussen S, Allentoft ME, Nielsen K, Orlando L, Sikora M, Sjögren K-G, Pedersen AG, Schubert M, Van Dam A, Kapel CMO, et al.: **Early Divergent Strains of *Yersinia pestis* in Eurasia 5,000 Years Ago.** *Cell* 2015, **163**:571–582.

8. Waldor MKM, Mekalanos JJJ: **Lysogenic conversion by a filamentous phage encoding cholera toxin.** *Science* 1996, **272**:1910–1914.

9. Chun J, Grim CJ, Hasan NA, Lee JH, Choi SY, Haley BJ, Taviani E, Jeon Y-S, Kim DW, Lee J-H, et al.: **Comparative genomics reveals mechanism for short-term and long-term clonal transitions in pandemic *Vibrio cholerae*.** *Proc. Natl. Acad. Sci. USA* 2009, **106**:15442–15447.

10. Rosen MJ, Davison M, Bhaya D, Fisher DS: **Fine-scale diversity and extensive recombination in a quasisexual bacterial population occupying a broad niche.** *Science* 2015, **348**:1019–1023.

* Deep sequencing of 90 cyanobacterial marker genes reveals extensive recombination, with each genome composed of a random mix of alleles from the gene pool.

11. Cui Y, Yang X, Didelot X, Guo C, Li D, Yan Y, Zhang Y, Yuan Y, Yang H, Wang J, et al.: **Epidemic Clones, Oceanic Gene Pools, and Eco-LD in the Free Living Marine Pathogen *Vibrio parahaemolyticus*.** *Mol Biol Evol* 2015, **32**:1396–1410.

* In-depth population genomic evidence of a panmictic ocean gene pool containing non-clonal ecological populations.

12. Shapiro BJ, Friedman J, Cordero OX, Preheim SP, Timberlake SC, Szabo G, Polz MF, Alm EJ: **Population Genomics of Early Events in the Ecological Differentiation of Bacteria.** *Science* 2012, **336**:48–51.

13. Vulić M, Dionisio F, Taddei F, Radman M: **Molecular keys to speciation: DNA**
polymorphism and the control of genetic exchange in enterobacteria. *Proc. Natl.*
Acad. Sci. USA 1997, **94**:9763–9767.

14. Majewski JJ, Cohan FMF: **DNA sequence similarity requirements for interspecific**
recombination in *Bacillus*. *Genetics* 1999, **153**:1525–1533.

15. Wiedenbeck J, Cohan FM: **Origins of bacterial diversity through horizontal genetic**
transfer and adaptation to new ecological niches. *FEMS Microbiology Reviews* 2011,
35:957–976.

16. Shapiro BJ, Polz MF: **Ordering microbial diversity into ecologically and genetically**
cohesive units. *Trends Microbiol* 2014, **22**:235–247.

* A comprehensive discussion of the roles of selection and recombination in structuring microbial
diversity.
316
317 17. Takeuchi N, Cordero OX, Koonin EV, Kaneko K: **Gene-specific selective sweeps in**
318 **bacteria and archaea caused by negative frequency-dependent selection**. *BMC*
319 *Biology* 2015, **13**:20.

320 * Mathematical modeling shows how individual genes can sweep through populations with low
321 recombination rates in the presence of negative frequency-dependent selection.
322
18. Niehus R, Mitri S, Fletcher AG, Foster KR: **Migration and horizontal gene transfer**
divide microbial genomes into multiple niches. *Nature Communications* 2015, **6**:1–9.

19. Mutreja A, Kim DW, Thomson NR, Connor TR, Lee JH, Kariuki S, Croucher NJ, Choi
SY, Harris SR, Lebens M, et al.: **Evidence for several waves of global transmission in**
the seventh cholera pandemic. *Nature* 2011, **477**:462–U111.

20. Orata FD, Kirchberger PC, Méheust R, Barlow EJ, Tarr CL, Boucher Y: **The Dynamics**
of Genetic Interactions between *Vibrio metoecus* and *Vibrio cholerae*, Two Close
Relatives Co-Occurring in the Environment. *Genome Biol Evol* 2015, **7**:2941–2954.

* A clear demonstration that genes in the integron are recombined between species just as often
as within species, and that 24% of 'core' genes have crossed species boundaries.
333
21. Boucher Y, Cordero OX, Takemura A, Hunt DE, Schliep K, Bapteste E, Lopez P, Tarr
CL, Polz MF: **Local Mobile Gene Pools Rapidly Cross Species Boundaries To Create**
Endemicity within Global Vibrio cholerae Populations. *mBio* 2011, **2**:e00335–10.

22. Boucher Y, Orata FD, Alam M: **The out-of-the-delta hypothesis: dense human**
populations in low-lying river deltas served as agents for the evolution of a deadly
pathogen. *Front Microbiol* 2015, **6**:1120.

23. Mentzer von A, Connor TR, Wieler LH, Semmler T, Iguchi A, Thomson NR, Rasko DA,
Joffre E, Corander J, Pickard D, et al.: **Identification of enterotoxigenic *Escherichia coli***
(ETEC) clades with long-term global distribution. *Nature Genetics* 2014, **46**:1321–
1326.

24. Cao Q, Didelot X, Wu Z, Li Z, He L, Li Y, Ni M, You Y, Lin X, Li Z, et al.: **Progressive**
genomic convergence of two *Helicobacter pylori* strains during mixed infection of a
patient with chronic gastritis. *Gut* 2015, **64**:554–561.

25. Vos M, Didelot X: **A comparison of homologous recombination rates in bacteria and**
archaea. *The ISME journal* 2009, **3**:199–208.

26. Baines SL, Holt KE, Schultz MB, Seemann T, Howden BO, Jensen SO, van Hal SJ,
Coombs GW, Firth N, Powell DR, et al.: **Convergent adaptation in the dominant global**
hospital clone ST239 of methicillin-resistant *Staphylococcus aureus*. *mBio* 2015,
6:e00080.

27. Hershberg R, Lipatov M, Small PM, Sheffer H, Niemann S, Homolka S, Roach JC,
Kremer K, Petrov DA, Feldman MW, et al.: **High Functional Diversity in**
***Mycobacterium tuberculosis* Driven by Genetic Drift and Human Demography**. *Plos*
Biol 2008, **6**:e311.

28. Namouchi A, Didelot X, Schock U, Gicquel B, Rocha EPC: **After the bottleneck:**
Genome-wide diversification of the *Mycobacterium tuberculosis* complex by mutation,
recombination, and natural selection. *Genome Research* 2012, **22**:721-734.

29. Casali N, Nikolayevskyy V, Balabanova Y, Harris SR, Ignatyeva O, Kontsevaya I,
Corander J, Bryant J, Parkhill J, Nejentsev S, et al.: **Evolution and transmission of drug-**
resistant tuberculosis in a Russian population. *Nature Genetics* 2014, **46**:279–286.

30. Yahara K, Didelot X, Jolley KA, Kobayashi I, Maiden MCJ, Sheppard SK, Falush D: **The**
landscape of realized homologous recombination in pathogenic bacteria. *Mol Biol*
Evol 2015, doi:10.1093/molbev/msv237.

31. Cordero OX, Polz MF: **Explaining microbial genomic diversity in light of evolutionary**
ecology. *Nature Reviews Microbiology* 2014, **12**:263-273.

32. Cadillo-Quiroz H, Didelot X, Held NL, Herrera A, Darling A, Reno ML, Krause DJ,
Whitaker RJ: **Patterns of Gene Flow Define Species of Thermophilic Archaea**. *Plos*
Biol 2012, **10**:e1001265.

33. Croucher NJ, Didelot X: **The application of genomics to tracing bacterial pathogen**
transmission. *Current Opinion in Microbiology* 2015, **23**:62–67.

34. Bendall ML, Stevens SL, Chan L-K, Malfatti S, Schwientek P, Tremblay J, Schackwitz
W, Martin J, Pati A, Bushnell B, et al.: **Genome-wide selective sweeps and gene-specific**
sweeps in natural bacterial populations. *The ISME journal* 2016,
doi:10.1038/ismej.2015.241

* A pioneering application of time-course metagenomics to infer genome-wide and gene-specific
selective sweeps in natural lake bacterial populations.
379
35. Achtman M, Wagner M: **Microbial diversity and the genetic nature of microbial**
species. *Nature Reviews Microbiology* 2008, **6**:431–440.

36. Snitkin ES, Zelazny AM, Thomas PJ, Stock F, NISC Comparative Sequencing Program,

Henderson DK, Palmore TN, Segre JA: **Tracking a Hospital Outbreak of Carbapenem-**
Resistant *Klebsiella pneumoniae* **with Whole-Genome Sequencing**. *Science*
Translational Medicine 2012, **4**:148ra116.

37. Comas I, Coscolla M, Luo T, Borrell S, Holt KE, Kato-Maeda M, Parkhill J, Malla B,
Berg S, Thwaites G, et al.: **Out-of-Africa migration and Neolithic coexpansion of**
Mycobacterium tuberculosis **with modern humans.** *Nature Genetics* 2013, **45**:1176–
1182.

38. Croucher NJ, Finkelstein JA, Pelton SI, Mitchell PK, Lee GM, Parkhill J, Bentley SD,
Hanage WP, Lipsitch M: **Population genomics of post-vaccine changes in**
pneumococcal epidemiology. *Nature Genetics* 2013, **45**:656–663.

39. Chewapreecha C, Harris SR, Croucher NJ, Turner C, Marttinen P, Cheng L, Pessia A,
Aanensen DM, Mather AE, Page AJ, et al.: **Dense genomic sampling identifies**
highways of pneumococcal recombination. *Nature Genetics* 2014, **46**:305–309.

40. Marttinen P, Croucher NJ, Gutmann MU, Corander J, Hanage WP: **Recombination**
produces coherent bacterial species clusters in both core and accessory genomes.
Microbial Genomics 2015, **DOI: 10.1099/mgen.0.000038**.

* The first simulation of bacterial evolution including mutation, allelic exchange (homologous
recombination) and gene gain/loss (non-homologous recombination).
401
402 41. Rocha EP, Smith JM, Hurst LD, Holden MT, Cooper JE, Smith NH, Feil EJ:
403   **Comparisons of dN/dS are time dependent for closely related bacterial genomes**. *J*
404   *Theor Biol* 2006, **239**:226–235.

405 42. Kryazhimskiy S, Plotkin JB: **The Population Genetics of dN/dS**. *PLoS Genet* 2008,
406   **4**:e1000304.

Figure 1

(a) Genome-wide sweep

(b) Gene-specific sweeps

Legend

Genetic diversity

high

low

Position in the genome

time (years) ⟶

time (years) ⟶

← gene-specific sweep completed **during** the period of study

← gene-specific sweep completed **before** the period of study

Figure 2