1 **Title: How clonal are bacteria over time?**

3 **Author: B. Jesse Shapiro[1]**

5 **Affiliation**

6   1.   Département de sciences biologiques, Université de Montréal, Montréal, QC H3C 3J7,

7   Canada

* jesse.shapiro@umontreal.ca

Abstract

Bacteria and archaea reproduce clonally (vertical descent), but exchange genes by recombination

(horizontal transfer). Recombination allows adaptive mutations or genes to spread rapidly within

(or even between) species, and reduces the burden of deleterious mutations. Clonality – defined

here as the balance between vertical and horizontal inheritance – is therefore a key microbial trait,

determining how quickly a population can adapt and the size of its gene pool. Here, I discuss

whether clonality varies over time and if it can be considered a stable trait of a given population. I

show that, in some cases, clonality is clearly not static. For example, non-clonal (highly

recombining) populations can give rise to clonal expansions, often of pathogens. However, an

analysis of time-course metagenomic data from a lake suggests that a bacterial population's past

clonality (as measured by its genetic diversity) is a good predictor of its future clonality. Clonality

therefore appears to be relatively – but not completely – stable over evolutionary time.

Introduction

Here, I revisit the question posed in the title of a classic paper by John Maynard Smith and

colleagues [1]: How clonal are bacteria, and more specifically how does clonality vary among

different microbial populations and over time? First, what do we mean by clonality? Perfectly

clonal bacteria replicate by cell division (vertical descent) and evolve by random mutations that

occur during DNA replication. In this theoretical population, there is negligible horizontal

transfer of DNA by recombination across the resulting tree of vertical descent. Very few (if any)

natural bacterial populations fit this idealized, theoretical definition of clonality. Or, as discussed

below, they might only fit it for a short amount of time. However, knowing where a bacterial

population of interest happens to fall along a spectrum of clonality can help us understand its

biology, and even make predictions about its evolution.

The opposite of clonality is panmixis – a situation in which the rate of horizontal transfer

is higher than the rate of vertical cell division, resulting in random association (linkage

equilibrium) among loci in the genome [1,2]. However, rates of horizontal transfer

(recombination) vary widely across the genome, such that a population can be mostly clonal,

except for a few loci in the genome [3]. These loci came to be termed genomic islands – a

metaphor I will build upon below. Some of the first islands identified were called pathogenicity

islands because they contained virulence factors [4]. However, non-pathogenic environmental

bacteria also contain islands, conferring adaptation to different ecological niches. For example,

genes in *Prochlorococcus* genomic islands confer adaptation to light and nutrient conditions

[5,6]. But islands need not confer niche adaptation to their host genome; they can be neutral to

host fitness or even detrimental, selfish parasites. Here, I define genomic islands broadly as any

piece of DNA that is transferred horizontally (by either homologous or nonhomologous

recombination) from cell to cell and therefore evolves independently (*i.e.* is unlinked) from the

rest of the genome.

I will begin by extending the use of island analogies to include continents, peninsulas and

archipelagos (**Table 1**). I will then use these analogies to discuss to what extent microbial

populations are clonal or panmictic, and how often they transition between the two regimes.

Are some islands really peninsulas?

In the classic analogy, an island is totally disconnected from the mainland, meaning that genes in

the island evolve independently of the genome (**Table 1**). Examples of islands that fit this strict

independence might include integrated phages and other "selfish" elements, or genes that reside

in a particular niche but not in a particular genome (*e.g.* a gene ecology model [7]). Peninsulas

provide an analogy that might better describe how islands are related to microbial genomes. A

peninsula (or "presque-ile," from the French for "almost island") is a geographic term for a very

narrow strip of land connected to (but distinct from) the mainland. In my analogy, an island is

evolutionarily independent of the mainland genome, but their fates may become linked, forming a

peninsula. For example, a bacterium may acquire a gene from a vast microbial gene pool. This

gene allows the bacterium to invade a new ecological niche, triggering a clonal expansion in

which the fate of the gene and its new host genome are linked, at least for the duration of the

clonal expansion. One such example could be *Yersinia pestis,* which acquired a single gene

allowing flea-borne transmission and triggering a clonal expansion in the form of Plague

pandemics [8]. Another peninsula, the prophage-encoded cholera toxin, and its links to the

mainland *Vibrio cholerae* genome [9,10], is discussed below.

Are some genomes archipelagos?

The very concept of one or a few islands implies a contrast with the large, clonal genomic

mainland or continent. But some microbial genomes may contain so many islands that there is no

mainland, only a vast archipelago (**Table 1**). A striking recent example is a population of

hotspring cyanobacteria in which virtually every gene in the genome evolved independently due

to frequent recombination [11], leading the authors to call the population "quasi-sexual" (in other

words, panmictic). Frequent recombination was confirmed by another group, using different

methods to study the same cyanobacteria [12]. This group also found that despite a history of

panmixis over long time scales, populations are more clonal over shorter time scales. Similarly,

the Asian ocean population of *Vibrio parahaemolyticus* also forms a panmictic gene pool, with

each recombination block of ~1.8 kbp evolving independently [13]. However, the panmictic gene

pool occasionally gives rise to pandemic clones. In another example, we found that almost every

gene in a population of *Vibrio cyclitrophicus* genomes showed signs of recombination over

relatively recent time scales [14]. Such apparently high rates of recombination in natural

populations were mysterious at first, contradicting recombination rates measured in the lab

[15,16] and predicted by theory [7,17]. However, theoretical models (discussed below) suggest

mechanisms capable of explaining how genes can spread through populations more rapidly by

recombination than by clonal expansion [18-20].

Clonal expansions from panmictic pools

I propose that archipelagos are not necessarily static over time, and that archipelagos can

sometimes coalesce into continents. Given the right ecological opportunity, a genome from a

panmictic gene pool can escape the "gravitational pull" of recombination and take off into a

clonal expansion. An example mentioned earlier is *V. cholerae*, a genetically diverse group of

coastal marine bacteria, some of which cause cholera. Virulence is mainly determined by two loci

in the genome: the cholera toxin and the toxin-coregulated pilus. Both genes are frequently gained

and lost by recombination [21,22], but are always found in one lineage of *V. cholerae* – the

lineage causing severe disease with pandemic potential, known as the phylocore genome (PG)

group [10]. It remains a mystery why the PG lineage evolved once, and only once. If PG *V.*

cholerae really did evolve just once, this would be surprising because *V. cholerae* draws on a

diverse, global gene pool and can be considered panmictic [23]. Therefore multiple different

lineages would be expected to acquire the two (or perhaps a handful of) genetic elements required

for pandemic disease. This leads to the hypothesis that pandemic cholera emergence is *selection*

limited rather than *diversity limited*. In other words, benign *V. cholerae* strains constantly acquire

virulence genes. However, these strains rarely encounter the right ecological niche to flourish,

e.g. a human population consuming brackish water. "The right niche" has appeared a few times in

human history: for example in India in the 1800s, when the Classical lineage evolved, and again

in Indonesia in the 1950s, when the El Tor lineage evolved [24]. When the right conditions

appear, the PG lineage, along with its virulence factors, takes off in a clonal expansions which

continue to wreak havoc today (*e.g.* cholera pandemics from the 1800s to today, all caused by the

PG clonal group). The virulence factors, previously islands in an archipelago, became a peninsula

connected to the PG mainland. The linkage between virulence factors and PG remains imperfect

because different variants of the cholera toxin continue to flow in and out of the PG continent

[10,21]; hence the toxin remains a peninsula, not firmly part of the mainland.

V. cholerae is a particularly well-characterized example of a panmictic gene pool giving

rise to a clonal expansion, but similar evolutionary dynamics are seen in other pathogens as well

(*e.g. V. parahaemolyticus* described above [13]). Enterotoxigenic *Escherichia coli* (ETEC) seems

to behave similarly, with deep branches of the phylogeny obscured by frequent recombination

and plasmid exchange, but more recent branches experiencing mostly clonal descent, with tight

linkage between virulence factors and the genomic mainland [25]. These observations are

consistent with an ancient, panmictic gene pool giving rise to clonal expansions, which can last

for decades or centuries.

The balance between recombination and selection

Let us consider the evolutionary forces that determine clonality: natural selection and

recombination. The effect of recombination on clonality is straightforward: more recombination

means less clonality. The effect of natural selection is more complex, but is defined here simply

as a force which favors clonal expansions of adaptive mutants within an ecological niche.

Selection, as defined here, therefore includes ecological effects. When driven by ecological

selection, clonal expansions are called selective sweeps, in which one clone outcompetes all

others, purging genetic diversity in the population.

Recombination and selection interact to determine the clonality of a population.

Recombination rates depend both on the ability of DNA to enter a cell and be incorporated into

the genome (the baseline rate) and the ability of that DNA to be retained by a balance of genetic

drift and natural selection (the realized rate). Some bacteria, such as *Helicobacter pylori,* have

realized recombination rates that are much higher than point mutation rates, exchanging at least
10% of their genome within a single four-year human infection [26]. Others, such as
Staphylococcus aureus [27,28] and *Mycobacterium tuberculosis* are decidedly more clonal [29-
31]. Recombination rates (both realized and baseline) vary widely across the genome. Of 10
pathogenic bacterial species studied, all had identifiable recombination 'hot' regions, although
their length, genomic location and gene content varied [3]. Genes of different functions had
different realized recombination rates, implying a role for natural selection on gene function in
determining whether newly acquired genes are retained.
144
Modeling the recombination-selection balance
When rates of recombination are relatively low compared to rates of natural selection on adaptive
genes within niches, entire genomes will sweep to fixation before they can be shuffled by
recombination. Following previous modelling work, s is defined here as the selective coefficient
of a niche-adaptive allele, and r is the recombination rate, per locus per generation [7,32,33].The
$s \gg r$ regime is well described in the Stable Ecotype Model [17], which predicts that most of the
genome will follow a single, clonal phylogeny. Genome-wide sweeps thus increase clonality and
can be considered a hallmark of clonal populations (**Table 1, Figure 1A**). In the $r \gg s$ regime,
individual genes (rather than entire genomes) will sweep to fixation (*i.e.* reach 100% frequency)
in ecological niches to which they are adapted, without affecting genetic diversity elsewhere in
the genome (**Figure 1B**). The first serious theoretical attempt to reconcile the observations of
gene-specific sweeps with low recombination rates was the "Adapt Globally, Act Locally" model
[18,20], in which globally adaptive genes (adaptive in multiple different niches) trigger genome-
wide sweeps within a niche before being transferred to the next niche. When viewed across
niches, the result is a gene-specific sweep. A recent model suggested another mechanism by
which these gene sweeps can occur at moderate (not unrealistically high) rates of recombination
[20]. In this model, a microbial habitat is bombarded with genetically maladapted migrants,
allowing gene sweeps to occur, although the adaptive allele never reaches 100% frequency due to
the constant input of migrants. In another model, Takeuchi et al. [19] show that gene sweeps can
occur when r is either very high or – counter-intuitively – when r is very low, but only when
negative frequency-dependent selection (NFDS) reduces the rate of genome-wide selective
sweeps. NFDS might be commonly imposed on bacteria and archaea by viral (phage) predation,
providing a selective advantage to rare alleles of phage receptor genes [34].
One parameter not thoroughly explored in any of these models is the effective population
size (N_e). Populations with small N_e are dominated by drift, and natural selection is inefficient.

They may experience genome-wide sweeps independently of *r* and *s*. As discussed below, N_e is

probably an important determinant of clonality.

Genome-wide and gene-specific sweeps in nature

To date, empirical evidence for gene-specific and genome-wide sweeps has come mostly from

cross-sectional studies of a single population of genomes at a single point in time, with

recombination and selection inferred backward in time [11,13,14,35]. Sequencing microbial

genomes or metagenomes sampled over time – already a typical practice in genomic

epidemiology (e.g. [28,36]) – promises to elucidate the rates of gene-specific and genome-wide

sweeps in nature (**Figure 1**).

In a pioneering study, Bendall et al. [37] sampled a lake over nine years and followed

single-nucleotide polymorphism (SNP) and gene frequencies in 30 bacterial populations by

metagenomic sequencing. They inferred that one of the populations (*Chlorobium*-111) had

undergone a near-complete genome-wide sweep over the nine-year study, with most SNP

diversity purged genome-wide (**Figure 1A**). By identifying regions of the genome with

unexpectedly low diversity compared to the genome-wide average, they inferred that gene-

specific sweeps had occurred in six of the populations, but these sweeps occurred before the

beginning of the nine-year study. During the nine-year study, they observed examples "where a

few adjacent SNPs trended toward fixation while genome-wide diversity was maintained"

(**Figure 1B**). They took this observation as consistent with gene-specific selective sweeps, but did

not attempt to determine whether the sweeps were due to selection or drift. Similarly, the inferred

genome-wide sweep could have been caused by selection or drift.

Is the frequency of genome-wide sweeps controlled mainly by the balance of

recombination and selection, or could drift (controlled by the effective population size, N_e) play

an important role as well? In their Figure 2 legend, Bendall et al. observe that "populations with

many SNPs were not necessarily sequenced deeper than those with few SNPs." This statement

was simply meant to assure the reader that SNP calling was not biased by sequence coverage, but

it also suggests that microbial populations tend to be far from a standard neutral model with

constant population size (*i.e.* the Wright-Fisher coalescent [38]). Taking SNP density as a rough

measure of N_e and sequence coverage as a rough measure of the census population size, the data

show that census population size is a poor predictor of N_e, and thus that populations are likely

impacted by population bottlenecks and/or selection – although it is difficult to distinguish

between them.

As a whole, the study showed that both genome-wide and gene-specific sweeps can occur

in different microbial populations from the same environment. Whether microbial populations

behaved differently due to differences in their ecology (*i.e.* regime of natural selection) or in their

baseline recombination rates remains a question for future study; Cohan recently suggested that

ecological differences could be more important [39]. Specifically, he suggests that ecological

"generalists" could have more opportunities for diversification, and thus be relatively immune to

genome-wide sweeps, compared to ecological specialists. I suggest that these generalists may

simply have larger N_e, such that diversity is rarely purged by drift (*e.g.* bottlenecks), and that

diversity is mainly governed by selection and recombination. It is possible that ecological

generalists tend to have large N_e, or that N_e and ecology exert independent effects (*e.g.* ecological

generalists can have low N_e and still resist genome-wide sweeps). The fact that one genome-wide

sweep was observed over a nine year period suggests that such events might be relatively rapid

but rare (only observed in one of 30 populations). Meanwhile, gene-sweeps might be more

common historically (affecting six of 30 populations), but could take longer to proceed to

completion.

Is clonality a stable trait?

As described in the *V. cholerae* example, some pathogenic bacterial populations can switch

between panmictic and clonal lifestyles [11,13,14,19,25,34,35]. Therefore clonality can vary over

time, but how much and how often? To quantify the stability of clonality over time, not just in

pathogens but in free-living environmental bacteria, I re-analyzed the lake time-course of Bendall

et al. [37]. Because estimates of selection and recombination rates were not readily available for

this dataset, I defined clonality based on the total genetic diversity (SNP density) in a population,

which scales with N_e, and includes the effects of both drift and of genome-wide selective sweeps.

Frequent genome-wide selective sweeps ($s >> r$) and/or small N_e result in clonality. I identified

20 "old, diverse" populations as those with a high density of SNPs (>1500 SNPs/Mbp) at the

beginning of the time-course. These populations are likely "old" because they have gone a

relatively long time since the last genome-wide purge of genetic diversity and are relatively non-

clonal. They include the six populations inferred to have undergone gene-specific sweeps [37].

The remaining ten populations were defined as "young, low-diversity," having more recently

experienced a genome-wide purge of diversity. The "old, diverse" populations have a relatively

low ratio of nonsynonymous (N) to synonymous (S) SNPs, suggesting large N_e and ample time

for purifying selection to remove (mostly deleterious) nonsynonymous mutations (**Figure 2**). In

contrast, the "young, low-diversity" populations are more likely to have high N:S ratios,

suggesting smaller N_e and less time for purifying selection to act. The cutoff of 1500 SNPs/Mbp
(0.15% SNP density) is somewhat arbitrary, but seems to correspond to an inflection point in
Figure 2, and also falls squarely within the 0-0.30% SNP density, above which *E. coli* evolution
appears to transition from clonal to recombining [40].
With young (presumably clonal, or low N_e) and old (less clonal, higher N_e) populations
thus defined, I asked whether "old, diverse" populations tended to maintain their diversity
through the 9-year period of the study. In their paper, Bendall et al. defined two alternative
population types: 1) those that maintained stable SNP diversity over nine years, and 2) those that
experienced significant fluctuations in diversity due to clonal expansions – defined when one, but
not all timepoints are dominated by a single allele (≥95% frequency) at >40% of SNP sites in the
genome [37]. By definition, the 19 populations of the first type did not experience genome-wide
sweeps during the study, while the 11 populations of the second type did experience genome-
wide purges of diversity, which were transient in ten cases and apparently permanent in one case
(*Chlorobium*-111). Strikingly, 17 out of 20 "old, diverse" populations maintained their diversity
over the nine-year study, compared to only two out of ten "young, low-diversity" populations
(Fisher test, Odds Ratio = 19.4, $P < 0.001$). This result suggests that populations with a history of
genome-wide sweeps tend to experience subsequent genome-wide sweeps, and those that have
maintained genetic diversity in the past tend to maintain their diversity into the future. In other
words, clonality can be considered a relatively stable microbial trait. However, we must take care
in taking a genome-wide sweep as evidence for clonality ($s \gg r$). In small effective population
sizes, genome-wide sweeps can occur due to drift and bottlenecks, independently of s and r.
Therefore, many of "young, low diversity" populations (**Figure 2**) may have lost diversity over
nine years due to low N_e, not due to low r.
260
History repeats itself
It appears that pathogens are more likely than free-living bacteria to undergo clonal expansions,
due in part to their ecology and transmission dynamics [41]. Free-living aquatic bacteria, on the
other hand, seem to be more likely to live in large, panmictic populations and behave like
archipelagos [11,13,14,37]. If clonality is indeed a stable trait, this implies that history will repeat
itself, and that the future behavior of microbial populations can be predicted with some
confidence from their past behavior. Diverse populations tend to stay diverse. Clonal populations
(that experience frequent genome-wide sweeps) tend to stay clonal. But history is not doomed to
repeat itself forever. As we have seen, clonal expansions, such as pandemic *V. cholerae,* may
originate when a panmictic gene pool (an archipelago) coalesces into a clonal continent, with

virulence factors linked as peninsulas. Many such pathogenic clones have been documented, with

life spans of decades to thousands of years [25,28,42,43]. Other pathogens, such as *Streptococcus*

pneumoniae, may retain their panmictic population structure throughout an outbreak [44-46].

Why some pathogens are clonal and others are panmictic is an open question, but surely depends

on the balance between recombination and selection, on the effective population size, and on the

time scales considered.

Acknowledgments. I am grateful to Rex Malmstrom, Yan Boucher, and Salvador Almagro-

Moreno for their thoughtful comments which improved the manuscript. I was supported by a

Canada Research Chair.

Table 1. Extended island metaphors of microbial genome evolution.

Geographic metaphor	Genetic unit to which the metaphor applies	Type of selective sweep experience by the unit	Dominant mode of genetic transmission	Example
Island	Gene	Gene-specific	horizontal	genes in the *V. cholerae* integron [22,23]
Peninsula	Gene	Genome-wide	vertical (clonal)	the cholera toxin gene, acquired horizontally, then linked to a clonal *V. cholerae* genome [9,21]
Continent	Genome	Genome-wide	vertical (clonal)	clonal expansions of *S. aureus* [28], *M. tuberculosis* [31,43]
Archipelago	Genome	Gene-specific	horizontal	hotspring cyanobacteria [11], ocean vibrios [13,14], pneumococcus [44,46]

Figures

Figure 1. Temporal dynamics of genome-wide and gene-specific selective sweeps inferred from metagenomic data. Genetic diversity can be measured by mapping metagenomic reads to a reference genome, identifying SNPs, and calculating the allele frequencies at each SNP position in the genome over time. The lowest possible genetic diversity occurs when a single allele is present in 100% of metagenomic reads. Alternatively, diversity could be defined in terms of gene presence/absence, based on relative coverage of a gene in the reference genome by metagenomic reads. **(a)** In a hypothetical genome-wide selective sweep, all positions in the genome tend toward low diversity. **(b)** In a hypothetical gene-specific selective sweep, only one or a few positions in the genome tend toward low diversity, while the rest of the genome maintains high or intermediate diversity.

Figure 2. Past diversity predicts future diversity. Based on data from Table 2 of Bendall et al. [37], the nonsynonymous to synonymous (N:S) SNP ratio was plotted against the total SNP density (SNPs per megabasepair) for each of 30 bacterial populations. A pseudocount of 1 was added to both N and S counts. These 30 populations were divided into 20 "old, diverse" populations (>1500 SNPs/Mbp) and 10 "young, low-diversity" populations (<1500 SNPs/Mbp). Each point represents one of the 30 populations, colored in black if diversity was maintained over a 9-year metagenomic time-course, or in red if it was not. Seventeen out of 20 "old, diverse" populations maintained their diversity over the 9-year study, compared to only 2 out of 10 "young, low-diversity" populations (Fisher test, Odds Ratio = 19.4, *P* < 0.001). The same result is obtained drawing a cutoff based on N:S rather than SNPs/Mbp: populations with N:S < 0.5 tend to maintain their diversity, whereas those with N:S > 0.5 tend to be purged of diversity over the 9 years (Fisher test, Odds Ratio = 5.94, *P* = 0.042). Consistent with previous observations that N:S depends on the evolutionary time available for purifying selection to act [47,48], N:S is negatively correlated with SNPs/Mbp, a proxy for evolutionary time or the time since the last genome-wide purge of genetic diversity in this dataset (Pearson's correlation of $\log_{10}$ transformed data, *r* = -0.81, *P* = 5.6e-8).

References

1. Smith JM, Smith NH, O'Rourke M, Spratt BG: **How clonal are bacteria?** *Proc Natl Acad Sci USA*. 1993, 90: 4384–4388.

2. Guttman DS, Dykhuizen DE: **Clonal Divergence in Escherichia coli as a Result of Recombination, Not Mutation**. *Science*. 1994, 266: 1380–1383.

3. Yahara K, Didelot X, Jolley KA, Kobayashi I, Maiden MCJ, Sheppard SK, et al.: **The landscape of realized homologous recombination in pathogenic bacteria**. *Mol Biol Evol*. 2016, 33: 456–471. doi:10.1093/molbev/msv237

4. Hacker J, Blum-Oehler G, Mühldorfer I, Tschäpe H: **Pathogenicity islands of virulent bacteria: structure, function and impact on microbial evolution**. *Mol Microbiol*. 1997, 23: 1089–1097.

5. Coleman ML, Sullivan MB, Martiny AC, Steglich C, Barry K, Delong EF, et al.: **Genomic islands and the ecology and evolution of *Prochlorococcus***. *Science*. 2006, 311: 1768–1770.

6. Coleman ML, Chisholm SW: **Ecosystem-specific selection pressures revealed through comparative population genomics**. *Proc Natl Acad Sci USA*. 2010, 107: 18634–18639. doi:10.1073/pnas.1009480107

7. Shapiro BJ, Polz MF: **Ordering microbial diversity into ecologically and genetically cohesive units**. *Trends Microbiol*. 2014, 22: 235–247. doi:10.1016/j.tim.2014.02.006

* A comprehensive discussion of the roles of selection and recombination in structuring microbial diversity.

8. Rasmussen S, Allentoft ME, Nielsen K, Orlando L, Sikora M, Sjögren K-G, et al.: **Early Divergent Strains of Yersinia pestis in Eurasia 5,000 Years Ago**. *Cell*. 2015, 163: 571–582. doi:10.1016/j.cell.2015.10.009

9. Waldor MK, Mekalanos JJ: **Lysogenic conversion by a filamentous phage encoding cholera toxin**. *Science*. 1996, 272: 1910–1914.

10. Chun J, Grim CJ, Hasan NA, Lee JH, Choi SY, Haley BJ, et al.: **Comparative genomics reveals mechanism for short-term and long-term clonal transitions in pandemic *Vibrio cholerae***. *Proc Natl Acad Sci USA*. 2009, 106: 15442–15447. doi:10.1073/pnas.0907787106

11. Rosen MJ, Davison M, Bhaya D, Fisher DS: **Fine-scale diversity and extensive recombination in a quasisexual bacterial population occupying a broad niche**. *Science*. 2015, 348: 1019–1023. doi:10.1126/science.aaa4456

* Deep sequencing of 90 cyanobacterial marker genes reveals extensive recombination, with each genome composed of a random mix of alleles from the gene pool.

353   12.   Melendrez MC, Becraft ED, Wood JM, Olsen MT, Bryant DA, Heidelberg JF, et al.:
354   **Recombination Does Not Hinder Formation or Detection of Ecological Species of**
355   ***Synechococcus* Inhabiting a Hot Spring Cyanobacterial Mat**. *Front Microbiol*. 2016,
356   6: 544. doi:10.1128/AEM.72.1.723-732.2006

357   13.   Cui Y, Yang X, Didelot X, Guo C, Li D, Yan Y, et al.: **Epidemic Clones, Oceanic Gene**
358   **Pools, and Eco-LD in the Free Living Marine Pathogen *Vibrio parahaemolyticus***. *Mol*
359   *Biol Evol*. 2015, 32: 1396–1410. doi:10.1093/molbev/msv009

360   * In-depth population genomic evidence of a panmictic ocean gene pool containing non-clonal
361   ecological populations.
362

363   14.   Shapiro BJ, Friedman J, Cordero OX, Preheim SP, Timberlake SC, Szabo G, et al.:
364   **Population Genomics of Early Events in the Ecological Differentiation of Bacteria**.
365   *Science*. 2012, 336: 48–51. doi:10.1126/science.1218198

366   15.   Vulić M, Dionisio F, Taddei F, Radman M: **Molecular keys to speciation: DNA**
367   **polymorphism and the control of genetic exchange in enterobacteria**. *Proc Natl Acad*
368   *Sci USA*. 1997, 94: 9763–9767.

369   16.   Majewski J, Cohan FM: **DNA sequence similarity requirements for interspecific**
370   **recombination in *Bacillus***. *Genetics*. 1999, 153: 1525–1533.

371   17.   Wiedenbeck J, Cohan FM: **Origins of bacterial diversity through horizontal genetic**
372   **transfer and adaptation to new ecological niches**. *FEMS Microbiology Reviews*. 2011,
373   35: 957–976. doi:10.1111/j.1574-6976.2011.00292.x

374   18.   Majewski J, Cohan FM: **Adapt globally, act locally: the effect of selective sweeps on**
375   **bacterial sequence diversity**. *Genetics*. 1999, 152: 1459–1474.

376   19.   Takeuchi N, Cordero OX, Koonin EV, Kaneko K: **Gene-specific selective sweeps in**
377   **bacteria and archaea caused by negative frequency-dependent selection**. *BMC*
378   *Biology*. 2015, 13: 20. doi:10.1186/s12915-015-0131-7

379   * Mathematical modeling shows how individual genes can sweep through populations with low
380   recombination rates in the presence of negative frequency-dependent selection.
381

382   20.   Niehus R, Mitri S, Fletcher AG, Foster KR: **Migration and horizontal gene transfer**
383   **divide microbial genomes into multiple niches**. *Nature Communications*. 2015, 6: 8924.
384   doi:10.1038/ncomms9924

385   21.   Mutreja A, Kim DW, Thomson NR, Connor TR, Lee JH, Kariuki S, et al.: **Evidence for**
386   **several waves of global transmission in the seventh cholera pandemic**. *Nature*. 2011,
387   477: 462–U111. doi:10.1038/nature10392

388   22.   Orata FD, Kirchberger PC, Méheust R, Barlow EJ, Tarr CL, Boucher Y: **The Dynamics**
389   **of Genetic Interactions between *Vibrio metoecus* and *Vibrio cholerae*, Two Close**
390   **Relatives Co-Occurring in the Environment**. *Genome Biology and Evolution*. 2015, 7:
391   2941–2954. doi:10.1093/gbe/evv193/-/DC1

392   * A clear demonstration that genes in the integron are recombined between species just as often
393   as within species, and that 24% of 'core' genes have crossed species boundaries.
394

23.   Boucher Y, Cordero OX, Takemura A, Hunt DE, Schliep K, Bapteste E, et al.: **Local**
Mobile Gene Pools Rapidly Cross Species Boundaries To Create Endemicity within
Global *Vibrio cholerae* Populations. *mBio*. 2011, 2: e00335–10.
doi:10.1128/mBio.00335-10

24.   Boucher Y, Orata FD, Alam M: **The out-of-the-delta hypothesis: dense human**
populations in low-lying river deltas served as agents for the evolution of a deadly
pathogen. *Front Microbiol*. 2015, 6: 1120. doi:10.3389/fmicb.2015.01120

25.   Mentzer von A, Connor TR, Wieler LH, Semmler T, Iguchi A, Thomson NR, et al.:
Identification of enterotoxigenic *Escherichia coli* (ETEC) clades with long-term
global distribution. *Nature Genetics*. 2014, 46: 1321–1326. doi:10.1038/ng.3145

26.   Cao Q, Didelot X, Wu Z, Li Z, He L, Li Y, et al.: **Progressive genomic convergence of**
two *Helicobacter pylori* strains during mixed infection of a patient with chronic
gastritis. *Gut*. 2015, 64: 554–561. doi:10.1136/gutjnl-2014-307345

27.   Vos M, Didelot X: **A comparison of homologous recombination rates in bacteria and**
archaea. *The ISME journal*. 2009, 3: 199–208. doi:10.1038/ismej.2008.93

28.   Baines SL, Holt KE, Schultz MB, Seemann T, Howden BO, Jensen SO, et al.:
Convergent adaptation in the dominant global hospital clone ST239 of methicillin-
resistant *Staphylococcus aureus*. *mBio*. 2015, 6: e00080. doi:10.1128/mBio.00080-15

29.   Hershberg R, Lipatov M, Small PM, Sheffer H, Niemann S, Homolka S, et al.: **High**
Functional Diversity in *Mycobacterium tuberculosis* Driven by Genetic Drift and
Human Demography. *PLOS Biol*. 2008, 6: e311.
doi:10.1371/journal.pbio.0060311.st006

30.   Namouchi A, Didelot X, Schock U, Gicquel B, Rocha EPC: **After the bottleneck:**
Genome-wide diversification of the *Mycobacterium tuberculosis* complex by mutation,
recombination, and natural selection. *Genome Research*. 2012, 22: 721–734.
doi:10.1101/gr.129544.111

31.   Casali N, Nikolayevskyy V, Balabanova Y, Harris SR, Ignatyeva O, Kontsevaya I, et al.:
Evolution and transmission of drug-resistant tuberculosis in a Russian population.
Nature. 2014, 46: 279–286. doi:10.1038/ng.2878

32.   Shapiro BJ, David LA, Friedman J, Alm EJ: **Looking for Darwin's footprints in the**
microbial world. *Trends in Microbiology*. 2009, 17: 196–204.
doi:10.1016/j.tim.2009.02.002

33.   Friedman J, Alm EJ, Shapiro BJ: **Sympatric Speciation: When Is It Possible in**
Bacteria? *PLOS ONE*. 2013, 8: e53539. doi:10.1371/journal.pone.0053539.g004

34.   Cordero OX, Polz MF: **Explaining microbial genomic diversity in light of evolutionary**
ecology. *Nature Reviews Microbiology*. 2014, 12: 263–273. doi:10.1038/nrmicro3218

35.   Cadillo-Quiroz H, Didelot X, Held NL, Herrera A, Darling A, Reno ML, et al.: **Patterns**
of Gene Flow Define Species of Thermophilic Archaea. *PLOS Biol*. 2012, 10:
e1001265. doi:10.1371/journal.pbio.1001265.t001

36.   Croucher NJ, Didelot X: **The application of genomics to tracing bacterial pathogen**
transmission. *Current Opinion in Microbiology*. 2015, 23: 62–67.
doi:10.1016/j.mib.2014.11.004

37.   Bendall ML, Stevens SL, Chan L-K, Malfatti S, Schwientek P, Tremblay J, et al.:
Genome-wide selective sweeps and gene-specific sweeps in natural bacterial
populations. *The ISME journal*. 2016, advance online publication, 8 January 2016;
doi:10.1038/ismej.2015.241

** A pioneering application of time-course metagenomics to infer genome-wide and gene-
specific selective sweeps in natural lake bacterial populations.
443

38.   Kingman J: **On the genealogy of large populations**. *J Appl Prob*. 1982, 19: 27–43.

39.   Cohan FM: **Bacterial Speciation: Genetic Sweeps in Bacterial Species**. *Curr Biol*. 2016,
26: R112–R115. doi:10.1016/j.cub.2015.10.022

40.   Dixit PD, Pang TY, Studier FW, Maslov S: **Recombinant transfer in the basic genome**
of *Escherichia* coli. *Proc Natl Acad Sci USA*. 2015, 112: 9070–9075.
doi:10.1073/pnas.1510839112

41.   Achtman M, Wagner M: **Microbial diversity and the genetic nature of microbial**
species. *Nature Reviews Microbiology*. 2008, 6: 431–440. doi:10.1038/nrmicro1872

42.   Snitkin ES, Zelazny AM, Thomas PJ, Stock F, NISC Comparative Sequencing Program,
Henderson DK, et al.: **Tracking a Hospital Outbreak of Carbapenem-Resistant**
***Klebsiella pneumoniae* with Whole-Genome Sequencing.** *Science Translational*
Medicine. 2012, 4: 148ra116–148ra116. doi:10.1126/scitranslmed.3004129

43.   Comas I, Coscolla M, Luo T, Borrell S, Holt KE, Kato-Maeda M, et al.: **Out-of-Africa**
migration and Neolithic coexpansion of *Mycobacterium tuberculosis* with modern
humans. *Nature Genetics*. 2013, 45: 1176–1182. doi:10.1038/ng.2744

44.   Croucher NJ, Finkelstein JA, Pelton SI, Mitchell PK, Lee GM, Parkhill J, et al.:
Population genomics of post-vaccine changes in pneumococcal epidemiology. *Nature*
Genetics. 2013, 45: 656–663. doi:10.1038/ng.2625

45.   Chewapreecha C, Harris SR, Croucher NJ, Turner C, Marttinen P, Cheng L, et al.: **Dense**
genomic sampling identifies highways of pneumococcal recombination. *Nature*
Genetics. 2014, 46: 305–309. doi:10.1038/ng.2895

46.   Marttinen P, Croucher NJ, Gutmann MU, Corander J, Hanage WP: **Recombination**
produces coherent bacterial species clusters in both core and accessory genomes.
Microbial Genomics. 2015, 1. doi:10.1099/mgen.0.000038.

* The first simulation of bacterial evolution including mutation, allelic exchange (homologous
recombination) and gene gain/loss (non-homologous recombination).
470

471 47. Rocha EP, Smith JM, Hurst LD, Holden MT, Cooper JE, Smith NH, et al.: **Comparisons
472   of dN/dS are time dependent for closely related bacterial genomes**. *J Theor Biol*. 2006,
473   239: 226–235.

474 48. Kryazhimskiy S, Plotkin JB: **The Population Genetics of dN/dS**. *PLOS Genet*. 2008, 4:
475   e1000304.

Figure 1

(a) Genome-wide sweep   **(b) Gene-specific sweeps**   **Legend**

Position in the genome

Genetic diversity

high

low

← gene-specific
sweep completed
during the period
of study

← gene-specific
sweep completed
before the period
of study

time (years) ⟶   time (years) ⟶

Figure 2