Differential methylation between ethnic sub-groups reflects the effect of genetic ancestry and environmental exposures

Joshua M. Galanter, MD, MAS[1,2,3], Christopher R. Gignoux, PhD[4], Sam S. Oh, PhD[1], Dara Torgerson, PhD[2], Maria Pino-Yanes, PhD[5,6], Neeta Thakur, MD, MPH[1], Celeste Eng, BS[1], Donglei Hu, PhD[1], Scott Huntsman, MS[1], Harold J. Farber, MD[7], Pedro C Avila, MD[8], Emerita Brigino-Buenaventura, MD[9], Michael A LeNoir, MD[10], Kelly Meade, MD[11], Denise Serebrisky, MD[12], William Rodríguez-Cintrón, MD[13], Rajesh Kumar, MD[14], Jose R Rodríguez-Santana, MD[15], Max A. Seibold, PhD[16], Luisa N. Borrell, DDS, PhD[17], Esteban G. Burchard, MD, MPH[1,2*], Noah Zaitlen, PhD[1*]

1. Department of Medicine, University of California, San Francisco, CA

2. Department of Bioengineering and Therapeutic Sciences, University of California, San Francisco, CA

3. Department of Epidemiology and Biostatistics, University of California, San Francisco, CA

4. Department of Genetics, Stanford University, Stanford, CA

5. Hospital Universitario Nuestra Señora de Candelaria, Tenerife, Spain

6. CIBER de Enfermedades Respiratorias, Instituto de Salud Carlos III, Madrid, Spain

7. Department of Pediatrics, Baylor College of Medicine and Texas Children's Hospital, Houston, Texas

8. Division of Allergy and Immunology, Feinberg School of Medicine, Northwestern University, Chicago, IL

9. Kaiser Permanente-Vallejo Medical Center, Vallejo, CA

10. Bay Area Pediatrics, Oakland, CA

11. Department of Pediatrics, Children's Hospital and Research Center, Oakland, CA

12. Jacobi Medical Center, Bronx, NY

13. Veterans Caribbean Health System, San Juan, Puerto Rico

14. Division of Allergy and Immunology, The Ann and Robert H. Lurie Children's Hospital of Chicago, Chicago, IL

15. Centro de Neumología Pediátrica, San Juan, Puerto Rico

16. Center for Genes, Environment, and Health, Department of Pediatrics, National Jewish Health, Denver, CO

17. Graduate School of Public Health and Health Policy, City University of New York, New York, NY

*   These authors contributed equally to this work

Please address correspondence to:

Esteban G. Burchard, MD, MPH

University of California, San Francisco

Departments of Bioengineering & Therapeutic Sciences and Medicine

UCSF Box 2911

San Francisco, CA 94143-2911

Ph: (415) 514-9677

Fax: (415) 514-4365

e-mail: esteban.burchard@ucsf.edu

or

Noah Zaitlen, PhD

University of California, San Francisco

Department of Medicine

UCSF Box 2552

San Francisco, CA 94143-2552

Ph: (415) 502-2027

e-mail: noah.zaitlan@ucsf.edu

or

Joshua Galanter, MD, MAS

University of California, San Francisco

Departments of Medicine and Epidemiology & Biostatistics

UCSF Box 2911

San Francisco, CA 94143-2552

Ph: (415) 514-9931

e-mail: galanter@gmail.com

Sources of Funding:

This research was supported in part by the Sandler Family Foundation, the American Asthma Foundation, National Institutes of Health (P60 MD006902, R01 HL117004, R21ES24844, U54MD009523, R01 ES015794, R01 HL088133, M01 RR000083, R01 HL078885, R01 HL104608, U19 AI077439, M01 RR00188), ARRA grant RC2 HL101651, and TRDRP 24RT-0025; EGB was supported in part through grants from the Flight Attendant Medical Research Institute (FAMRI), and NIH (K23 HL004464); NZ was supported in part by an NIH career development award from the NHLBI (K25HL121295). JMG was supported in part by NIH Training Grant T32 (T32GM007546) and career development awards from the NHLBI (K23HL111636) and NCATS (KL2TR000143) as well as the Hewett Fellowship; N.T. was supported in part by an institutional training grant from the NIGMS (T32-GM007546) and career development awards from the NHLBI (K12-HL119997 and K23-HL125551), Parker B. Francis Fellowship Program, and the American Thoracic Society; CRG was supported in part by NIH Training Grant T32 (GM007175) and the UCSF Chancellor's Research Fellowship and Dissertation Year Fellowship; RK was supported with a career development award from the NHLBI (K23HL093023); HJF was supported in part by the GCRC (RR00188); PCA was supported in part by the Ernest S. Bazley Grant; MAS was supported in part by 1R01HL128439-01. This publication was supported by various institutes within the National Institutes of Health. Its contents are solely the responsibility of the authors and do not necessarily represent the official views of the NIH.

Abstract

In clinical practice and biomedical research populations are often divided categorically

into distinct racial and ethnic groups. In reality, these categories, which are based on

social rather than biological constructs, comprise diverse groups with highly

heterogeneous histories, cultures, traditions, religions, as well as social and

environmental exposures. While the factors captured by these categories contribute to

clinical practice and biomedical research, the use of race/ethnicity is widely debated. As

a response to this debate, genetic ancestry has been suggested as a complement or

alternative to this categorization. However, few studies have examined the effect of

genetic ancestry, racial/ethnic identity, and environmental exposures on biological

processes. Herein, we examine the associations between self-identified ethnicity and

epigenetic modification of DNA methylation, a phenomenon affected by both genetic

and environmental factors. We also assess the relative contributions of genetic ancestry

and environmental factors on these associations. We typed over 450,000 variably

methylated CpG sites in primary whole blood of 573 individuals of Mexican and Puerto

Rican descent who also had high-density genotype data. We found that both genetically

determined ancestry and self-identified ethnicity were significantly associated with

methylation levels at a large number of CpG sites. In addition, we found an enrichment

of ethnicity-associated sites amongst loci previously associated with environmental and

social exposures, particularly maternal smoking during pregnancy. This suggests that

ethnic self-identification may function as a proxy for environmental exposures. Overall

we conclude that race and ethnicity provide important and relevant clinical and

biomedical information above and beyond and individual's genetic ancestry.

Significance Statement

Race, ethnicity, and genetic ancestry have had a complex and often controversial history

within biomedical research and clinical practice. For example, race- and ethnicity-

specific clinical reference standards are based on an average derived from statistical

modeling applied to population-based sampling on a given physical trait such as

pulmonary function. However, because race and ethnicity are social constructs, they

ignore the heterogeneity within the categories. To account for these heterogeneities and

avoid social and political controversies, the genetics community has integrated the use

of genetic ancestry as a proxy because genetic sequence is not altered by environmental

or social factors. We explore the relative contributions of ancestry, ethnicity and

environment to variation in methylation, a fundamental biological process.

Introduction

Race, ethnicity, and genetic ancestry have had a complex and often controversial history

within biomedical research and clinical practice[1-3]. For example, race- and ethnicity-

specific clinical reference standards are based on an average derived from statistical

modeling applied to population-based sampling on a given physical trait such as

pulmonary function[4,5]. However, because race and ethnicity are social constructs and

poor markers for genetic diversity, they ignore the heterogeneity within the categories[6].

To account for these heterogeneities and avoid social and political controversies, the

genetics community has integrated the use of genetic ancestry instead of race and

ethnicity[3] because genetic sequence is not altered by environmental or social factors,

such as those related to racial or ethnic self-identity. Indeed, recent work from our

group and others have demonstrated that genetic ancestry improves diagnostic

precision compared to crude racial/ethnic categorizations for specific medical

conditions and clinical decisions[7-9].

Epigenetic modification of the genome through methylation plays a key role in the

regulation of diverse cellular processes[10]. Changes in DNA methylation patterns have

been associated with complex diseases, including various cancers[11], cardiovascular

disease[12,13], obesity[14], diabetes[15], autoimmune and inflammatory diseases[16], and

neurodegenerative diseases[17]. Epigenetic changes are thought to reflect influences of

both genetic[18] and environmental factors[19]. The discovery of methylation quantitative

trait loci (meQTL's) across populations by Bell *et al.* established the influence of genetic

factors on methylation levels in a variety of tissue types[18], with meQTL's explaining

between 22% and 63% of the variance in methylation levels. Multiple environmental

factors have also been shown to affect methylation levels, including endocrine

disruptors, tobacco smoke[20,21], polycyclic aromatic hydrocarbons, infectious pathogens,

particulate matter, diesel exhaust particles[22], allergens, heavy metals, and other indoor

and outdoor pollutants[23]. Psychosocial factors, including measures of traumatic

experiences[24-26], socioeconomic status[27,28], and general perceived stress[29], also affect

methylation levels. Racial and ethnic categories reflect the shared experiences and

exposures to known risk factors for disease, such as air pollution and tobacco smoke,

poverty, and inadequate access to medical services, which have all contributed to worse

disease outcomes in certain populations[30,31]. Since environmental exposures affect

methylation, they may be reflected in systematic differences in methylation levels

between racial or ethnic groups that cannot be captured by using genetically defined

ancestry alone.

Given the roles of both genetic and environmental influences upon methylation, we

leveraged genome-wide methylation data as an intermediate phenotype to examine the

degree to which self-identified ethnicity and genetic ancestry are reflected in differences

in methylation. We hypothesized that while genetic ancestry can explain many of the

differences in methylation between these groups, some ethnic-specific methylation

differences reflecting social and environmental differences between groups, might

remain. If our hypothesis is correct, our findings would have important implications for

both the use of ancestry to capture the biological changes and of race/ethnicity to

account for social and environmental exposures. Epigenome-wide association studies in

diverse populations may be susceptible to confounding due to environmental exposures

in addition to confounding due to population stratification[32].

We also examined the relationship between genome-wide (global) estimates of ancestry

and locus-specific (local) ancestry to determine the extent to which associations between

global ancestry and methylation are reflective of genetic factors acting in -cis. Finally, by

using dense genotyping arrays, we queried whether methylation differences associated

with ancestry can be traced back to meQTLs whose allele frequencies differ by ancestry.

To address these aims, we analyzed data from 573 Latino children according to their

national origin identity or ethnic subgroup (such as Puerto Rican and Mexican),

enrolled in the Genes-Environments and Admixture in Latino Americans (GALA II)

study of childhood asthma[33].

Results

The study included 573 participants, the majority of whom self-identified as being either

of Puerto Rican (n = 220) or Mexican origin (n = 276). Table 1 displays baseline

characteristics of the GALA II study participants with methylation data included in this

study, stratified by ethnic subgroups (Puerto Rican, Mexican, Other Latino, and Mixed

Latinos who had grandparents of more than one national origin). SI Appendix Figure 1

shows the distribution of African, European, and Native American ancestry among the

524 participants with genomic ancestry estimates.

Differences in ethnicity and ancestry resulted in discernible patterns in the global

methylation profile as demonstrated in a multidimensional scaling analysis (SI

Appendix Figure 2A). As expected[27,34], the first few principal coordinates are strongly

correlated to imputed cell composition (SI Appendix Figure 2B and C). There are also

significant associations of self-identified sub-ethnicity with PC2 (p-ANOVA = 0.003),

PC3 (p-ANOVA = 0.004), PC6 (p-ANOVA = 0.0001), PC7 (p-ANOVA = 0.0003) [SI

Appendix Figure 3A], and PC8 (p-ANOVA = 0.0003), after adjusting for age, sex,

disease status, cell components, and technical factors (plate and position). Genetic

ancestry was associated with PC3 (p-ANOVA = 0.002), PC7 (p-ANOVA = 0.0004) [SI

Appendix Figure 3B] and PC8 (p-ANOVA = 0.001) in a two degree of freedom ANOVA

test, adjusting for age, sex, disease status, cell components, technical factors, and

ethnicity. SI Appendix Table 1 summarizes the results of the simple correlation analysis

of methylation with ethnicity and ancestry, as well as the adjusted nested ANOVA

models described above and the mediation results described below.

A mediation analysis[35] revealed that the associations between ethnicity and PCs 3, 7,

and 8 were significantly mediated by Native American ancestry (mediation p = 0.01,

<0.001, and <0.001, respectively) and inclusion of Native American ancestry in the

regression model of PCs 3, 7, and 8 caused the ethnicity associations to be non-

significant. However, the associations of ethnicity with PCs 2 and 6 were not explained

by Native American, African or European ancestry (mediation p > 0.05), suggesting that

ethnic differences are associated with global methylation patterns not captured by

genetic differences. When genetic ancestry was regressed on the methylation data with

the principal coordinates recalculated using the residuals of the regression between

methylation and ancestry, there was an association between ethnicity and PC6 (p-

ANOVA = 0.003). However, there was no association with any of the other principal

coordinates. These observations suggest that while genetic ancestry can explain some of

the association between ethnicity and global methylation patterns, other non-genetic

factors, such as environmental and social exposure differences associated with ethnicity,

influence methylation independent of genetic ancestry.

An epigenome-wide association study of self-identified ethnicity (see methods for

details of ascertainment of ethnicity) and methylation identified a significant difference

in methylation M-values between ethnic groups at 916 CpG sites at a Bonferroni-

corrected significance level of less than 1.6×10^{-7} [Figure 1A and SI Appendix Table 2].

The most significant association with ethnicity occurred at cg12321355 in the ABO blood

group gene (*ABO*) on chromosome 3 (p-ANOVA 6.7×10^{-22}) [Figure 1B]. A two degree of

freedom ANOVA test for genomic ancestry was also significantly associated with

methylation level at this site (p = 2.3×10^{-5}) [Figure 1C], and when the analysis was

stratified by ethnic sub-group, showed an association in both Puerto Ricans and

Mexicans (p = 0.001 for Puerto Ricans, p = 0.003 for Mexicans). Although adjusting for

genomic ancestry attenuated the effect of ethnicity, a significant association between

ethnicity and methylation remained (p = 0.04). Recruitment site, an environmental

exposure proxy, was not significantly associated with methylation at this locus (p = 0.5),

suggesting that environmental differences associated with ethnicity beyond geography

and ancestry are driving the association.

When we repeat the analysis adjusting for ancestry, a significant association remained

in 314 of the 834 (37.8%, $p < 2.2 \times 10^{-16}$ for enrichment) CpG sites associated with

ethnicity [SI Appendix Figure 4A and SI Appendix Table 2] (82 sites were excluded

because they demonstrated unstable coefficient estimates and inflated standard errors

due to strong correlations between ethnicity and ancestry, especially Native American

ancestry [see SI Appendix Figure 1]). Genomic ancestry explained a median of 4.2%

(IQR 1.8% to 8.3%) of the variance in methylation at these loci and accounts for a

median of 75.7% (IQR 45.8% to 92%) of the total variance in methylation explained

jointly by ethnicity and ancestry [SI Appendix Figure 4B]. Sensitivity tests for

departures from linearity, fine scale population substructure and the exclusion of the 16

participants who self-identified as "Mixed Latino" sub-ethnicity, did not meaningfully

affect our results [See SI Appendix Results and SI Appendix Tables 2-6]. We conclude

that genetic ancestry explains much but not all of the association between ethnicity and

methylation. We hypothesize that other non-genetic factors associated with ethnicity

could explain the ethnicity-associated methylation changes that cannot be accounted for

by genomic ancestry alone.

Methylation at CpG loci that had previously been reported to be associated with

environmental exposures whose exposure prevalence differs between ethnic groups

were tested for association with ethnicity in this study. A recent meta-analysis of

maternal smoking during pregnancy, an exposure that varies significantly by ethnicity[33],

identified associations with methylation at over 6,000 CpG loci[21]. We found 1341 of

4404 that passed QC in our own study (30.4%) were nominally associated with ethnicity

($p < .05$), which represented a highly significant ($p < 2 \times 10^{-16}$) enrichment. Using a

Bonferroni correction for the 4404 loci tested, 126 maternal-smoking related loci were

associated with ethnicity ($p < 1.1 \times 10^{-5}$), and 27 loci were among the 916 CpG's reported

above as associated with ethnicity [SI Appendix Table 7]. We also examined methylation

loci from an earlier study of maternal smoking in Norwegian newborns[20] as well as

studies of diesel exhaust particles[22] and exposure to violence[24]. These results are

supportive of our hypothesis that environmental exposures may be responsible for the

observed differences in methylation between ethnic groups and are presented in the SI

Appendix Text and in SI Appendix Table 8.

We also found 194 loci with a significant association between global genetic ancestry

and methylation levels (after adjusting for ethnicity) at a Bonferroni corrected

association p-value of less than 1.6×10^{-7} [SI Appendix Figure 5 and SI Appendix Table

9], including 48 that were associated with ethnicity in our earlier analysis. Of these

significant associations, 55 were driven primarily by differences in African ancestry, 94

by differences in Native American ancestry, and 45 by differences in European ancestry.

The most significant association between methylation and ancestry occurred at

cg04922029 in the Duffy antigen receptor chemokine gene (DARC) on chromosome 1

(ANOVA p-value 3.1×10^{-24}) [SI Appendix Figure 5B]. This finding was driven by a strong

association between methylation level and global African ancestry; each 25 percentage

point increase in African ancestry was associated with an increase in M-value of 0.98,

which corresponds to an almost doubling in the ratio of methylated to unmethylated

DNA at the site (95% CI 0.72 to 1.06 per 25% increase in African ancestry, $p = 1.1\times10^{-21}$).

The distribution of methylation M-values at cg04922029 is tri-modal, raising the

possibility that a SNP whose allele frequency differs between African and non-African

populations may be driving the association. There was no significant heterogeneity in

the association between genetic ancestry and methylation between Puerto Ricans and

Mexicans (p-het = 0.5). Mexicans have a mean unadjusted methylation M-value 0.48

units lower than Puerto Ricans (95% CI 0.35 to 0.62 units, $p = 1.1\times10^{-11}$). However,

adjusting for African ancestry accounts for the differences in methylation level between

the two sub-groups (p-adjusted = 0.4), demonstrating that ethnic differences in

methylation at this site are due to differences in African ancestry.

A substantial proportion of the effect of global ancestry on local methylation levels is

due to local ancestry acting in -cis. Among the 194 CpG sites associated with global

ancestry, local ancestry at the CpG site explained a median of 10.4% (IQR 3.0% to

19.4%) of the variance in methylation, accounting for a median of 52.8% (IQR 20.3% to

84.9%) of the total variance explained jointly by local and global ancestry [SI Appendix

Figure 5].

In an admixture mapping study, we find that methylation at 3,694 of 321,503 CpG's

(1.1%), was significantly associated with ancestry at the CpG site at a Bonferroni

corrected association p-value of less than 1.6×10^{-7} [Figure 2A and SI Appendix Table

10]. This included 118 of the 194 loci identified above (61%), where global ancestry was

associated with methylation. The most significant CpG site was again cg04922029,

which was almost perfectly correlated with African ancestry at the locus (p = 6×10^{-162})

[Figure 2B]. Each African haplotype at the CpG site was associated with an increase in

methylation M-value of 2.7, corresponding to a 6.5-fold increase in the ratio of

methylated to unmethylated DNA per African haplotype at that locus. The second most

significant association occurred at cg06957310 on chromosome 17; each increase in

African ancestry at the locus was associated with a decrease in M-value of 1.7 (a 3.2-fold

decrease in the ratio of methylated to unmethylated DNA; p = 3.7×10^{-75}). We obtained

similar results when performing the analysis using methylation ß values instead of M-

values [See SI Appendix text and SI Appendix Table 11]

For each of the admixture mapping loci, we tested whether a single nucleotide

polymorphism (SNP) within 1 Mb of the CpG was associated with methylation. We

found 3637 loci out of the 3694 (98.5%) admixture mapping findings with at least one

SNP within 1 Mb that was significantly associated with methylation levels (after

adjustment of the number of SNPs in cis-) [SI Appendix Table 12]

The SNP/CpG pair were separated by a median distance of 10.9 kb (interquartile range

2.9 kb to 35.1 kb). The furthest SNP/CpG pair were 998 kb apart. The most significant

SNP/CpG pair were cg25134647/rs4963867, on chromosome 12, which are separated by

412 base pairs [SI Appendix Figure 6A/B]. Each copy of the T allele was associated with

a decrease in M-value of 3.58, corresponding to a nearly 12-fold decrease in the ratio of

methylated to unmethylated DNA at the site [SI Appendix Figure 6C]

(p < 10^{-370}). We found that CpG cg04922029 (our top admixture mapping association)

was significantly correlated with SNP rs2814778 [Figure 2C], the Duffy null mutation,

212 base pairs away; each copy of the C allele was associated with an increase in M-value

of 1.5, or a 2.9-fold increase in the ratio of methylated to unmethylated DNA (p =

3.8×10^{-90}) [Figure 2D]. We obtained similar results when performing the analysis using

methylation ß values instead of M-values [See SI Appendix text and SI Appendix Table

13]

Discussion

We have shown that both genomic ancestry and self-identified ethnicity are

independently associated with methylation levels throughout the genome. While

genomic ancestry can explain a portion of the association between ethnicity and

methylation, genomic ancestry inadequately accounts for the association between

ethnicity and methylation at 34% (314/916) of loci. Our results suggest ethnic self-

identification is unlikely to have a direct causal relationship with methylation but rather,

that other non-genetic factors associated with self-identified ethnicity may affect

differences in methylation patterns between Latino subgroups. These factors may

include social, economic, cultural, and environmental exposures.

We conclude that systematic environmental differences between ethnic subgroups likely

play an important role in shaping the methylome for both individuals and populations.

Loci previously associated with diverse environmental exposures such as *in utero*

exposure to tobacco smoke[20,21], as well as diesel exhaust particles[22] and psychosocial

stress[24] were enriched in our set of loci where methylation was associated with ethnicity.

Twenty-seven of the loci associated with maternal smoking during pregnancy in a large

consortium meta-analysis[21] were associated with methylation at a genome-wide

significance threshold of 1.6×10^{-7}. Thus, inclusion of relevant social and environmental

exposures in studies of methylation may help elucidate racial/ethnic disparities in

disease prevalence, health outcomes and therapeutic response. However, in many cases,

a detailed environmental exposure history is unknown, unmeasurable or poorly

quantifiable, and race/ethnicity becomes a useful, albeit imperfect proxy.

Our comprehensive analysis of high-density methyl- and genotyping from genomic DNA

allowed us to investigate the genetic control of methylation in great detail and without

the potential destabilizing effects of EBV transformation and culture in cell lines[36]. The

strongest patterns of methylation are associated with cell composition in whole blood[27].

However, the specific type of Latino ethnic-subgroups (Puerto Rican, Mexican, other, or

mixed) is also associated with principal coordinates of genome-wide methylation.

Our approach has some potential limitations. It is possible that fine-scale population

structure (sub-continental ancestry) within European, African, and Native American

populations may contribute to ethnic differences in methylation, as we had previously

reported in the case of lung function[37]. However, despite the presence of additional

substructure among the GALA II participants, PC's 3-10 explained the association

between ethnicity and ancestry at only 51 loci. PCs from chip-based genotypes will not

capture all forms of genetic variation. Clusters of ethnicity specific rare variants of large

effect or strong ethnicity-specific selective sweeps in the last 8-12 generations[38] could

also give rise to methylation differences, but these are inconsistent with existing rare

variant and selection analyses[39,40]. Our models of genetic ancestry assumed a linear

effect of ancestry on methylation, whereas a nonlinear association or other model

misspecification could have led to incomplete adjustment for genetic ancestry, and thus,

led to a residual association between ethnicity and methylation. However, when we

added second and third order polynomials or cubic splines to our models, we found

evidence for a nonlinear association between ancestry and methylation at only 25 and

26 loci, respectively, and it did not affect the association between ethnicity and

methylation. Although it is impossible to account for all types of non-linearity and non-

additivity (such as gene by gene or gene by environment interaction), our analysis

suggests that non-linear effects are unlikely to be significant. To rule out any residual

confounding due to recruitment sites, we conducted an additional analysis on the effect

of recruitment site on methylation both for the overall study and for the Mexican

participants (the largest study population in this analysis). We observed no significant

independent effect of recruitment site suggesting that confounding due to recruitment

region was limited, at least within the United States. We were unable to test for the

effect of geographic differences between the United States and Puerto Rico because our

study included relatively few Puerto Ricans recruited outside of Puerto Rico.

The presence of a strong association between genetic ancestry and methylation raises

the possibility that epigenetic studies can be confounded by population stratification,

similar to genetic association studies, and that adjustment for either genetic ancestry or

selected principal components is warranted. This possibility was first demonstrated in a

previous analysis of the association between self-described race and methylation[41].

However, the study only evaluated two distinct racial groups (African Americans and

Whites), while the present study demonstrates the possibility of population

stratification in an admixed and heterogeneous population with participants from

diverse Latino national origins. The tendency to consider Latinos as a homogenous or

monolithic ethnic group makes any analysis of this population particularly challenging.

Our finding of loci whose methylation patterns differed between Latino ethnic

subgroups, even after adjusting for genetic ancestry, suggests that any analysis of these

populations in disease-association studies without adjusting for ethnic heterogeneity is

likely to result in spurious associations even after controlling for genomic ancestry.

Our analysis of local genetic ancestry and methylation demonstrates that loci associated

with genome-wide ancestry are driven primarily by allele frequency differences between

ancestral populations in 118 out of 194 loci, suggesting that in most cases global ancestry

is acting in -cis. In addition, methylation-QTLs whose allele frequencies differ between

ethnic groups are found in 95% (3637/3694) of loci associated with local ancestry. Of

particular interest, the most significant ancestry-associated locus, the *DARC* gene,

harbors an association between ancestry and methylation at cg04922029, which can be

entirely explained by the genotype at rs2814778, the Duffy null mutation. This mutation,

which confers resistance to *P. vivax* malaria, has an allele frequency of 100% in

individuals in the five 1000 genomes populations[42-44] from Africa (Esan in Nigeria,

Gambian in the Western Division of Gambia, Luhya in Webuye, Kenya, Mende in Sierra

Leone, and Yoruban in Ibadan, Nigeria), and 80% to 90% in admixed populations of

African origin in the Americas (89% in Afro Caribbeans in Barbados, and 80% in

African Americans in the Southwest U.S.). In contrast, the allele frequency is 1% in the

five European populations and 0% in the five Asian populations [Figure 2E], consistent

with the known high level of ancestry information at the locus. Latinos, being admixed,

have intermediate and more varied minor allele frequencies, ranging from 3% in

Mexicans to 14% in Puerto Ricans.

In summary, this study provides a framework for understanding how genetic, social and

environmental factors can contribute to systematic differences in methylation patterns

between ethnic subgroups, even between presumably closely related populations such as

Puerto Ricans and Mexicans. Methylation QTL's whose allele frequency varies by

ancestry lead to an association between local ancestry and methylation level. This, in

turn, leads to systematic variation in methylation patterns by ancestry, which then

contributes to ethnic differences in genome-wide patterns of methylation. However,

although genetic ancestry has been used to adjust for confounding in genetic studies,

and can account for much of the ethnic differences in methylation in this study, ethnic

identity is associated with methylation independent of genetic ancestry. This is likely

due to social and environmental effects captured by ethnicity. Indeed, we find that CpG

sites known to be influenced by social and environmental exposures are also

differentially methylated between ethnic subgroups. These findings called attention to a

more complete understanding of the effect of social and environmental variables on

methylation in the context of race and ethnicity to fully understanding this complex

process.

Our findings have important implications for the independent and joint effects of race,

ethnicity, and genetic ancestry in biomedical research and clinical practice, especially in

studies conducted in diverse or admixed populations. Our conclusions may be

generalizable to any population that is racially mixed such as those from South Africa,

India, and Brazil, though we would encourage further study in diverse populations. As

the National Institutes of Health (NIH) embarks on a precision medicine initiative, this

research underscores the importance of including diverse populations and studying

factors capturing the influence of social, cultural, and environmental factors, in addition

to genetic ones, upon disparities in disease and drug response.

Methods

All research on human subjects was approved by the Institutional Review Board at the

University of California and each of the recruitment sites (Kaiser Permanente Northern

California, Children's Hospital Oakland, Northwestern University, Children's Memorial

Hospital Chicago, Baylor College of Medicine on behalf of the Texas Children's Hospital,

VA Medical Center in Puerto Rico, the Albert Einstein College of Medicine on behalf of

the Jacobi Medical Center in New York and the Western Review Board on behalf of the

Centro de Neumologia Pediatrica), and all participants/parents provided age-

appropriate written assent/consent. Latino children were enrolled as a part of the

ongoing GALA II case-control study, where details of recruitment can be found[33] as well

as in the SI Appendix text.

Genomic DNA (gDNA) was extracted from whole blood using Wizard® Genomic DNA

Purification Kits (Promega, Fitchburg, WI). A subset of 573 participants (311 cases with

asthma and 262 healthy controls) was selected for methylation. Methylation was

measured using the Infinium HumanMethylation450 BeadChip (Illumina, Inc., San

Diego, CA) following the manufacturer's instructions. Details of methylation measures

and quality control are described in the SI Appendix text.

Details of genotyping and quality control procedures for single nucleotide

polymorphisms (SNPs) and individuals have been described elsewhere[45] and

summarized in the SI Appendix Text.

Unless otherwise noted, all regression models were adjusted for case status, age, sex,

estimated cell counts, and plate and position. To account for possible heterogeneity in

the cell type makeup of whole blood we inferred white cell counts using the method by

Houseman et al[34]. Indicator variables were used to code categorical variables with more

than two categories, such as ethnicity. In these cases, a nested analysis of variance

(ANOVA) was used to compare models with and without the variables to obtain an

omnibus p-value for the association between the categorical variable and the outcome.

For analyses of dependent beta-distributed variables (such as African, European, and

Native American ancestries), or cell proportion, k-1 variables were included in the

analysis, and a nested analysis of variance (ANOVA) was used to compare models with

and without the variables to obtain an k-1 degree of freedom omnibus p-value for the

association between predictor (such as ancestry) and the outcome variable.

The Bonferroni method was used to adjust for multiple comparisons. For methylome-

wide associations, the significance threshold was adjusted for 321,503 probes, resulting

in a Bonferroni threshold of 1.6×10^{-7}. Analyses were performed using R version 3.2.1

(The R Foundation for Statistical Computing)[46] and the Bioconductor package version

2.13. Further details on the models used for specific statistical analyses is in the SI

Appendix Text.

Acknowledgements

The authors acknowledge the families and patients for their participation and thank the

numerous health care providers and community clinics for their support and

participation in GALA II. In particular, the authors thank study coordinator Sandra

Salazar; the recruiters who obtained the data: Duanny Alva, MD, Gaby Ayala-Rodriguez,

Lisa Caine, Elizabeth Castellanos, Jaime Colon, Denise DeJesus, Blanca Lopez , Brenda

Lopez, MD, Louis Martos, Vivian Medina, Juana Olivo, Mario Peralta, Esther Pomares,

MD, Jihan Quraishi, Johanna Rodriguez, Shahdad Saeedi, Dean Soto, Ana Taveras. We

also thank Sasha Gusev for helpful discussion. Computations in this manuscript were

performed using the UCSF Biostatistics High Performance Computing System.

References

1.   Risch, N., Burchard, E., Ziv, E. & Tang, H. Categorization of humans in biomedical
research: genes, race and disease. *Genome Biology* **3,** comment2007 (2002).

2.   Cooper, R. S., Kaufman, J. S. & Ward, R. Race and genomics. *N. Engl. J. Med.*
348, 1166–1170 (2003).

3.   Yudell, M., Roberts, D., Desalle, R. & Tishkoff, S. SCIENCE AND SOCIETY.
Taking race out of human genetics. *Science* **351,** 564–565 (2016).

4.   Hankinson, J. L., Odencrantz, J. R. & Fedan, K. B. Spirometric reference values
from a sample of the general U.S. population. *Am. J. Respir. Crit. Care Med.* **159,**
179–187 (1999).

5.   Quanjer, P. H. *et al.* Multi-ethnic reference values for spirometry for the 3-95-yr
age range: the global lung function 2012 equations. *European Respiratory*
Journal **40,** 1324–1343 (2012).

6.   Borrell, L. N. Racial identity among Hispanics: implications for health and well-
being. *Am J Public Health* **95,** 379–381 (2005).

7.   Kumar, R. *et al.* Genetic Ancestry in Lung-Function Predictions. *N. Engl. J. Med.*
363, 321–330 (2010).

8.   Udler, M. S. *et al.* Effect of Genetic African Ancestry on eGFR and Kidney Disease.
J. Am. Soc. Nephrol. **26,** 1682–1692 (2015).

9.   Nalls, M. A. *et al.* Admixture Mapping of White Cell Count: Genetic Locus
Responsible for Lower White Blood Cell Count in the Health ABC and Jackson

Heart Studies. *The American Journal of Human Genetics* **82,** 81–87 (2008).

10.   Smith, Z. D. & Meissner, A. DNA methylation: roles in mammalian development.
Nat Rev Genet **14,** 204–220 (2013).

11.   Kulis, M. & Esteller, M. DNA methylation and cancer. *Adv. Genet.* **70,** 27–56
(2010).

12.   Udali, S., Guarini, P., Moruzzi, S., Choi, S.-W. & Friso, S. Cardiovascular
epigenetics: from DNA methylation to microRNAs. *Mol. Aspects Med.* **34,** 883–
901 (2013).

13.   Kato, N. *et al.* Trans-ancestry genome-wide association study identifies 12 genetic
loci influencing blood pressure and implicates a role for DNA methylation. *Nat.*
Genet. (2015). doi:10.1038/ng.3405

14.   Bell, C. G. *et al.* Integrated genetic and epigenetic analysis identifies haplotype-
specific methylation in the FTO type 2 diabetes and obesity susceptibility locus.
PLoS ONE **5,** e14040 (2010).

15.   Chambers, J. C. *et al.* Epigenome-wide association of DNA methylation markers
in peripheral blood from Indian Asians and Europeans with incident type 2
diabetes: a nested case-control study. *Lancet Diabetes Endocrinol* **3,** 526–534
(2015).

16.   Liu, Y. *et al.* Epigenome-wide association data implicate DNA methylation as an
intermediary of genetic risk in rheumatoid arthritis. *Nat. Biotechnol.* **31,** 142–147
(2013).

17.   Lardenoije, R. *et al.* The epigenetics of aging and neurodegeneration. *Prog.*
Neurobiol. **131,** 21–64 (2015).

18.   Bell, J. T. *et al.* DNA methylation patterns associate with genetic and gene
expression variation in HapMap cell lines. *Genome Biology* **12,** R10 (2011).

19.   Feil, R. & Fraga, M. F. Epigenetics and the environment: emerging patterns and
implications. *Nat Rev Genet* **13,** 97–109 (2011).

20.   Joubert, B. R. *et al.* 450K epigenome-wide scan identifies differential DNA

methylation in newborns related to maternal smoking during pregnancy. *Environ.*
Health Perspect. **120,** 1425–1431 (2012).

21. Joubert, B. R. *et al.* DNA Methylation in Newborns and Maternal Smoking in
Pregnancy: Genome-wide Consortium Meta-analysis. *Am. J. Hum. Genet.* **98,**
680–696 (2016).

22. Jiang, R., Jones, M. J., Sava, F., Kobor, M. S. & Carlsten, C. Short-term diesel
exhaust inhalation in a controlled human crossover study is associated with
changes in DNA methylation of circulating mononuclear cells in asthmatics. *Part*
Fibre Toxicol **11,** 71 (2014).

23. Ho, S.-M. *et al.* Environmental epigenetics and its implication on disease risk and
health outcomes. *ILAR J* **53,** 289–305 (2012).

24. Chen, W. *et al.* ADCYAP1R1 and asthma in Puerto Rican children. *Am. J. Respir.*
Crit. Care Med. **187,** 584–588 (2013).

25. Ressler, K. J. *et al.* Post-traumatic stress disorder is associated with PACAP and
the PAC1 receptor. *Nature* **470,** 492–497 (2011).

26. van der Knaap, L. J. *et al.* Glucocorticoid receptor gene (NR3C1) methylation
following stressful events between birth and adolescence. The TRAILS study.
Transl Psychiatry **4,** e381 (2014).

27. Lam, L. L. *et al.* Factors underlying variable DNA methylation in a human
community cohort. *Proceedings of the National Academy of Sciences* **109 Suppl**
2, 17253–17260 (2012).

28. Borghol, N. *et al.* Associations with early-life socio-economic position in adult
DNA methylation. *International Journal of Epidemiology* **41,** 62–74 (2012).

29. Vidal, A. C. *et al.* Maternal stress, preterm birth, and DNA methylation at imprint
regulatory sequences in humans. *Genet Epigenet* **6,** 37–44 (2014).

30. Nguyen, A. B., Moser, R. & Chou, W. Y. Race and health profiles in the United
States: an examination of the social gradient through the 2009 CHIS adult survey.
Public Health **128,** 1076–1086 (2014).

31. Evans, G. W. & Kantrowitz, E. Socioeconomic Status and Health: The Potential
Role of Environmental Risk Exposure. *Annual Review of Public Health* **23,** 303–
331 (2002).

32. Michels, K. B. *et al.* Recommendations for the design and analysis of epigenome-
wide association studies. *Nat. Methods* **10,** 949–955 (2013).

33. Oh, S. S. *et al.* Effect of secondhand smoke on asthma control among black and
Latino children. *J. Allergy Clin. Immunol.* **129,** 1478–83.e7 (2012).

34. Houseman, E. A. *et al.* DNA methylation arrays as surrogate measures of cell
mixture distribution. *BMC Bioinformatics* **13,** 86 (2012).

35. Tingley, D., Yamamoto, T., Hirose, K., Keele, L. & Imai, K. mediation: R package
for causal mediation analysis. *UCLA Statistics/American Statistical Association*
(2014).

36. Grafodatskaya, D. *et al.* EBV transformation and cell culturing destabilizes DNA
methylation in human lymphoblastoid cell lines. *Genomics* **95,** 73–83 (2010).

37. Moreno-Estrada, A. *et al.* Human genetics. The genetics of Mexico recapitulates
Native American substructure and affects biomedical traits. *Science* **344,** 1280–
1285 (2014).

38. Galanter, J. M. *et al.* Development of a panel of genome-wide ancestry informative
markers to study admixture throughout the Americas. *PLoS Genet.* **8,** e1002554
(2012).

39. Hernandez, R. D. *et al.* Classic selective sweeps were rare in recent human
evolution. *Science* **331,** 920–924 (2011).

40. Tang, H. *et al.* Recent Genetic Selection in the Ancestral Admixture of Puerto
Ricans. *The American Journal of Human Genetics* **81,** 626–633 (2007).

41. Barfield, R. T. *et al.* Accounting for population stratification in DNA methylation
studies. *Genet. Epidemiol.* **38,** 231–241 (2014).

42. 1000 Genomes Project Consortium *et al.* A map of human genome variation from
population-scale sequencing. *Nature* **467,** 1061–1073 (2010).

43.   1000 Genomes Project Consortium *et al*. An integrated map of genetic variation
499   from 1,092 human genomes. *Nature* **491,** 56–65 (2012).

500   44.   Sudmant, P. H. *et al*. An integrated map of structural variation in 2,504 human
501   genomes. *Nature* **526,** 75–81 (2015).

502   45.   Galanter, J. M. *et al*. Genome-wide association study and admixture mapping
503   identify different asthma-associated loci in Latinos: the Genes-environments &
504   Admixture in Latino Americans study. *J. Allergy Clin. Immunol.* **134,** 295–305
505   (2014).

46.   Team, R. C. R: A language and environment for statistical computing.

Figure Legends

Figure 1: [A] Manhattan plot showing the associations between ethnicity and
methylation at individual CpG loci. [B] Violin plot showing one such locus, cg19145607.
Mexicans are relatively hypermethylated compared to Puerto Ricans ($p = 1.4 \times 10^{-19}$). [C]
Plot showing the association between Native American ancestry at the locus and
methylation levels at the locus colored by ethnicity; Native American ancestry accounts
for 58% of the association between ethnicity and methylation at the locus.

Figure 2: [A] Manhattan plot showing the association between local ancestry and
methylation at individual CpG loci. [B] Association between cg04922029 on the *DARC*
locus and African ancestry, color coded by ethnic group. There is near perfect
correlation between the two. [C] Association between SNPs located within 1Mb of
cg04922029 and methylation levels at that CpG. [D] Association between rs2814778
(Duffy null) genotype and methylation at cg04922029, color coded by the number of
African alleles present. There is near perfect correlation between genotype, ancestry and
methylation at the locus. [E] Allele frequency of rs2814778 by 1000 Genomes
population. The C allele is nearly ubiquitous in African populations and nearly absent
outside of African populations and their descendants.

TABLE 1: Baseline characteristics of GALA II participants with methylation data, stratified by ethnicity.

	Mexican	Puerto Rican	Mixed Latino	Other Latino
n	276	220	16	61
Males (%)	125 (45.3%)	127 (57.7%)	6 (37.5%)	28 (45.9%)
Age	11.4 [9.3: 14.7]	12.3 [10.4: 14.2]	11.8 [10.7: 14.9]	11.8 [10: 15.7]
Asthma cases (%)	124 (44.9%)	147 (66.8%)	9 (56.3%)	31 (50.8%)
Ancestry (n = 524)				
African	4.3% [2.9%: 6.0%]	22.8% [16.6%: 29.4%]	8.5% [5.6%: 19.2%]	12.3% [6.3%: 25.8%]
Native American	55.4% [44.5%: 65.7%]	11.2% [9.8%: 13%]	31.5% [20.9%: 45.6%]	32.8% [10.4%: 49.3%]
European	40.5% [29.9%: 50.2%]	65.7% [59.2%: 71%]	50.5% [44.6%: 57.6%]	48.9% [40%: 58.5%]
Recruitment Site				
Chicago	140 (50.7%)	15 (6.8%)	11 (68.9%)	15 (24.6%)
New York	18 (6.5%)	10 (4.5%)	1 (6.3%)	23 (37.7%)
Puerto Rico	0	193 (87.7%)	0	0
San Francisco	78 (28.3%)	0	2 (12.5%)	23 (37.7%)
Houston	40 (14.5%)	2 (0.9%)	2 (12.5%)	5 (8.2%)
Cell Counts (estimated)				
Granulocytes	51.2% [46.0%: 55.7%]	51.6% [46.8%: 57%]	51% [43.6%: 57.2%]	49.1% [43.8%: 55.8%]
Lymphocytes	41.9% [36.9%: 46.6%]	41.8% [36.9%: 46.5%]	41.9% [36.1%: 51.6%]	43.9% [36.8%: 49.6%]
Monocytes	7.1% [5.8%: 8.3%]	6.74% [5.74%: 8.24%]	6.6% [5.7%: 7.6%]	7.4% [6.2%: 8.6%]

Figure 1

Figure 2

A

B

Methylation by Local African Ancestry

Ethnicity
- Mexican
- Mixed Latino
- Other Latino
- Puerto Rican

C

Methylation QTLs for cpg cg04922029 in chromosome 1

DARC

rs10021525   Distance = -212

D

Methylation at cg04922029 by rs2814778 genotype

E