Ageing, TOR and amino acid restriction: a cross-tissue transcriptional network connects GATA factors to *Drosophila* longevity

Adam J Dobson[1], Xiaoli He[1], Eric Blanc[2], Ekin Bolukbasi[1], Yodit Feseha[1], Mingyao Yang[1,2], Matthew DW Piper[1,4*]*

1 - Institute of Healthy Ageing, Department of Genetics, Evolution and Environment, University College London, Gower Street, London WC1E 6BT, UK.

2 - Berlin Institute of Health, Luisenstraße 56, 10117 Berlin, Germany.

3 - Current address: Institute of Animal Genetics and Breeding, Sichuan Agricultural University, Chengdu, Sichuan, China.

4 – Current address: School of Biological Sciences, Monash University, Victoria 3800, Australia.

* For correspondence: yangmingyao@sicau.edu.cn, matthew.piper@monash.edu

Keywords: Ageing, TOR, transcriptional networks, transcription factors, systems biology.

Abstract

Animal lifespan can be extended by dietary restriction (DR), but at a cost to fitness. This phenomenon depends on essential amino acids (EAAs) and TOR signalling, but roles of specific tissues and downstream transcriptional regulators are poorly characterised. Manipulating relevant transcription factors (TFs) specifically in lifespan-limiting tissues may ameliorate ageing without costs of DR. Here we identify TFs which regulate the DR phenotype in *Drosophila*, analysing organs as an interacting system and reducing its transcriptional complexity by two orders of magnitude. Evolutionarily conserved GATA TFs are predicted to regulate the overlapping effects of DR and TOR on organs, and genetic analyses confirmed that these TFs interact with diet to determine lifespan. Importantly, *Srp* knockdown insulated fly lifespan from the pernicious effects of EAAs, but tissue-specific knockdown reduced the corrolary costs. These results provide the first indication that benefits of EAAs for early-life fitness can be decoupled from longevity by tissue-specific transcriptional reprogramming.

Background

How can lifelong health be maximised? Answering this question is a major goal, as ever-increasing human lifespans outpace advances in gerontology, at great social, personal and financial cost [1]. Dietary restriction (DR) has the evolutionarily conserved capacity to improve lifelong health by reducing nutrient intake, but at a cost of reduced biological fitness and vigour in youth [2]. Despite having been discovered 80 years ago [3], the molecular mechanisms underpinning lifespan extension by DR remain elusive. Defining these mechanisms could ameliorate the burden of ageing without the costs of DR.

Calories do not fully account for the benefits of DR: specific nutrients and their relative ratios are key [4–6]. In *Drosophila*, the ratio of dietary sugar to yeast modulates lifespan, which is explained by essential amino acids (EAAs) from the yeast [7], and importantly the same mechanism is conserved in mice [8,9]. Recent evidence indicates that the phenotype of EAA-restricted *Drosophila* is recapitulated by pharmacologically suppressing the Target of Rapamycin (TOR) pathway [10,11], consistent with molecular evidence that EAAs positively regulate TOR [12]. Understanding of how TOR curtails lifespan is incomplete, although maintenance of proteome quality likely plays a role [13–15]. TOR also affects transcription [16–18], but to date this effect has been relatively poorly studied.

In *Drosophila,* transcriptomic responses to DR and TOR have been characterised at the cellular and organismal levels [18], but information on tissue-specific and organ-specific transcription is a requisite advance for many reasons. Primarily, an animal's overall phenotype is determined by tissues coordinating to match their collective functions to the environment. Consequently, to fully understand organismal phenotypes we must account for tissues as an interacting and integrated system. By quantifying the changes to that system in low-TOR or DR states, we gain a complete view of how tissues collectively mediate the DR phenotype. Such an understanding is likely to help identify manipulations of regulators of the DR transcriptional state, which may impart longevity even under *ad libitum* feeding. Minimising the number of tissues in which these regulators are manipulated may ameliorate associated costs, if costs and benefits of DR are mediated by distinct tissues. This approach recognises the findings of studies of insulin signalling, which indicate that tissue-restricted manipula-

tions are sufficient to extend lifespan, and therefore that transcriptional mechanisms

affecting longevity are incompletely described by organismal analyses [19,20]. Altogeth-

er, these observations suggest that understanding longevity via DR requires (1)

modelling of transcription across systems of interacting tissues, (2) identification of

changes to that system under DR, (3) quantification of the TOR-dependence of these

changes, (4) prediction and testing of tissue-specific regulators of DR-like gene ex-

pression.

Recent analytical advances facilitate prediction of transcriptional regulators

from patterns of gene expression. Regulatory elements, such as transcription factors,

can be predicted by enrichment analysis amongst groups of genes [21,22]. Transcrip-

tional network analysis, also known as coexpression analysis, identifies transcripts

which are expressed together [23,24]. The patterns in a transcriptional network therefore

reflect the tendency of genes to be switched on and off together, which can be con-

sidered a reflection of the underlying gene-regulatory machinery. Consequently, tran-

scriptional network analyses provide a strong basis from which to predict regulatory

elements. Network analysis can also answer the need to understand tissues as a

complete system, by identifying how patterns of expression in one tissue which cor-

respond to that in others. Therefore, by defining transcriptional networks across tis-

sues and how they change under DR, we can distill the complex interactions between

transcripts and tissues that define the phenotype of the DR animal, and systematical-

ly reflect the status of the underlying gene-regulatory machinery.

Here, we address a key question: By studying transcription across tissues,

can we identify manipulations of transcriptional regulators to recapitulate the DR

phenotype? This study is facilitated by dietary manipulations that offer a precise tool-

set to dissect DR. Previously, we developed a semi-defined *Drosophila* diet, which is

optimal for early-life egg laying, in which 50% of available EAAs are provided as a

supplement to yeast-based medium [7,10]. Against this "fully fed" control, lifespan can

be extended by two interventions, both with correlated fecundity costs to egg laying in

early life: either omission of the EAA supplement (i.e. DR); or the addition of rapamy-

cin (EAA+rapamycin), which extends lifespan in the presence of EAAs by suppress-

ing TOR pharmacologically. These tools allow us to establish the effects of EAA dilu-

tion, and the TOR-dependence of these effects. Taking these diets and capitalising

on orthology between *Drosophila* and vertebrate organs, we studied transcriptomes

in the brain, fat body (the analogue of the vertebrate liver and adipose), gut, ovary

and thorax (which largely comprises muscle). We then used transcriptional network analysis to analyse these tissues integratively, reflecting underlying regulatory machinery. Promoter analysis was then applied to identify candidate transcription factor regulators. Crucially, when we tested these transcription factors genetically, we were able to validate their role as mediators of dietary effects on lifespan, and also demonstrate that costs of lifespan extension can be mitigated by targeting interventions to specific tissues.

Results

The effects of DR on ageing are thought to be mediated by the TOR pathway. Here, by characterising the organ-specific transcriptional changes caused by EAA restriction (DR) and pharmacological TOR suppression, we isolate changes that are associated with longevity in both experimental conditions, and address whether the effects of DR mimic the effects of low TOR. We also associate *cis*-regulatory elements with these changes, to predict relevant transcription factors. The study design is presented in Figure 1. We term DR as a diet-induced longevity condition, and rapamycin supplementation as a drug-induced longevity condition.

Figure 1. Study design. Against a control in which short lifespan is determined by enrichment of essential amino acids (EAAs), lifespan is extended by DR of the EAAs (diet-induced longevity), or pharmacologically, by administration of the TOR-suppressive drug rapamycin (drug-induced longevity). This lifespan extension comes at a biological cost of reduced early-life fitness (i.e. fecundity). Tissue-specific transcriptional changes associated with both conditions are therefore longevity-

associated, EAA-dependent and TOR-dependent. Identifying *Cis*-regulatory elements associated with these transcriptional networks predicts mechanisms by which DR affects transcription via TOR.

Tissue-specific transcriptomic effects of DR are recapitulated by TOR suppression

To assess whether the conditions of diet-induced longevity and drug-induced longevity have equivalent transcriptional effects, we first studied the equivalence of these treatments within tissues. We compared the changes in expression of all genes in the transcriptome under these two conditions, with the prediction that the induced changes would be positively correlated. As predicted, in whole flies, brains, guts, ovaries and thoraces, the diet-induced and drug-induced longevity conditions had positively correlated effects on gene expression (Figure 2). The transcriptional changes observed in these organs therefore mirror the associated changes to lifespan [10]. However, the two long-lived conditions did not have equivalent effects in the fat body, indicating that the coordination of function is largely but not obligately coupled across organs. To test explicitly the overlapping effects of DR and TOR suppression at the level of individual transcripts, differential expression analysis was performed. Surprisingly, no genes were differentially expressed in the ovary under DR, despite the robust decrease in egg laying in this condition [10]. However, in each other tissue, both lifespan-enhancing treatments had overlapping transcriptional signatures (Table 1). Overall, these analyses demonstrate that the transcriptional effects of DR largely mirror those of pharmacological TOR suppression.

Table I. Overlapping effects of the diet-induced longevity (DR) and drug-induced longevity (rapamycin) conditions on changes in gene expression within specific tissues.

Gene set		Number of genes[†]			
Tissue	Expression change*	DR	Rapamycin	Overlap	P-value[§]
Whole fly	Up	254	2239	203	5.55e-121
	Down	40	2159	17	3.99e-06
Brain	Up	154	22	1	0.02
	Down	439	82	63	8.44e-82
Fat body	Up	221	160	6	0.02
	Down	408	156	56	2.75e-46
Gut	Up	8	49	6	7.60e-17
	Down	72	52	6	1.78e-08
Ovary	Up	0	30	NA	NA
	Down	0	47	NA	NA
Thorax	Up	141	39	10	1.32e-13
	Down	153	25	6	8.11e-09

* Relative to EAA diet
† Number of genes showing given change in expression
§ Hypergeometric test

125
126

Figure 2. Changes in expression induced by DR correlate those induced by TOR suppression. Log_2 fold-changes in expression in the diet-induced longevity and drug-induced longevity conditions, relative to the EAA-enriched control (DR/EAA fold-change and rapa/EAA fold-change, respectively), were calculated for every gene in the transcriptome. Fold-changes were calculated by DESeq2. Text above panels indicates correlations (Kendall's Tau) between fold-changes induced by the two long-lived conditions. All p-values $< 2.2e^{-16}$

To look for ubiquitous molecular signatures of lifespan extension, we exam-
ined overlaps between the transcriptional changes observed between different tis-
sues. Surprisingly, no one transcript responded to DR/TOR in all organs, and the
whole-fly samples captured only a small portion of the organ-specific changes asso-
ciated with lifespan extension (Figure S1). Therefore, DR cannot be understood in
terms of any one tissue. To address whether this was functionally relevant, we ana-
lysed enrichment of GO terms amongst the transcripts associated with longevity.
Most GO terms (83%) were associated with specific organs, showing that there is al-
so no ubiquitous functional signature of DR/TOR (Figure S2). Together these anal-
yses show that the DR regulon can largely be accounted for by TOR (excepting the
ovary), but that the identity of the responsive transcripts is tissue-dependent. The or-
gan-specificity of longevity-associated changes indicates parallel responses amongst
organs to EAAs and rapamycin, suggesting the coordination of organ-specific func-
tions changes under DR and low systemic TOR.

Diet and TOR suppression have overlapping effects on the orchestration of
expression across organs

Discrete tissues compartmentalise functions, and an individual's fitness de-
pends on correctly orchestrating these tissue-specific functions to match the nutri-
tional environment. It is therefore possible that the lifespan benefits of DR and TOR
suppression may be mediated by certain tissues, and the biological costs by others.
Therefore we investigated how DR changes the orchestration of gene expression
across organs, studied as an integrated system. To identify interdependent gene ex-
pression (i.e. gene coexpression) amongst organs, we employed Weighted Gene
Coexpression Network Analysis (WGCNA; [23]). This approach can identify codepend-
ent gene expression and reduce dimensionality of complex data: therefore, applying
it in our study allowed us to identify codependent gene expression across organs,
and changes in those codependencies in the long-lived conditions. After applying
quality controls and removing genes with zero variance, 11164 of 13442 analysed
genes clustered into 14 modules (Figure 3a, Figure 3b, Supplementary Materials),
representing groups of genes that share expression patterns across organs. By de-
scribing modularity in our data, this network provides a complete description of the
structure of the transcriptome across all organs and diets under study.

The table within the figure:

Module colour code	Module number	N. genes in module	Variance explained by Eigengene		
			DR	Control	Rapamycin
	1	4228	0.784	0.802	0.793
	2	2283	0.852	0.826	0.825
	3	1181	0.719	0.738	0.736
	4	1000	0.757	0.813	0.785
	5	873	0.840	0.821	0.816
	6	775	0.790	0.782	0.762
	7	200	0.849	0.838	0.805
	8	124	0.824	0.777	0.760
	9	118	0.835	0.857	0.805
	10	100	0.845	0.874	0.839
	11	83	0.845	0.874	0.839
	12	83	0.831	0.836	0.798
	13	65	0.850	0.894	0.844
	14	51	0.880	0.881	0.872

Figure 3. A network description of transcription across *Drosophila* organs and long-lived conditions. (a) Consensus transcriptional modules found across organs in all three experimental conditions. Genes were clustered according to their expression (log$_2$ RPKM) across organs by Weighted Gene Coexpression Network Analysis (WGCNA), revealing modules of genes showing similar coexpression across organs. Leaves of the tree indicate genes. The tree was cut by hybrid tree cutting to define transcriptional modules (clusters). Tree cutting assigned genes to one of 14 consensus transcriptional modules, indicated by colour-coded vertical lines that together form the horizontal bar beneath the dendrogram (see Supplementary Spreadsheets

for assignments of genes to modules). Genes not assigned to any module are colour-coded grey. **(b)** Module assignments identify clear structure in the transcriptional network, using the EAA-fed control as an example. Modules are indicated by coloured side bars. The heatmap indicates gene-gene adjacency (i.e. signed suared correlations) in expression. **(c)** Representative expression (Eigengenes) of all modules (y-axis showing Eigengene values, A.U.). These data summarise changes in transcriptional networks across organs and experimental conditions. Data are plotted by experimental condition in columns ("DR" = diet-induced longevity, "Rapamycin" = drug-induced longevity, short-lived control = "EAA"), and transcriptional module in rows. Within each plot, boxes are colour-coded by tissue (see key). Boxplots show medians (horizontal midline), 1st and 3rd quartiles (hinges), and range of data points.

We built on the transcriptional network analysis to isolate significant changes

in network structure in the two long-lived conditions, by identifying pairs of modules

exhibiting significantly changed coregulation. Quantifying module coregulation re-

quired reduction of the complexity of the data. Therefore, we summarised gene ex-

pression in each module with a single "Eigengene" vector per module (Figure 3c),

calculated as the first principal component of expression of the genes in each module

[25]. These Eigengenes accounted for between 72% and 89% of the variance in each

module (weighted average = 79.86%, Figure 3b), thus simplifying the description of

the organ system by two orders of magnitude, from $11.1e^3$ transcripts to 14

Eigengenes, whilst retaining ~80% of total information. We then used these

Eigengenes to quantify changes in module-module coregulation. For all pairs of

Eigengenes, correlations were calculated in each dietary condition then, for each

long-lived condition, the correlation coefficients were subtracted from the correspond-

ing coefficient in the EAA-fed control. This quantified changes in module-module

coregulation induced in the two long-lived conditions. Repeating the same procedure

on random permutations of the data generated null distributions for each pair of mod-

ules. Pairs of modules which showed significantly altered coregulation were identified

by comparing the observed changes in correlation coefficients to the null distribu-

tions. This analysis thereby systematically identified pairs of modules whose coregu-

lation was significantly altered by the conditions causing longer life (Figure 4). The

diet-induced longevity condition significantly changed the coregulation of three pairs

of modules (4 & 13, 4 & 3, 14 & 13). All significant changes in the drug-induced lon-

gevity condition were paired with module 11 (relative to modules 1, 4, 7, 8, 9, 12, 13).

This correlation-permutation approach identified changes in coregulation of pairs of

modules, but did not reveal whether those changes were driven by one or both mem-

bers of each pair. Therefore, by three separate methods, we asked which individual

modules' Eigengenes were most strongly perturbed across the experimental condi-

tions (Figure S3). These three analyses identified modules 4, 11 and 14, which were

all identified by the correlation-permutation analysis. Therefore, the Eigengene anal-

yses collectively indicated that altered regulation of modules 4, 11 and 14 changes

the coordination of functions across organs in the long-lived conditions. Module 11

was associated specifically with drug-induced longevity, module 14 was associated

specifically with diet-induced longevity, and module 4 was associated with both diet-

induced and drug-induced longevity. These results indicate that the coexpression of

module 4 with other modules may be associated generally with longevity by TOR

suppression, whilst modules 11 and 14 exhibit intervention-specific changes in coex-

pression.

Figure 4. Structure of longevity-associated changes in the transcriptional network. The heatmap shows the observed change in correlation (Spearman's rho) between pairs of modules in response to DR and in response to TOR suppression (Rapamycin), relative to the fully EAA-fed condition. Row and column labels represent transcriptional modules. Values indicate the difference in correlation between the given condition and the EAA-fed condition, and are given when the change was significant (p≤0.05) according to permutation testing.

To identify functions that are likely affected by changes to coexpression

across organs, we analysed enrichment of Gene Ontology (GO) terms within each

module (Supplementary Materials). Distinct transcriptional modules were associated

with non-overlapping functions, indicating that these modules represent functionally

distinct suites of genes, and therefore changes to their coexpression is likely physio-

logically relevant. Of the modules implicated in responses to DR, Module 4 was en-

riched in extracellular metabolic enzymes - particularly peptidases - and also lipases

and carbohydrases. Module 14 was enriched in transmembrane sugar transporters

and polysaccharide/carbohydrate binding. Module 11 was enriched in ATP-binding

cassette (ABC) transporter activity. Taken together, the transcriptional network and

GO analyses suggest that longevity is associated with altered orchestration of meta-

bolic functions across organs, and not changes to the same metabolic program in all

tissues.

Predicting transcription factors to regulate longevity-associated transcriptional

changes

We sought to identify candidate regulators of transcriptional changes associ-

ated with diet-induced longevity and drug-induced longevity. We predict that manipu-

lating transcription factors (TFs) that regulate DR/TOR-dependent transcriptional

regulation will mimic DR by correctly coordinating diverse transcriptional targets. We

identified transcription factor binding site motifs (TFBSs) associated with transcrip-

tional variation across organs in the long-lived conditions. This process associated

each transcriptional module with a potentially unique set of TFs, but clustering mod-

ules by these TF sets grouped modules 4, 11 and 14 (Figure 5a), which the transcrip-

tional network analyses had implicated in longevity. This clustering therefore sug-

gested that these longevity-associated modules share common upstream regulators.

Indeed, the sets of TFs identified in the longevity-associated modules overlapped

substantially (Figure 5b), and these shared motifs were also the most significantly

enriched; for each longevity-associated module, the most strongly enriched motifs

bind GATA family TFs (*GATAd, GATAd, grn, pnr, srp*) (Table II). Furthermore, the

most significant association observed in the entire analysis was between GATA-

binding motifs and Module 4, the only module associated with both diet-induced and

drug-induced longevity. Other motifs associated with all three longevity-associated

modules were annotated with *Bx* - which physically binds the GATA TF *pnr*; *Ham*,

which has roles in cell fate determination [26,27]; and *CG10348*, which has no known
regulatory function. However, the association with these latter three transcription fac-
tors was less significant than the association with GATA factors. Looking across all
modules, all but 2 and 5 were associated with some GATA motifs, however, the
strength of enrichment (Escore) for the GATA factors was significantly greater for the
longevity-associated modules 14 and 4 (Figure 5c). The final evidence of association
between GATA factors and longevity-associated modules was that Module 4 was not
only highly enriched in GATA binding sites, but also contained GATAe (Supplemen-
tary Files). Experimental studies have shown correlated expression of a TF and its
putative targets is a strong predictor of causal relationships [21,22]. Altogether, these
findings strongly implicated GATA factors in the tissue-specific transcriptional chang-
es observed under low TOR.
244

Table II. Annotations for top five-ranked motifs enriched in association with transcriptional modules. Unannotated motifs are excluded.

Longevity-associated	Module number	TF*	Escore[†]
Diet-induced & Drug-induced	4	grn srp pnr GATAd GATAe	11.58
		grn srp pnr GATAd GATAe	11.21
		grn srp pnr GATAd GATAe	10.07
		grn srp pnr GATAd GATAe	10.07
		grn srp pnr GATAd GATAe	10.07
Drug-induced	11	grn srp pnr GATAd GATAe	5.95
		grn srp pnr GATAd GATAe	5.88
		CG10348 ham	5.68
		grn srp pnr GATAd GATAe	5.56
		CG10348 ham	5.56
Diet-induced	14	grn GATAd pnr GATAe	10.63
		grn srp pnr GATAd GATAe	9.67
		grn GATAd pnr GATAe	9.05
		grn srp pnr GATAd GATAe	8.69
		grn GATAd pnr GATAe	8.60
No	1	Dref	8.50
		Dref	7.28
		Dref	5.66
		crp	4.78
		CG7928	4.47
No	2	lola	4.57
		lola	4.44
		Trl	4.43
		Trl	4.43
		Trl	4.42
No	3	grh	7.30
		grh	6.64
		GATAe srp pnr GATAd	6.24
		grn srp pnr GATAd GATAe	5.91
		grn srp pnr GATAd GATAe	5.91
No	5	CG9906 CG1924 Cnx99A	4.94
		jim	4.21
		nerfin-1 nerfin-2	4.01
		slp1 slp2 foxo fd19B	3.88
		Dref	3.87
No	6	vri	6.88
		gt	5.73
		Mef2	5.54
		Mef2	5.43
		vri	5.18
No	7	crp	5.08
		pnr	3.82
		HLH106	3.80
		CrebB-17A	3.61
		Hnf4	3.43
No	8	CG10348 ham	5.63
		CG10348 ham	5.44
		scrt	5.35
		scrt	5.00
		CG12605	4.91
No	9	crp	6.92
		Trl	5.44
		Trl	5.44
		Trl	4.96
		Trl CG33260	4.68
		Atf6	5.60
		grn GATAd pnr GATAe	4.37

No	10	CG11071	4.35
		grn GATAd pnr GATAe	4.34
		srp	4.31
No	12	crp	9.27
		CG6792	6.20
		kay Atf3	5.58
		Dref	4.91
		vri	4.74
No	13	HLH106	8.86
		HLH106	7.99
		GATAd	6.58
		srp	6.50
		GATAe	6.37

245 * Genes annotated as associated with the motif
246 † Escore for the motif
247

Figure 5. A signature of GATA transcription factors as regulators of the transcriptional response to DR and/or TOR. (a) Lifespan-associated transcriptional modules are grouped by sharing of transcription factor binding site motifs. Each module was tested for enrichment of *cis*-regulatory elements, and clustered by the

presence/absence of associated transcription factor binding site motifs. Dendrogram labels represent module numbers. This analysis clusters together transcriptional modules associated with longevity (4, 11, 14, also clustered with module 8). **(b)** Longevity-associated modules are associated with overlapping sets of transcription factor binding sites. Text below the Venn diagram names the transcription factors associated with binding sites enriched in all three longevity-associated modules. **(c)** Distribution per module of enrichment (Escore) of motifs annotated as binding GATA transcription factors. The box plots show medians (horizontal midline), 1st and 3rd quartiles (hinges), and range of data points. The notches equate to ~95% confidence intervals for the medians [(1.58 * interquartile range) / (square root n)]. The plot shows that enrichment of GATA binding sites is particularly strong for modules 4 and 14, both of which are associated with lifespan extension by DR. **(d)** GATA factor binding sites are enriched across multiple gene sets regulated by DR/TOR. Columns represent sets of genes according to tissue and the sign of expression change per treatment. Rows represent TFs associated with the sets of differentially expressed genes by enrichment of binding site motifs in the gene set. The axes are ordered by clustering the presence (blue) and absence (grey) of TF binding sites, using binary distance. Names of transcription factors occurring in a cluster across multiple gene sets have been magnified. **(e)** The bar plot shows the frequency of associations with longevity-associated transcriptional changes within organs, for the most frequently associated TFs. TFs associated with <10 gene sets are excluded.

248   To complement the analysis of transcriptional network regulators, we repeated

249 the same approach on sets of genes which showed changed regulation (differential

250 expression) within organs in one or both of the long-lived conditions (Table III). Con-

251 sistent with the absence of an organ-nonspecific transcriptional signature of DR/TOR,

252 most TFBSs identified by this analysis were associated with specific tissues. Howev-

253 er, one cluster of TFs was associated with multiple gene sets across organs, and that

254 cluster contained the GATA factors (Figure 5d). Across all gene sets, each GATA

255 factor was associated with at least 32% of diet/tissue-specific gene sets (Figure 5e).

256 GATA factors were therefore associated with DR/TOR-dependent transcriptional

257 changess within organs, recapitulating the parallel observations across organs.

Table III. Annotations for top five ranked motifs enriched in association with differentially expressed gene sets. Unannotated motifs are excluded.

Gene set				
Tissue	DR*	Rapamycin*	TFs**	Escore[†]
Whole fly	-	-	Tbp,Taf1	11.11
			E2f,E2f2,Dp	10.22
			Tbp	8.98
			E2f,E2f2,Dp	8.17
			E2f,E2f2,Dp	7.96
Whole fly	+	+	Cf2	6.25
			Mef2	5.41
			bin	5.40
			Cf2	5.37
			Cf2	5.22
Whole fly	-		E2f,E2f2,Dp	6.55
			Tbp,Taf1	6.44
			lola	6.27
			Blimp-1	5.98
			Blimp-1	5.98
Whole fly	+		Cf2	5.84
			Mef2	5.23
			Mef2	5.22
			Optix	5.19
			bin	5.07
Whole fly		-	Top2	4.32
			Dref	4.05
			Trl,CG33260	3.93
			Dref	3.92
			Dref	3.87
Whole fly		+	lola	8.29
			lola	8.27
			lola	7.61
			Optix	4.70
			bin	4.45
Brain	-	-	Tbp	8.40
			Tbp,Taf1	8.35
			grn,pnr	6.85
			pnr	6.21
			lola	6.13
Brain	-		pnr	11.03
			grn	10.04
			ham	10.01
			grn,pnr	9.95
			pnr	9.55
Brain	+		Jra	4.31
			Jra	4.29
			Atf3	4.02
			sna	3.77
			ato	3.68
Brain		-	Tbp,Taf1	9.58
			Tbp	7.63
			srp	5.33
			lola	5.02
			grn,pnr	4.76
			Tbp	8.07
			Tbp,Taf1	7.42

Tissue			Gene	Value
Brain		+	gem	4.89
			pnr	4.63
			tin	4.57
Fat body	-	-	Rel	8.75
			Rel	7.10
			Tbp	6.94
			pnr	5.50
			Tbp,Taf1	4.99
Fat body	+	+	hb	6.89
			hb	6.34
			sqz,rn,Lin29	5.90
			hb	5.83
			sqz,rn,Lin29	5.70
Fat body	-		pnr	5.45
			grn	5.34
			pnr	4.85
			grn	4.77
			GATAe	4.65
Fat body	+		Hsf	6.54
			Hsf	5.75
			Hsf	5.10
			ovo	4.77
			bs	4.49
Fat body		-	GATAd	6.61
			srp	5.77
			pnr	5.53
			GATAe	5.46
			Tbp	5.38
Fat body		+	lola	6.46
			grn	6.45
			lola	6.15
			grn	5.87
			pnr	5.62
Gut	-	-	cic	10.44
			Tbp	8.89
			Tor	7.88
			Tbp,Taf1	6.90
			Tbp	6.81
Gut	+	+	scro	13.34
			oc	12.11
			scro	10.44
			ttk	10.16
			vnd	9.38
Gut	-		gem	6.79
			grn	6.49
			pnr	6.37
			pnr	5.53
			grn,pnr	4.92
Gut	+		oc	8.97
			fru	7.82
			scro	7.61
			fru	7.51
			ken	7.28
Gut		-	zfh1	8.56
			Tbp,Taf1	8.19
			da	8.18
			Tbp	7.93

Tissue	Sign*	Sign*	Gene**	Escore†
			da	7.41
Gut		+	Hsf	4.58
			pnr	4.46
			ERR	4.11
			br	3.77
			grh	3.71
Ovary		-	E2f,E2f2,Dp	4.56
			Myb	4.55
			Myb	4.46
			vri	4.41
			vri	4.26
Ovary		+	twi	5.19
			HLH106	4.18
			CG13897	3.89
			bin	3.87
			foxo	3.81
Thorax	-	-	grn	9.63
			grn	8.02
			pnr	7.72
			pnr	7.71
			pnr	7.63
Thorax	+	+	zfh1	6.02
			sna	5.55
			sna	5.41
			ham	5.32
			Rfx	4.24
Thorax	-		Tbp	9.37
			grn,pnr	9.02
			Tbp,Taf1	8.47
			GATAd	7.43
			GATAe	7.07
Thorax	+		Tbp,Taf1	10.54
			Tbp	8.63
			lola	7.43
			grn	6.99
			Hnf4	6.86
Thorax		-	ttk	7.43
			crp	7.19
			ttk	6.93
			Fer1	6.46
			ase	6.34
Thorax		+	Rel	7.19
			Tbp,Taf1	7.02
			Tbp	6.95
			Rel	5.03
			Rfx	4.68

* sign of expression change, relative to EAA diet. The gene set is defined by only the effect of one treatment when only one symbol is given.

** Genes annotated as associated with the motif

† Escore for the motif

The TFBS enrichment analysis associated GATA factors both with transcrip-

tional changes within organs and altered functions of the organs as a collective

whole. This association was consistent with prior knowledge of the evolutionarily

conserved biology of these TFs. GATA factors play known roles in signalling amino

acid availability via TOR in evolutionarily diverse eukaryotes (e.g. yeast: [28]; and mos-

quitos: [29]), as well as being required for lifespan extension by some tissue-specific

genetic lesions in worms [20]. To test for evolutionarily conserved connections between

GATA factors and DR-responsive transcripts, we established the association be-

tween TFBSs and orthologs of DR-responsive genes across 12 *Drosophila* species.

These 12 *Drosophila* species span ~40 of million years of evolution, which due to

their rapid development time, equates to a greater evolutionary time than that for the

mammalian radiation. The method used for this analysis was independent of that

used for *D. melanogaster*, but also showed that the association between GATA mo-

tifs and orthologs of DR-responsive genes from *D. melanogaster* was robust across

Drosophila species (Supplementary Spreadsheets). Together, these data and previ-

ous findings strongly implicate GATA factors as evolutionarily-conserved regulators

of tissue-specific transcriptional responses to EAAs.

Functional roles for GATA factors in mediating dietary effects on lifespan

Of the five *D. melanogaster* GATA TFs, *GATAe* and *srp* were of particular inte-

rest for a role in longevity via DR/TOR. *GATAe* was associated more strongly than

any other TF with longevity-associated transcriptional changes (see above), and *srp*

has known roles in the fat body in regulating oogenesis via yolk proteins [30]. Conse-

quently we focussed on testing the roles of *srp* and *GATAe* in dietary regulation of

longevity. We knocked down *srp* and *GATAe* with tissue-specific RNAi, and evaluat-

ed whether these knockdowns altered the effects of dietary EAAs on egg laying and

lifespan. *GATAe* and *srp* were most highly expressed in the gut and fat body, respec-

tively (Figure S4), so we targeted *GATAe* in the midgut epithelium (with the TiGS

driver: [31]) and *srp* in the fat body (with the $S_1 106$ driver, which drives in both fat body

and gut [32], however fat body-specific expression of *srp* (Supplementary Materials)

suggests that RNAi phenotypes can be attributed solely to the fat body). These

GeneSwitch drivers [33] are activated by feeding flies the RU_{486} inducer (RU).

Figure 6. Tissue-restricted knockdown of *GATAe* and *srp* interact with dietary EAAs to determine egg laying and lifespan. Survival curves **(A & C)** and egg laying indices (overnight egg laying by vials of 10 flies; **B & D**) of flies expressing *GATAe*[RNAi] in the gut (A, B), or *srp*[RNAi] in the gut and fat body.

In *GATAe^RNAi* flies there was a significant interaction between the effect of feeding the transgene-inducing drug and EAA on the lifespan of GATAe knockdown flies (Cox Proportional Hazards Regression, $P < 0.005$). Specifically, knocking down *GATAe* in the gut accelerated the onset of mortality in early life but, surprisingly, this lifespan shortening was partially rescued by feeding dietary EAAs (Figure 6a). However GATAe^RNAi affected neither egg laying nor altered the effect of EAAs on egg laying (Figure 6b), indicating that the regulation of egg laying is independent of the lifespan-limiting process induced by knocking down GATAe in the gut.

Figure 7. Ubiquitous knockdown of *srp* reverses lifespan-shortening effects of EAAs and desensitises egg laying to EAAs. (A) Survival curves and **(B)** egg laying indices of flies ubiquitously expressing RNAi against *srp*.

In contrast to *GATAe* knockdown, *srp* knockdown had positive effects on lon-

gevity (Figure 6c). Fat body-specific expression of *srp*[RNAi] protected flies from the

lifespan-shortening effects of EAAs (log-rank test, p<0.001), but there was no effect

on lifespan in the absence of EAAs (p=0.57). Egg laying was not affected by srp

knockdown (p=0.08). Therefore, knocking down *srp* in the fat body insulates lifespan

against the pernicious effects of EAAs, but does not preclude flies from the early-life

benefits of EAA feeding for egg laying. Together, the GATAe[RNAi] and srp[RNAi] experi-

ments validate our bioinformatic analysis, by linking age-dependent physiology with

interactions between dietary EAAs and the tissue-specific regulation of GATA factors.

Our expression data showed that *srp* is expressed most strongly in the fat

body, but is also expressed in other tissues e.g. the ovary. Therefore, we asked

whether systemic *srp* knockdown using a *Daughterless* GeneSwitch driver further al-

tered fly phenotype. Ubiquitous knockdown of *srp* was sufficient to neutralise the

lifespan-shortening effects of EAAs, restoring lifespan to that of flies fed DR food

(Figure 7a). Furthermore, ubiquitous *srp* RNAi extended lifespan of flies on DR food,

meaning that transgene expression resulted in a median 7 day lifespan extension

whether or not the flies were fed high levels of dietary EAAs. Whilst systemic *srp*

knockdown extended lifespan more strongly than the tissue-specific knockdown, the

tradeoff with egg laying was much greater (Figure 7b). Without induction of *srp*[RNAi],

the flies laid ~25% more eggs when EAAs were supplemented to the food, but *srp*

knockdown arrested egg laying (ANOVA, $p < 2.2^{-16}$) and rendered it insensitive to

EAA feeding (EAA * RU interaction, p<0.005). The egg laying phenotype thus con-

trasted the lifespan phenotype, since egg laying was entirely desensitised to diet,

whereas median lifespan remained sensitive. Together, these data indicate that the

relationship between lifespan and egg laying is set by an interaction of dietary EAAs

and *srp*, and the integration of egg laying and lifespan depends on the tissues in

which this interaction occurs.

Discussion

Dietary restriction improves lifelong health and extends lifespan in a range of

organisms, from yeast to mammals. A growing body of evidence shows the particular

importance of dietary nutrient balance for ageing, particularly lowered pro-
tein:carbohydrate ratio [6,34], which implicates amino-acid sensitive TOR signalling [11].
This implicit role of TOR was recently validated in *Drosophila*, by the demonstration
that active TOR is required for EAA enrichment to shorten lifespan [10]. These results
have the important implication that the lifespan effects of EAA restriction are not ac-
counted for solely by metabolic effects, but also by cellular signalling. Using the same
experimental conditions, our present study predicts lifespan-relevant and EAA-
dependent signalling mechanisms, by showing that EAA restriction has transcription-
al effects which are congruent - both within and between organs - with those of
pharmacological TOR suppression, indicating that the transcriptional effects of DR
are indeed mediated in large part by TOR. Furthermore, these overlapping transcrip-
tional changes are also associated with overlapping sets of *cis* regulatory elements.
Thus, DR and rapamycin have overlapping effects on lifespan, on transcription within
organs, on the coordination of functions across organs, and their transcriptional tar-
gets share predicted regulators.

Our results connect lifespan regulation by DR and low TOR with GATA tran-
scription factors. The GATA factor knockdown experiments validate the bioinformatic
connections between dietary EAAs, transcription factors and lifelong physiology.
Specifically, these experiments show that genetic knockdown of either *GATAe* or *srp*
substantially alters the effect of diet on fly survival and, in the case of srp, knockdown
both extended lifespan and dictated the influence of diet on egg laying, thus pheno-
copying flies treated with rapamycin. Curiously, *GATAe* knockdown in the gut
changed the sign of the effect of EAAs on lifespan, suggesting either a different opti-
mal dietary balance in these flies, or that elevated EAAs rescued the pathological ef-
fects of GATA knockdown. By contrast, *srp* knockdown with two different drivers can-
celled the lifespan-shortening effects of EAA enrichment. Furthermore the GATA
knockdown experiments show that, whilst lifespan and egg laying normally show
coupled responses to diet, they can be decoupled by genetic manipulation of GATA
factors, depending on which tissues are targeted. The phenotypic differences be-
tween flies in which srp is locally or systemically knocked down provide key evidence
that tissue-targeted manipulation of TFs associated with DR can recapitulate the
lifespan benefits of DR, and that the associated costs depend on the tissue-
specificity of the intervention. This supports the notion that ageing can be ameliorated
by specifically targeting pathologies in lifespan-limiting tissues. Whilst there is some

existing evidence of a role for GATA factors in lifespan regulation [20], to our knowledge they have not yet been associated explicitly with DR, nor has there been a prior demonstration of any transcriptional intervention blocking the phenotypic effects of EAAs. The indication that costs and benefits of dietary alteration are mediated by distinct tissues has the important biomedical implication that lifelong health may be improved by altering the coordination of function across multiple tissues, and the evolutionary implication that phenotypic tradeoffs evolve due to coupled functions of tissues.

The GATA factors are an ancient family of transcription factors, which have well-characterised and essential roles in coordinating development and growth in organisms from yeast to mice. In multicellular differentiated organisms, GATA factors are required in the development of multiple tissue types, which in *Drosophila* includes the heart [35], fat body [36] and gut [37]. GATA factors are also expressed tissue-specifically in adulthood (Figure S4), but their roles in coordinating adult functions are not well characterised. In *Drosophila*, recent studies have shown a role of *GATAe* in maintenance of cell identity and tissue function in the adult gut, but it is not yet clear from which molecular pathways *GATAe* integrates information [38–40]. One of the better-described roles for GATA factors in adult animals is in nutrient regulation of oogenesis in mosquitos [29,41–43]. In *Aedes aegypti*, oogenesis requires a blood meal, which contains the mosquito's only source of protein. Egg production is suppressed before feeding, due in part to GATA-mediated repression in the fat body of the major yolk precursor protein gene *Vg* [44]. After the blood meal, *Vg* expression is activated by TOR enhancing expression of the transcriptional activator *AaGATAa* [43]. In *Drosophila*, there is evidence that this regulatory circuit may be conserved, since *srp* regulates expression of yolk proteins [30]. Our new and recent data show that rapamycin abrogates higher egg laying caused by EAA enrichment [10], corresponding to changes in the fat body in expression of yolk proteins (Supplementary Materials). The connection in *Drosophila* between dietary nitrogen and GATA transcriptional control is consistent with mechanisms in the yeast *Saccharomyces cerevisiae*, in which selective amino acid catabolism is controlled by a circuit known as Nitrogen Catabolite Repression. In this system, when the available nitrogen sources only support poor growth, TOR-dependent nuclear localisation of a GATA transcription factor triggers the expression of genes involved in the transport and metabolism of less-preferred nitrogen sources

[28]. Together, these data point to highly evolutionarily conserved connections between

protein uptake, growth and reproduction, TOR signaling and transcriptional control by

GATA factors. Theory suggests that ageing results from antagonistic pleiotropic ef-

fects of mechanisms that promote growth and reproduction in the young [45]. GATA

factors fit these criteria, as nutrient-responsive regulators of growth that we associate

with molecular responses to lifespan-extending regimes. Due to the evolutionarily

conserved connections between dietary nitrogen and lifespan on one hand, and die-

tary nitrogen and GATA factor-regulated transcription on the other, we anticipate new

interest in the role that GATA factors play in lifespan regulation.

Our experiments demonstrate that the tissue-restricted knockdown of GATA

factors is sufficient to modify the lifespan-shortening effect of EAAs. Previous studies

have shown that tissue-targeted interventions to lower insulin signalling are sufficient

to extend lifespan in worms and flies [46], and GATA factors are required for some

such effects. It is well-established that the balance of dietary nutrients has the evolu-

tionarily conserved capacity to determine lifelong health [11], and to recognise this bal-

anced supply, insulin and TOR signalling must coordinate. Reducing either insulin or

TOR signalling is sufficient to extend lifespan, suggesting that the lifespan-extending

effects of these two interventions may be mediated by the nexus of their signalling

effects. In *C. elegans*, the GATA factor *ELT-2* is required for longevity following die-

tary restriction or mutation of the insulin receptor [20], and GATA factor overexpression

extends lifespan [47]. Additionally, previous transcriptional studies in both flies and

worms have uncovered enrichment of GATA motifs in the insulin regulon [48,49]. To-

gether with our data, these results suggest that signals from TOR and other nutrient

sensing pathways are mediated at least in part by GATA factors. In light of these da-

ta, our demonstration that transcriptional effects of diet are largely tissue-specific

suggests that TOR and the GATA factors mediate cell-autonomous interpretations of

global signals (e.g. insulins or bioamines), into a local language that dictates physio-

logical change appropriate to the tissue in question.

Transcription factors known to be downstream of TOR signalling and insulin

signalling are candidates for interpretation by GATA factors in *Drosophila. REPTOR*

and *REPTOR-BP* have recently been discovered by Tiebe and colleagues as a novel

TOR-dependent transcription factor complex [18]. Given the complementary nature of

Tiebe *et al*'s cell culture experiments and our study of dissected tissues, an emergent

hypothesis is that GATA factors determine organ-specific responses to TOR by modi-

fying effects of systemic *REPTOR* signalling. Downstream of insulin signalling, mo-

lecular interactions between *FoxO* and TOR signalling are already established: TOR

lies in a network of interacting signaling pathways including insulin [50], and *FoxO* is

required for reduced insulin signalling to extend lifespan [51]. However, FoxO binds

TOR's promoter and is required for normal *TOR* expression, and GATA motifs are

associated with genes that are not bound by *FoxO* but show *FoxO*-dependent regula-

tion by IIS [49], suggesting a regulatory circuit from IIS to GATA factors via *FoxO* and

TOR. Given that *FoxO* is not required for the lifespan-extending effects of DR [52] and

that different insulin-like peptides are produced in response to different nutritional

stimuli [53], we suggest that signals from IIS via FoxO may be modified in peripheral

tissues by TOR-modulated GATA activity, which shapes tissue-specific responses to

diet that ultimately modify physiology and longevity (Figure 8).

Figure 8. Predicted molecular integration of GATA factors in nutrient-sensing and lifespan-relevant transcriptional networks in *Drosophila*. Our transcriptomic analysis associates transcriptional effects of EAA restriction with TOR, and the TOR-dependent EAA regulon with GATA factors. Previous studies have shown that (1) TOR directly represses *REPTOR*, (2) The insulin pathway bifurcates via Pi3K and Ras to regulate lifespan via *FOXO* and *Aop*, and (3) GATA binding sites are associated with genes that are differentially expressed in *FOXO* mutants but not bound by *FOXO*, that *FOXO* both binds *TOR*'s promoter and *TOR* transcription is reduced in *FOXO* mutants. Together, these results suggest that *TOR* integrates signals from IIS/Pi3K signalling via *FOXO*, that regulons of DR, TOR and *FOXO* are all enriched in GATA factor binding sites, and therefore that GATA factors mediate transcriptional effects of both EAAs and *FOXO* via TOR.

The gut appears to be a particularly effective target for interventions to extend
lifespan [46]. The adult gut has a critical role in the ongoing health of organisms, bal-
ancing the passage of nutrients whilst resisting environmental stresses [54]. In flies,
there are complex relationships between age, gut maintenance, diet, metabolism,
resident microbiota and expression of antimicrobial genes [55-62]. Recent work sug-
gests a central role for the gut in mediating lifespan extension by DR in female flies,
revealing sexually dimorphic pathologies in aged female guts [63], dependent on cell-
autonomous expression of sex determination pathways [64]. Failure of gut integrity is
associated with changes in microbiota and greater antimicrobial peptide expression,
and appears to be a marker of imminent death [58,65]. In worms, the GATA factor ELT-2
interacts with p38 transcriptional regulators to modify adult gut immunity [66], whilst
GATA factors are required for normal gut development and maintenance in *Drosophi-*
la and mice [38,67]. It is thus tempting to speculate that increased risk of death in late
life may be preciptated by a loss of gut integrity enhancing exposure to environmental
microbes and toxins. Our results are directly relevant to these issues, because coor-
dination across organs of a transcriptional module (module 4) that was perturbed un-
der long-lived conditions strongly corresponded to *GATAe* expression, and *GATAe*
and paracrine insulin signalling have each been shown to be required for intestinal
stem cell proliferation [38,68]. Our lifespan data are consistent with *GATAe* knockdown
in the gut increasing mortality, but since this effect appears to be stochastic and not
necessarily connected to age, the role of *GATAe* regulation in late-life mortality re-
mains to be established. However, fly lifespan is not necessarily extended by en-
hancing expression of components of the immune system [69], antibiotic treatment [55] or
genetically reducing gut dysplasia [59]. Thus, although compromised gut integrity is
clearly a marker of frailty in late life, it may not be obligately linked to death.

This study reveals how the fly's tissues interact as a system, and how that
system responds to EAA dilution and low TOR. The results showed that the effects of
EAA restriction on transcription are likely mediated by TOR via GATA transcription
factors. Entirely consistent with this prediction, genetic analysis confirmed that the
tissue-specific activity of GATA factors dictate the effect of dietary nutrient balance on
phenotype. Importantly, these experiments also suggest that the costs and benefits
of dietary variation may be mediated by different tissues, and therefore that benefits
may be reaped without fitness tradeoffs by targeting specific tissues. The evolution-
ary conservation of GATA factors, of their connection to regulating amino acid me-

tabolism, and of the capacity of TOR to mediate lifespan extension, suggests that
GATA factors may be relevant to ameliorating ageing by DR in a broad range of or-
ganisms, including humans.

Materials & methods

Drosophila melanogaster and diets were prepared according to (10). 1SY me-
dium is a dietary restriction (DR: diet-induced longevity) regime that extends lifespan
in wild-type and laboratory-maintained flies [71,72], containing 100 g/l autolysed yeast
(MP Biomedicals, OH, USA), 50 g/l sucrose (Tate & Lyle, London, UK), 15 g/l agar
(Sigma-Aldrich, Dorset, UK), 30 ml/l nipagin (Chemlink Specialities, Dorset, UK), and
3 ml/l propionic acid (Sigma-Aldrich, Dorset, UK). EAA food comprised 1SY with the
addition of cocktail of EAAs dissolved in pH 4.5 water (final concentrations in fly me-
dia: L-arginine 0.43 g/l, L-histidine 0.21 g/l, L-isoleucine 0.34 g/l, L-leucine 0.48 g/l, L-
lysine 0.52 g/l, L-methionine 0.1 g/l, L-phenylalanine 0.26 g/l, L-threonine 0.37 g/l, L-
tryptophan 0.09 g/l, L-valine 0.4 g/l: all suppled by Sigma). EAA+rapamycin (drug-
induced longevity) food consisted of EAA food with the addition of rapamycin (LC la-
boratories, MA, USA) dissolved in ethanol, to a final concentration of 200 µM in the
diet. For RNAi experiments, RU_{486} (Sigma M8046) dissolved in ethanol was added to
1SY or EAA food to a final concentration of 200 µm. Vehicle controls were added to
media as appropriate.

Outbred wild-type Dahomey flies bearing the endosymbiont *Wolbachia* were
cultured on a 12:12 light cycle at $25^{o}C$ and 60% humidity. For RNAi experiments, the
TiGS and S_1106 drivers and RNAi constructs (UAS-srp[RNAi]: Vienna Stock Center
#33748; UAS-GATAe[TRiP]: Bloomington Stock Center #33748) were backcrossed into
flies bearing the *w* mutation for at least 6 generations (UAS-GATAe[TRiP] was back-
crossed by PCR), and maintained at large population sizes to maintain outbred ge-
netic diversity. For all experiments, parents of experimental flies oviposited onto
grape juice agar for 18h. Eggs were washed from this agar, added to 1SY and cul-
tured to adulthood at standardised density. Newly emerged flies were allowed to ma-
te *ad libitum* for 48h before being lightly anaesthetised with CO_2. Males were re-
moved, and female flies were allocated to experimental diets. For RNA sequencing,
females were maintained on 1SY, EAA or EAA+rapamycin for six days before dissec-

tion. The RNA sequencing experiment was independently replicated three times,

generating three samples per organ or per whole fly, per diet (with the exception of

the gut on 1SY: see Supplementary Text). For egg laying and lifespan experiments,

survival was scored three times per week. Egg laying was scored on day 10 after

eclosion, after 18h egg laying. Fecundity indices were calculated as number of eggs

per female (n=10 females per vial).

RNA was collected 6-10h into the flies' light cycle. To prepare RNA for se-

quencing, whole flies were flash-frozen. Brains, abdominal fat bodies, ovaries, guts

and thoraces were micro-dissected in ice-cold RNAlater solution and frozen at -80°C.

RNA was extracted using the QIAGEN total RNA isolation kit and quantified on an

Agilent 2100 bioanalyser. Sequencing was performed by the high throughput ge-

nomics services center at the Huntsman Cancer Institute (University of Utah). Sam-

ple concentration and purity of RNA was measured on a NanoDrop spectrophotome-

ter, and RNA integrity was assessed on an Agilent 2200 TapeStation. Illumina

TruSeq libraries were prepared from this RNA with the Illumina TruSeq Stranded

mRNA Sample Prep kit and sequenced on an Illumina HiSeq2000 101 v3 platform

using paired-end sequencing.

Reads were aligned to the *D. melanogaster* genome annotation 5.57 using

TopHat2 2.0.14 and counted using HTSeq 0.5.4p3 [73,74]. Non-protein coding genes

were retained. Unmapped reads were discarded. Enumerated reads were then ana-

lysed in R (3.0 & 3.1) using BioConductor. RPKM was calculated from read counts

generated by HTSeq, using the EdgeR library.

Differential expression across the three experimental conditions was deter-

mined with a negative binomial GLM fitted by DESeq2 (1.8.1, [75]), without rejection

based on Cook's distance, calculating P-values with a two-sided Wald test, and cal-

culating false discovery rate by Storey's method. Intersections between gene sets

and enriched GO terms were visualised with the upset package in R [76]. GO term en-

richent was analysed using GOrilla (http://cbl-gorilla.cs.technion.ac.il/).

Unsigned gene coexpression networks were determined using data from all

organs, excluding the whole-fly samples, using the WGCNA package in R [23]. Con-

sensus modules were determined automatically using the *blockwiseConsensusMod-*

ules function with default settings and a power of 26, stipulating a minimum module

size of 50 genes. Eigengene determination, variance explained by Eigengenes and

clustering were performed using internal WGCNA functions, as per the package tuto-

rial

(http://labs.genetics.ucla.edu/horvath/CoexpressionNetwork/Rpackages/WGCNA/Tut

orials/index.html). Changes in between-module correlations by experimental condi-

tion were calculated by custom R functions: correlation matrices (Spearman's rho) for

module Eigengenes were calculated in each experimental condition, and changes

were calculated by subtracting the DR matrix and the EAA+rapamycin matrix from

the EAA matrix, to calculate observed changes in correlations between pairs of

Eigengenes under DR and under rapamycin administration, respectively. To generate

null distributions for changes in correlations for each pair of modules in each of the

two long-lived conditions, the same procedure was repeated 10,000 times, permuting

each Eigengene and calculating changes in correlation for each pair. Observed

changes were considered significant when they did not fall between the 2.5^{th} and

97.5^{th} percentiles of their respective null distribution.

The most likely acyclic network between modules was determined by Additive

Bayesian Network analysis of module Eigengenes using the R package ABN with

10000 iterations. This approach determines the likelihood of inter-dependencies be-

tween variables by randomly simulating data and comparing to the observed inter-

dependencies. ABN found a consensus structure to the data after only ~1000 itera-

tions, indicating that the structure is robust to further simulation. The ABN was plotted

in Cytoscape, using hierarchical network ordering. Meta-module analysis and

Eigengene perturbation analysis were performed according to [25].

Enrichment of cis-regulatory motifs was analysed using i-Cis target [77], exclud-

ing genes for which DEseq2 models did not converge. Unannotated CRMs were ex-

cluded from further analysis. Samples were hierarchically clustered according to the

presence/absence of transcription factor binding sites with the R hclust function, us-

ing a binary distance metric. Notches on boxplots, approximating 95% confidence

intervals of medians were produced using the "notch" argument to the R boxplot

function. Heatmaps of TF:gene-set associations were plotted using the heatmap.2

function from the R gplots package, and ordered by heirarchical clustering using bi-

nary distance.

Analysis of evolutionary conservation of the association between TF binding

motifs and differentially expressed genes was conducted per tissue. Groups of differ-

entially expressed genes (according to DEseq2 analysis) showing the same signs of

fold-change in response to EAA restriction were scanned for significantly over-

represented motifs in their promoter regions. 706 motifs were included in the search,

from OnTheFly (2014 release), flyreg v2, dmmpmm2009 and idmmpmm2009 data-

bases, all provided by MEME 4.10 (http://meme-suite.org). Pscan version 1.2.2 was

used to score the motifs for the gene list, using the sets of genes that either didn't

show any diet-induced differential expression in any tissue, or were not expressed,

as a 'null hypothesis' background. For each gene list, the 706 scores (one per motif)

produced by pscan were normalized to z scores, and P values were computed, cor-

rected for multiple testing (Benjamini-Hochberg).

The same procedure was repeated for the evolutionarily conserved regulation

analysis, except that the genes considered were obtained from the 12 *Drosophila*

species (*D.melanogaster* and *D.ananassae*, *D.erecta*, *D.grimshawi*, *D.mojavensis*,

D.persimilis, *D.pseudoobscura*, *D.sechellia*, *D.simulans*, *D.virilis*, *D.willistoni* &

D.yakuba) whose genomic sequences are part of FlyBase, taking only 1:1 orthologs

of genes which DESeq2 analysis had shown to be differentially expressed in *D. mel-*

anogaster. Gene selection was done in two steps: for all gene groupings (differential

expression and background), genes for which OrthoDB version 7 didn't predict a one-

to-one ortholog in all other 11 species were removed. Then, each grouping was ex-

panded with all the relevant orthologs predicted by OrthoDB, such that each gene

group contained 12 times as many genes at after the first step.

Survival data were analysed in Microsoft Excel for log-rank statistics, or in R

for Cox Proportional Hazards analysis, using the CoxPH function from the Survival

package. Fecundity indices were analysed with ANOVAs fitted with the lm function in

R.

Bibliography

1. Niccoli T, Partridge L. Ageing as a risk factor for disease. *Curr Biol.* **22,** R741–
752 (2012)

2. Piper MD, Partridge L, Raubenheimer D, Simpson SJ. Dietary Restriction and
Aging: A Unifying Perspective. *Cell Metab.* **14 PAGE**(2011).

3. McCay CM, Crowell MF, Marnard LA. The effect of retarded growth upon the
length of life span and upon the ultimate body size. *J Nutr.* **10**, 63-59 (1935).

4. Mair W, Piper MD, Partridge L. Calories Do Not Explain Extension of Life Span
by Dietary Restriction in Drosophila. *PLoS Bio.* **3**, e223 (2005).

5. Lee K, Simpson SJ, Clissold FJ, Brooks R, Ballard WJ, Taylor PW, et al.
Lifespan and reproduction in Drosophila: New insights from nutritional geome-
try. *Proc Natl Acad Sci.* **7**, 2498–2503 (2008).

6. Skorupa DA, Dervisefendic A, Zwiener J, Pletcher SD. Dietary composition
specifies consumption, obesity, and lifespan in Drosophila melanogaster. *Ag-*
ing Cell **7**, 478–490 (2008).

7. Grandison RC, Piper MD, Partridge L. Amino-acid imbalance explains exten-
sion of lifespan by dietary restriction in Drosophila. *Nature* **462**, 1061–1064
(2009)

8. Solon-Biet SM, Aisling CM, Ballard WJ, Ruohonen K, Wu LE, Cogger VC, et
al. The Ratio of Macronutrients, Not Caloric Intake, Dictates Cardiometabolic
Health, Aging, and Longevity in Ad Libitum-Fed Mice. *Cell Metab.* **19**, 418–430
(2014).

9. Solon-Biet SM, Mitchell SJ, Coogan SC, Cogger VC, Gokarn R, C M Aisling, et
al. Dietary Protein to Carbohydrate Ratio and Caloric Restriction: Comparing
Metabolic Outcomes in Mice. *Cell Reports* **11**, 1529-1534 (2015).

10. Emran S, Yang M, He X, Zandveld J, Piper MD. Target of rapamycin signalling
mediates the lifespan-extending effects of dietary restriction by essential ami-
no acid alteration. *Aging (Albany)* **6**, 390–8 (2014).

11. Simpson SJ, Couteur DG, Raubenheimer D. Putting the Balance Back in Diet.
Cell **161**, 1823 (2015).

12. Bar-Peled L, Sabatini D. Regulation of mTORC1 by amino acids. *Trends Cell*
Biol. **24**, 400-406 (2014)

13. Taylor RC, Dillin A. XBP-1 is a cell-nonautonoous regulator of stress re-
sistance and longevity. *Cell* **153**, 1435–1447 (2013).

14. Katewa S, Kapahi P. Role of TOR signaling in aging and related biological
processes in Drosophila melanogaster. *Exp Gerontol.* **46**, 382-390 (2011).

15. Vilchez D, Simic M, Dillin A. Proteostasis and aging of stem cells. *Trends Cell*
Biol. **24**, 161-171 (2014).

16. Bülow MH, Aebersold R, Pankratz MJ, Jünger MA. The Drosophila FoxA
Ortholog Fork Head Regulates Growth and Gene Expression Downstream of
Target of Rapamycin. *PLoS ONE* **5**, e15171 (2010).

17. Robida-Stubbs S, Glover-Cutter K, Lamming D, Mizunuma M, Narasimhan S,
Neumann-Haefelin E, et al. TOR signaling and rapamycin influence longevity
by regulating SKN-1/Nrf and DAF-16/FoxO. *Cell Metab.* **15**, 713-724 (2012).

18. Tiebe M, Lutz M, Garza A, Buechling T, Boutros M, Teleman AA. REPTOR
and REPTOR-BP Regulate Organismal Metabolism and Transcription Down-
stream of TORC1. *Dev Cell* **33**, 272–84 (2015).

19. Alic N, Tullet JM, Niccoli T, Broughton S, Hoddinott MP, Slack C, et al. Cell-nonautonomous effects of dFOXO/DAF-16 in aging. *Cell Rep.* **6**, 608–616 (2014).

20. Zhang P, Judy M, Lee S-J, Kenyon C. Direct and Indirect Gene Regulation by a Life-Extending FOXO Protein in C. elegans: Roles for GATA Factors and Lipid Gene Regulators. *Cell Metab.* **17**, 85–100 (2013).

21. Potier D, Davie K, Hulselmans G, Sanchez M, Haagen L, Vân H-T, et al. Mapping Gene Regulatory Networks in Drosophila Eye Development by Large-Scale Transcriptome Perturbations and Motif Inference. *Cell Reports* **9**, 2290-2303 (2014).

22. Dutta D, Dobson AJ, Houtz PL, Gläßer C, Revah J, Korzelius J, et al. Regional Cell-Specific Transcriptome Mapping Reveals Regulatory Complexity in the Adult Drosophila Midgut. *Cell Reports* **12**, 346–358 (2015).

23. Langfelder P, Horvath S. WGCNA: an R package for weighted correlation network analysis. BMC *Bioinformatics.* **9**, 559-PAGE (2008).

24. Mccormick M, Promislow D. Networks in the Biology of Aging: Powerful Tools for a Complex Process. *Annu Rev Gerontol Geriat.* **34**, 243-266 (2014).

25. Langfelder P, Horvath S. Eigengene networks for studying the relationships between co-expression modules. *Bmc Syst Biol.* **1**, 54 PAGE (2007).

26. Moore A, Jan L, Jan Y. Hamlet, a binary genetic switch between single-and multiple-dendrite neuron morphology. *Science.* **297**, 1355-1358 (2002).

27. Moore AW, Roegiers F, Jan LY, Jan Y-NN. Conversion of neurons and glia to external-cell fates in the external sensory organs of Drosophila hamlet mutants by a cousin-cousin cell-type respecification. *Genes Dev.* **18**, 623–628 (2004).

28. Cooper T. Transmitting the signal of excess nitrogen in Saccharomyces cerevisiae from the Tor proteins to the GATA factors: connecting the dots. *FEMS Microbiol Rev.* **26**, 223-238 (2002).

29. Attardo GM, Higgs S, Klingler KA, Vanlandingham DL, Raikhel AS. RNA interference-mediated knockdown of a GATA factor reveals a link to anautogeny in the mosquito Aedes aegypti. *Proc Natl Acad Sci.* **100**, 13374–13379 (2003).

30. Lossky M, Wensink P. Regulation of Drosophila yolk protein genes by an ovary-specific GATA factor. *Mol Cell Biol.* **15**, 6943–6952 (1995).

31. Alic N, Giannakou ME, Papatheodorou I, Hoddinott MP, Andrews DT, Bolukbasi E, et al. Interplay of dFOXO and Two ETS-Family Transcription Factors Determines Lifespan in Drosophila melanogaster. *Plos Genet.* **10**, e1004619 (2014).

32. Giannakou ME, Goss M, Jünger MA, Hafen E, Leevers SJ, Partridge L. Long-Lived Drosophila with Overexpressed dFOXO in Adult Fat Body. *Science* **305**, 361 PAGE (2004).

33. Poirier L, Shane A, Zheng J, Seroude L. Characterization of the Drosophila
gene-switch system in aging studies: a cautionary tale. *Aging Cell.* **7**, 758–770
(2008).

34. Couteur DG, Samantha S-B, Cogger VC, Mitchell SJ, Senior A, de Cabo R, et
al. The impact of low-protein high-carbohydrate diets on aging and lifespan.
Cell Mol Life Sci **73**, 1237–1252 (2016).

35. Sorrentino RP, Gajewski KM, Schulz RA. GATA factors in Drosophila heart
and blood cell development. *Semin Cell Dev Biol.* **16**, 107–116 (2005).

36. Sam S, Leise W, Hoshizaki D. The serpent gene is necessary for progression
through the early stages of fat-body development. *Mech Dev.* **60**, 197–205
(1996).

37. Murakami R, Okumura T, Uchiyama H. GATA factors as key regulatory mole-
cules in the development of Drosophila endoderm. *Dev Growth Differ.* **47**,
581–589 (2005).

38. Buchon N, Osman D, David F, Fang H, Boquete J-P, Deplancke B, et al. Mor-
phological and Molecular Characterization of Adult Midgut Compartmentaliza-
tion in Drosophila. *Cell Reports* **3**, 1725-1738 (2013).

39. Okumura T, Takeda K, Kuchiki M, Akaishi M. GATAe regulates intestinal stem
cell maintenance and differentiation in Drosophila adult midgut. *Dev Biol.* **410**,
24-35 (2016).

40. Hansen IA, Attardo GM, Park J-H, Peng Q, Raikhel AS. Target of rapamycin-
mediated amino acid signaling in mosquito anautogeny. *Proc Natl Acad Sci.*
101, 10626–10631 (2004).

41. Hansen IA, Attardo GM, Roy SG, Raikhel AS. Target of Rapamycin-dependent
Activation of S6 Kinase Is a Central Step in the Transduction of Nutritional
Signals during Egg Development in a Mosquito. *J Biol Chem.* **280**, 20565–
20572 (2005).

42. Park J-H, Attardo GM, Hansen IA, Raikhel AS. GATA Factor Translation Is the
Final Downstream Step in the Amino Acid/Target-of-Rapamycin-mediated Vi-
tellogenin Gene Expression in the Anautogenous Mosquito Aedes aegypti. *J.*
Biol. Chem. **281**, 11167–11176 (2006).

43. Martín D, Piulachs M, Raikhel A. A novel GATA factor transcriptionally re-
presses yolk protein precursor genes in the mosquito Aedes aegypti via inter-
action with the CtBP corepressor. *Mol Cell Biol.* **21**, 164–174 (2001).

44. Williams GC. Pleiotropy, natural selection, and the evolution of senescence.
Evolution **11**, 398–411 (1957).

45. Rera M, Azizi M, Walker D. Organ-specific mediation of lifespan extension:
More than a gut feeling? *Ageing Res Rev* **12**, 436–444 (2013).

46. Budovskaya YV, Wu K, Southworth LK, Jiang M, Tedesco P, Johnson TE, et
al. An elt-3/elt-5/elt-6 GATA Transcription Circuit Guides Aging in C. elegans.
Cell., **134**, 1–13 (2007).

47. Murphy CT, A M Steven, Bargmann CI, Fraser A, Kamath RS, Ahringer J, et
al. Genes that act downstream of DAF-16 to influence the lifespan of Caeno-
rhabditis elegans. *Nature.* **424**, 277–284 (2003).

48. Alic N, Andrews DT, Giannakou ME, Papatheodorou I, Slack C, Hoddinott MP,
et al. Genome wide dFOXO targets and topology of the transcriptomic re-
sponse to stress and insulin signalling. *Mol Syst Biol.* **7**, 502 PAGE (2011).

49. Teleman AA. miR-200 De-FOGs Insulin Signaling. *Cell Metab.* **11**, 8–9 (2010).

50. Slack C, Giannakou ME, Foley A, Goss M, Partridge L. dFOXO-independent
effects of reduced insulin-like signaling in Drosophila. *Aging Cell* **10**, 735–748
(2011).

51. Min K, Yamamoto R, Buch S, Pankratz M, Tatar M. Drosophila lifespan control
by dietary restriction independent of insulin-like signaling. *Aging Cell.* **7**, 199-
206 (2008).

52. Kim J, Neufeld T. Dietary sugar promotes systemic TOR activation in Dro-
sophila through AKH-dependent selective secretion of Dilp3. *Nat Comms* **6**,
6846 PAGE (2015).

53. Buchon N, Osman D. All for one and one for all: Regionalization of the Dro-
sophila intestine. *Insect Biochem Mol Biol.* **67**:2-8 (2015).

54. Ren C, Webster P, Finkel SE, Tower J. Increased internal and external bacte-
rial load during Drosophila aging without life-span trade-off. JOURNAL **6**, 144–
152 (2007).

55. Biteau B, Karpac J, Supoyo S, Matthew D, Lehmann R, Jasper H. Lifespan
Extension by Preserving Proliferative Homeostasis in Drosophila. *PLoS Genet.*
6, e1001159 (2010).

56. Rera M, Bahadorani S, Cho J, Koehler C, Ulgherait M. Modulation of longevity
and tissue homeostasis by the Drosophila PGC-1 homolog. *Cell Metab.* **14,**
623–634 (2011).

57. Rera M, Clark R, Walker D. Intestinal barrier dysfunction links metabolic and
inflammatory markers of aging to death in Drosophila. *Proc Natl Acad Sci.*
109, 21528–21533 (2012).

58. Ayyaz A, Li H, Jasper H. Haemocytes control stem cell activity in the Dro-
sophila intestine. *Nat Cell Biol.* **17**, 736–748 (2015).

59. Carlson K, Gardner K, Pashaj A, Carlson D. Genome-wide gene expression in
relation to Age in large laboratory cohorts of Drosophila melanogaster. *Genet*
Res Int, 835624 (2015)

60. Petkau K, Parsons B, Duggal A, Foley E. A deregulated intestinal cell cycle
program disrupts tissue homeostasis without affecting longevity in Drosophila.
J Biol Chem. **289**, 28719-28729 (2014).

61. Wong A, Dobson AJ, Douglas AE. Gut microbiota dictates the metabolic re-
sponse of Drosophila to diet. *J Exp Biol.* **217**, 1894–1901 (2014).

62. Regan JC, Khericha M, Dobson AJ, Bolukbasi E, Rattanavirotkul N, Partridge
L. Sex difference in pathology of the ageing gut mediates the greater response
of female lifespan to dietary restriction. *Elife.* **5**, e10956 (2016).

63. Hudry B, Khadayate S, Miguel-Aliaga I. The sexual identity of adult intestinal
stem cells controls organ size and plasticity. *Nature* **530**, 344–348 (2016).

64. Clark RI, Salazar A, Yamada R, Sorel F-G, Morselli M, Alcaraz J, et al. Distinct
Shifts in Microbiota Composition during Drosophila Aging Impair Intestinal
Function and Drive Mortality. *Cell Rep.* **12**, 1656-1667 (2015).

65. Block DH, Kwame T-B, Kang H, Carlisle JA, Hanganu A, Lai T, et al. The De-
velopmental Intestinal Regulator ELT-2 Controls p38-Dependent Immune Re-
sponses in Adult C. elegans. *PLoS Genet.* **11**, e1005265 (2015).

66. Aronson BE, Stapleton KA, Krasinski SD. Role of GATA factors in develop-
ment, differentiation, and homeostasis of the small intestinal epithelium. *Am J*
Physiol Gastrointest Liver Physiol. **306**, 474–490 (2014).

67. O'Brien LE, Soliman SS, Li X, Bilder D. Altered Modes of Stem Cell Division
Drive Adaptive Intestinal Growth. *Cell* **147**, 603–614 (2011).

68. Libert S, Chao Y, Chu X, Pletcher SD. Trade-offs between longevity and path-
ogen resistance in Drosophila melanogaster are mediated by NFκB signaling.
Aging Cell **5**, 533–43 (2006).

69. Bass T, Grandison RC. Wong R, Martinez P, Partridge L, Piper MDW. Optimi-
zation of dietary restriction protocols in Drosophila. *J Gerontol A Biol Sci Med*
Sci **62**, 1071-1081 (2007).

70. Metaxakis A, Partridge L. Dietary Restriction Extends Lifespan in Wild-Derived
Populations of Drosophila melanogaster. *PLoS ONE* **8**, e74681 (2013).

71. Kim D, Pertea G, Trapnell C, Pimentel H, Kelley R, Salzberg SL. TopHat2: ac-
curate alignment of transcriptomes in the presence of insertions, deletions and
gene fusions. *Genome Biol* **14**, R36 PAGE (2013).

72. Anders S, J M Davis, Chen Y, Okoniewski M, Smyth GK, Huber W, et al.
Count-based differential expression analysis of RNA sequencing data using R
and Bioconductor. *Nat Protoc.* **8**, 1765–1786 (2013).

73. Love MI, Huber W, Anders S. Moderated estimation of fold change and dis-
persion for RNA-seq data with DESeq2. JOURNAL **15** 550 PAGE (2014).

74. Lex A, Gehlenborg N, Strobelt H. UpSet: visualization of intersecting sets. *IEEE Trans Vis Comput Graph* **20**, 1983-1992 (2014).

75. Herrmann C, de Sande B, Potier D, Aerts S. i-cisTarget: an integrative genomics method for the prediction of regulatory features and cis-regulatory modules. JOURNAL **40**, e114–e114 THIS ISN'T RIGHT (2012).

Acknowledgments

We thank Cathy Slack, Nazif Alic, Angela Douglas and Virginia Howick for comments which improved the manuscript. This study was funded by grants to MP from the Biotechnology and Biological Sciences Research Council (BBI011544/1) and the Royal Society (UF100158 & RG110303).

Authors' Contributions

AJD analysed data, performed experiments, designed the study and wrote the manuscript. XH performed experiments. E Blanc analyed data. E Bolukbasi contributed transgenic flies and performed experiments. YF performed experiments. MY performed experiments and designed the study. MDWP designed the study and wrote the paper.

Competing financial interests

The authors declare they have no competing financial interests.

Data availability

RNAseq data have been deposited with NCBI. Custom R scripts have been uploaded to GitHub.