1 Ecological factors rather than barriers to dispersal shape genetic structure of algal
2 symbionts in horizontally-transmitting corals
3

4 Davies SW[1,2*], Wham FC[3], Kanke MR[1,4] and MV Matz[1]
5
6 [1] The University of Texas at Austin, Department of Integrative Biology, Austin, TX
7
8 [2] The University of North Carolina at Chapel Hill, Department of Marine Sciences,
9 Chapel Hill, NC
10
11 [3] Pennsylvania State University, Department of Biology, University Park, PA
12
13 [4] The University of North Carolina at Chapel Hill, Department of Genetics, Chapel Hill,
14 NC
15
* Corresponding author:
Sarah W. Davies, 512-609-9134. daivessw@gmail.com
Department of Marine Sciences, 123 South Road, The University of North Carolina at
Chapel Hill, Chapel Hill, NC 27599-3300

KEYWORDS: coral, symbiosis, *Symbiodinium,* population structure, local adaptation,
host-specificity, ecological partitioning

Abstract

Many reef-building corals acquire their algal symbionts (*Symbiodinium* sp.) from the local environment upon recruitment. This horizontal transmission strategy where hosts pair with locally available symbionts could serve to increase coral fitness across diverse environments, as long as hosts maintain high promiscuity and symbionts adapt locally. Here, we tested this hypothesis in two coral species by comparing host and symbiont genetic structures across different spatial scales in Micronesia. Each host species associated with two genetically distinct *Symbiodinium* lineages, confirming high promiscuity in broadly dispersing hosts. However, contrary to our initial expectation, symbiont genetic structure was independent of physical barriers to dispersal between islands, unlike genetic structure of their hosts that was nearly perfectly explained by ocean currents. Instead, *Symbiodinium* consistently demonstrated genetic divergence among local reefs and between the two host species at each island, although not necessarily between distant islands. These observations indicate that *Symbiodinium* lineages disperse much more broadly than previously thought and continuously adapt to specific hosts and reef environments across their range, following the classical Baas Becking's hypothesis: "Everything is everywhere, but the environment selects". Overall, our findings confirm that horizontal transmission could be a mechanism for broadly dispersing coral species to enhance their local fitness by associating with locally adapted symbionts. Dramatic differences in factors driving the genetic structures of horizontally-transmitting corals and their *Symbiodinium* imply that viewing their combined genomes as a single entity ('hologenome') would not be useful in the context of their evolution and adaptation.

Introduction

Many well-known symbioses involve the passing of symbionts from parents to offspring

(vertical transmission), fully aligning the evolutionary trajectories of symbiotic partners

and typically lead to their deep integration at biochemical and genomic levels (i.e.

Buchnera in aphids (1, 2)). The result of such symbiosis is essentially a novel composite

organism. In other types of symbioses, the association between partners must be newly

established in every generation (horizontal transmission), which allows for the

maintenance of each partner's species identity. In theory, this kind of relationship should

generate novel ecological opportunities for both symbiotic partners through their mixing

and matching across environments. For example, association with ecologically

specialized algal photobionts can lead to distinct ecological guilds of lichens (3) or allow

a fungal partner to expand its geographic range across a broader climatic envelope (4).

Similarly, in aphids, association with various horizontally transmitted bacterial symbionts

allows these insects to colonize novel host plants across climatic zones (5). Considering

these and other examples of ecological adaptation based on varying symbiotic

associations, it has been argued that the joint genomic content of symbiotic systems

should be studied as a single unit of evolution, the 'hologenome' (6, 7). However, the

usefulness of this concept depends on whether the evolutionary trajectories of both

symbiotic partners are sufficiently aligned to present a unified target of selection (8).

Here, we explore this question in the symbiosis between a horizontally transmitting reef-

building coral and dinoflagellate algae of the genus *Symbiodinium* (9).

Association with *Symbiodinium* is obligatory for many coral hosts that rely on

algal photosynthesis for energy, while the algae benefit from protected and light-exposed

residence as well as inorganic nutrients and CO_2 concentration regulatory mechanisms

provided by the host (10-13). Given the obligatory nature of this symbiosis for the host, it

is somewhat surprising that in the majority of coral species (~85%) *Symbiodinium* are not

transmitted vertically, but rather must be acquired by the juvenile coral from its local

environment (14, 15). One possible explanation is that dispersal ranges of aposymbiotic

coral larvae typically extend over hundreds of kilometers (i.e. (16), while the

environmental variation corals must deal with exists on much smaller spatial scales: reef

environments with varying light, thermal and nutrient conditions can be separated by

meter-scale distances (i.e. (17). Under such circumstances, coral hosts would benefit

from the mixing and matching strategy, improving their fitness by associating with the

locally available, and putatively ecologically specialized, algal strains (18-20).

Establishing the relative roles of these symbiotic partners in adaptation to variable

environments is essential for better prediction of coral reefs' future under climate change

(i.e. (21).

The working hypothesis for this study, the 'global host, local symbiont'

hypothesis, posits that (i) coral host disperses widely and is able to establish symbiosis

with diverse algal strains across locations; (ii) *Symbiodinium* algae are poor dispersers

(i.e., 21), which results in strong divergence among locations aligning with physical

barriers and facilitates their local environmental specialization; (iii) local *Symbiodinium*

strains also diverge with respect to the host species as a result of selective pressure

towards higher host specificity (22, 23). To validate all three components of our

hypothesis, we examined multi-locus genotypes (MLG) of clade C *Symbiodinium* in two

species of *Acropora – A. hyacinthus* and *A. digitifera –* collected from the same reef

locations across the Micronesian Pacific (Fig. 1A). Our previous work has shown that

both host species exhibit extensive genetic connectivity and their genetic structure is

nearly perfectly explained by the patterns of regional surface currents (16). By using two

coral species that co-occur across the same locations as well as local reef environments

we aimed to disentangle the roles of environmental specialization, host specialization,

and physical barriers to dispersal in driving the fine-scale genetic structure of

Symbiodinium.

Figure 1: Geographic location of the Micronesian islands where *Acropora hyacinthus* and *A.*
digitifera coral samples were collected (A). Top: Map of the Micronesian Pacific with an inset of
the Pacific Ocean for reference. Islands where samples were collected and analyzed for
Symbiodnium genetics are designated with grey boxes. Detailed information on each sampling
site is located in Table S3. No *A. hyacinthus* were present in Guam and *A. digitifera* were not
collected in Ngulu. Analyses of microsatellite loci data for *Symbiodinium* hosted by *Acropora*
hyacinthus and *Acropora digitifera* using multilocus genotyping data. B. STRUCTURE population
assignment for *Symbiodinium* from two *Acropora* host species across greater Micronesia at an
optimal population number (K=2), corresponding to C40 (pink) and C3 (turquoise) clade C
subclades. Colors in the bottom panels correspond to host species and shades of grey correspond
to different sites within each island. C. Within-subclade STRUCTURE analysis.

Results

Two *Symbiodinium* clade C lineages

To enable standard population genetic analysis, data analysis was restricted to corals

hosting a single diploid *Symbiodinium* clone, which encompassed the majority (69% *A.*

digitifera and 64% *A. hyacinthus*) of our samples. Across the two coral species in

Micronesia (Fig. 1A), two distinct *Symbiodinium* lineages were observed, which were

most clearly discriminated by two of the six microsatellite loci (Fig. S1). Sequencing of

the psbA$_{ncr}$ gene for a subset samples classified these lineages as C40 and C3 (39, (24)

(Fig. S2). Corals of both species from Palau and Ngulu hosted exclusively C40 (Fig. 1B,

pink bars). This lineage was also prevalent in *A. digitifera* at one reef site on Yap and was

rarely found in *A. digitifera* across the rest of Micronesia. All other coral hosts of either

species associated with C3 (Fig. 1B, turquoise bars). Both *Symbiodinium* lineages

possessed high genetic diversity, with a total of 70 unique alleles across three islands in

C40 (Table S1A) and 130 unique alleles across five islands in C3 (Table S1B). Mean

numbers of alleles per island for each locus for C40 ranged from 3.00-4.67 with the

highest numbers of private alleles in Palau (N=7) (Table S1A). Mean allele numbers per

island for each locus for C3 ranged from 3.83-5.17 and numbers of private alleles ranged

from 1-4 (Table S1B).

***Symbiodinium* genetic structure**

All pairwise between-island F_{ST}'s for C40 were significant (Table 1A). C3 had one non-

significant F_{ST} (Yap-Pohnpei) while all others were significant and ranged from 0.058

(Guam-Chuuk) to 0.078 (Yap-Kosrae) (Table 1B). Notably, *Symbiodinium* C3

differentiation did not show the isolation-by-distance (IBD) pattern observed in both

coral hosts (Fig. 2). IBD could not be computed for C40 due to too few between-island

comparisons. C3 F_{ST} values generally exceeded F_{ST} values for the host (Fig. 2A),

however, this result did not imply stronger symbiont genetic differentiation: higher F_{ST} in

the symbionts was a consequence of higher mean heterozygosity in *Symbiodinium*

markers (e.g. (25). Indeed, an alternative measure of genetic differentiation controlling

for mean heterozygosity - Jost's D (26) - demonstrated that symbiont genetic divergence

was in fact intermediate between the two hosts (Fig. 2B).

Table 1. Summary of pairwise F_{ST} values between all islands for *Symbiodinium* C40 (A) and C3
(B). Permutations were run 9999 times. All significant comparisons are shaded in grey.
A. *Symbiodinium* C40

	PAL	NGU	YAP
PAL	0.000	***	0.004
NGU	0.279	0.000	***
YAP	0.056	0.326	0.000

145 B. *Symbiodinium* C3

	YAP	GUA	CHU	POH	KOS
YAP	0.000	***	***	0.059	***
GUA	0.062	0.000	***	***	***
CHU	0.066	0.058	0.000	***	***
POH	0.009	0.071	0.063	0.000	***
KOS	0.078	0.077	0.076	0.067	0.000

147
Figure 2: Comparison of two host species and *Symbiodinium* C3 differentiation. A. Pairwise
genetic differentiation [(F_{ST} /(1-F_{ST})] of two species of *Acropora* coral and *Symbiodinium* C3
across linear distances (km) demonstrating significant isolation by distance for the two host
species but no correlation for the symbiont. B. Pairwise Jost's D for the same two host species
and *Symbiodinium* C3 across linear distances (km) demonstrating no isolation by distance and no
difference in overall divergence between the host and symbiont.

Host specificity and environmental specialization

Discriminant analysis of principal components (DAPC) strongly differentiated between host species for both *Symbiodinium* C40 and C3 (Table S2, Fig. S3A, B), suggesting that host specificity drives symbiont diversity. In addition, DAPC demonstrated consistent differences among islands for each *Symbiodinium* lineage irrespective of host species: strong per-island assignment proportions were observed for C40 (Fig. S3C, 82-100%) and C3 (Fig. S3D, 51-98%), consistent with both STRUCTURE (Fig. 1B) and F_{ST} results (Table 1). Moreover, nearly all pairwise F_{ST} values between different reef sites within islands were significant for both lineages (Table 2), suggesting environmental partitioning of symbionts. In accord with these results, of the two top eigenvalues in DAPC of individual within islands, one explained divergence by host species (host specificity) and the other explained differences between reef sites (environmental specialization) (Fig. 3).

Table 2. Summary of pairwise F_{ST} values for *Symbiodinium* C40 (A) and C3 (B) between all sites pooled across host species. Permutations were run 9999 times. All significant comparisons are shaded in grey.

A. *Symbiodinium* C40

	PAL1	PAL2	NGU	YAP2
PAL1	0.000	0.009	***	0.007
PAL2	0.031	0.000	***	0.007
NGU	0.283	0.303	0.000	***
YAP2	0.057	0.074	0.326	0.000

B. *Symbiodinium* C3

	YAP1	YAP2	GUA1	GUA2	CHU1	CHU2	POH1	POH2	KOS1	KOS2
YAP1	0.000	0.139	0.002	0.009	***	***	0.002	0.055	***	***
YAP2	0.011	0.000	***	0.003	***	***	0.027	0.064	***	***
GUA1	0.059	0.107	0.000	0.045	***	***	***	***	***	***
GUA2	0.045	0.062	0.025	0.000	0.002	***	***	0.008	***	***
CHU1	0.071	0.073	0.104	0.065	0.000	***	***	***	***	***
CHU2	0.079	0.112	0.070	0.086	0.068	0.000	***	***	***	***
POH1	0.053	0.028	0.126	0.111	0.085	0.126	0.000	***	***	***
POH2	0.018	0.018	0.093	0.039	0.067	0.092	0.063	0.000	***	***
KOS1	0.154	0.146	0.170	0.110	0.163	0.176	0.168	0.119	0.000	***
KOS2	0.092	0.121	0.116	0.131	0.129	0.090	0.136	0.100	0.183	0.000

Figure 3: Discriminant analysis of principal components (DAPC) of MLG data for *Symbiodinium* C40 and C3 hosted by *Acropora hyacinthus* and *Acropora digitifera* at twelve sites across six islands (Ngulu not included) in Micronesia. DAPC analysis on two discriminant functions demonstrating host species assignments and site assignments. DAPC scatter plots for individual samples from within A. Palau for *Symbiodinium* C40. B. Yap for *Symbiodinium* C3, C. Chuuk for *Symbiodinium* C3, D. Pohnpei for *Symbiodinium* C3, E. Kosrae for *Symbiodinium* C3, and F. Guam for *Symbiodinium* C3 for *A. digitifera* hosts only (1 DF axis). Proportions of assignments are indicated in the clusters. Information on the DAPC models can be found in Table S2.

Discussion

Promiscuity of the coral host

Clade C *Symbiodinium* are considered to be the most derived lineage within the genus

Symbiodinium and exhibit significantly higher within-clade diversity when compared to

other, more basal, clades (9, 27, 28). Across the Micronesian Pacific (Fig. 1A), both coral

hosts associated exclusively with clade C *Symbiodinium*, which was nevertheless

represented by two distinct lineages– C40 and C3 (Fig. S2). This observation confirms

the first prediction of the 'global host, local symbiont' hypothesis: coral hosts show

considerable flexibility in their symbiotic association across their range and within their

habitat.

192
Lack of dispersal limitation in <u>Symbiodinium</u> *across Micronesia*

Initially, we expected to find strong isolation by distance in *Symbiodinium*, since the

prevailing view of the *Symbiodinium* life cycle involves symbiotic existence in sedentary

hosts alternating with a free-living form that largely exists in the benthos where dispersal

by ocean currents must be limited (29-32). In contrast, we observed that genetic

differentiation between islands of *Symbiodinium* C3 across Micronesia did not exceed

F_{ST} = 0.078, which contrasts studies from other locations reporting symbiont F_{ST} as high

as 0.54 in clade A (33) and Φ_{ST} as high as 0.468 in clade C (34). This lack of physical

dispersal limitation is best demonstrated by comparing *Symbiodinium* C3 divergence to

divergence of their coral hosts, which has been shown to nearly perfectly follow

biophysical barriers (16). If *Symbiodinium* dispersal were limited by the same physical

barriers and to a greater extent, as our hypothesis posited, we would have observed

steeper increases in divergence with distance relative to the host; instead, no increase of

F_{ST} with distance was observed (Fig. 2A). Moreover, after correcting for mean

heterozygosity, genetic divergence (Jost's D) in *Symbiodinium* C3 was found to be

intermediate between *A. hyacinthus* and *A. digitifera* (Fig. 2B), once again demonstrating

that symbiont dispersal is less physically limited than their hosts. This finding contradicts

the second prediction of our 'global host, local symbiont' hypothesis and is contrary to

our expectations based on earlier F_{ST}-based reports of stronger *Symbiodinium*

differentiation compared to hosts (20-22, 34).

Host specificity

The majority of reef-building coral species associate with a specific strain (termed

"clade" or "subclade") of *Symbiodinium* broadly defined based on ribosomal and/or

chloroplast markers (22, 35-37). Previous *Symbiodinium* multilocus genotyping

studies revealed that each of these strains harbors greater diversity, both in terms of

genetic and functional diversity (20, 34, 38). Our data indicate that the local association

of hosts and symbionts of the same genotypic cluster is due to pervasive evolution of host

specificity in *Symbiodinium* (Fig. S3 & 3). As our study includes two coral species, we

also observe that this specificity is not perfect: at every location there were symbionts

that would have been assigned to another coral host based on their multilocus genotype

(Fig. 3). This suggests that host specialization in *Symbodinium* arises in the face of

considerable exchange between symbiont communities hosted by different coral species

at the same location.

Environmental partitioning

The most ecologically relevant postulate of the 'global host, local symbiont' hypothesis is

that locally available symbionts are also locally adapted, giving rise to a locally adapted

holobiont. Recurrent genetic divergence of symbionts between reef sites within the same

island (Fig. 3) despite the lack of physical dispersal limitation across much larger

distances (Fig. 2) suggests that symbiont genetic divergence is likely due to poor survival

of immigrants rather than to physical barriers to migration. Given that hosts are available

at every site, this leaves other ecological parameters of local reef environment as the most

likely barrier-forming force, preventing survival of immigrants adapted to a different

environment– a situation termed "phenotype-environment mismatch" (39). This in turn

implies environmental specialization (i.e., local adaptation) in the symbionts, which

supports our hypothesis. Interestingly, just like host specificity, this environmental

specialization is imperfect: there are several *Symbiodinium* genotypes that appear to be

successful migrants between reef sites (particularly clearly visible for Kosrae, Fig. 3E),

indicating that environmental partitioning also arises in the face of considerable gene

flow.

Conclusions

Both coral host species associated with two divergent *Symbiodinium* lineages, indicating

that coral hosts are promiscuous in their symbiont associations across the seascape. In

contrast, within each *Symbiodinium* lineage, strong associations with particular host

species were observed, suggesting that host-specificity is an important driver of

Symbiodinium diversification. Unexpectedly, *Symbiodinium* genetic divergence did not

correlate with physical distance between islands, but was driven by the combined effects

of local (within-island) environment and host identity. This indicates that *Symbiodinium*

assemblages across Micronesia are comprised by broadly dispersing lineages

dynamically adapting to specific coral hosts and reef environments across their range.

Assuming that host species can be regarded as part of *Symbiodinium*'s environment, this

pattern clearly follows the classical hypothesis of Baas Becking (originally proposed for

soil microbes): "Everything is everywhere, but the environment selects" (40). Overall,

our observations support the view that horizontal transmission might be a strategy for

broadly dispersing coral hosts to improve their local fitness by associating with locally

available, environmentally specialized, *Symbiodinium*. Finally, dramatic differences in

factors that structure genetic diversity in the coral host and *Symbiodinium* indicate that

viewing their combined genomes as a single 'hologenome' would not be useful in the

context of their evolution and adaptation.

Materials and Methods

Sampling: This study comprised a subset of samples previously analyzed for coral host

genetics in Davies *et al.* (16) (Table S3, Fig. 1A). Twenty-five individuals of each coral

host species (*Acropora hyacinthus* and *A. digitifera*) were examined at two reef sites

within seven islands, with the exception of Ngulu (only collected *A. hyacinthus*) and

Guam (no *A. hyacinthus* found), for a total of thirteen sites.

Laboratory Procedures: Holobiont DNA was isolated following Davies *et al.* (41).

Microsatellite primers consisted of six previously published clade C loci (42, 43) and one

novel locus mined using Msatcommander (44) from nucleotide EST data for

Symbiodinium sp. clade C3 (45) (Table S4). Loci were multiplexed in 20 µl polymerase

chain reaction (PCR) mixtures containing 10 ng of template, 0.1 µM of each forward and

reverse primers, 0.2 mM dNTP, 1 µl 10X *ExTaq* buffer, 0.025 U *ExTaq* Polymerase and

0.0125 U *Pfu* Polymerase. Amplifications began at 94°C for 5 min, followed by 35 cycles

of 94°C for 40 s, annealing temperature for 120 s, and 72°C for 60 s and a 10 minute

72°C extension period. Molecular weights were analyzed using the ABI 3130XL

capillary sequencer. Data were binned by repeat size and individuals failing to amplify at

≥3 loci were excluded from analyses.

Data Analysis: *Symbiodinium* were scored as diploid based on available information on

their ploidy (22, 43, 46) Since each host could potentially contain populations of

genetically distinct *Symbiodinium*, multi-locus genotypes (MLG) were only considered if

they contained two or fewer alleles across all loci, suggesting that all alleles originate

from a single, clonally replicated, genome (46), which was the case for 190 of 277 *A.*

digitifera hosts and 178 of 278 *A. hyacinthus* hosts. STRUCTURE v2.3.3 (47) assigned

similar MLGs to populations (10^6 iterations, burn in = 300,000) across ten replicate runs

for each K (1-10) using an admixture model with no location prior. Following (48), ΔK

was calculated in STRUCTURE Harvester (49) and CLUMPP (50) and DISTRUCT (51)

produced all graphics. We observed high percent assignment to two clusters among

sympatric individuals. Following (46), we investigated the relationship between these

assignments and allele identities and found strong relationships between assignments and

allele identity, particularly at Sgr_21 and Spl_33 (Fig. S1), and we concluded that two

distinct *Symbiodinium* lineages were present. MLG data were then split into lineages and

additional STRUCTURE runs were completed.

Partitioned lineage data were investigated for genetic divergences among islands

and sites using F_{ST} (AMOVA, 9999 permutations in GENALEX v6.5, (52). Pairwise

differentiations (F_{ST} and Jost's D) and pairwise island distances were correlated to test

for isolation by distance (IBD) in *Symbiodinium* C3 (insufficient C40 data) and these

trends were compared to results from the host species (53). Mean allelic diversities per

island and numbers of private alleles per island were also calculated in GENALEX v6.5.

To resolve differences between islands, between host species and between sites

and host species within each island, assignment of samples to genetic clusters using

discriminant analysis of principal components (DAPC) was performed in R (54) using the

ADEGENET package (55, 56). Data were converted into principle components (PCs) and

then a-scores were used to determine trade-offs between power of discrimination and

model over-fitting. Relationships were examined by DAPC and model information are

listed in Table S2.

Sequencing Analysis of *Symbiodinium* psbA^{ncr}

To confirm individual assignments to different *Symbiodinium* lineages, the non-coding

region of the circular plastid (psbA^{ncr}) was amplified in a subset of samples using the

methods described by (24). Amplified products were directly sequenced and aligned with

ClustalW2 (http://www.ebi.ac.uk/Tools/msa/clustalw2/) to clade C psbA^{ncr} sequences

from common Indo-Pacific *Symbiodinium* types (Table S5). The phylogeny was

reconstructed using the default settings of Phylogeny.fr (57) and identities

of *Symbiodinium* types were interpreted based on percent sequence identity with

previously identified *Symbiodinium* types.

Acknowledgements

Thanks to field assistants Carly Kenkel, Tim Kiett, Irina Yakushenok and David Stump.
Nida Khawaja Rahman was integral in molecular work and James Derry assited with data
management. We are grateful to Ulrich Mueller for numerous suggestions. We
acknowledge all permit authorities. This study was supported by the Coral Reef
Conservation program of the National Oceanic and Atmospheric Administration
(administered through the Hawaiian Undersea Research Laboratory) and the National
Science Foundation grant DEB-1054766 to M.V.M. DW was funded in part by
Pennsylvania State University and grants from the National Science Foundation (OCE-
0928764 and IOS-1258058).

References

1. Nakabachi A, Ishida K, Hongoh Y, Ohkuma M, & Miyagishima SY (2014) Aphid gene of bacterial origin encodes a protein transported to an obligate endosymbiont. *Curr Biol* 24(14):R640-641.

2. Shigenobu S & Wilson AC (2011) Genomic revelations of a mutualism: the pea aphid and its obligate bacterial symbiont. *Cellular and molecular life sciences : CMLS* 68(8):1297-1309.

3. Peksa O & Skaloud P (2011) Do photobionts influence the ecology of lichens? A case study of environmental preferences in symbiotic green alga Asterochloris (Trebouxiophyceae). *Mol Ecol* 20(18):3936-3948.

4. Fernandez-Mendoza F, *et al.* (2011) Population structure of mycobionts and photobionts of the widespread lichen Cetraria aculeata. *Mol Ecol* 20(6):1208-1232.

5. Henry LM, *et al.* (2013) Horizontally transmitted symbionts and host colonization of ecological niches. *Curr Biol* 23(17):1713-1717.

6. Brucker RM & Bordenstein SR (2014) Response to Comment on "The hologenomic basis of speciation: Gut bacteria cause hybrid lethality in the genus Nasonia". *Science* 345(6200).

7. Zilber-Rosenberg I & Rosenberg E (2008) Role of microorganisms in the evolution of animals and plants: the hologenome theory of evolution. *Fems Microbiol Rev* 32(5):723-735.

8. Moran NA & Sloan DB (2015) The Hologenome Concept: Helpful or Hollow? *PLoS biology* 13(12):e1002311.

9. Lesser MP, Stat M, & Gates RD (2013) The endosymbiotic dinoflagellates (Symbiodinium sp.) of corals are parasites and mutualists. *Coral Reefs*:603-611.

10. Muscatine L (1990) The role of symbiotic algae in carbon and energy flux in reef corals. *Ecosystems of the World* 25:75-87.

11. Barott KL, Venn AA, Perez SO, Tambutte S, & Tresguerres M (2015) Coral host cells acidify symbiotic algal microenvironment to promote photosynthesis. *Proc Natl Acad Sci U S A* 112(2):607-612.

12. Muscatine L & Cernichiari E (1969) Assimilation of photosynthetic products of zooxanthellae by a reef coral. *The Biological Bulletin* 137(3):506-523.

13. Trench RK & Blank RJ (1987) Symbiodinium microadriaticum Fredenthal, S. Goreau sp. nov., S. Kawagut2 sp. nov., and S. pilosum sp. nov.: Gymnodinoid dinoflagellate symbionts of marine invertebrates. *J Phycol* 23(3):469-481.

14. Fadlallah YH (1983) Sexual reproduction, development and larval biology of scleractinian corals. A review. *Coral Reefs* 2:129-150.

15. Harrison PLW, C.C. (1990) *Reproduction, dispersal and recruitment of scleractinian corals* (elsevier Science Publications Amsterdam).

16. Davies SW, Treml EA, Kenkel CD, & Matz MV (2015) Exploring the role of Micronesian islands in the maintenance of coral genetic diversity in the Pacific Ocean. *Mol Ecol* 24(1):70-82.

17. Gorospe KD & Karl SA (2011) Small-Scale Spatial Analysis of In Situ Sea Temperature throughout a Single Coral Patch Reef. *Journal of Marine Biology* 2011:12.

18. Byler KA, Carmi-Veal M, Fine M, & Goulet TL (2013) Multiple symbiont
acquisition strategies as an adaptive mechanism in the coral Stylophora pistillata.
Plos One 8(3):e59596.
19. Rowan R & Knowlton N (1995) Intraspecific diversity and ecological zonation in
coral-algal symbiosis. *Proc Natl Acad Sci U S A* 92(7):2850-2853.
20. Howells EJ*, et al.* (2012) Coral thermal tolerance shaped by local adaptation of
photosymbionts. *Nat Clim Change* 2:116-120.
21. Baums IB, Devlin-Durante MK, & LaJeunesse TC (2014) New insights into the
dynamics between reef corals and their associated dinoflagellate endosymbionts
from population genetic studies. *Mol Ecol* 23(17):4203-4215.
22. Thornhill DJ, Lewis AM, Wham DC, & LaJeunesse TC (2014) Host-specialist
lineages dominate the adaptive radiation of reef coral endosymbionts. *Evolution*
68(2):352-367.
23. Coffroth MA, Santos SR, & Goulet TL (2001) Early ontogenetic expression of
specificity in a cnidarian–algal symbiosis. *Mar Ecol Prog Ser* 222:85-96.
24. LaJeunesse TC & Thornhill DJ (2011) Improved resolution of reef-coral
endosymbiont (Symbiodinium) species diversity, ecology, and evolution through
psbA non-coding region genotyping. *Plos One* 6(12):e29013.
25. Jakobsson M, Edge MD, & Rosenberg NA (2013) The Relationship Between F-
ST and the Frequency of the Most Frequent Allele. *Genetics* 193(2):515-528.
26. Jost L (2008) G(ST) and its relatives do not measure differentiation. *Mol Ecol*
17(18):4015-4026.
27. Pochon X & Gates RD (2010) A new Symbiodinium clade (Dinophyceae) from
soritid foraminifera in Hawai'i. *Molecular phylogenetics and evolution* 56(1):492-
497.
28. Pochon X, Montoya-Burgos JI, Stadelmann B, & Pawlowski J (2006) Molecular
phylogeny, evolutionary rates, and divergence timing of the symbiotic
dinoflagellate genus Symbiodinium. *Molecular phylogenetics and evolution*
38(1):20-30.
29. Fitt WK, Chang SS, & Trench RK (1981) Motility Patterns of Different Strains of
the Symbiotic Dinoflagellate Symbiodinium (=Gymnodinium) Microadriaticum
(Freudenthal) in Culture. *Bulletin of Marine Science* 31(2):436-443.
30. Fitt WK & Trench RK (1983) The Relation of Diel Patterns of Cell-Division to
Diel Patterns of Motility in the Symbiotic Dinoflagellate Symbiodinium-
Microadriaticum Freudenthal in Culture. *New Phytol* 94(3):421-432.
31. Yacobovitch T, Benayahu Y, & Weis VM (2004) Motility of zooxanthellae
isolated from the Red Sea soft coral Heteroxenia fuscescens (Cnidaria). *J Exp*
Mar Biol Ecol 298(1):35-48.
32. Littman RA, van Oppen MJH, & Willis BL (2008) Methods for sampling free-
living Symbiodinium (zooxanthellae) and their distribution and abundance at
Lizard Island (Great Barrier Reef). *J Exp Mar Biol Ecol* 364(1):48-53.
33. Andras JP, Kirk NL, & Harvell CD (2011) Range-wide population genetic
structure of Symbiodinium associated with the Caribbean Sea fan coral, Gorgonia
ventalina. *Mol Ecol* 20(12):2525-2542.

34.   Howells EJ, van Oppen MJ, & Bay LK (2009) High genetic differentiation and
424   cross-shelf patterns of genetic diversity among Great Barrier Reef populations of
425   Symbiodinium. *Coral Reefs* 28(1):215-225.
426   35.   Fabina NS, Putnam HM, Franklin EC, Stat M, & Gates RD (2013) Symbiotic
427   specificity, association patterns, and function determine community responses to
428   global changes: defining critical research areas for coral-Symbiodinium
429   symbioses. *Global Change Biol* 19(11):3306-3316.
430   36.   Rodriguez-Lanetty M, Krupp DA, & Weis VM (2004) Distinct ITS types of
431   Symbiodinium in Clade C correlate with cnidarian/dinoflagellate specificity
432   during onset of symbiosis. *Mar Ecol Prog Ser* 275(97-102).
433   37.   Weis VM, Reynolds WS, deBoer MD, & Krupp DA (2001) Host-symbiont
434   specificity during onset of symbiosis between the dinoflagellates Symbiodinium
435   spp. and planula larvae of the scleractinian coral Fungia scutaria. *Coral Reefs*
436   20(3):301-308.
437   38.   Santos SR, Shearer TL, Hannes AR, & Coffroth MA (2004) Fine-scale diversity
438   and specificity in the most prevalent lineage of symbiotic dinoflagellates
439   (Symbiodinium, Dinophyceae) of the Caribbean. *Mol Ecol* 13(2):459-469.
440   39.   Marshall DJ, Monro K, Bode M, Keough MJ, & Swearer S (2010) Phenotype-
441   environment mismatches reduce connectivity in the sea. *Ecol Lett* 13(1):128-140.
442   40.   Baas Becking LGM (1934) Geobiologie of inleiding tot de milieukunde. *Den*
443   *Haag [Netherlands]: W.P. Van Stockum & Zoon, N. V.*
444   41.   Davies SW*, et al.* (2013) Novel polymorphic microsatellite markers for
445   population genetics of the endangered Caribbean star coral, Montastraea
446   faveolata. *Mar Biodivers* 43(2):167-172.
447   42.   Bay LK, Howells EJ, & van Oppen MJ (2009) Isolation, characterisation and
448   cross amplification of thirteen microsatellite loci for coral endo-symbiotic
449   dinoflagellates (Symbiodinium clade C). *Conserv Genet Resour* 1(1):199-203.
450   43.   Wham DC, Carmichael M, & LaJeunesse TC (2014) Microsatellite loci for
451   Symbiodinium goreaui and other Clade C Symbiodinium. *Conserv Genet Resour*
452   6(1):127-129.
453   44.   Faircloth BC (2008) MSATCOMMANDER: detection of microsatellite repeat
454   arrays and automated, locus-specific primer design. *Molecular ecology resources*
455   8(1):92-94.
456   45.   Leggat W, Hoegh-Guldberg O, Dove S, & Yellowlees D (2007) Analysis of an
457   EST library from the dinoflagellate (Symbiodinium sp.) symbiont of reef-building
458   corals. *J Phycol* 43(5):1010-1021.
459   46.   Traylor-Knowles N*, et al.* (2011) Production of a reference transcriptome and
460   transcriptomic database (PocilloporaBase) for the cauliflower coral, Pocillopora
461   damicornis. *BMC genomics* 12:585.
462   47.   Pritchard JK, Stephens M, & Donnelly P (2000) Inference of population structure
463   using multilocus genotype data. *Genetics* 155(2):945-959.
464   48.   Evanno G, Regnaut S, & Goudet J (2005) Detecting the number of clusters of
465   individuals using the software STRUCTURE: a simulation study. *Mol Ecol*
466   14(8):2611-2620.

49.   Earl DA & Vonholdt BM (2012) STRUCTURE HARVESTER: a website and
program for visualizing STRUCTURE output and implementing the Evanno
method. *Conserv Genet Resour* 4(2):359-361.
50.   Jakobsson M & Rosenberg NA (2007) CLUMPP: a cluster matching and
permutation program for dealing with label switching and multimodality in
analysis of population structure. *Bioinformatics* 23(14):1801-1806.
51.   Rosenberg NA (2004) DISTRUCT: a program for the graphical display of
population structure. *Mol Ecol Notes* 4(1):137-138.
52.   Peakall R & Smouse PE (2006) GENALEX 6: genetic analysis in Excel.
Population genetic software for teaching and research. *Mol Ecol Notes* 6(1):288-
295.
53.   Davies SW, Treml EA, Kenkel CD, & Matz MV (2015) Exploring the role of
Micronesian islands in the maintenance of coral genetic diversity in the Pacific
Ocean. *Mol Ecol* 24(1):70-82.
54.   R Development Core Team (2015) R: A language and environment for statistical
computing. (R Foundation for Statistical Computing, Vienna, Austria).
55.   Jombart T (2008) adegenet: a R package for the multivariate analysis of genetic
markers. *Bioinformatics* 24(11):1403-1405.
56.   Jombart T, Devillard S, & Balloux F (2010) Discriminant analysis of principal
components: a new method for the analysis of genetically structured populations.
BMC genetics 11:94.
57.   Dereeper A*, et al.* (2008) Phylogeny.fr: robust phylogenetic analysis for the non-
specialist. *Nucleic Acids Res* 36(Web Server issue):W465-469.
490
491

492 **Supporting Information**

493
494 Figure S1: Bubble plot of the SSR allele frequencies of C3 (red) and C40 (blue) at the two loci
495 that showed the largest discrimination power, Sgr_21 (A) and Spl_33 (B). The size of each
496 bubble is proportional to the allele frequency within each *Symbiodinium* type.

498
499 Figure S2. Phylogenetic analysis of psbA^{ncr} sequences from representative samples from this
500 study (labeled branch tips) along with publically available psbA^{ncr} sequences from known
501 *Symbiodinium* subclades identifies two lineages present (C3 and C40). Bootstrap support values
502 are shown at the partitions that define known subclades. Scale bar: replacements per nucleotide
503 site. Sequence accession numbers for reference sequences are given in Table S1.

Figure S3: Discriminant analysis of principal components (DAPC) of MLG data for *Symbiodinium* C40 and C3 hosted by *Acropora hyacinthus* and *Acropora digitifera* at thirteen sites across seven islands in Micronesia. A. DAPC analysis on two discriminant functions demonstrating strong host species assignments across all islands for *Symbiodinium* C40 and B. *Symbiodinium* C3. Numbers overlaying the curves indicate successfully assigned fraction of samples. C. DAPC scatter plot for individual samples from *Symbiodinium* C40 represented by colored dots clustered by islands. D. DAPC scatter plot for individual samples from *Symbiodinium* C3 represented by colored dots clustered by islands. Proportions of assignments are indicated in the clusters or in the legends. Information on the DAPC models can be found in Table S2.

515 Table S1: Summary of allelic diversity measures for *Symbiodinium* C40 (A) and *Symbiodinium*
516 C3 (B). Na: number of alleles, He: expected heterozygosity, Ho: observed heterozygosity, PA:
517 number of private alleles
518 A. *Symbiodinium* C40

	Na	**He**	**Ho**	**PA**
PAL	4.67	0.52	0.64	7
NGU	4.00	0.42	0.55	6
YAP	3.00	0.43	0.51	2

519 B. *Symbiodinium* C3

	Na	**He**	**Ho**	**PA**
YAP	4.33	0.47	0.56	1
GUA	3.83	0.42	0.46	3
CHU	5.17	0.58	0.51	3
POH	4.33	0.47	0.56	1
KOS	4.00	0.48	0.51	4

521 Table S2: Discriminant analysis of principle component (DAPC) model information including the
522 number of principle components ("PC") and discriminant functions ("DF") retained, the
523 proportion of conserved variance by the clustering model ("var"), and the overall assignment
524 proportions across the model ("assign"). A. DAPC information for *Symbiodinium* from different
525 host species, B. islands, and C. sites and host species within each island.
526
527 Model Information

A. Host Species	**PC**	**DF**	**var**	**assign**
Symbiodinium C40	7	1	0.814	0.879
Symbiodinium C3	14	1	0.922	0.846
B. Islands	**PC**	**DF**	**var**	**assign**
Symbiodinium C40	16	4	0.970	0.981
Symbiodinium C3	19	4	0.966	0.796
C. Within Island	**PC**	**DF**	**var**	**assign**
Palau C40	14	3	0.984	0.735
Yap C3	7	3	0.844	0.778
Guam C3	3	3	0.647	0.860
Chuuk C3	12	3	0.933	0.927
Pohnpei C3	8	3	0.906	0.900
Kosrae C3	7	3	0.878	0.888

529   Table S3: Reef Site Collections. GPS coordinates, main island group, number of *A. digitifera* and
530   *A. hyacinthus* hosts genotyped. The numbers in brackets are individuals hosting a single
531   *Symbiodinium* MLG, which were included in all analyses presented here. Site letter corresponds
532   to island insets in Figure 1A.

Site	Island	GPS	*A. digitifera*	*A. hyacinthus*
WC. West Channel Reef	Palau	7°31'55.7 N, 134°29'42.8 E	24 (20)	25 (17)
LH. Lighthouse Reef	Palau	7°16'62.4 N, 134°27'61.9 E	23 (15)	24 (16)
NG. Ngulu	Ngulu Atoll	8°18'12.0 N, 137°29'18.7 E	0[1]	39 (25)
ST. South Tip Reef	Yap	9°26'05.4 N, 138°02'10.4 E	25 (10)	25 (11)
GO. Goofnuw Channel	Yap	9°34'26.4 N, 138°12'19.2 E	24 (15)[m]	25 (20)
PB. Pago Bay	Guam	13°25'66.6 N, 144°47'94.3 E	26 (20)	0*
TG. Tanguisson	Guam	13°32'61.1 N, 144°48'52.6 E	21 (17)	0*
WP. West Polle	Chuuk	7°19'69.7 N, 151°33'21.1E	15 (6)[m]	24 (18)
SE. South East Pass	Chuuk	7°14'60.3 N, 152°01'29.1 E	21 (17)[m]	22 (16)
AN. Ant Atoll (East)	Pohnpei	6°47'42.3 N, 158°01'20.7 E	24 (16)	22 (13)
RO. Roj	Pohnpei	6°46'37.7 N, 158°12'24.1 E	24 (16)	23 (15)
CG. Coral Garden	Kosrae	5°18'47.2 N, 162°53'01.8 E	25 (15)[m]	24 (13)
HP. Hiroshi Point	Kosrae	5°15'88.0 N, 162°59'01.8 E	25 (23)[m]	25 (14)
TOTAL			**277 (190)**	**278 (178)**

* indicates that no individuals of this species were found
[1] indicates that individuals were not collected from this site but are likely present
[m] indicated sites where multiple *Symbiodinium* species were detected

Table S4: Summary of six polymorphic microsatellite loci used to assess genetic variation
Symbiodinium clade C hosted by *A. hyacinthus* and *A. digitifera* and their corresponding
multiplexing groups.

Multiplex Group	Locus (Repeat)	Primer Sequence 5'-3'	Repeat	Annealing Temperature	Source
A	*SgrSpl_30*	F: FAM-ccgaactacctttggtcaac R: aaaagacaaggacatctcgg	TA	53	Wham et al., 2014
B	*SgrSpl_78*	F: FAM-tgaaattcggtgttcattgt R:ctcagatgtttccgacgagt	TA	54	Wham et al., 2014
	Sgr_21	F: HEX-tgctgagtggcgtgtatatc R: tgatggtacttgatggtg	TCA	54	Wham et al., 2014
	Spl_33	F: HEX-acttgcaaagtccaagatcg R: gaacggtgaaaggaaaatga	CAT	54	Wham et al., 2014
C	*C784*	F: Hadp-ctccttaggactggactcgc R: agaagtcaaatcgtcaccatcg	ATC	60	This Study
D	*C105*	F: FAM-tttcgttgttggacattgttatg R: ggactgaaaggtgcttgagg	complex	55	Bay *et al.,* 2009

539   Fadp- labeled primers were indirectly labeled in each PCR reaction with an additional FAM labeled adapter
540   tag sequence: FAM: agcagcgaactcagtacaaca
541   Hadp- labeled primers were indirectly labeled in each PCR reaction with an additional FAM labeled
542   adapter tag sequence: HEX: tcgtcgcttgagtcatcgtta
543

544 Table S5: Accession number and corresponding Clade C *Symbiodinium* lineage for each unique
545 sequence identifier from Figure S2.

Sequence ID	Accession Number	Lineage
1	JQ043642	C3
2	KF572372	C3
3	JQ043643	C3
4	JQ043641	C3
5	JQ043640	C3
6	JQ043644	C3
7	JQ043635	C3
8	JQ043638	C3
9	JQ043637	C3
10	JQ043636	C3
11	KF572331	C3
12	KF572332	C3
13	KF572318	C3
14	KF572319	C3
15	KF572373	C3
16	KF572374	C3
17	KF572358	C3
18	KF572368	C40
19	KF572359	C40
20	KF572367	C40
21	KF572366	C40
22	KF572360	C40
23	KF572357	C40
24	KF572363	C40
25	KF572365	C40
26	KF572364	C40
27	KF572370	C40
28	JQ043603	C31
29	JQ043599	C31
30	JQ043604	C31
31	JQ043671	C27
32	JQ043673	C27
33	JQ043670	C27
34	JQ043669	C27
35	JQ043676	C27

546