ELF3 polyQ variation in *Arabidopsis thaliana* reveals PIF4-independent role in

thermoresponsive flowering*.*

Maximilian O. Press[1], Amy Lanctot[2], Christine Queitsch[1]*

1: University of Washington Department of Genome Sciences

2: University of Washington Molecular and Cellular Biology Program, University of

Washington Department of Biology

Keywords: flowering, thermomorphogenesis, ELF3, PIF4, temperature sensing,

Arabidopsis thaliana, polyglutamine

*: correspondence to queitsch@uw.edu

Short title: ELF3/PIF4 independence in plant adult thermal responses

ABSTRACT

Plants have evolved elaborate mechanisms controlling developmental responses to

environmental stimuli. A particularly important stimulus is temperature. Previous work

has identified the interplay of PIF4 and ELF3 as a central circuit underlying thermal

responses in *Arabidopsis thaliana.* However*,* thermal responses vary widely among

strains, possibly offering mechanistic insights into the wiring of this circuit. ELF3

contains a polyglutamine (polyQ) tract that is crucial for ELF3 function and varies in

length across strains. Here, we use transgenic analysis to test the hypothesis that

natural polyQ variation in ELF3 is associated with the observed natural variation in

thermomorphogenesis. We found little evidence that the polyQ tract plays a specific role

in thermal responses beyond modulating general ELF3 function. Instead, we made the

serendipitous discovery that ELF3 plays a crucial, PIF4-independent role in

thermoresponsive flowering under conditions more likely to reflect field conditions. We

present evidence that ELF3 acts through the photoperiodic pathway, pointing to a

previously unknown symmetry between low and high ambient temperature responses.

Moreover, in analyzing two strain backgrounds with vastly different thermal responses,

we demonstrate that responses may be shifted rather than fundamentally rewired

across strains. Our findings tie together disparate observations into a coherent

framework in which multiple pathways converge in accelerating flowering in response to

temperature, with some such pathways modulated by photoperiod.

INTRODUCTION

The responses of plants to temperature variation are of central importance to food

security in a changing world [1]. Therefore, the elucidation of the genetic pathways

underlying these responses has been a key mission of plant science [2]. Many previous

studies examined the phenomena of circadian temperature compensation [3–5],

thermoresponsive flowering [6–10], and temperature effects on plant morphology [11–

16]. Several have converged on PIF4 as a master regulator of temperature responses,

and ELF3 as an input to PIF4 integration, among many other genes and pathways [14–

16]. Given known regulatory interactions between ELF3 and PIF4 [17–20], it is

reasonable to predict that both operate in the same pathway for thermal response

phenotypes [21]. Recent reports focusing on one such phenotype, hypocotyl elongation,

support this expectation [14–16].

ELF3 serves to repress hypocotyl elongation by reducing PIF4 levels. This

repression of PIF4 occurs at both the transcriptional level, through the role of ELF3 in

the Evening Complex (EC) [17,19], and at the post-translational level, through PIF4

destabilization by phytochrome phyB [22]. Light sensing enforces circadian oscillations

of the EC and other components, leading to calibration of the circadian clock [23,24],

resulting in diurnal repression of hypocotyl elongation through repression of PIF4 and

PIF5 [17,19]. ELF3 also plays a crucial role as a flowering repressor [25]. Consequently,

elf3 null mutants show elongated hypocotyls even in the light, and flower early.

PIF4 is one of a family of basic helix-loop-helix (bHLH) "phytochrome-interacting

factors" (PIFs), transcription factors with overlapping functions promoting

skotomorphogenesis. Under dark conditions, the PIFs act to target phyB for ubiquitin-

mediated degradation by the E3 ubiquitin ligase COP1, thereby repressing

photomorphogenesis [26]. Under light conditions, degradation of PIFs is mediated by

direct interactions with photoactivated phyB [22]. PIF4 is distinct from the other PIFs in

having specific roles in temperature sensing and flowering [27]. *pif4* null mutants show

short hypocotyls with photomorphogenic attributes even in the dark [28].

At elevated ambient temperatures (27º-29º) the wiring of these signaling

pathways changes. Several independent studies have recently found that elevated

temperatures, specifically during dark periods [29], inhibit the activity of the EC by an

unknown mechanism [14–16], leading to increased expression of *PIF4* and its targets

[11,27]. This increased PIF4 activity leads to several morphological temperature

responses through various signaling pathways [13,27]. PIF4 is also required for the

acceleration of flowering at 27ºC under short photoperiods [9,29], though these

observations have been disputed [30,31]. While PIF4 action alone (among PIFs) is

essentially sufficient for most described thermomorphogenic responses [11,27], there is

evidence for a limited role of PIF5 (though not other PIFs) in thermoresponsive

flowering under short days (SDs) [30,31]. In contrast, under continuous light, *pif4* null

mutants have an intact temperature-dependent acceleration of flowering [11]. Lastly,

pif4 null mutants lose the normal elongation of petioles under high temperatures [11]. It

is unclear why PIF4 does not affect thermoresponsive flowering under continuous light;

yet, this phenomenon may reflect low PIF4 levels under these conditions due to

inhibition by phyB. Under longer photoperiods and higher temperature a flowering

acceleration still exists [7,11], which suggests a PIF4-independent thermoresponsive

flowering pathway. Nonetheless, recent reviews of the literature tend to emphasize the

primacy of PIF4 in this response [10,32,33], although the condition of elevated

temperature with short photoperiods is probably rare in the field.

Recent studies have identified ELF3 as a plausible upstream regulator of PIF4 in

thermal responses [14–18]. However, others have implicated different candidates, such

as FCA [13], and mathematical modeling has suggested that ELF3/EC complex

regulation alone is insufficient to explain PIF4 thermal regulation [14,34]. The exact

mechanisms of this response have yet to be unraveled.

Specifically, the mechanism by which EC/ELF3 activity is reduced under elevated

temperatures ("temperature sensing") is not known. We recently used transgenic

experiments to demonstrate that ELF3 function is dependent on the unit copy number of

its C-terminal polyglutamine (polyQ) tract [35]. This domain is likely disordered, and

disordered domains evince structural changes in response to physical parameters such

as temperature [36]. Thermal remodeling of this polyQ tract is a plausible mechanism by

which ELF3 activity could be modulated through temperature. This polyQ tract also

shows substantial natural variation [35], potentially serving as a factor underlying natural

variation in thermoresponsive phenotypes. For example, in flies, variable repeats are

associated with local temperature compensation adaptations [37]. In short, the ELF3-

polyQ is an attractive candidate for adaptive variation in the ecologically relevant trait of

temperature response [38].

In this study, we used transgenic polyQ variants of ELF3 in two *A. thaliana*

genetic backgrounds to dissect the contribution of the polyQ tract to temperature

response. We show that polyQ repeat copy number modulates temperature sensing by

affecting overall ELF3 function. Surprisingly, we found that ELF3's role in

thermoresponsive flowering appears to be entirely independent of PIF4. We postulate

that ELF3's primary role in thermoresponsive flowering is PIF4-independent and occurs

through the photoperiodic pathway, and that this role is in turn dependent on the genetic

background.

RESULTS

The hypocotyl elongation temperature response is modulated by the ELF3 polyQ tract

affecting overall gene function.

Many recent studies noted the involvement of ELF3 in temperature-dependent

hypocotyl elongation [14–16,39], concluding that ELF3 protein activity is reduced under

elevated temperatures, thereby relieving ELF3 repression of *PIF4. PIF4* up-regulation

then leads to the observed hypocotyl elongation. We examined whether polyQ tract

variation in ELF3 in two backgrounds affects hypocotyl elongation at 27º (Fig. 1). We

previously showed that ELF3 polyQ variation has pleiotropic background-dependent

effects, with nonlinear associations between polyQ tract length and quantitative

phenotypes (including hypocotyl elongation at 22ºC; ref. 33). Certain variants (16Q for

Ws, >20Q for Col) generally complemented *elf3* null mutant phenotypes in Col and Ws

A. thaliana strains, whereas other variants complemented only specific phenotypes or

behaved as hypomorphs across all tested phenotypes. Here, we observed similar

trends for thermoresponsive hypocotyl elongation (Fig. 1). For example, in the Ws

background (Fig. 1A), the endogenous ELF3 variant (16Q) partially complements the

elf3 null mutant; another variant (9Q) fully complements the hypocotyl temperature

response. Other polyQ variants behaved as hypomorphs in Ws. In the Col background

(Fig. 1B), the endogenous 7Q variant, among other variants, failed to rescue the

response, agreeing with our previous observation that these transgenic lines are

hypomorphic in this background [35]. Deleting the entire polyQ tract eliminated

thermoresponsive hypocotyl elongation in both Col and Ws backgrounds. We next

addressed whether the observed phenotypic variation among polyQ variants was due to

variation in thermosensing or variation in general ELF3 function. We found that robust

thermal responses were strongly correlated with the overall functionality of each ELF3

variant in hypocotyl elongation (Fig. 1C), such that variants with intact thermal

responses exhibited short hypocotyls at 22ºC, whereas ELF3 variants with defective

thermal responses exhibited elongated hypocotyls regardless of temperature.

Furthermore, this ELF3 functionality effect is dependent on genetic background

(comparing for instance the 16Q and 20Q responses). Together, these results suggest

that the ELF3 polyQ tract controls repression of hypocotyl elongation regardless of

temperature, rather than sensing temperature specifically. Nonetheless, our transgenic

ELF3 polyQ lines remain informative as an allelic series of ELF3 function to understand

the role of ELF3 in the de-repression of PIF4, which is thought to underlie

thermomorphogenesis [14–16,39–41].

Figure 1

Fig. 1. Response to elevated temperature (27º, relative to 22º) among transgenic lines

expressing ELF3-polyQ variants. Mean response and error were estimated by

regression, based on two independently-generated transgenic lines for each genotype,

with n >= 30 seedlings of each genotype in each condition (Table S1). WT = Ws, *elf3* =

elf3 mutant+vector control, 0Q = *elf3* mutant+*ELF3* transgene lacking polyQ, etc. Error

bars indicate standard error of the mean. (A): Ws (Wassilewskija) strain background.

Lines are generated in an *elf3-4* background. (B): Response in the Col (Columbia) strain

background, lines were generated in an *elf3-200* background. In both (A) and (B), **:

Bonferroni-corrected $p < 0.01$, **: Bonferroni-corrected $p < 0.05$, . : Bonferroni-corrected

$p < 0.1$ in testing the interaction term (different response from WT, Ws or Col). (C):

Temperature response is a function of ELF3 functionality (repression of hypocotyl

elongation at 22º). Simple means of 22º hypocotyl length, regression estimates of

temperature response. PCC = Pearson correlation coefficient; p-value is from a Pearson

correlation test.

Expression of PIF4 and PIF4 targets as a function of temperature and ELF3.

To evaluate the hypothesis that the thermal response defects in the transgenic lines

was due to up-regulation of PIF4 and PIF4 targets, we measured transcript levels of

PIF4 and its target *AtHB2* in seedlings of selected lines from both backgrounds at 22ºC

and 27ºC (Fig. S1). Like others [15,16], we observed an inverse relationship between

ELF3 functionality and transcript levels of *PIF4* and *AtHB2*, with larger effects on *PIF4*

expression. The ELF3 lines with the strongest thermal response (*e.g.* 16Q in the Ws

background) showed the most robust de-repression of *PIF4* at elevated temperature.

However, *elf3* null mutants retained some *PIF4* up-regulation under these conditions,

especially in the Ws background. We conclude that ELF3-mediated de-repression of

PIF4 is involved in thermal responses as suggested by prior studies [15,16]; however,

de-repression of PIF4 and its targets may not be sufficient to explain the entirety of

thermal response defects in *elf3* null mutants.

ELF3 polyQ variation affects thermoresponsive adult morphology and flowering time.

Following the expectation that ELF3's thermal response acts through PIF4, we

reasoned that ELF3 should also play a role in other PIF4-dependent thermal responses.

One well-known response to elevated temperature is adult petiole elongation. *pif4*

mutants fail to show this response when grown at elevated temperatures [11]. We

measured petiole length in the ELF3 polyQ transgenic lines, expecting that, due to

general PIF4 de-repression, poorly-functioning ELF3 polyQ lines would show no

response (perhaps due to constitutively elongated petioles, similar to hypocotyls; Fig.

2). In stark contrast to this expectation, we found that all lines had a robust petiole

response to temperature (Fig. 2A, B). This effect was apparent in both Ws (Fig. 2A) and

Col backgrounds (Fig. 2B). Moreover, this response was actually accentuated in *elf3*

null mutants and in poorly-functioning ELF3 polyQ variants (Fig. 2A, B).

Figure 2

10   **Fig. 2.** Adult plant responses to elevated temperature (27º, relative to 22º) in long days

11   among transgenic lines expressing different ELF3-polyQ variants. (A) and (C):

12   Response in the Ws (Wassilewskija) strain background. Lines are in an *elf3-4*

13   background. (B) and (D): Response in the Col (Columbia) strain background, lines are

in an *elf3-200* background. (A) and (B) display PL:LL temperature response, (C) and (D)

display RLN temperature response. Average responses and errors were estimated in a

regression model accounting for variation between experiments (Table S2), based on

two to three independently-generated transgenic lines for each genotype. n >= 24 plants

overall for each genotype in each condition. PL:LL = petiole to leaf length ratio at 25

days post germination, RLN = rosette leaf number at flowering, WT = wild type, *elf3* =

elf3 mutant+vector control, 0Q = *elf3* mutant+*ELF3* transgene with entire polyglutamine

removed, etc. Error bars indicate standard error. In each case, **: Bonferroni-corrected

$p < 0.01$, **: Bonferroni-corrected $p < 0.05$, . : Bonferroni-corrected $p < 0.1$ in testing the

interaction term (different response from WT, Ws or Col).

Further, we measured flowering time in transgenic lines as the number of rosette

leaves at flowering (Fig. 2C, D). PIF4 is not required for the accelerated flowering

temperature response under longer photoperiods [11]. Hence, we expected that loss of

ELF3 function should not affect thermoresponsive flowering if ELF3's thermal signaling

role acts through PIF4. In contrast to this expectation, in the Col background, *elf3*

mutants had an abrogated flowering response to elevated temperature (Fig. 2D).

Moreover, most variants in the Col background entirely failed to rescue this phenotype,

and even the endogenous 7Q shows only a partial rescue. While *elf3* mutants show

abrogated flowering response to lower temperatures [8], it was not expected that a

similar role would extend to the elevated temperature flowering response, which is

usually considered to be dominated by PIF4 [9], gibberellin signaling [30], and other

transcriptional regulators of *FT* such as SVP [31].

Unlike Col, Ws lacks a robust flowering response to elevated temperature under

these conditions [42], and indeed, variants in the Ws background generally showed no

thermoresponsive flowering (Fig. 2C). Thus, ELF3 polyQ variation does not suffice to

enhance the negligible thermoresponsive flowering in the Ws background under these

conditions. In light of this data, the roles of ELF3 and PIF4 in the elevated temperature

response appear to be independent of one another under these experimental conditions

and for these traits. These results are intriguing, given that the PIF4 pathway is the

best-recognized mechanism for thermoresponsive flowering at high temperatures

[9,10,32,33]. Therefore, we suggest that ELF3 acts in a PIF4-independent pathway for

thermoresponsive flowering at high temperatures.

ELF3 regulates thermoresponsive flowering under long days, and is not required for

PIF4-dependent adult thermomorphogenesis.

Our results with ELF3-polyQ variants suggested that ELF3 dysfunction does not

meaningfully affect PIF4-dependent traits, but does affect PIF4-independent traits in

adult plants in long days. However, these results may be due to subtle differences in

conditions between our approach and those used by previous investigators. We

therefore directly addressed the relationship of ELF3 and PIF4 in adult

thermoresponsive phenotypes by growing *pif4* and *elf3* mutants with various thermal

treatments. Previous experiments with *pif4* mutants used different conditions from ours,

specifically a later transfer to elevated temperature [11]. Hence, it was possible that the

observed inconsistencies between *elf3* and *pif4* effects on adult thermoresponsive

phenotypes were a trivial consequence of experimental conditions. Specifically, the

effects of elevated temperature during the early seedling stages (the conditions we use)

may induce pathways irrelevant to treatments at later, vegetative stages. Thus, we

tested both transfer conditions under long days (Fig. 3). We found that the effect of

different experimental conditions is negligible, though the earlier 27ºC treatment showed

a slightly stronger morphological response (Fig. 3A, B). Thus, the timing of the 27ºC

treatment (early seedling vs. vegetative stage) does not substantially affect adult

thermoresponsive traits. Further, our results under long days were similar to previous

observations under continuous light [11], showing that PIF4 is essential for petiole

elongation (Fig. 3B), but dispensable for thermoresponsive flowering (Fig. 3C). Our

PIF4 results were in direct contrast to ELF3, which was dispensable for petiole

elongation (Fig. 3B), but essential for thermoresponsive flowering (Fig. 3C). These

results indicate the apparent independence of ELF3 and PIF4 in these specific

responses, and suggest that seedling thermomorphogenesis, adult

thermomorphogenesis, and thermoresponsive flowering constitute three independent

developmental responses.

Figure 3

Fig. 3. *elf3* and *pif4* null mutant phenotypes are independent under LD treatments and robust to conditions. (A), (B), and (C): 22º: constant 22º LD growth; 27º 14d: transfer from 22º to 27º at 14 days post-germination; 27º 1d: transfer from 22º to 27º at 1 day post-germination. (A): Col (WT), *elf3-200*, and *pif4-2* plants grown under long days with three different temperature regimes were photographed at 20 days post germination. Experiment was repeated with similar results. (B and D): Petiole elongation responses of the indicated genotypes, measured by ratio of petiole to whole leaf length at 25 days post germination. Regression analysis of data in Table S3. In each case, **: Bonferroni-corrected p < 0.01, **: Bonferroni-corrected p < 0.05, . : Bonferroni-corrected p < 0.1 in testing the interaction term (different response from Col).

One open question was whether the dispensability of ELF3 for petiole elongation reflected increased importance of other inputs to PIF4, such as FCA, which is involved

in PIF4-dependent thermoresponsive petiole elongation in 7-day-old seedlings [13]. We

therefore measured adult thermoresponsive petiole elongation in *fca* mutants (Fig.

S2A), and unexpectedly found no substantial difference between *fca* mutants and WT

Col. Regulatory rewiring across development may remove FCA and ELF3 as inputs to

PIF4-dependent thermomorphogenesis in 25-day-old adult plants.

A second question was whether loss of *ELF3* function can affect

thermoresponsive flowering in the Ws strain under other temperature conditions. We

therefore assayed flowering in Ws and the Ws-derived null mutant *elf3-4* at 16ºC and

22ºC (Fig. S2B). Under these conditions, Ws robustly accelerated flowering at 22ºC,

whereas *elf3-4* showed no perceptible difference in flowering between the two

temperatures. Thus, ELF3's role in thermoresponsive flowering is not restricted to the

Col strain or a certain temperature, but rather is necessary for whatever

thermoresponsive reaction norm a strain may have for flowering.

ELF3 and PIF4 regulate adult thermoresponsive phenotypes independently.

If ELF3 and PIF4 were independent in controlling thermal responses of adult

phenotypes under long days, then *elf3 pif4* double mutants would show approximately

additive phenotypes. We generated *elf3 pif4* double mutants and subjected them to the

same experiments as above. Our results indicated that flowering and petiole elongation

constitute independent temperature responses, with ELF3 controlling the former and

PIF4 controlling the latter in additive fashions (Fig. 4). That is, *elf3 pif4* double mutants

showed negligible thermoresponsive flowering like *elf3*, and a negligible petiole

response like *pif4*. Additionally, *elf3 pif4* flowered slightly later than *elf3* at 22º, while

maintaining a negligible thermal response in flowering, indicating that *elf3* mutants are

not simply restricted by a physiological limit of early flowering. The additivity of these

phenotypes establishes that, under these conditions, ELF3 and PIF4 likely operate in

separate thermal response pathways.

Figure 4

6 **Fig. 4.** Double mutant analysis confirms PIF4 and ELF3 independence in adult

7 temperature responses and non-redundancy of PIF4 with PIF5. (A): Col, *elf3-200*, *pif4-*

8 *2,* and *elf3-200 pif4-2* plants grown under long days with two different temperature

9 regimes were photographed at 25 days post germination. (B): Petiole elongation

10 responses of the indicated genotypes, measured by ratio of petiole to whole leaf length

at 25 days post germination. (C): Flowering temperature response of indicated

genotypes, measured by rosette leaf number (RLN) at flowering. (B) and (C): n > 8

plants for each genotype in each treatment. All "27°" plants were seeded and incubated

one day at 22° before transfer to 27°. Experiments were repeated with similar results.

Regression analysis of data reported in Tables S6 and S7. In each case, **: Bonferroni-

corrected $p < 0.01$, **: Bonferroni-corrected $p < 0.05$, . : Bonferroni-corrected $p < 0.1$ in

testing the interaction term (different response from Col).

Previous studies have indicated that other members of the PIF family have

negligible or minor (in the case of PIF5) roles in these same thermal response

phenotypes [11,27,43]. For instance, under SD, *pif1 pif3 pif4 pif5* mutants behave

essentially identical to *pif4 pif5* mutants in flowering response and *FT* expression, which

in turn show only a very slight abrogation of these responses relative to *pif4* mutants

[31]. *pif4 pif5* double mutants do show slightly abrogated thermoresponsive flowering

under 12 hour light : 12 hour dark photoperiods relative to single mutants [29], similar to

other thermally responsive phenotypes [11,29–31]. These previous findings, combined

with the completely intact flowering response of *pif4* mutants, suggest that redundancy

between PIFs plays little meaningful role in this response. However, to directly address

this possibility, we evaluated thermoresponsive flowering in *pif5* and *pif4 pif5* mutants

(Fig. 4D), because PIF5 is most often considered to act redundantly with PIF4

[20,29,31,44], and the only other PIF to show any small contribution to

thermoresponsive flowering [29–31]. As expected, both *pif5* single mutants and *pif4 pif5*

double mutants demonstrate intact thermoresponsive flowering. These observations

indicate that redundancy with other PIFs is not responsible for the apparent

independence of PIF4 and ELF3. Notably, petiole elongation at elevated temperatures

is equally disrupted in *pif4* and *pif4 pif5* mutants, but intact in *pif5* single mutants (Figure

S3), reproducing the known dependence of this trait upon PIF4 alone [11].

Consequently, our results support the previously-suggested dominance of

thermomorphogenesis by PIF4 rather than other PIFs, and the irrelevance of PIF4 (and

most likely other PIFs as well) to thermoresponsive flowering under LD.

Overall, the strong photoperiod-dependence of PIF4-related thermoresponsive

flowering necessitates the existence of some pathway or pathways independent of PIF4

under long days, given the persistence of the phenomenon under these conditions.

Based on our data, ELF3 acts in one such pathway.

Thermoresponsive flowering under long days can operate through the photoperiodic

pathway.

ELF3 operates in thermoresponsive flowering at low ambient temperatures via the

photoperiodic pathway, through repressing *GI* expression, after which GI in turn directly

activates FT [45,46]. To evaluate whether this pathway might explain our results, we

measured transcript levels of *GI* and *CO* in wild-type and *elf3* mutants under 22ºC and

27ºC (Fig. 5A). We found that *GI* is strongly up-regulated in *elf3* null mutants of Col and

Ws backgrounds, confirming previous reports in Col [39,46]. Further, wild-type Ws

showed higher basal *GI* levels compared to Col, which did not increase at higher

temperatures. In contrast, Col showed very low basal *GI* levels that increased at higher

temperatures to approximately the same levels as Ws. *CO* levels, however, were not

substantially increased by either *elf3* mutation or increased temperature, consistent with

previous reports [8,46]. Thus, robust thermoresponsive flowering was correlated with

low basal levels of *GI,* and with temperature-dependent *GI* up-regulation, as observed in

Col. The ELF3-dependent thermal responsiveness of *GI* expression confirms previous

reports [15], though the among-strain variation in responsiveness appears to be novel

and correlated specifically with flowering induction (but not hypocotyl or petiole

elongation, Figures 1 and 2). High basal *GI* levels in Ws may be associated with other

thermoresponsive deficiencies at high temperatures in this strain [42,47,48]. These

observations support the model under which ELF3 acts in the photoperiodic pathway to

engender thermoresponsive flowering, just as it does in response to lower ambient

temperatures [8,46].

We attempted to measure *FT* transcript levels in these samples, expecting that

they would be elevated in the early-flowering *elf3* and 27ºC conditions (Figure S4).

Unexpectedly, we observed that, while *FT* levels may increase slightly in the *elf3*

mutants, *FT* appears dramatically down-regulated in all 27ºC samples. This finding is

difficult to interpret in light of the phenotypic data, as most models of thermoresponsive

flowering agree that signaling operates through *FT* [7–9,29–31], suggesting rather that

these 7-day-old seedlings may be too young to measure physiologically relevant *FT*

expression, at least under 27ºC conditions.

Figure 5

Fig. 5. ELF3 and GI regulate thermoresponsive flowering. (A): Temperature-responsive

expression of photoperiodic pathway components at ZT0. Expression of each gene is

quantified relative to levels in Col at 22° (Col 22 = 1.0). Error bars represent SEM

across three biological replicates. *elf3-4*: *elf3* null in Ws background; *elf3-200*: *elf3* null

in Col background. (B): Thermoresponsive flowering in various flowering mutants. LD

RLN = rosette leaf number at flowering under long days. * : interaction term for

genotype by environment at Bonferroni-corrected p < 0.05; details of regression model

in Table S9. (C) Thermoresponsive petiole elongation in various flowering mutants. For

(B) and (C), n >= 8 plants of each genotype in each condition; white boxes indicate

measurements at 22º, red boxes indicate measurements at 27º. *gi: gi-2*, *co*: *co-101*,

spy: *spy-3*, *soc1*: *soc1* T-DNA insertion, *elf3: elf3-200*. This experiment was repeated

with similar results. (D): Models of thermoresponsive flowering under long and short

photoperiods. Dashed edges indicate speculated temperature sensing mechanisms.

Edges with increased weight indicate relative increases of influence between conditions.

Pathways are indicated, along with other important actors reported elsewhere.

If the photoperiodic pathway contributes to thermoresponsive flowering at

elevated ambient temperatures in long days (LD), we would expect mutants in this

pathway to show abrogated thermal responses, as they do under short days (SD), along

with members of the autonomous pathway [7]. These two pathways also contribute

independently to thermoresponsive flowering at low temperatures (16ºC vs. 23ºC) [6,8].

Altogether, we would expect that a photoperiodic thermoresponsive flowering pathway

would operate independently of both PIF4 and the autonomous pathways in long days.

It is not clear whether the autonomous pathway would be independent of PIF4, given

known interactions between FCA and PIF4 [13].

To evaluate whether these past results under other conditions also apply to long

days and elevated temperatures, we measured flowering time at 22ºC and 27ºC in

mutants in the photoperiodic pathway (*gi*, *co,* Fig. 5B). We also tested mutants of the

gibberellin pathway (*spy*), and a terminal floral integrator (*soc1*), which we do not expect

to be necessary for thermoresponsive flowering. We found robust thermal responses in

all mutants except *elf3* and *gi*, similar to previous results under different conditions

[7,8,45,46]. These results emphasize once again that differences in thermoresponsive

flowering are not generalizable between photoperiods, as it has recently been shown

that *co* mutants have a partial flowering acceleration defect under SD. These results

implicate GI (but not CO) as an actor in thermoresponsive flowering at elevated

temperatures. Collectively, these experiments suggest that the photoperiod pathway is

necessary in promoting thermoresponsive flowering in long days, and expression data

in this and other studies suggests that ELF3 is likely to act within this pathway.

DISCUSSION

ELF3 and PIF4 are both crucial integrators of temperature and light signaling in

controlling *A. thaliana* development. Recent literature has emphasized the centrality of

PIF4-dependent thermoresponsive regulation in a variety of phenotypes, including in

flowering [9,10,32]. Here, we show that PIF4 is dispensable for thermoresponsive

flowering under long photoperiod conditions [11], and that ELF3 is essential for

thermoresponsive flowering under these conditions. Our results integrate previous

knowledge about thermoresponsive flowering, and identify at least one pathway for this

response that does not involve PIF4. Moreover, we show that while polyQ variation in

ELF3 affects ELF3 function, the polyQ tract is unlikely a temperature-responsive

component in itself. Our results allow us to integrate the many disparate findings of

current studies into classic models of thermal responses in *A. thaliana*, allowing a

comprehensive view of the genetic underpinnings of this agronomically crucial plant

trait.

ELF3 polyglutamine variation appears to affect thermoresponsive traits by modulating

overall ELF3 activity.

In previous work, we demonstrated that polyQ variation in ELF3 is (i) common, (ii)

affects many known ELF3-dependent phenotypes, and (iii) is dependent on the genetic

background [35]. Following the recent discoveries that ELF3 is involved with thermal

response [14–16], we confirmed that ELF3 polyQ variation also affects thermal

response phenotypes in a background-dependent fashion. However, we found little

support for the hypothesis that the polyQ tract has a special role in temperature

sensing. Instead, as was the case for other ELF3-dependent phenotypes, ELF3 polyQ

variation appeared to affect overall ELF3 functionality, with less functional ELF3 variants

lacking robust temperature responses. However, a more exhaustive series of polyQ

variants may be required for revealing polyQ-specific effects, in particular because the

molecular mechanism(s) by which polyQ variation affects ELF3 functionality remain

unknown.

ELF3-PIF4 relationship in thermomorphogenesis.

One question that remains unanswered is to what extent ELF3 participates in PIF4-

dependent thermoresponsive morphologies. While our study and previous work

[14,16,39] support a PIF4-ELF3 link in thermoresponsive hypocotyl elongation, this

relationship disappears in the analogous case of thermoresponsive petiole elongation.

These results can be explained by many hypotheses. For instance, it is possible that

ELF3 regulation of PIF4 is only relevant at the early seedling stage. Another possible

hypothesis is that ELF3 regulation of PIF4 in some instances is sufficient but not

necessary for thermal responses. More studies are needed to understand the

mechanistic details of the ELF3 and PIF4 relationship in thermomorphogenesis.

Natural variation in temperature response.

Several studies have found that different *A. thaliana* strains respond to temperature

differently, either shifting or inverting the reaction norm of the phenotype in question

[42,47,48]. Ws has a shifted reaction norm with respect to temperature compared to Col

for photoperiod-related phenotypes, including flowering. For instance, Ws displays

accelerated flowering at 23ºC vs. 16ºC [42], but accelerates flowering no further at

27ºC. Here, we show that this acceleration requires ELF3, like the elevated temperature

acceleration in Col. Another example of differential mutational effects among strains is

that *gi* mutants in the L*er* background display robust thermoresponsive flowering [6,7]. It

is unclear whether this finding is due to altered wiring of pathways between these

backgrounds.

Thermoresponsive flowering requires either PIF4 or ELF3, depending on photoperiod.

Under various conditions, both ELF3 and PIF4 have been found to be crucial for

thermoresponsive flowering. Other members of the autonomous and the photoperiodic

pathways have also been implicated in thermoresponsive flowering [6–8] (besides other

pathways, [49]). Consequently, some combination of these pathways, modulated by

experimental conditions, must require ELF3 and/or PIF4. We and others [11,29] have

observed that PIF4 and its paralogs are not required for proper thermoresponsive

flowering under longer photoperiods. Furthermore, we and others [8,46] have shown

that ELF3 and the photoperiod pathway (excluding CO) are essential for proper

thermoresponsive flowering under long days. It has been previously shown that PIF4

and the photoperiodic pathway contribute to thermoresponsive flowering via

independent pathways [9], suggesting that under longer photoperiods PIF4 activity is

inhibited, allowing other mechanisms to dominate thermoresponsive flowering.

We propose a model of thermoresponsive flowering, in which PIF4, ELF3, the

photoperiodic pathway, and other pathways interact depending upon condition and

genetic background (Fig. 5D). Under short days or other short photoperiods, phyB

activity is down-regulated, leading to up-regulation of PIF4 [22,50–52], which at high

levels occupies the promoter of the flowering integrator *FT* and induces flowering [9].

However, under longer photoperiods, phyB up-regulation leads to an attenuation of

PIF4 activity, and consequently the role of PIF4 and other PIFs becomes negligible [11].

This allows canonical ambient temperature responses (such as the photoperiodic

pathway, including ELF3, [8,46]) to take a dominant role in thermoresponsive flowering.

Constitutive overexpression of PIF4, PIF5, and PIF3 under long day conditions induces

early flowering [30], supporting the hypothesis that differences in PIF levels underlie the

photoperiod-dependence of PIF4's role. We have not formally excluded the possibility

that members of the large PIF family other than PIF4 and PIF5 might contribute to the

phenotype; however, there is no evidence at present to suggest that they might

[11,27,30,31]. Several reports have indicated that GI and COP1, but not CO, are

involved in thermoresponsive flowering [7,8,46], with GI directly binding the *FT* promoter

[46]. Under each of these conditions, *FT*-induced flowering is activated by a different

signaling cascade. This interpretation leads to a coherent view of how light and

temperature responses are integrated in this important plant trait.

To summarize, at least three independent mechanisms have been described that

promote thermoresponsive flowering in any context. These include the photoperiodic

pathway (PHYB/ELF3/GI/COP1), the autonomous pathway

(PHYA/FCA/FVE/TFL1/FLC), and the PIF4-dependent pathway

(PIF4/H2A.Z/gibberellin), all of which converge by regulating *FT* (although the last

pathway may also act through other integrators [29,30]). Our results showing a down-

regulation of *FT* at elevated temperatures in young seedlings may also hold clues

regarding the interaction of these pathways, but are difficult to interpret in the context of

the other evidence. The collective results of our experiments and previous work suggest

that the first two pathways are necessary but not sufficient for thermoresponsive

flowering, and that the third (PIF4) is sufficient but not necessary for thermoresponsive

flowering. Further study will be necessary in understanding the interdependencies of the

three pathways. For instance, it has been suggested that PIF4 binding to the *FT*

promoter is dependent on cooperativity with a second photoperiod-controlled actor [34].

In conclusion, we observe that ELF3 is involved in the hypocotyl response to

elevated temperature as reported previously, and that this response can be abrogated

by poorly-functioning ELF3 polyQ variants. We further demonstrate that ELF3 has little

effect on the petiole temperature response, and is necessary for the flowering

temperature response, suggesting that it functions independently of PIF4, potentially in

the photoperiodic pathway. These results reiterate the complexity of these crucial

environmental responses in plants, and can serve as a basis for further development of

our understanding of how plants respond to elevated temperatures. In the context of

climatic changes, this understanding will serve those attempting to secure the global

food supply.

MATERIALS AND METHODS

Plant materials and growth conditions. All mutant lines (except *pif4-2 elf3-200*) were

either described previously or obtained as T-DNA insertions from the Arabidopsis

Biological Resources Center at Ohio State University [53,54], and are described in

Table S11. *pif4-2 elf3-200* was obtained via crossing and genotyping. T-DNA insertions

were confirmed with primers described in Table S10. For hypocotyl assays, seedlings

were grown for 15d in incubators set to SD (8h light : 16h dark days, with light supplied

at 100 µmol· $m^{-2}·s^{-1}$ by cool white fluorescent bulbs) on vertical plates as described

previously [35]. All plates were incubated at 22º for one day, after which one replicate

arm was transferred to an incubator set to 27º, with another replicate arm maintained at

22º. For flowering time assays, plants were stratified 3-5d at 4º in 0.1% agarose and

seeded into Sunshine #4 soil in 36-pot or 72-pot flats to germinate at 22º under LD (16h

light : 8 hr dark days, with light supplied at 100 µmol·$m^{-2}·s^{-1}$ by cool white fluorescent

bulbs). Replicate arms were subsequently transferred to 27º LD conditions as indicated,

with others remaining at 22º. Different temperature treatments of the same experiment

were identical with respect to randomization, setup, and format. At 25d, petiole length

and whole leaf length (including petiole) of the third leaf were measured, and the ratio of

these values was further analyzed. Flowering was defined as an inflorescence ≥1cm

tall; at this point, date and rosette leaf number were recorded.

Trait data analysis. All data analysis was performed using R v3.2.1 [55]. Where

indicated, temperature responses were modeled using multiple regression in the form

Phenotype ~ μ + β_G*Genotype* + β_T*Temperature* + β_{GxT}(*Genotype* x *Temperature*) +

β_E*Experiment* + *Error*. All experiments were included in models for transgenic

experiments, and thus the β_E term describes systematic variation between experiments,

whereas line-specific effects among transgenics should be modeled in the error term. Where temperature responses are reported, they consist of the $\beta_T + \beta_{GxT}$ terms and associated errors ($\sqrt{\sigma_T^2 + \sigma_{GxT}^2}$) where σ_T is the standard error for β_T and σ_{GxT} is the standard error for β_{GxT}), and thus are corrected for systematic experimental variation and temperature-independent genotype effects. Where p-values are reported for GxT interaction effects, they have been subjected to a Bonferroni correction to adjust for multiple comparisons. Analysis scripts and data are provided at

https://figshare.com/articles/elf3_pif4_data_code_v2/3398353.

Gene expression analyses. Seedlings were grown for 1d under LD at 22°, after which one replicate arm was transferred to LD at 27°, with another replicate arm remaining at 22°, and all seedlings were harvested 6d later at indicated times. At harvest, ~30mg aerial tissue of pooled seedlings was flash-frozen immediately in liquid nitrogen and stored at -80°. RNA extraction, cDNA synthesis, and real-time quantitative PCR were performed as described previously [35], using primers in Table S10. Transcript levels were quantified from technical triplicate across at least two biological replicates using the $\Delta\Delta C_t$ method, assuming 100% primer efficiency [56].

ACKNOWLEDGMENTS

We thank Philip Wigge and Jaehoon Jung for ideas, helpful conversations, sharing unpublished data, and comments on this manuscript. We thank Evan Eichler for use of the LightCycler instrument. We thank members of the Queitsch lab for helpful discussions. This work was supported by National Institutes of Health New Innovator Award DP2OD008371 to CQ.

REFERENCES

1.   Battisti DS, Naylor RL (2009) Historical warnings of future food insecurity with unprecedented seasonal heat. Science 323: 240–244. doi:10.1126/science.1164363.

2.   Quint M, Delker C, Franklin KA, Wigge PA, Halliday KJ, et al. (2016) Molecular and genetic control of plant thermomorphogenesis. Nat Plants 2: 15190. doi:10.1038/nplants.2015.190.

3.   Gould PD, Locke JCW, Larue C, Southern MM, Davis SJ, et al. (2006) The molecular basis of temperature compensation in the Arabidopsis circadian clock. Plant Cell 18: 1177–1187. doi:10.1105/tpc.105.039990.

4.   Edwards KD, Anderson PE, Hall A, Salathia NS, Locke JCW, et al. (2006) FLOWERING LOCUS C mediates natural variation in the high-temperature response of the Arabidopsis circadian clock. Plant Cell 18: 639–650. doi:10.1105/tpc.105.038315.

5.   Thines B, Harmon FG (2010) Ambient temperature response establishes ELF3 as a required component of the core Arabidopsis circadian clock. Proc Natl Acad Sci U S A 107: 3257–3262. doi:10.1073/pnas.0911006107.

6.   Blázquez MA, Ahn JH, Weigel D (2003) A thermosensory pathway controlling flowering time in Arabidopsis thaliana. Nat Genet 33: 168–171. doi:10.1038/ng1085.

7.   Balasubramanian S, Sureshkumar S, Lempe J, Weigel D (2006) Potent induction of Arabidopsis thaliana flowering by elevated growth temperature. PLoS Genet 2: e106. doi:10.1371/journal.pgen.0020106.

8.   Strasser B, Alvarez MJ, Califano A, Cerdán PD (2009) A complementary role for ELF3 and TFL1 in the regulation of flowering time by ambient temperature. Plant J 58: 629–640. doi:10.1111/j.1365-313X.2009.03811.x.

9.   Kumar SV, Lucyshyn D, Jaeger KE, Alós E, Alvey E, et al. (2012) Transcription factor PIF4 controls the thermosensory activation of flowering. Nature 484: 242–245. doi:10.1038/nature10928.

10.   Song YH, Ito S, Imaizumi T (2013) Flowering time regulation: photoperiod- and temperature-sensing in leaves. Trends Plant Sci 18: 575–583. doi:10.1016/j.tplants.2013.05.003.

11.   Koini MA, Alvey L, Allen T, Tilley CA, Harberd NP, et al. (2009) High temperature-mediated adaptations in plant architecture require the bHLH transcription factor PIF4. Curr Biol 19: 408–413. doi:10.1016/j.cub.2009.01.046.

12. Johansson H, Jones HJ, Foreman J, Hemsted JR, Stewart K, et al. (2014) Arabidopsis cell expansion is controlled by a photothermal switch. Nat Commun 5: 4848. doi:10.1038/ncomms5848.

13. Lee H-J, Jung J-H, Cortés Llorca L, Kim S-G, Lee S, et al. (2014) FCA mediates thermal adaptation of stem growth by attenuating auxin action in Arabidopsis. Nat Commun 5: 5473. doi:10.1038/ncomms6473.

14. Box MS, Huang BE, Domijan M, Jaeger KE, Khattak AK, et al. (2014) ELF3 Controls Thermoresponsive Growth in Arabidopsis. Curr Biol 25: 194–199. doi:10.1016/j.cub.2014.10.076.

15. Mizuno T, Nomoto Y, Oka H, Kitayama M, Takeuchi A, et al. (2014) Ambient temperature signal feeds into the circadian clock transcriptional circuitry through the EC night-time repressor in Arabidopsis thaliana. Plant Cell Physiol 55: 958–976. doi:10.1093/pcp/pcu030.

16. Raschke A, Ibañez C, Ullrich KK, Anwer MU, Becker S, et al. (2015) Natural variants of ELF3 affect thermomorphogenesis by transcriptionally modulating PIF4-dependent auxin response genes. BMC Plant Biol 15: 197. doi:10.1186/s12870-015-0566-6.

17. Nusinow DA, Helfer A, Hamilton EE, King JJ, Imaizumi T, et al. (2011) The ELF4-ELF3-LUX complex links the circadian clock to diurnal control of hypocotyl growth. Nature 475: 398–402. doi:10.1038/nature10182.

18. Nieto C, López-Salmerón V, Davière J-M, Prat S (2014) ELF3-PIF4 Interaction Regulates Plant Growth Independently of the Evening Complex. Curr Biol 25: 187–193. doi:10.1016/j.cub.2014.10.070.

19. Nozue K, Covington MF, Duek PD, Lorrain S, Fankhauser C, et al. (2007) Rhythmic growth explained by coincidence between internal and external cues. Nature 448: 358–361. doi:10.1038/nature05946.

20. Filo J, Wu A, Eliason E, Richardson T, Thines BC, et al. (2015) Gibberellin driven growth in elf3 mutants requires PIF4 and PIF5. Plant Signal Behav. doi:10.4161/15592324.2014.992707.

21. Lorenzo CD, Sanchez-Lamas M, Antonietti MS, Cerdán PD (2016) Emerging Hubs in Plant Light and Temperature Signaling. Photochem Photobiol 92: 3–13. doi:10.1111/php.12535.

22. Lorrain S, Allen T, Duek PD, Whitelam GC, Fankhauser C (2008) Phytochrome-mediated inhibition of shade avoidance involves degradation of growth-promoting bHLH transcription factors. Plant J 53: 312–323. doi:10.1111/j.1365-313X.2007.03341.x.

23. McWatters HG, Bastow RM, Hall A, Millar AJ (2000) The ELF3 zeitnehmer regulates light signalling to the circadian clock. Nature 408: 716–720. doi:10.1038/35047079.

24. Covington MF, Panda S, Liu XL, Strayer CA, Wagner DR, et al. (2001) ELF3 modulates resetting of the circadian clock in Arabidopsis. Plant Cell 13: 1305–1315.

25. Zagotta MT, Hicks KA, Jacobs CI, Young JC, Hangarter RP, et al. (1996) The Arabidopsis ELF3 gene regulates vegetative photomorphogenesis and the photoperiodic induction of flowering. Plant J 10: 691–702. doi:10.1046/j.1365-313X.1996.10040691.x.

26. Leivar P, Quail PH (2011) PIFs: pivotal components in a cellular signaling hub. Trends Plant Sci 16: 19–28. doi:10.1016/j.tplants.2010.08.003.

27. Proveniers MCG, van Zanten M (2013) High temperature acclimation through PIF4 signaling. Trends Plant Sci 18: 59–64. doi:10.1016/j.tplants.2012.09.002.

28. Bai M-Y, Shang J-X, Oh E, Fan M, Bai Y, et al. (2012) Brassinosteroid, gibberellin and phytochrome impinge on a common transcription module in Arabidopsis. Nat Cell Biol 14: 810–817. doi:10.1038/ncb2546.

29. Thines BC, Youn Y, Duarte MI, Harmon FG (2014) The time of day effects of warm temperature on flowering time involve PIF4 and PIF5. J Exp Bot 65: 1141–1151. doi:10.1093/jxb/ert487.

30. Galvão VC, Collani S, Horrer D, Schmid M (2015) Gibberellic acid signaling is required for ambient temperature-mediated induction of flowering in Arabidopsis thaliana. Plant J 84: 949–962. doi:10.1111/tpj.13051.

31. Fernández V, Takahashi Y, LeGourrierec J, Coupland G (2016) Photoperiodic and thermosensory pathways interact through CONSTANS to promote flowering at high temperature under short days. Plant J. doi:10.1111/tpj.13183.

32. Wigge PA (2013) Ambient temperature signalling in plants. Curr Opin Plant Biol 16: 661–666. doi:10.1016/j.pbi.2013.08.004.

33. Liu J, Feng L, Li J, He Z (2015) Genetic and epigenetic control of plant heat responses. Front Plant Sci 06. doi:10.3389/fpls.2015.00267.

34. Seaton DD, Smith RW, Song YH, MacGregor DR, Stewart K, et al. (2015) Linked circadian outputs control elongation growth and flowering in response to photoperiod and temperature. Mol Syst Biol 11: 776. doi:10.15252/msb.20145766.

35. Undurraga SF, Press MO, Legendre M, Bujdoso N, Bale J, et al. (2012) Background-dependent effects of polyglutamine variation in the Arabidopsis thaliana gene ELF3. Proc Natl Acad Sci U S A 109: 19363–19367. doi:10.1073/pnas.1211021109.

36. Uversky VN (2009) Intrinsically disordered proteins and their environment: effects of strong denaturants, temperature, pH, counter ions, membranes, binding partners, osmolytes, and macromolecular crowding. Protein J 28: 305–325. doi:10.1007/s10930-009-9201-4.

37. Sawyer LA (1997) Natural Variation in a Drosophila Clock Gene and Temperature Compensation. Science (80-) 278: 2117–2120. doi:10.1126/science.278.5346.2117.

38. Gemayel R, Vinces MD, Legendre M, Verstrepen KJ (2010) Variable tandem repeats accelerate evolution of coding and regulatory sequences. Annu Rev Genet 44: 445–477. doi:10.1146/annurev-genet-072610-155046.

39. Mizuno T, Kitayama M, Oka H, Tsubouchi M, Takayama C, et al. (2014) The EC night-time repressor plays a crucial role in modulating circadian clock transcriptional circuitry by conservatively double-checking both warm-night and night-time-light signals in a synergistic manner in Arabidopsis thaliana. Plant Cell Physiol 55: 2139–2151. doi:10.1093/pcp/pcu144.

40. Nomoto Y, Nomoto Y, Kubozono S, Miyachi M, Yamashino T, et al. (2012) A circadian clock- and PIF4-mediated double coincidence mechanism is implicated in the thermosensitive photoperiodic control of plant architectures in Arabidopsis thaliana. Plant Cell Physiol 53: 1965–1973. doi:10.1093/pcp/pcs141.

41. Mizuno T, Kitayama M, Takayama C, Yamashino T (2015) Insight into a Physiological Role for the EC Night-Time Repressor in the Arabidopsis Circadian Clock. Plant Cell Physiol 56: 1738–1747. doi:10.1093/pcp/pcv094.

42. Ibañez C, Poeschl Y, Peterson T, Bellstädt J, Denk K, et al. (2015) Developmental plasticity of Arabidopsis thaliana accessions across an ambient temperature range. bioRxiv: 017285. doi:10.1101/017285.

43. Stavang JA, Gallego-Bartolomé J, Gómez MD, Yoshida S, Asami T, et al. (2009) Hormonal regulation of temperature-induced growth in Arabidopsis. Plant J 60: 589–601. doi:10.1111/j.1365-313X.2009.03983.x.

44. Sun J, Qi L, Li Y, Zhai Q, Li C (2013) PIF4 and PIF5 transcription factors link blue light and auxin to regulate the phototropic response in Arabidopsis. Plant Cell 25: 2102–2114. doi:10.1105/tpc.113.112417.

45. Sawa M, Kay SA (2011) GIGANTEA directly activates Flowering Locus T in Arabidopsis thaliana. Proc Natl Acad Sci U S A 108: 11698–11703. doi:10.1073/pnas.1106771108.

46. Jang K, Gil Lee H, Jung S-J, Paek N-C, Joon Seo P (2015) The E3 Ubiquitin Ligase COP1 Regulates Thermosensory Flowering by Triggering GI Degradation in Arabidopsis. Sci Rep 5: 12071. doi:10.1038/srep12071.

47. Edwards KD, Lynn JR, Gyula P, Nagy F, Millar AJ (2005) Natural allelic variation in the temperature-compensation mechanisms of the Arabidopsis thaliana circadian clock. Genetics 170: 387–400. doi:10.1534/genetics.104.035238.

48. Edwards KD, Guerineau F, Devlin PF, Millar AJ (2015) Low-temperature-specific effects of PHYTOCHROME C on the circadian clock in Arabidopsis suggest that PHYC underlies natural variation in biological timing. bioRxiv: 030577. doi:10.1101/030577.

49. Lee JH, Yoo SJ, Park SH, Hwang I, Lee JS, et al. (2007) Role of SVP in the control of flowering time by ambient temperature in Arabidopsis. Genes Dev 21: 397–402. doi:10.1101/gad.1518407.

50. Duek PD, Fankhauser C (2005) bHLH class transcription factors take centre stage in phytochrome signalling. Trends Plant Sci 10: 51–54. doi:10.1016/j.tplants.2004.12.005.

51. Huq E, Quail PH (2002) PIF4, a phytochrome-interacting bHLH factor, functions as a negative regulator of phytochrome B signaling in Arabidopsis. EMBO J 21: 2441–2450. doi:10.1093/emboj/21.10.2441.

52. Luo Q, Lian H-L, He S-B, Li L, Jia K-P, et al. (2014) COP1 and phyB Physically Interact with PIL1 to Regulate Its Stability and Photomorphogenic Development in Arabidopsis. Plant Cell 26: 2441–2456. doi:10.1105/tpc.113.121657.

53. Alonso JM, Stepanova AN, Leisse TJ, Kim CJ, Chen H, et al. (2003) Genome-wide insertional mutagenesis of Arabidopsis thaliana. Science 301: 653–657. doi:10.1126/science.1086391.

54. Kleinboelting N, Huep G, Kloetgen A, Viehoever P, Weisshaar B (2012) GABI-Kat SimpleSearch: new features of the Arabidopsis thaliana T-DNA mutant database. Nucleic Acids Res 40: D1211–D1215. doi:10.1093/nar/gkr1047.

55. R Core Team (2016) R: A Language and Environment for Statistical Computing.

56. Pfaffl MW (2001) A new mathematical model for relative quantification in real-time RT-PCR. Nucleic Acids Res 29: e45.

Supporting Information Captions

Fig. S1. Expression analysis of *PIF4* and *AtHB2* depends on temperature, genetic background, and ELF3 functionality. Error bars represent the standard deviation across two biological replicates. White bars represent 22º expression, red bars 27º expression for each line. Tissue was collected from 7d seedlings at ZT0.

Fig. S2. Regulation of adult thermoresponsive traits by ELF3 and FCA is independent of PIF4 and modulated by genetic background. Flowering temperature response of indicated genotypes under indicated conditions, measured by petiole length to leaf length ratio at 25 days or rosette leaf number (RLN) at flowering. For each experiment, n > 10 plants for each genotype in each treatment. Regression analysis of data in Tables S4 and S5.

Fig. S3. Regulation of adult thermoresponsive petiole elongation traits occurs principally through PIF4. Petiole elongation temperature response of indicated genotypes under indicated conditions, measured by ratio of petiole length to leaf length at 25d. For each experiment, n > 10 plants for each genotype in each treatment. This experiment was repeated with similar results. Regression analysis of data in Table S8.

Fig. S4. Expression of *FT* in 7d seedlings responds to temperature and *elf3* status. White bars represent 22º expression, red bars 27º expression for each line.

Tissue was collected from 7d seedlings at ZT0. Error bars indicate SEM across three biological replicates.

Table S1. Regression analysis of hypocotyl elongation temperature response among Col and Ws transgenic lines.

Table S2. Regression analysis of petiole : leaf length ratio and rosette leaf number at flowering temperature response among Col and Ws transgenic lines.

Table S3. Regression analysis of rosette leaf number at flowering and petiole : leaf length ratio temperature responses in *elf3* and *pif4*.

Table S4. Regression analysis of rosette leaf number at flowering temperature response in Ws and *elf3-4*.

Table S5. Regression analysis of petiole : leaf length ratio temperature response in Col and *fca* mutants.

Table S6. Regression analysis of rosette leaf number at flowering temperature response in *elf3 pif4* double mutants.

Table S7. Regression analysis of rosette leaf number at flowering temperature response in *pif4 pif5* double mutants.

Table S8. Regression analysis of the petiole elongation temperature response in *pif4 pif5* double mutants.

Table S9. Regression analysis of rosette leaf number at flowering and petiole : leaf length ratio temperature responses in flowering pathway mutants.

Table S10. Primers used in this study.

Table S11. Mutant lines used in this study.