1 **Title** (88 chars < 120): Punctuated evolution and transitional hybrid network in an
2 ancestral cell cycle of fungi
3

4 **Authors**: Edgar M. Medina[1,2], Jonathan J. Turner[3], Jan M. Skotheim[3], and Nicolas E.
5 Buchler[1,2]
6
7 [1] Department of Biology, Duke University, Durham, NC, 27708, USA
8 [2] Center for Genomic and Computational Biology, Duke University, Durham, NC,
9 27710, USA
10 [3] Department of Biology, Stanford University, Stanford, CA, 94305, USA
11
12 **Contact:** nicolas.buchler@duke.edu

13
14 **Abstract:** (130 words < 150)
15

Although cell cycle control is an ancient, conserved, and essential process, some core

animal and fungal cell cycle regulators share no more sequence identity than non-

homologous proteins. Here, we show that evolution along the fungal lineage was

punctuated by the early acquisition and entrainment of the SBF transcription factor.

Cell cycle evolution in the fungal ancestor then proceeded through a hybrid network

containing both SBF and its ancestral animal counterpart E2F, which is still

maintained in many basal fungi. We hypothesize that a viral SBF may have initially

hijacked cell cycle control by activating transcription via the *cis*-regulatory elements

targeted by the ancestral cell cycle regulator E2F, much like extant viral oncogenes.

Consistent with this hypothesis, we show that SBF can regulate promoters with E2F

binding sites in budding yeast.

Impact statement (15-30 words): Cell cycle network evolution in a fungal ancestor

was punctuated by arrival of a viral DNA-binding protein that was permanently

incorporated into the G1/S regulatory network controlling cell cycle entry.

Keywords: cell cycle / evolution / fungi / chytrid / virus

INTRODUCTION

The networks regulating cell division in yeasts and animals are highly similar in both

physiological function and network structure (Figure 1) (Cross et al., 2011; Doonan

and Kitsios, 2009). For example, the cell cycle controls proliferation in response to a

variety of internal and external signals during the G1 phase, between cell division

and DNA replication. These input signals, including cell growth, are integrated into a

gradual increase in cyclin dependent kinase (Cdk) activity, which triggers a feedback

loop at the basis of the all-or-none irreversible decision to proliferate (Bertoli et al.,

2013).

Many of the molecular mechanisms underlying G1 regulation are highly conserved.

In animal cells, Cyclin D, in complex with either Cdk4 or Cdk6, initiates cell cycle

entry by phosphorylating the retinoblastoma protein, pRb. This begins the

inactivation of pRb and the concomitant activation of the E2F transcription factors

that induce transcription of downstream cyclins E and A, which complete the

inhibition of pRb thereby forming a positive feedback loop (Bertoli et al., 2013).

Similarly, in budding yeast, the G1 cyclin Cln3-Cdk1 complex initiates the transition

by phosphorylating and partially inactivating Whi5, an inhibitor of the SBF

transcription factor (Costanzo et al., 2004; de Bruin et al., 2004; Nasmyth and

Dirick, 1991; Ogas et al., 1991; Sidorova and Breeden, 1993). This allows for SBF-

dependent transcription of the downstream G1 cyclins *CLN1* and *CLN2*, which also

inactivate Whi5 to complete a positive feedback loop (Skotheim et al., 2008). Thus,

both the biochemical function of G1 regulators and their specific targets are highly

conserved (Figure 1).

Many of the individual proteins performing identical roles are unlikely to be true

orthologs, *i.e.*, it cannot be inferred from sequence identity that the proteins evolved

from a common ancestral gene. In yeast, a single cyclin-dependent kinase, Cdk1,

binds distinct cyclin partners to perform all the functions of three non-orthologous

animal Cdks (Cdk2, 4 and 6) during cell cycle entry (Liu and Kipreos, 2000).

Furthermore, no member of the transcription factor complex SBF-Whi5 exhibits

amino acid sequence identity or structural similarity to any member of the E2F-pRb

complex (Cross et al., 2011; Hasan et al., 2013; Taylor et al., 1997). Finally, Cdk

inhibitors such as Sic1 and p27 play analogous roles in yeast and mammals despite a

total lack of sequence identity (Cross et al., 2011). Taken together, these examples

imply significant evolution of cell cycle regulatory proteins in fungi and/or animals

while the network topology remains largely intact. While identification of network

topology is restricted to a few model organisms and is not as broad as sequence

analysis, the similar network topology in budding yeast and animals suggests that

this feature is more conserved than the constituent regulatory proteins (Cross et al.,

2011; Doonan and Kitsios, 2009).

The shared presence of E2F-pRb within plants (Archaeaplastida) and animal

(Metazoa) lineages indicates that this regulatory complex, rather than the fungal

SBF-Whi5 complex, was present in the last eukaryotic common ancestor (Cao et al.,

2010; Doonan and Kitsios, 2009; Fang et al., 2006; Harashima et al., 2013). The

divergence of G1 regulator sequences is surprising because fungi and animals are

more closely related to one another than either is to plants. This fungal-metazoan

difference raises the question as to where the fungal components came from. Fungal

components could either be rapidly evolved ancestral regulators or have a distinct

evolutionary history, which would suggest convergent evolution of regulatory

networks.

Here, we examine conserved and divergent features of eukaryotic cell cycle

regulation. In contrast to previous work that considered a protein family of cell cycle

regulators in isolation (Cao et al., 2010; 2014; Eme et al., 2011; Gunbin et al.,

2011; Ma et al., 2013; Wang et al., 2004), we studied the evolutionary history of an

entire regulatory network. We examined a greater number of genomes covering

most of eukaryotic diversity, including Excavata, Haptophyta, Cryptophyta, SAR

(Stramenopiles, Alveolata & Rhizaria), Archaeaplastida (plants), Amoebozoa,

Apusozoa and the Opisthokonta (animals and fungi). This survey allowed us to

estimate the cell cycle repertoire of the last eukaryotic common ancestor (LECA), a

prerequisite to clarifying the evolutionary transitions of the cell cycle components of

both animals and fungi.

Our results indicate that LECA likely had complex cell cycle regulation involving at

least one Cdk, multiple cyclin families, activating and inhibitory E2F transcription

factors, and pRb-family pocket proteins. The LECA repertoire helps establish that the

emergence of SBF-Whi5 is abrupt and distinguishes fungi from all other eukaryotes.

We also show that basal fungi can have both ancestral E2F-pRb and fungal SBF-Whi5
components. Thus, fungal evolution appears to have proceeded through a hybrid
network before abruptly losing the ancestral components in the lineage leading to
Dikarya. This supports the hypothesis that network structure, but not the individual
components, has been conserved through the transition to fungi and argues against
the case of convergent evolution.
109
Finally, our data confirm that SBF shows homology to KilA-N, a poorly characterized
domain present in prokaryotic and eukaryotic DNA viruses. Thus, SBF is not derived
from E2F (its functional analog) and likely emerged through horizontal gene transfer
in the fungal ancestor. That SBF can still regulate promoters with E2F binding sites
in budding yeast suggests that a viral SBF may have initially hijacked cell cycle
control, activating transcription via the *cis*-regulatory elements targeted by the
ancestral cell cycle regulator E2F, much like extant viral oncogenes.
117
118 **RESULTS**
119
Reconstruction of complex cell cycle control in the ancestral eukaryote
Recent work shows that the last eukaryotic common ancestor (LECA) already had a
complex repertoire of proteins (Dacks and Field, 2007; Eichinger et al., 2005;
Merchant et al., 2007). Indeed, all sequenced eukaryotic lineages have lost entire
gene families that were present in LECA (Fritz-Laylin et al., 2010). In contrast to the
growing consensus that LECA had an extensive repertoire of proteins, the prevailing
view of the LECA cell cycle is that it was based on a simple oscillator constructed
with relatively few components (Coudreuse and Nurse, 2010; Nasmyth, 1995).
According to the 'simple' LECA cell cycle model, an ancestral oscillation in Cyclin B-
Cdk1 activity drove periodic DNA replication and DNA segregation, while other
aspects of cell cycle regulation, such as G1 control, may have subsequently evolved
in specific lineages. The model was motivated by the fact that Cdk activity of a single
Cyclin B is sufficient to drive embryonic cell cycles in frogs (Murray and Kirschner,
1989) and fission yeast (Stern and Nurse, 1996), and that many yeast G1 regulators
have no eukaryotic orthologs (Figure 1).
135
To determine the complexity of LECA cell cycle regulation, we examined hundreds of
diverse eukaryotic genomes (Fungi, Amoebozoa, Excavata, SAR, Haptophyta,

Cryptophyta). We first built sensitive profile Hidden Markov Models (Eddy, 2011) for

each of the gene families of cell cycle regulators from model organisms *Arabidopsis*

thaliana, Homo sapiens, Schizosaccharomyces pombe, and *Saccharomyces*

cerevisiae. These HHMs were then used to query the sequenced eukaryotic genomes

for homologs of both fungal and animal cell cycle regulators (Figures 2-3; see

Methods). Phylogenetic analyses were performed on the detected homologs for

accurate sub-family assignment of the regulators and infer their evolutionary history

(see Methods). If LECA regulation were simple, we would expect little conservation

beyond the Cyclin B-Cdk1 mitotic regulatory module. However, if LECA regulation

were more complex, we would expect to see broad conservation of a wider variety of

regulators.

We did not find either of the fungal regulators, SBF and Whi5, in any Archaeplastida,

Amoebozoa, SAR, Haptophyta, Cryptophyta, Excavata or Metazoa. However, we did

find the cyclin sub-families (A, B, D, and E) known to regulate the cell cycle in

metazoans across all major branches of eukaryotes (for cyclin phylogeny see Figure

3-figure supplement 1). We also found examples of all three sub-families of E2F

transcription factors (E2F1-6, DP, E2F7/8) and the pRb family of pocket proteins (for

E2F/DP and pRb phylogeny see Figure 3-figure supplement 2 and Figure 3-figure

supplement 3). Nearly all species contain the APC specificity subunits Cdc20 and

Cdh1/Fzr1, which regulate exit from mitosis and maintain low Cdk activity in G1 (for

Cdc20-family APC phylogeny see Figure 3-figure supplement 4). Taken together,

these data indicate that LECA cell cycle regulation was based on multiple cyclin

families, as well as regulation by the APC complex and members of the pRb and E2F

families.

Members of the Cdk1-3 family (i.e. CdkA in plants) are also broadly conserved across

eukaryotes, suggesting they were the primary LECA cell cycle Cdks (for CDK

phylogeny see Figure 3-figure supplement 5). Other cell cycle Cdk families in animal

(Cdk4/6) and plant (CdkB) are thought to be specific to those lineages. However, we

found CdkB in Stramenopiles, which may have arrived via horizontal transfer during

an ancient secondary endosymbiosis with algae as suggested by (Cavalier-Smith,

1999). We excluded from our analysis other families of cyclin-Cdks including Cdk7-9,

which regulate transcription and RNA processing, and Cdk5 (yeast Pho85), which

regulate cell polarity, nutrient regulation, and contribute to cell cycle regulation in

yeast (Cao et al., 2014; Guo and Stiller, 2004; Ma et al., 2013; Moffat and Andrews,

2004). While interesting and important, an extensive examination of these cyclin-

Cdk families is beyond the scope of this work.

A hybrid E2F-pRb-SBF-Whi5 network on the path of fungal evolution

To identify the possible origins of SBF and Whi5, we searched for sequence homologs

across eukaryotic genomes (see Methods). We were unable to find any eukaryotic

homologs of SBF or Whi5 outside of fungi, with one exception that we discuss in the

next section. The emergence of the new fungal components SBF, which includes the

large APSES family, and Whi5 is abrupt and occurs near the split of basal fungi from

metazoans (Figures 4-5; for SBF only, SBF+APSES, and Whi5 phylogeny see Figure

5-figure supplement 1, Figure 5-figure supplement 2, and Figure 5-figure

supplement 3). The precise location remains unclear because we have only 1

Nuclearid genome and Microsporidia are fast-evolving fungal parasites with reduced

genomes (Cuomo et al., 2012). Interestingly, the new regulators (SBF and Whi5)

and ancestral regulators (E2F and Rb) co-exist broadly across basal fungi and the

lineages formerly known as "zygomycetes". At one extreme, Chytridiomycota (*e.g.*,

Spizellomyces punctatus) can have both fungal and animal cell cycle regulators,

which likely represents the ancestral hybrid network. However, this hybrid network

was evolutionarily unstable, as different constellations of components are present in

the extant zygomycetes and basal fungi. For example, the zygomycetes have lost

pRb while retaining E2F, which was then abruptly lost in the transition to Dikarya.

The instability of the hybrid network suggests some functional redundancy between

the new SBF-Whi5 and ancient E2F-Rb pathway.

The SBF and E2F family of transcription factors are unlikely to be orthologs.

The DNA-binding domains of SBF/MBF (Taylor et al., 2000; Xu et al., 1997) and

E2F/DP (Zheng et al., 1999) are structurally classified as members of the winged-

helix-turn-helix (wHTH) family, which is found in both prokaryotes and eukaryotes

(Aravind and Koonin, 1999; Aravind et al., 2005; Gajiwala and Burley, 2000).

Although the DNA-binding domains of E2F/DP and SBF/MBF are both classified as

wHTH proteins, they show important differences in sequence, overall structure, and

mode of protein-DNA complex formation that lead us to conclude that it is highly

unlikely that they are orthologs.

The strongest arguments against SBF-E2F orthology are based on structural biology.

Many wHTH transcription factors, including the E2F/DP family, have a 'recognition

helix' that interacts with the major or minor grooves of the DNA. The E2F/DP family

has an RRXYD DNA-recognition motif in its helix that is invariant within the E2F/DP

family and is responsible for interacting with the conserved, core GCGC motif (Zheng

et al., 1999) (see Figure 6A: red structure). The RRXYD recognition motif is strikingly

conserved in E2F/DP across all eukaryotes, including the E2F/DP proteins uncovered

in basal fungi (Figure 6B, left). In contrast, the first solved SBF/MBF structure, Mbp1

from *S. cerevisiae* in the absence of DNA, suggested Mbp1 recognizes its MCB (Mlu I

cell cycle box, ACGCGT) binding site via a recognition helix (Taylor et al., 1997; Xu

et al., 1997). A recent structure of PCG2, an SBF/MBF homolog in the rice blast

fungus *Magnaporthe oryzae*, in complex with its MCB binding site does not support

this proposed mode of DNA binding (Liu et al., 2015). In striking contrast to many

wHTH structures, in which the recognition helix is the mediator of DNA binding

specificity, the wing of PCG2 binds to the minor groove to recognize the MCB binding

site. The two glutamines in the wing (Q82, Q89) are the key elements that recognize

the core MCB binding motif CGCG (Figure 6A, blue structure). Family-specific

conservation in the DNA-binding domain is observed for all members of the SBF and

APSES family, including basal fungal sequences (Figure 6B, right). In summary, the

incongruences in sequence, structure, and mode of DNA-interaction between E2F/DP

and SBF/MBF families strongly suggest that SBF is not derived from E2F.

Viral origin and evolution of the fungal SBF and APSES family

Since SBF is unlikely to be orthologous to the E2F family of transcription factors, we

therefore sought an alternative evolutionary origin. SBF is a member of a larger

family of transcription factors in yeast that includes Xbp1, Bqt4, and the APSES

family (Acm1, Phd1, Sok2, Egf1, StuA); see Figure 5. Previous work has shown that

the DNA-binding domain of the APSES and SBF proteins is homologous to a viral

KilA-N domain (Iyer et al., 2002). KilA-N is a member of a core set of "viral hallmark

genes" found across diverse DNA viruses that infect eubacteria, archaea, and

eukaryotes (Koonin et al., 2006). Outside the fungal SBF/APSES sub-family, little is

known about the KilA-N domain structure, its DNA-binding recognition sequence, and

function (Brick et al., 1998). The wide distribution of DNA viruses and KilA-N across

the three domains of life suggests that the fungal ancestor likely acquired SBF via
horizontal gene transfer.

We were unable to find ancestral SBF-like proteins in other eukaryotes—with the
notable exception of *Trichomonas vaginalis*, a parasitic excavate that has over 100
homologs. These potential SBF homologs were compared to the Uniprot protein
database, and their best sequence matches were to Mimivirus, a large double-
stranded DNA virus of the Nucleo-Cytoplasmic Large DNA Viruses (Yutin et al.,
2009). Upon closer inspection, both fungal SBF sequences and *T. vaginalis* Mimivirus
sequences share the viral KilA-N domain. To gain insight into the possible
evolutionary origins of the SBF subfamily, we aligned all KilA-N sequences from the
Uniprot and PFAM database (Finn et al., 2014) to all our fungal SBF sequences and
built a phylogenetic tree (Figure 7). There are three major phylogenetic groups of
KilA-N domains: those found in eukaryotic viruses, prokaryotic viruses, and the
fungal SBF family. Our current phylogeny is unable to distinguish whether the SBF
family arrived in a fungal ancestor through a eukaryotic virus or a phage-infected
bacterium. Structural and functional characterization of existing viral KilA-N domains
could help distinguish between these two hypotheses. However, our phylogeny does
show that *T. vaginalis* obtained its KilA-N proteins independently from the fungal
ancestor because its KilA-N domains are clearly placed within the eukaryotic DNA
viruses (*e.g.*, Mimivirus) rather than with the fungal subfamily (Figure 7).

SBF ancestor could regulate E2F-target genes

264
Of all the members of the SBF/APSES family, the most likely candidate to be a
"founding" TF is SBF, as it is the only member present in all fungi. In budding yeast
and other fungi, SBF functions in G1/S cell cycle regulation and binds a consensus
site CGCGAA (Gordân et al., 2011), which overlaps with the consensus site
GCGSSAAA for the E2F family (Rabinovich et al., 2008). The APSES regulators, Xbp1,
and MBF in budding yeast bind TGCA, TCGA, ACGCGT motifs, respectively. A viral
origin of the SBF/APSES family—with the founding member involved in cell cycle
control—suggests the hypothesis that perhaps the founder TF functioned like a DNA
tumor virus protein and hijacked cell cycle control to promote proliferation.

For the viral TF (SBF) to hijack cell cycle control in the fungal ancestor, it must have
been able to both bind E2F regulatory regions and then activate the expression of

genes under E2F in a cell cycle-regulated fashion. The overlap between the

conserved E2F and SBF consensus sites suggests that ancestral SBF could bind E2F

regulatory regions. However, a single base pair substitution in the SBF motif can

reduce gene expression by up to ~95% (Andrews and Moore, 1992) and flanking

regions outside the core are often important for binding affinity and gene expression

(Nutiu et al., 2011). To test whether yeast SBF can bind a canonical E2F binding site,

we inserted consensus E2F binding sites in the budding yeast genome; see Figure 8.

The hijacking hypothesis would be supported *in vivo* if E2F binding sites could

generate SBF-dependent cell cycle regulated gene expression. We used the well-

studied *CLN2* promoter, which has three binding sites for SBF (SCB, Swi6-dependent

cell cycle box) in a nucleosome-depleted region. Removal of these SCB sites is

known to eliminate cell cycle-dependent gene expression (Bai et al., 2010). We

changed all three SCBs to the E2F binding site variant GCGCGAAA known to regulate

the histone gene cluster in mammals (Rabinovich et al., 2008). We observed

significant oscillations in GFP expression, which were coordinated with the cell cycle.

Importantly, the amplitude of these oscillations was dependent on the budding yeast

SBF, but not MBF, and disappeared when the 3 binding sites were removed. Taken

together, this set of *in vivo* experiments lends support to the hijacking hypothesis,

where an ancestral SBF took control of several E2F-regulated genes.

DISCUSSION

298
Cell division is an essential process that has been occurring in an uninterrupted chain

for billions of years. Thus, one expects strong conservation in the regulatory network

controlling the eukaryotic cell division cycle. Consistent with this idea, cell cycle

network structure is highly similar in budding yeast and animal cells. However, many

components performing similar functions, such as the central SBF and E2F

transcription factors, lack sequence identity suggesting a significant degree of

evolution or independent origin. To identify axes of conservation and evolution in

eukaryotic cell cycle regulation, we examined a large number of genome sequences

in Archaeplastida, Amoebozoa, SAR, Haptophyta, Cryptophyta, Excavata, Metazoa

and Fungi. Across eukaryotes, we found a large number of proteins homologous to

metazoan rather than fungal G1/S regulators. Our analysis indicates that the last

eukaryotic common ancestor likely had complex cell cycle regulation based on Cdk1,

Cyclins D, E, A and B, E2F, pRb and APC family proteins.

In contrast, SBF was not present in the last common eukaryotic ancestor, and

abruptly emerged, with its regulator Whi5, in fungi likely due to the co-option of a

viral protein at the base of the fungal lineage. The origin of Whi5 is unclear because

we found no homologs outside of fungi. Whi5 is a mostly unstructured protein,

which, like pRb, recruits transcriptional inhibitor proteins to specific sites on DNA via

transcription factor binding (Huang et al., 2009; Wang et al., 2009). The relatively

simple structure of Whi5 suggests that it may have been subsequently co-opted as a

phosphopeptide to entrain SBF activity to cell cycle regulated changes in Cdk activity

(Figure 9).

The replacement of E2F-Rb with SBF-Whi5 at the core of the cell cycle along the

fungal lineage raises the question as to how such a drastic change to fundamental

regulatory network could evolve? One answer can be found in the evolution of

transcription factors. When the function of an essential transcription factor does

change, it often leaves behind a core part of its regulon for another factor (Brown et

al., 2009; Gasch et al., 2004; Lavoie et al., 2009). This process of handing off

transcription factor function has been observed to proceed through an intermediate

state, present in some extant genomes, in which both factors perform the function

(Tanay et al., 2005). The logic of proceeding through an intermediate state has been

well-documented for the regulation of genes expressed only in yeast of mating type

a (asgs) (Tsong et al., 2006). In the ancestral yeast and many extant species, asgs

expression is activated by a protein only present in **a** cells, while in other yeasts,

expression is repressed by a protein only present in α cells and **a**/α diploids. The

replacement of the ancestral positive regulation by negative regulation occurred via

yeast that contained both systems illustrating how an essential function can evolve

through a hybrid state (Baker et al., 2012).

Clearly, something similar happened during cell cycle evolution. It appears that the

replacement of the E2F-pRb transcription regulatory complex with the SBF-Whi5

complex proceeded via a hybrid intermediate that preserved its function. In the

hybrid intermediate, E2F-Rb and SBF-Whi5 may have evolved to be parallel

pathways whose functions overlapped to such an extent that the previously essential

E2F-Rb pathway could be lost in the transition to Dikarya. Interestingly, many basal

fungi have preserved rather than lost this hybrid intermediate, which suggests that

each pathway may have specialized functions. Chytrids exhibit both animal (e.g.

centrioles, flagella, amoeboid movement) and fungal features (e.g. cell walls, hyphal

morphology) whose synthesis needs to be coordinated with cell division. The

preservation of the hybrid network in chytrids could then be explained if animal and

fungal features are regulated by the E2F-Rb and SBF-Whi5 pathways respectively.

The origin of the hybrid network at the base of Fungi is abrupt and may have been

initiated by the arrival of SBF via virus. Many tumor viruses activate cell

proliferation. For example, the DNA tumor viruses Adenovirus and SV40 highjack cell

proliferation in part by activating the expression of E2F-dependent genes by binding

pRb to disrupt inhibition of E2F (DeCaprio, 2009). While the specific mechanisms

may differ, when SBF entered the fungal ancestor cell it might have activated the

transcription of E2F target genes. Rather than inhibiting the inhibitor of E2F, SBF

may have directly competed for E2F binding sites with transcriptionally inactive Rb-

E2F complexes (Figure 9). Consistent with this model, we have shown here that SBF

can directly regulate gene expression in budding yeast via a consensus E2F binding

site. Thus, the cooption of a viral protein generated a hybrid network to ultimately

facilitate dramatic evolution of the core cell cycle network in fungi.

MATERIALS AND METHODS

Identification of potential protein family homologs.

We used Profile-Hidden Markov Models (profile-HMMs) to detect homologs for each of

the families studied, using the HMMER 3 package (Eddy, 2011). Profile-HMMs are

sensitive tools for remote homology detection. Starting with a set of diverse yet

reliable protein homologs is fundamental for detecting remote protein homology and

avoiding "model poisoning" (Johnson et al., 2010). To this end, we used reliable

training-set homologs from the cell cycle model organisms *Arabidopsis thaliana,*

Homo sapiens, Schizosaccharomyces pombe, and *Saccharomyces cerevisiae*, to build

the profile-HMMs used to detect homologs. Our profile-HMM search used a stringent

e-value threshold of 1e-10 to detect putative homologs in the "best" filtered protein

sets (where available) of our 100+ eukaryotic genomes (see Supplementary File 1A

for genome details). All putative homologs recovered through a profile-HMM search

were further validated (or rejected) using an iterative search algorithm (Jackhmmer)

against the annotated SwissProt database using the HMMER web server (Finn et al.,
2011).

Our profile-HMM for E2F/DP family only detects E2F or DP, where as our profile-HMM
for SBF/MBF family only detects SBF/MBF (or APSES). The same protein was never
identified by both profile-HMMs because the sequence profiles and the structure are
different. In the case of basal fungi, which have both E2F/DP and SBF/MBF, all
proteins classified as an E2F/DP had clear homology to E2F or DP (see alignment in
new Figure 6B) and all proteins that we classified as SBF/MBF had clear homology to
SBF/MBF (see alignment in new Figure 6B). Thus, we have not misclassified E2Fs or
SBF/MBFs in basal fungi and we have not reached the limit of homology detection.

Phylogenetic-based classification of protein homologs in sub-families.

A phylogenetic analysis and classification was built in four stages. In the first stage,
we used MAFFT-L-INS-i (-maxiterate 1000) to align the sequences of eukaryotic
protein family members (Katoh and Standley, 2013). We then used probabilistic
alignment masking using ZORRO (Wu et al., 2012) to create different datasets with
varying score thresholds. Next, we used ProtTest 3 to determine the empirical
amino-acid evolutionary model that best fit each of our protein datasets using
several criteria: Akaike Information Criterion, corrected Akaike Information Criterion,
Bayesian Information Criterion and Decision Theory (Darriba et al., 2011). Last, for
each dataset and its best-fitting model, we ran different phylogenetic programs that
use maximum-likelihood methods with different algorithmic approximations (RAxML
and PhyML) and Bayesian inference methods (PhyloBayes-MPI) to reconstruct the
phylogenetic relationships between proteins.

404
405

For RAxML analyses, the best likelihood tree was obtained from five independent
maximum likelihood runs started from randomized parsimony trees using the
empirical evolutionary model provided by ProtTest. We assessed branch support via
rapid bootstrapping (RBS) with 100 pseudo-replicates. PhyML 3.0 phylogenetic trees
were obtained from five independent randomized starting neighbor-joining trees
(RAND) using the best topology from both NNI and SPR moves. Non-parametric
Shimodaira-Hasegawa-like approximate likelihood ratio tests (SH-aLRTs) and
parametric *à la Bayes* aLRTs (aBayes) were calculated to determine branch support
from two independent PhyML 3.0 runs. For Bayesian inference we used PhyloBayes

(rather than the more frequently used MrBayes) because it allows for site-specific

amino-acid substitution frequencies, which better models the level of heterogeneity

seen in real protein data (Lartillot and Philippe, 2004; Lartillot et al., 2009). We

performed Phylobayes analyses by running three independent chains under CAT and

the exchange rate provided by ProtTest 3 (e.g. CAT-LG), four discrete gamma

categories, and with sampling every 10 cycles. Proper mixing was initially confirmed

with Tracer v1.6 (Rambaut et al., 2014). The first 1000 samples were discarded as

burn-in, and convergence was assessed using bipartition frequencies and summary

statistics provided by bpcomp and tracecomp from Phylobayes. These were visually

inspected with an R version of AWTY (https://github.com/danlwarren/RWTY)

(Nylander et al., 2008). The best phylogenies are shown in Figure 3-figure

supplement 1-5 and Figure 5-figure supplement 1-3, and were used to tentatively

classify sequences into sub-families and create Figures 2-5.

We note that the confidence of each node in the phylogenetic trees was assessed

using multiple, but complementary support metrics: (1) posterior probability for the

Bayesian inference, (2) rapid bootstrap support (Stamatakis, 2006; Stamatakis et

al., 2008) for RAxML, and (3) non-parametric Shimodaira-Hasegawa-like

approximate likelihood ratio tests (SH-aLRTs) and parametric *à la Bayes* aLRTs

(aBayes) for PhyML. These different support metrics complement each other in their

advantages and drawbacks. SH-aLRT is conservative enough to avoid high false

positive rates but performs better compared to bootstrapping (Guindon et al., 2010;

Simmons and Norton, 2014). aBayes is powerful compared to non-parametric tests,

but has a tendency to increase false-positive rates under serious model violations,

something that can be balanced with SH-aLRTs (Anisimova and Gascuel, 2006;

Anisimova et al., 2011).

Strain construction:

Our *CLN2pr-GFP-CLN2PEST* constructs were all derived from pLB02-0mer (described

in (Bai et al., 2010) and obtained from Lucy Bai). To create pLB02-CLN2, a synthetic

DNA fragment (IDT, Coralville, IA) encompassing a region of the *CLN2* promoter

from 1,130 bp to 481 bp upstream of the *CLN2* ORF was digested with BamHI and

SphI and ligated into pLB02-0mer digested with the same enzymes. To create

pLB02-E2F, which contains E2F binding sites, the same procedure was applied to a

version of the promoter fragment in which the SCBs at 606bp, 581bp, and 538bp upstream of the ORF were replaced with the E2F binding site consensus sequence GCGCGAAA (Thalmeier et al., 1989). All these plasmids were linearized at the BbsI restriction site in the *CLN2* promoter and transformed. Both *swi4Δ* and *mbp1Δ* strains containing pLB02-0mer, pLB02-Cln2, pLB02-E2F fluorescent expression reporters were produced via mating lab stocks using standard methods. JE103 was a kind gift from Dr. Jennifer Ewald. Plasmids and strains are listed in Supplementary File 1B and 1C, respectively.

Imaging and analysis:

Imaging proceeded essentially as described in Bean *et al*., 2006. Briefly, early log-phase cells were pre-grown in SCD and gently sonicated and spotted onto a SCD agarose pad (at 1.5%), which was inverted onto a coverslip. This was incubated on a heated stage on a Zeiss Observer Z.1 while automated imaging occurred (3 minute intervals, 100-300 ms fluorescence exposures). Single-cell time-lapse fluorescence intensity measurements were obtained using software described in (Doncic and Skotheim, 2013; Doncic et al., 2011), and oscillation amplitudes were obtained manually from the resulting traces. The single-cell fluorescence intensity traces used mean cellular intensity with the median intensity of the entire field of view subtracted, to control for any fluctuations in fluorescent background. The resulting measurements were analyzed in R.

ACKNOWLEDGEMENTS
We thank F. Cross and A. Robinson-Mosher for extensive discussions, and R. Gordan, J. Heitman, D. Lew, J. Nevins, S. Rubin for critical comments on the manuscript. We thank J. Stajich for providing access to the Bioinformatics cluster at the Institute for Integrative Genome Biology at UC Riverside supported by the College of Natural and Agricultural Sciences, the National Science Foundation, and Alfred P Sloan Foundation.

References:

Adl, S.M., Simpson, A.G.B., Lane, C.E., Lukeš, J., Bass, D., Bowser, S.S., Brown, M.W., Burki, F., Dunthorn, M., Hampl, V., et al. (2012). The revised classification of eukaryotes. Journal of Eukaryotic Microbiology *59*, 429–493.

Andrews, B.J., and Moore, L. (1992). Mutational analysis of a DNA sequence involved
in linking gene expression to the cell cycle. Biochem. Cell Biol. *70*, 1073–1080.

Anisimova, M., and Gascuel, O. (2006). Approximate likelihood-ratio test for
branches: A fast, accurate, and powerful alternative. Systematic Biology *55*, 539–
552.

Anisimova, M., Gil, M., Dufayard, J.-F., Dessimoz, C., and Gascuel, O. (2011).
Survey of branch support methods demonstrates accuracy, power, and robustness of
fast likelihood-based approximation schemes. Systematic Biology *60*, 685–699.

Aravind, L., and Koonin, E.V. (1999). DNA-binding proteins and evolution of
transcription regulation in the archaea. Nucleic Acids Res *27*, 4658–4670.

Aravind, L., Anantharaman, V., Balaji, S., Babu, M.M., and Iyer, L.M. (2005). The
many faces of the helix-turn-helix domain: Transcription regulation and beyond⋆.
FEMS Microbiol Rev *29*, 231–262.

Bai, L., Charvin, G., Siggia, E.D., and Cross, F.R. (2010). Nucleosome-Depleted
Regions in Cell-Cycle-Regulated Promoters Ensure Reliable Gene Expression in Every
Cell Cycle. Dev Cell *18*, 544–555.

Baker, C.R., Booth, L.N., Sorrells, T.R., and Johnson, A.D. (2012). Protein
modularity, cooperative binding, and hybrid regulatory states underlie transcriptional
network diversification. Cell *151*, 80–95.

Bertoli, C., Skotheim, J.M., and de Bruin, R.A.M. (2013). Control of cell cycle
transcription during G1 and S phases. Nat Rev Mol Cell Biol *14*, 518–528.

Brick, D.J., Burke, R.D., Schiff, L., and Upton, C. (1998). Shope fibroma virus RING
finger protein N1R binds DNA and inhibits apoptosis. Virology *249*, 42–51.

Brown, V., Sabina, J., and Johnston, M. (2009). Specialized Sugar Sensing in Diverse
Fungi. Curr Biol.

Cao, L., Chen, F., Yang, X., Xu, W., Xie, J., and Yu, L. (2014). Phylogenetic analysis
of CDK and cyclin proteins in premetazoan lineages. BMC Evol Biol *14*, 10.

Cao, L., Peng, B., Yao, L., Zhang, X., Sun, K., Yang, X., and Yu, L. (2010). The
ancient function of RB-E2F pathway: insights from its evolutionary history. Biol
Direct *5*, 55–.

Cavalier-Smith, T. (1999). Principles of Protein and Lipid Targeting in Secondary
Symbiogenesis: Euglenoid, Dinoflagellate, and Sporozoan Plastid Origins and the
Eukaryote Family Tree. Journal of Eukaryotic Microbiology *46*, 347–366.

Costanzo, M., Nishikawa, J.L., Tang, X., Millman, J.S., Schub, O., Breitkreuz, K.,
Dewar, D., Rupes, I., Andrews, B., and Tyers, M. (2004). CDK activity antagonizes
Whi5, an inhibitor of G1/S transcription in yeast. Cell *117*, 899–913.

Coudreuse, D., and Nurse, P. (2010). Driving the cell cycle with a minimal CDK
control network. Nature *468*, 1074–1079.

Cross, F.R., Buchler, N.E., and Skotheim, J.M. (2011). Evolution of networks and
sequences in eukaryotic cell cycle control. Philosophical Transactions of the Royal
Society B: Biological Sciences *366*, 3532–3544.

Cuomo, C.A., Desjardins, C.A., Bakowski, M.A., Goldberg, J., Ma, A.T., Becnel, J.J.,
Didier, E.S., Fan, L., Heiman, D.I., Levin, J.Z., et al. (2012). Microsporidian genome
analysis reveals evolutionary strategies for obligate intracellular growth. Genome Res
22, 2478–2488.

Dacks, J.B., and Field, M.C. (2007). Evolution of the eukaryotic membrane-trafficking
system: origin, tempo and mode. J Cell Sci *120*, 2977–2985.

Darriba, D., Taboada, G.L., Doallo, R., and Posada, D. (2011). ProtTest 3: fast
selection of best-fit models of protein evolution. Bioinformatics *27*, 1164–1165.

de Bruin, R.A.M., McDonald, W.H., Kalashnikova, T.I., Yates, J., and Wittenberg, C.
(2004). Cln3 activates G1-specific transcription via phosphorylation of the SBF bound
repressor Whi5. Cell *117*, 887–898.

DeCaprio, J.A. (2009). How the Rb tumor suppressor structure and function was
revealed by the study of Adenovirus and SV40. Virology *384*, 274–284.

Doncic, A., and Skotheim, J.M. (2013). Feedforward Regulation Ensures Stability and
Rapid Reversibility of a Cellular State. Mol Cell.

Doncic, A., Falleur-Fettig, M., and Skotheim, J.M. (2011). Distinct interactions select
and maintain a specific cell fate. Mol Cell *43*, 528–539.

Doonan, J.H., and Kitsios, G. (2009). Functional evolution of cyclin-dependent
kinases. Mol Biotechnol *42*, 14–29.

Eddy, S.R. (2011). Accelerated profile HMM searches. PLoS Comput Biol *7*,
e1002195.

Eichinger, L., Pachebat, J.A., Glöckner, G., Rajandream, M.-A., Sucgang, R.,
Berriman, M., Song, J., Olsen, R., Szafranski, K., Xu, Q., et al. (2005). The genome
of the social amoeba Dictyostelium discoideum. Nature *435*, 43–57.

Eme, L., Trilles, A., Moreira, D., and Brochier-Armanet, C. (2011). The phylogenomic
analysis of the anaphase promoting complex and its targets points to complex and
modern-like control of the cell cycle in the last common ancestor of eukaryotes. BMC
Evol Biol *11*, 265.

Fang, S.-C., de los Reyes, C., and Umen, J.G. (2006). Cell size checkpoint control by
the retinoblastoma tumor suppressor pathway. PLoS Genet *2*, e167.

Finn, R.D., Bateman, A., Clements, J., Coggill, P., Eberhardt, R.Y., Eddy, S.R., Heger,
A., Hetherington, K., Holm, L., Mistry, J., et al. (2014). Pfam: the protein families
database. Nucleic Acids Res *42*, D222–D230.

Finn, R.D., Clements, J., and Eddy, S.R. (2011). HMMER web server: interactive
sequence similarity searching. Nucleic Acids Res *39*, W29–W37.

Fritz-Laylin, L.K., Prochnik, S.E., Ginger, M.L., Dacks, J.B., Carpenter, M.L., Field,
M.C., Kuo, A., Paredez, A., Chapman, J., Pham, J., et al. (2010). The genome of
Naegleria gruberi illuminates early eukaryotic versatility. Cell *140*, 631–642.

Gajiwala, K.S., and Burley, S.K. (2000). Winged helix proteins. Curr Opin Struct Biol
10, 110–116.

Gasch, A.P., Moses, A.M., Chiang, D.Y., Fraser, H.B., Berardini, M., and Eisen, M.B.
(2004). Conservation and evolution of cis-regulatory systems in ascomycete fungi.
PLoS Biol *2*, e398.

Gordân, R., Murphy, K.F., McCord, R.P., Zhu, C., Vedenko, A., and Bulyk, M.L.
(2011). Curated collection of yeast transcription factor DNA binding specificity data
reveals novel structural and gene regulatory insights. Genome Biol *12*, R125.

Guindon, S., Dufayard, J.-F., Lefort, V., Anisimova, M., Hordijk, W., and Gascuel, O.
(2010). New algorithms and methods to estimate maximum-likelihood phylogenies:
assessing the performance of PhyML 3.0. Systematic Biology *59*, 307–321.

Gunbin, K.V., Suslov, V.V., Turnaev, I.I., Afonnikov, D.A., and Kolchanov, N.A.
(2011). Molecular evolution of cyclin proteins in animals and fungi. BMC Evol Biol *11*,
224.

Guo, Z., and Stiller, J.W. (2004). Comparative genomics of cyclin-dependent kinases
suggest co-evolution of the RNAP II C-terminal domain and CTD-directed CDKs. BMC
Genomics *5*, 69.

Hallmann, A. (2009). Retinoblastoma-related proteins in lower eukaryotes. Commun
Integr Biol *2*, 538–544.

Harashima, H., Dissmeyer, N., and Schnittger, A. (2013). Cell cycle control across
the eukaryotic kingdom. Trends Cell Biol *23*, 345–356.

Hasan, M.M., Brocca, S., Sacco, E., Spinelli, M., Papaleo, E., Lambrughi, M.,
Alberghina, L., and Vanoni, M. (2013). A comparative study of Whi5 and
retinoblastoma proteins: from sequence and structure analysis to intracellular
networks. Front Physiol *4*, 315.

Huang, D., Kaluarachchi, S., Van Dyk, D., Friesen, H., Sopko, R., Ye, W., Bastajian,
N., Moffat, J., Sassi, H., Costanzo, M., et al. (2009). Dual regulation by pairs of
cyclin-dependent protein kinases and histone deacetylases controls G1 transcription
in budding yeast. PLoS Biol *7*, e1000188.

Iyer, L.M., Koonin, E.V., and Aravind, L. (2002). Extensive domain shuffling in
transcription regulators of DNA viruses and implications for the origin of fungal
APSES transcription factors. Genome Biol *3*, RESEARCH0012.

Johnson, L.S., Eddy, S.R., and Portugaly, E. (2010). Hidden Markov model speed
heuristic and iterative HMM search procedure. BMC Bioinformatics *11*, 431.

Katoh, K., and Standley, D.M. (2013). MAFFT Multiple Sequence Alignment Software
Version 7: Improvements in Performance and Usability. Mol Biol Evol *30*, 772–780.

Koonin, E.V., Senkevich, T.G., and Dolja, V.V. (2006). The ancient Virus World and
evolution of cells. Biol Direct *1*, 29.

Lartillot, N., and Philippe, H. (2004). A Bayesian mixture model for across-site
heterogeneities in the amino-acid replacement process. Mol Biol Evol *21*, 1095–1109.

Lartillot, N., Lepage, T., and Blanquart, S. (2009). PhyloBayes 3: a Bayesian
software package for phylogenetic reconstruction and molecular dating.
Bioinformatics *25*, 2286–2288.

Lavoie, H., Hogues, H., and Whiteway, M. (2009). Rearrangements of the
transcriptional regulatory networks of metabolic pathways in fungi. Curr Opin
Microbiol *12*, 655–663.

Liu, J., and Kipreos, E.T. (2000). Evolution of cyclin-dependent kinases (CDKs) and
CDK-activating kinases (CAKs): differential conservation of CAKs in yeast and
metazoa. Mol Biol Evol *17*, 1061–1074.

Liu, J., Huang, J., Zhao, Y., Liu, H., Wang, D., Yang, J., Zhao, W., Taylor, I.A., and
Peng, Y.-L. (2015). Structural basis of DNA recognition by PCG2 reveals a novel DNA
binding mode for winged helix-turn-helix domains. Nucleic Acids Res *43*, 1231–1240.

Ma, Z., Wu, Y., Jin, J., Yan, J., Kuang, S., Zhou, M., Zhang, Y., and Guo, A.-Y.
(2013). Phylogenetic analysis reveals the evolution and diversification of cyclins in
eukaryotes. Molecular Phylogenetics and Evolution *66*, 1002–1010.

Merchant, S.S., Prochnik, S.E., Vallon, O., Harris, E.H., Karpowicz, S.J., Witman,
G.B., Terry, A., Salamov, A., Fritz-Laylin, L.K., Maréchal-Drouard, L., et al. (2007).
The Chlamydomonas genome reveals the evolution of key animal and plant
functions. Science *318*, 245–250.

Moffat, J., and Andrews, B. (2004). Late-G1 cyclin-CDK activity is essential for
control of cell morphogenesis in budding yeast. Nat Cell Biol *6*, 59–66.

Murray, A.W., and Kirschner, M.W. (1989). Cyclin synthesis drives the early
embryonic cell cycle. Nature *339*, 275–280.

Nasmyth, K. (1995). Evolution of the cell cycle. Philos Trans R Soc Lond, B, Biol Sci
349, 271–281.

Nasmyth, K., and Dirick, L. (1991). The role of SWI4 and SWI6 in the activity of G1
cyclins in yeast. Cell *66*, 995–1013.

Nutiu, R., Friedman, R.C., Luo, S., Khrebtukova, I., Silva, D., Li, R., Zhang, L.,
Schroth, G.P., and Burge, C.B. (2011). Direct measurement of DNA affinity
landscapes on a high-throughput sequencing instrument. Nat Biotechnol *29*, 659–
664.

Nylander, J.A.A., Wilgenbusch, J.C., Warren, D.L., and Swofford, D.L. (2008). AWTY
(are we there yet?): a system for graphical exploration of MCMC convergence in
Bayesian phylogenetics. Bioinformatics *24*, 581–583.

Ogas, J., Andrews, B.J., and Herskowitz, I. (1991). Transcriptional activation of

CLN1, CLN2, and a putative new G1 cyclin (HCS26) by SWI4, a positive regulator of
G1-specific transcription. Cell *66*, 1015–1026.

Rabinovich, A., Jin, V.X., Rabinovich, R., Xu, X., and Farnham, P.J. (2008). E2F in
vivo binding specificity: Comparison of consensus versus nonconsensus binding sites.
Genome Res *18*, 1763–1777.

Rambaut, A., Suchard, M.A., Xie, D., and Drummond, A.J. (2014). Tracer v1.6,
Available from http://beast.bio.edu.ac.uk/Tracer.

Sidorova, J., and Breeden, L. (1993). Analysis of the SWI4/SWI6 protein complex,
which directs G1/S-specific transcription in Saccharomyces cerevisiae. Mol Cell Biol
13, 1069–1077.

Simmons, M.P., and Norton, A.P. (2014). Divergent maximum-likelihood-branch-
support values for polytomies. Molecular Phylogenetics and Evolution *73*, 87–96.

Skotheim, J.M., Di Talia, S., Siggia, E.D., and Cross, F.R. (2008). Positive feedback
of G1 cyclins ensures coherent cell cycle entry. Nature *454*, 291–296.

Stamatakis, A. (2006). RAxML-VI-HPC: maximum likelihood-based phylogenetic
analyses with thousands of taxa and mixed models. Bioinformatics *22*, 2688–2690.

Stamatakis, A., Hoover, P., and Rougemont, J. (2008). A rapid bootstrap algorithm
for the RAxML Web servers. Systematic Biology *57*, 758–771.

Stern, B., and Nurse, P. (1996). A quantitative model for the cdc2 control of S phase
and mitosis in fission yeast. Trends Genet *12*, 345–350.

Tanay, A., Regev, A., and Shamir, R. (2005). Conservation and evolvability in
regulatory networks: The evolution of ribosomal regulation in yeast. Proc Natl Acad
Sci USA.

Taylor, I.A., Treiber, M.K., Olivi, L., and Smerdon, S.J. (1997). The X-ray structure of
the DNA-binding domain from the Saccharomyces cerevisiae cell-cycle transcription
factor Mbp1 at 2.1 A resolution. J Mol Biol *272*, 1–8.

Thalmeier, K., Synovzik, H., and Mertz, R. (1989). Nuclear factor E2F mediates basic
transcription and trans-activation by E1a of the human MYC promoter. Genes & ….

Travesa, A., Kalashnikova, T.I., de Bruin, R.A.M., Cass, S.R., Chahwan, C., Lee, D.E.,
Lowndes, N.F., and Wittenberg, C. (2013). Repression of G1/S Transcription Is
Mediated via Interaction of the GTB Motifs of Nrm1 and Whi5 with Swi6. Mol Cell Biol
33, 1476–1486.

Tsong, A.E., Tuch, B.B., Li, H., and Johnson, A.D. (2006). Evolution of alternative
transcriptional circuits with identical logic. Nature *443*, 415–420.

van den Heuvel, S., and Dyson, N.J. (2008). Conserved functions of the pRB and E2F
families. Nat Rev Mol Cell Biol *9*, 713–724.

Wang, G., Kong, H., Sun, Y., Zhang, X., Zhang, W., Altman, N., DePamphilis, C.W.,
and Ma, H. (2004). Genome-wide analysis of the cyclin family in Arabidopsis and

comparative phylogenetic analysis of plant cyclin-like proteins. Plant Physiol *135*,
1084–1099.

Wang, H., Carey, L.B., Cai, Y., Wijnen, H., and Futcher, B. (2009). Recruitment of
Cln3 cyclin to promoters controls cell cycle entry via histone deacetylase and other
targets. PLoS Biol *7*, e1000189.

Wu, M., Chatterji, S., and Eisen, J.A. (2012). Accounting for alignment uncertainty in
phylogenomics. PLoS ONE *7*, e30288.

Xu, R.M., Koch, C., Liu, Y., Horton, J.R., Knapp, D., Nasmyth, K., and Cheng, X.
(1997). Crystal structure of the DNA-binding domain of Mbp1, a transcription factor
important in cell-cycle control of DNA synthesis. Structure *5*, 349–358.

Yutin, N., Wolf, Y.I., Raoult, D., and Koonin, E.V. (2009). Eukaryotic large nucleo-
cytoplasmic DNA viruses: clusters of orthologous genes and reconstruction of viral
genome evolution. Virol. J. *6*, 223.

Zheng, N., Fraenkel, E., Pabo, C.O., and Pavletich, N.P. (1999). Structural basis of
DNA recognition by the heterodimeric cell cycle transcription factor E2F-DP. Genes
Dev *13*, 666–674.

693
694

A. Mammalian cell cycle network

B. Budding yeast cell cycle network

695
696
Figure 1. Topology of G1/S regulatory network in mammals and budding
yeast is conserved, yet many regulators exhibit no detectable sequence
homology. Schematic diagram illustrating the extensive similarities between animal
(A) and budding yeast (B) G1/S cell cycle control networks. Similar coloring denotes
members of a similar family or sub-family. Fungal components colored white denote
proteins with no identifiable animal orthologs.

703
704
Figure 2. Animal and plant G1/S regulatory network was present in the last
eukaryotic common ancestor. Distribution of cell cycle regulators across the
eukaryotic species tree (Adl et al., 2012). Animals (Metazoa) and yeasts (Fungi) are
sister groups (Opisthokonta), and are distantly related to plants (Charophyta), which
are members of the Archaeplastida. Check marks indicate the presence of at least
one member of a protein family in at least one sequenced species from the
corresponding group. See Figure 3 for a complete list of components in all species
analyzed.

	(H. sapiens)	Cdk1-4,6	CycB (CycB3)	CycA	CycE/D (CycO/F/J/G/I)	E2F1-6	E2F7-8	DP	Rb	Cdc20	Fzr1
Metazoa	*H. sapiens (human)*	5	3	2	13	6	2	3	3	2	1
	G. gallus (chicken)	4	2	2	13	6	2	2	3	2	1
	D. rerio (zebrafish)	5	3	2	12	5	2	3	3	1	2
	B. floridae (lancelet)	5	1	1	6	2	2	1	2	1	2
	C. intestinalis (sea squirt)	5	1	2	5		1	1	1	1	1
	S. purpuratus (urchin)	3	2	1	6	2		1	3	2	1
	L. gigantea (sea snail)	3	2	1	7	2	1	1	2	2	1
	D. melanogaster (fly)	3	2	1	3	2		1	2	1	2
	C. elegans (nematode)	3	4	1	2	2	1	1	1	1	1
	N. vectensis (anemone)	3	3	1	5	5	3	1	1	2	1
	A. queenslandica (sponge)	2	1	1	5	2	1	1	3	2	1
	T. adhaerens (placozoa)	3	2	1	3	2	1	1	2	1	1
Choanomonada	*Sphaeroforma arctica*		1		3	2		2	1	1	1
	Capsaspora owczarzaki		1	1	4	1	1	1	1	1	1
	Monosiga brevicollis	1	1	1	2	1	1	1	1	1	1
	Salpingoeca rosetta	2	1	1	2	2	1	1	2	1	1
Apusozoa	*Thecamonas trahens*	1	1	1	3	1	1	1	1	1	1
Amoebozoa	*Dictyostelium discoideum*	1	1	1	1	1		1	1	1	1
	Dictyostelium purpureum	1	1	1	1	1		1	1	1	1
	Dictyostelium fasciculatum	1	1	1	1	1	1	1	1	1	1
	Polysphondylium pallidum	1	1	1		1	2		1	1	1
	Entamoeba histolytica	6	2		3						2
	Entamoeba nuttalli	6	1		1						2
Diplomonatida	*Giardia intestinalis*	2			2		1				
Heterolobosea	*Naegleria gruberi*	3	3	3		1	1	1	1	3	1
Euglenozoa	*Trypanosoma brucei*	3	1		2					3	
	Leishmania major	3	2		1					2	
	Leishmania donovani	2	2		1					2	
Apicomplexa	*Symbiodinium minutum*	7	2							1	1
	Plasmodium falciparum	2									1
	Cryptosporidium muris	3			2	1	1	1			1
	Toxoplasma gondii	2									1
Ciliophora	*Tetrahymena thermophila*	7	5		11	2	2	3	2	1	8
Labyrinthulomycetes	*Aurantiochytrium limacinum*	3	1	1	3			1	1	1	1
	Schizochytrium aggregatum	3	1	1	2					1	1
	Aplanochytrium kerguelense	3	1	1	2	1		1	1	2	1
Peronosporomycetes	*Phytophthora infestans*	5	1	1	3	1	1	1	1	1	1
	Pythium ultimatum	1	1	1	4	1	1	1	1	1	1
	Hyaloperonospora parasitica	3		2	4			2	1	1	
	Saprolegnia parasitica	3	1	2	4	1	1	2	1	4	1
Diatomea	*Fragilariopsis cylindrus*	2	4	1	10	1	1	1		2	1
	Phaeodactylum tricornutum	2	1	1	10	2	1	1	1	2	1
	Thalassiosira pseudonana	2	3	1	26	1	1	1	1	2	2
Eustigmatales	*Nannochloropsis gaditana*	3	2	1	2			1	1	1	
Phaeophyceae	*Ectocarpus siliculosus*	3	2	1	2	1	1	1	1	2	1
Pelagophyceae	*Aureococcus anophaefferens*	2	2		2	1		1		2	1
Rhizaria	*Bigelowiella natans*	1	2	1	3	1	2			2	1
Cryptophyta	*Guillardia theta*	4	3	1	3	1	1	2	1	3	
Haptophyta	*Emiliania huxleyi*	2	2	3	1	2			1	6	2
Glaucophyta	*Cyanophora paradoxa*	1	1	1	1		1	1		1	2
Rhodophyceae	*Porphyridium cruentum*	3	1	1	1	1	1	1		1	1
	Cyanidioschyzon merolae	2		1	2	1	1	2	1	1	1
Chlorophyta	*Ostreococcus tauri*	2	1	1	2	1	1	1	1	1	1
	Ostreococcus lucimarinus	2	1	1	2	1	1	1	1	1	1
	Micromonas pusilla	2	1	1	2	1	1	1	1	1	1
	Coccomyxa subellipsoidea	4	1	1	1	1	1	1	1	1	1
	Volvox carteri	3	1	1	4	1		1	1	1	1
	Chlamydomonas reinhardtii	3	1	1	4	1		1	1	1	1
Charophyta	*Physcomitrella patens*	9	2	8	2	3	2	2	3	6	4
	Selaginella moellendorffii	3	1	3	3	2	1	1	2	3	2
	Brachypodium distachyon	5	5	6	14	3	2	2	2	5	2
	Oryza sativa	5	6	6	12	4	2	2	2	3	2
	Arabidopsis thaliana	5	11	10	11	3	3	2	1	6	3
	(A. thaliana)	CdkA/B	CycB	CycA	CycD (CycSDS)	E2FA-C	DEL	DP	RBR	Cdc20	Fzr1

713
714
Figure 3. Comparative genomic data of G1/S regulators across eukaryotes.
We developed profile-HMMs to detect cell division cycle regulators in eukaryotic
genomes. For each cell cycle regulatory family (*e.g.*, cyclins), we used molecular
phylogeny to classify eukaryotic sequences into sub-families (*e.g.*, Cyclins B, Cyclin
A, Cyclins E/D). See Methods for details and Figure 3-supplement 1 (Cyclin), Figure
3-supplement 2 (E2F/DP), Figure 3-supplement 3 (pRb), Figure 3-supplement 4
(Cdc20-family), and Figure 3-supplement 5 (CDK) for final phylogenies. Each entry
lists the number of sub-family members (column) for each eukaryotic genome (row).
Grey rows list the sub-family gene names in *H. sapiens* and *A. thaliana*. Additional
cyclin sub-family members are listed in parentheses.
725

Figure 4. Fungal ancestor evolved novel G1/S regulators, which eventually replaced ancestral cyclins, transcription factors, and inhibitors in Dikarya.
Basal fungi and "Zygomycota" contain hybrid networks comprised of both ancestral and fungal specific cell cycle regulators. Check marks indicate the presence of at least one member of a protein family in at least one sequenced species from the group. See Figure 5 for a complete list of components in all fungal species analyzed.

735
736
Figure 5. Comparative genomic data of G1/S regulators across fungi. We
developed profile-HMMs to detect fungal-specific cell division cycle regulators in
eukaryotic genomes. For each cell cycle regulatory family (*e.g.*, SBF/APSES), we
used molecular phylogeny to classify eukaryotic sequences into sub-families (*e.g.*,
SBF/MBF, APSES, Xbp1, Bqt4). Corresponding table shows the number of regulators
of each class for all species analyzed. See Methods for details and Figure 5-
supplement 1 (SBF only), Figure 5-supplement 2 (SBF+APSES), and Figure 5-
supplement 3 (Whi5) for final phylogenies. Grey rows list the sub-family gene
names in *S. cerevisiae*, *S. pombe*, and *H. sapiens*

746
Figure 6. E2F and SBF show incongruences in sequence, structure, and
mode of DNA binding. (A) Although both proteins share a winged helix-turn-helix
(wHTH) domain, the E2F/DP and SBF/MBF superfamilies do not exhibit significant
sequence identity or structural similarity to suggest a common recent evolutionary
origin according to CATH or SCOP databases. Furthermore, each wHTH has a
different mechanism of interaction with DNA: the arginine and tyrosine side-chains
of recognition helix-3 of E2F (E2F4 from *Homo sapiens* (Zheng et al., 1999)) interact
with specific CG nucleotides, where as the glutamine side-chains of the "wing" of
SBF/MBF (PCG2 from *Magnaporthe oryzae* (Liu et al., 2014)) interact with specific
CG nucleotides. (B) Sequence alignment of the DNA binding domain of
representative eukaryotic E2F/DP (left) and fungal SBF/MBF (right). The
corresponding secondary structure is above the sequence alignment. Evolutionary
conserved residues of sequence aligned DNA binding domains are highlighted in
black. Bold sequence names correspond to E2F/DP and SBF/MBF sequences from
basal fungi. Colored sequence names correspond to sequences of the structures
shown in panel A. PDB IDs for the structures used are shown in parentheses. W =
wing; T= turn.
764

765
Figure 7. Viral origin of yeast cell cycle transcription factor SBF. Maximum
Likelihood unrooted phylogenetic tree depicting relationships of fungal SBF-family
proteins (blue), KilA-N domains in prokaryotic (black, green), and eukaryotic DNA
viruses (orange, red). The particular origin of the HGT event that originated the
SBF/APSES family is unclear. We found no significant rapid bootstrapping support
(RBS ≥80%) for internal branches. Inference was performed under GTR+G model of
evolution. Scale bar in substitutions per site; (See Methods).
773

774
Figure 8. Yeast cell cycle transcription factor SBF can regulate cell cycle-
dependent transcription via E2F binding sites (A) Fluorescence images of cells
expressing a destabilized GFP from the SBF-regulated *CLN2* promoter. (B) Oscillation
of a transcriptional reporter in budding yeast. Characteristic time series of GFP
expression from a *CLN2* promoter (SCB), a *CLN2* promoter where the SBF binding
sites were deleted (ΔSCB), or a *CLN2* promoter where the SBF binding sites were
replaced with consensus E2F binding sites (E2F). Oscillation amplitudes were
quantified by scaling the mean fluorescence intensity difference from peak to trough
divided by the trough intensity $(S_{max} - S_{min})/S_{min}$. Circles denote time points
corresponding to (b). Triangles denote budding events. (C) Distribution of oscillation
amplitudes for different genotypes and GFP reporters. *swi4Δ* and *mbp1Δ* strains have
deletions of the SBF and MBF DNA-binding domain subunits respectively. t-test
comparisons within and across red and blue categories yield p-values > 0.3 or <
0.01 respectively. Boxes contain 25^{th}, median and 75^{th} percentiles, while whiskers
extend to 1.5 times this interquartile range.

Basal pathway

Addition of previously exogenous regulator

Entrainment of exogenous component to form hybrid network

Basal components lost, but pathway topology/function conserved

Figure 9. Punctuated evolution of a conserved regulatory network. Evolution can replace components in an essential pathway by proceeding through a hybrid intermediate. Once established, the hybrid network can evolve dramatically and lose previously essential regulators, while sometimes retaining the original network topology.

798
799
Figure 3 - supplement 1: Reduced phylogeny of eukaryotic cell cycle cyclins.
The cell division cycle (CDC) cyclin family consists of several sub-families with a well-
characterized cyclin box: (1) CycA (*H. sapiens, A. thaliana*), (2) CycB and CLB (*H.
sapiens, A. thaliana, S. cerevisiae, S. pombe*), (3) CycD (*H. sapiens, A. thaliana*), (4)
CycE (*H. sapiens*), and (5) CLN (*S. cerevisiae, S. pombe*). We combined CycA,
CycB, CycD, CycE, CLB, and CLN sequences from *H. sapiens* (10 cyclins)*, A. thaliana*
(13 cyclins)*, S. cerevisiae* (9 cyclins)*,* and *S. pombe* (5 cyclins) to create a
eukaryotic CDC cyclin profile-HMM (pCYC.hmm) following the procedure outlined in
the Methods. Our HMM profile was sensitive enough to discover known, but
uncharacterized cyclin sub-families (CycO, CycF, CycG, CycI, CycJ, CycSDS) as *bona*
fide CDC cyclins. A domain threshold of E-20 was used to identify potential CDC
cyclin homologs. We first made a phylogeny of all cyclins to classify them. This
dataset was then manually pruned to remove long-branches and problematic
lineages. Our reduced CDC cyclin dataset has a total of 499 sequences. Columns with
the top 10% Zorro score (496 positions) were used in our alignment. Confidence at
nodes was assessed with multiple support metrics using different phylogenetic
programs under LG model of evolution (aBayes and SH-aLRT metrics with PhyML,
RBS with RAxML, Bayesian Posterior Probability with Phylobayes (23,163 sampled
trees, meandiff= 0.01, maxdiff= 0.5)). Colored dots in branches indicate
corresponding branch supports. Thick branches indicate significant support by at
least two metrics, one parametric and one non-parametric; branch support
thresholds are shown in the center of the figure; see Methods.

822
823
Figure 3 - supplement 2: Phylogeny of eukaryotic E2F/DP transcription
factors. E2F-DP is a winged helix-turn-helix DNA-binding domain that is conserved
across eukaryotes (van den Heuvel and Dyson, 2008). There are three sub-families
within the E2F-DP family: (1) the E2F subfamily, (2) the E2F7-8/DEL subfamily, and
(3) the DP subfamily. The E2F family consists of E2F1-6 (*H. sapiens*) and E2FA-C (*A.
thaliana*). The E2F7-8/DEL family consists of E2F7-8 (*H. sapiens*) and DEL1-3 (*A.
thaliana*). The DP family consists of DP1-3 (*H. sapiens*) and DPA-B (*A. thaliana*). The
members of E2F form heterodimers with DP, whereas the DEL family has two DNA-
binding domains and does not require DP to bind DNA. We used the E2F_TDP.hmm
profile from PFAM to uncover members of the E2F/DP family across eukaryotes. A
domain threshold of E-10 was used to identify potential E2F/DP homologs. Our
E2F/DP dataset (pE2FDP.fasta) has 248 sequences. Columns with the top 8% Zorro
score (284 positions) were used in our alignment. Confidence at nodes was assessed
with multiple support metrics using different phylogenetic programs under LG model
of evolution (aBayes and SH-aLRT metrics with PhyML, RBS with RAxML, Bayesian
Posterior Probability with Phylobayes (53,009 sampled trees, meandiff=0.0064,
maxdiff=0.18)). Colored dots in branches indicate corresponding branch supports.
Thick branches indicate significant support by at least two metrics, one parametric
and one non-parametric; branch support thresholds are shown in the center of the
figure; see Methods.

844
845
Figure 3 - supplement 3: Phylogeny of eukaryotic Rb inhibitors. *H. sapiens*
has Rb1, RBL1 (p107), and RBL2 (p130), and *A. thaliana* has RBR1. The model fungi
S. cerevisiae and *S. pombe* do not have any obvious retinoblastoma pocket proteins.
We needed more eukaryotic sequences to create a robust HMM profile (pRb.hmm)
for the pRb family. Based on the pRb sequences collected in (Hallmann, 2009), we
built a profile-HMM using putative pRb homologs from *H. sapiens, G. gallus, C.*
intestinalis, D. melanogaster, C. elegans, N. vectensis, T. adhaerens (metazoa)*; B.*
dendrobatidis (fungi); *D. discoideum, D. purpureum, T. pseudonana, P. tricornutum,*
N. gruberi, E. huxleyi (protists)*; C. merolae, O. tauri, O. lucimarinus, M. pusilla, V.*
carteri, C. reinhartdii, P. patens, S. moellendorfii, A. thaliana (plants)*. A domain
threshold of E-20 was used to identify pRB homologs. Our pRB dataset (pRb.fasta)
has 72 sequences. Columns with the top 15% Zorro score (566 positions) were used
in our alignment. Confidence at nodes was assessed with multiple support metrics
using different phylogenetic programs under LG model of evolution (aBayes and SH-
aLRT metrics with PhyML, RBS with RAxML, Bayesian Posterior Probability with
Phylobayes (23,219 sampled trees, meandiff=0.0035, maxdiff=0.067)). Colored dots
in branches indicate corresponding branch supports. Thick branches indicate
significant support by at least two metrics, one parametric and one non-parametric;
branch support thresholds are shown in the center of the figure; see Methods.
865
866

867
868
Figure 3 - supplement 4: Reduced phylogeny of eukaryotic Cdc20-family APC
regulators. We combined CDC20 and CDH1/FZR1 sequences from *H. sapiens* (3
members)*, A. thaliana* (9 members)*,* and *S. cerevisiae* (3 members) to create a
eukaryotic CDC20-family APC regulator profile-HMM (pCDC20.hmm) following the
procedure outlined in the Methods. A domain threshold of E-50 was used to identify
CDC20 homologs. Our pCDC20 dataset (pCDC20.fasta) has 350 sequences. This
dataset was manually pruned to remove long-branches and problematic lineages.
Our reduced CDC20 dataset has a total of 289 sequences. Columns with the top 20%
Zorro score (608 positions) were used in our alignment. Confidence at nodes was
assessed with multiple support metrics using different phylogenetic programs under
LG model of evolution (aBayes and SH-aLRT metrics with PhyML, RBS with RAxML,
Bayesian Posterior Probability with Phylobayes (13,638 sampled trees,
meandiff=0.015, maxdiff=0.37)). Colored dots in branches indicate corresponding
branch supports. Thick branches indicate significant support by at least two metrics,
one parametric and one non-parametric; branch support thresholds are shown in the
center of the figure; see Methods.
885

886
887
Figure 3 - supplement 5: Phylogeny of eukaryotic cyclin-dependent kinases.
To create a profile-HMM (pCDCCDK.hmm) for eukaryotic cell cycle CDK, we
combined Cdk1-3, Cdk4, Cdk6 sequences from *H. sapiens,* CdkA and CdkB from *A.*
thaliana, Cdc28 from *S. cerevisiae,* and Cdc2 from *S. pombe*. A domain threshold of
E-20 was used to identify CDK homologs. Our cell cycle CDK dataset
(pCDCCDK.fasta) has 272 sequences. Columns with the top 15% Zorro score (473
positions) were used in our alignment. Confidence at nodes was assessed with
multiple support metrics using different phylogenetic programs under LG model of
evolution (aBayes and SH-aLRT metrics with PhyML, RBS with RAxML, Bayesian
Posterior Probability with Phylobayes (28,193 sampled trees, meandiff=0.015,
maxdiff=0.53)). Colored dots in branches indicate corresponding branch supports.
Thick branches indicate significant support by at least two metrics, one parametric
and one non-parametric; branch support thresholds are shown in the center of the
figure; see Methods.
902
903

Figure 5 - supplement 1: Phylogeny of fungal SBF transcription factors. SBF and MBF are transcription factors that regulate G1/S transcription in budding and fission yeast. To detect SMRC (Swi4/6 Mbp1 Res1/2 Cdc10) across fungi, we built a sensitive profile-HMM (pSMRC.hmm) by combining well-characterized SMRC sequences from *S. cerevisiae, C. albicans, N. crassa, A. nidulans,* and *S. pombe*. A domain threshold of E-10 was used to identify SMRC homologs. Our SMRC dataset (pSMRC.fasta) has 147 sequences. Columns with the top 20% Zorro score (709 positions) were used in our alignment. Confidence at nodes was assessed with multiple support metrics using different phylogenetic programs under LG model of evolution (aBayes and SH-aLRT metrics with PhyML, RBS with RAxML, Bayesian Posterior Probability with Phylobayes (19,457 sampled trees, meandiff=0.0056, maxdiff=0.145)). Colored dots in branches indicate corresponding branch supports. Thick branches indicate significant support by at least two metrics, one parametric and one non-parametric; branch support thresholds are shown in the center of the figure; see Methods.

923
924
Figure 5 - supplement 2: Phylogeny of fungal SBF and APSES transcription
factors. SBF and APSES transcription factors (Asm1, Phd1, Sok2, Efg1, StuA) share
a common DNA-binding domain (KilA-N), which is derived from DNA viruses. During
our search for SBF and APSES homologs, we consistently detected two additional
fungal sub-families with homology to KilA-N: XBP1 (family name taken from *S.
cerevisiae*) and BQT4 (family name taken from *S. pombe*). To detect APSES, XBP1,
and BQT4 homologs, we built profile-HMMs (APSES.hmm, XBP1.hmm, and
BQT4.hmm) by combining APSES, XBP1, and BQT4 homologs from *S. cerevisiae, C.
albicans, N. crassa, A. nidulans,* and *S. pombe*. A domain threshold of E-10 was used
to identify APSES, XBP1, and BQT4 homologs. Our final dataset (pKILA.fasta)
contains all fungal KILA sub-families (SBF, APSES, XBP1, BQT4) and has a total of
301 sequences. Columns with the top 10% Zorro score (447 positions) were used in
our alignment. Confidence at nodes was assessed with multiple support metrics
using different phylogenetic programs under LG model of evolution (aBayes and SH-
aLRT metrics with PhyML, RBS with RAxML, Bayesian Posterior Probability with
Phylobayes (15,251 sampled trees, meandiff=0.012, maxdiff=0.25)). Colored dots in
branches indicate corresponding branch supports. Thick branches indicate significant
support by at least two metrics, one parametric and one non-parametric; branch
support thresholds are shown in the center of the figure; see Methods.
944

945
946
Figure 5 - supplement 3: Phylogeny of fungal Whi5 inhibitors. WHI5 and
NRM1 are a yeast-specific protein family that has been identified and functionally
characterized across *S. cerevisiae*, *C. albicans*, and *S. pombe.* Both WHI5 and NRM1
are fast evolving proteins. There is a small conserved region of 25 amino-acids
(known as the GTB domain) that is responsible for interacting with Swi6/Cdc10
(Travesa et al., 2013). Unfortunately, the Whi5.hmm profile from PFAM is unable to
detect an SRL3 paralogue in *S. cerevisiae* or the NRM1 orthologues in *A. gossypii* or
C. albicans. We built a more sensitive profile-HMM (pWHI5.hmm) by combining
WHI5/NRM1 sequences across ascomycetes (including SRL3 from Saccharomyces
genomes and NRM1 from Candida genomes). A domain threshold of E-05 was used
to identify WHI5 homologs. Our WHI5 dataset (pWHI5.fasta) has 98 sequences.
Columns with the top 15% Zorro score (260 positions) were used in our alignment.
Confidence at nodes was assessed with multiple support metrics using different
phylogenetic programs under the LG model of evolution (aBayes and SH-aLRT
metrics with PhyML, RBS with RAxML, Bayesian Posterior Probability with Phylobayes
(77,696 sampled trees, meandiff=0.0068, maxdiff=0.11)). Colored dots in branches
indicate corresponding branch supports. Thick branches indicate significant support
by at least two metrics, one parametric and one non-parametric; branch support
thresholds are shown in the center of the figure; see Methods.
966

Supplementary File 1A: List of eukaryotic genomes. We downloaded and
analyzed the following annotated genomes using the "best" filtered protein sets when
available. We gratefully acknowledge the Broad Institute, the DOE Joint Genome
Institute, Génolevures, PlantGDB, SaccharomycesGD, AshbyaGD, DictyBase, JCV
Institute, Sanger Institute, TetrahymenaGD, PythiumGD, AmoebaDB,
NannochloroposisGD, OrcAE, TriTryDB, GiardiaDB, TrichDB, CyanophoraDB, and
CyanidioschizonDB for making their annotated genomes publicly available. We
especially thank D. Armaleo, I. Grigoriev, T. Jeffries, J. Spatafora, S. Baker, J.
Collier, and T. Mock for allowing us to use their unpublished data.
976
977 **Supplementary File 1B: Plasmids.**

978 **Supplementary File 1C: Strains.** All yeast strains were derived from W303 and
979 constructed using standard methods.