Getting the Hologenome Concept Right:

An Eco-Evolutionary Framework for Hosts and Their Microbiomes

Kevin R. Theis[1,*], Nolwenn M. Dheilly[2], Jonathan L. Klassen[3], Robert M. Brucker[4], John F.

Baines[5,6], Thomas C.G. Bosch[7], John F. Cryan[8], Scott F. Gilbert[9], Charles J. Goodnight[10],

Elisabeth A. Lloyd[11], Jan Sapp[12], Philippe Vandenkoornhuyse[13], Ilana Zilber-Rosenberg[14],

Eugene Rosenberg[15], Seth R. Bordenstein[16,*]

[1]Department of Immunology and Microbiology, Wayne State University, Detroit, MI 48201, USA.
[2]School of Marine and Atmospheric Sciences, Stony Brook University, Stony Brook, NY 11794-5000, USA.
[3]University of Connecticut, Department of Molecular & Cell Biology, Storrs, CT 06269, USA.
[4]The Rowland Institute at Harvard, Harvard University, Cambridge, MA 02142, USA.
[5]Max Planck Institute for Evolutionary Biology, August-Thienemann-Strasse 2, D-24306 Plön, Germany.
[6]Institute for Experimental Medicine, Christian-Albrechts-University of Kiel, Arnold-Heller-Strasse 3, D-
24105 Kiel, Germany.
[7]Zoological Institute, Christian-Albrechts-University, D-24118 Kiel, Germany.
[8]Department of Anatomy and Neuroscience, University College Cork, Cork, Ireland; Alimentary
Pharmabiotic Centre, University College Cork, Cork, Ireland.
[9]Department of Biology, Swarthmore College, Swarthmore, PA 19081, USA.
[10]Department of Biology, University of Vermont, Burlington, VT 05405, USA.
[11]History and Philosophy of Science Department, Indiana University, Bloomington, IN 47405, USA.
[12]Department of Biology, York University, 4700 Keele Street, Toronto, Ontario, M3J 1P3, Canada.
[13]Université de Rennes 1, CNRS, UMR6553 EcoBio, Rennes, France.
[14]Independent Scholar, Rachavat Ilan St., Givat Shmuel 51905, Israel.
[15]Department of Molecular Microbiology and Biotechnology, Tel-Aviv University, Tel Aviv, 69978, Israel.
[16]Departments of Biological Sciences and Pathology, Microbiology and Immunology, Vanderbilt
University, Nashville, TN 37225, USA.
28

30 *Corresponding authors:

31 Kevin Theis (ktheis@med.wayne.edu), Seth Bordenstein (s.bordenstein@vanderbilt.edu)

Abstract

Given the recently appreciated complexity of symbioses among hosts and their

microbes, significant rethinking in biology is occurring today. Scientists and philosophers

are asking questions at new biological levels of hierarchical organization - What is a

holobiont and hologenome? When should this vocabulary and associated concepts apply?

Are these points of view a null hypothesis for host-microbe systems or limited to a certain

spectrum of symbiotic interactions such as host-microbial coevolution? Legitimate

questions, advancements and revisions are warranted at this nascent stage of the field.

However, a productive and meaningful discourse can only commence when skeptics and

proponents alike use the same definitions and constructs. For instance, critiquing the

hologenome concept is not synonymous with critiquing coevolution, and arguing that an

entity is not necessarily the primary unit of selection is not synonymous with arguing that

it is not a unit of selection in general. Here, we succinctly deconstruct and clarify these

recent misconceptions. Holobionts (hosts and their microbes) and hologenomes (all

genomes of the holobiont) are multipartite entities that result from ecological, evolutionary

and genetic processes. They are not restricted to one special process but constitute a wider

vocabulary and framework for host biology in light of the microbiome. We invite the

community to consider these new perspectives in biology.

Main Text

The holobiont is a term used to describe an individual host and its microbial community, including viruses and cellular microorganisms [1-6] (Figure 1). It is derived from the Greek word *holos* which means whole or entire. Microbial symbionts can be constant or inconstant and can interact with a host in a context-dependent manner as harmful, harmless or helpful. In most cases, the net outcome of these interspecies relationships often varies with the presence of other symbionts. The term holobiont distinguishes itself by not only recognizing hosts and their obligate symbionts, but also emphasizing the diverse and complex array of symbionts and their dynamic associations within a host. In contrast to binary host-microbial interactions, the emergent properties of complex microbial communities and their hosts are newly appreciated and potentially universal. The host and microbial genomes of a holobiont are collectively defined as its hologenome [1,2], and the pluralistic attributes of a holobiont scale directly to the hologenome (Figure 1). This pluralism is an important point that is frequently missed. Microbial genomes can be stable or labile components of the hologenome, vertically or horizontally transmitted, and the functional traits that they encode are context dependent and may result in damage, benefit, or indifference to the holobiont [7].

The hologenome concept of evolution proposes that evolutionary forces acting at the phenotypic level of an individual organism (e.g., a single plant or animal) are instead acting at the holobiont level in many cases. For a given plant or animal, phenotype and fitness can often be deeply dependent on its microbial community [8], in both cooperative and competitive manners. Hologenomic evolution is a relatively new concept and is therefore liable to be interpreted in ways that misrepresent its original conception.

Importantly, a recent paper used different definitions for the holobiont and hologenome,

and restricted the term hologenome to those situations when the host and microbes

coevolve to form a primary unit of selection [9]. The result is a "straw man" conception of the

holobiont and hologenome. The hologenome concept is relatively new to biology and thus

requires critical evaluation; but to have a robust debate, skeptics and proponents must use

the same terminology. Our central goals in this brief essay are (a) to highlight errors in

these recent narrow definitions of the holobiont and hologenome, keeping them consistent

with their original pluralistic definitions, (b) to encourage productive discussion and

critical inquiry, and (c) to stimulate new ideas and understanding considering the link

between phenotype and genotype.

The first argument proposed by skeptics is that if X did not coevolve *sensu stricto*

with Y, then the hologenome is not real [9-11]. In this case, X and Y are respectively a

microbe/microbial community and a host. However, this argument is not relevant to the

hologenome concept. Hologenome is a term that encompasses all of the genomes of the

host and its microbes at a given point in time. How the symbiotic community was

assembled is independent of how selection can act upon the holobiont. Thus, holobionts

can be formed through neutral processes, selection at the level of the host, symbiont or

both (Figure 1). Although a component of it, coevolution is not the sole feature of the

hologenome and its associated concepts. By way of caricatural illustration, one would not

similarly say that if genes X and Y did not coevolve in a host, then they are not part of the

genome. Evolution of genomes and hologenomes is not a monolithic process. Genetic

conflict, epistasis, selection, drift, etc. are all operational [1,2]. Thus, objections to the

hologenome concept based on a lack of coevolution misrepresent what constitutes a

hologenome. To put it simply, coevolution is a process; the hologenome is an entity.

It was recently suggested that a non-coevolutionary application of the word

hologenome "would make it sufficiently general that it can be interpreted in any number of

ways" [9]. This comment refers to the original and more generally accepted definition of the

hologenome as all of the genomes in the holobiont, all of which in turn are evolving in that

context [6]. However, using this logic, the word genome would be as unhelpful to biology as

the term hologenome because it would be an insufficiently general definition of the types of

evolutionary processes occurring within the genome. As noted above, coevolution, genetic

conflict, selection, and drift all occur in the genome and hologenome. For this reason,

genetic interactions or epistasis between genes in the same genome follow a conceptual

continuum to intergenomic and interspecies interactions underlying community

phenotypes in hologenomes [1,12].

Prevalent misuse of coevolution to explain host-microbe associations was the

impetus for some of us coining the word "phylosymbiosis" [13]. It describes the concordance

between a host phylogeny and microbial community dendrogram that depicts community

relationships based on the degree of shared taxonomy and/or abundance of members of

the community [14-17]. Phylosymbiosis does not *a priori* imply coevolution or

codiversification to the observed pattern of symbiotic microbial community relationships

recapitulating host phylogenetics. Moreover, it "does not presume that microbial

communities are stable or even vertically transmitted from generation to generation" [15,16].

Rather, it refers to a pattern in which changes in separate parts of the holobiont (host

genome and microbiota) are related in a concordant manner across lineages of holobionts.

It is also a stepping-stone from population genetics to community genetics because when

phylosymbiosis is observed under strictly controlled conditions, it tests whether variation

in holobiont assembly is primarily stochastic or deterministic [13,15,18]. Stochastic assembly

means that each microbe has an equal opportunity of colonizing a host. Deterministic

assembly reflects ecological selection of a particular non-random microbial community and

its host, without reference to which partner, or potentially both, is doing the selection, and

it can be affected by genetic variation in the host or microbial species. In this context, when

genetic variation in the interacting species affects community assembly, it has been defined

as broad sense "community heritability," or H^2_C [19,20]. H^2_C measures a "heritable basis to

trophic-level interactions" [21]. Controlled studies of microbial community assembly across

different species of *Nasonia* wasps and *Hydra* have yielded such phylosymbiotic patterns

[13,14]. If there is a significant H^2_C, natural selection can act on ecological communities [18],

including organization of the holobiont and its emergent phenotypes [20].

Discussion of evolutionary processes brings forth a second unfounded argument

against the hologenome concept, namely that holobionts and their hologenomes must be

the "primary" unit of selection for the concept to be informative [9]. This strict claim leads

biologists into error, as multiple levels of selection can operate simultaneously. For

example, selfish genetic elements can be selected within a genome that is in turn selected

for any number of phenotypes that affect fitness—this is uncontroversial. While the

holobiont is posited to be "a unit of selection in evolution" [2,22-24], it is critically not proposed

as the only or primary unit of selection [1,2]. Primariness varies with what traits are targeted

by natural selection. From a phenotypic perspective, selection works on trait variation that

influences which entities reproduce and in what relative proportions they do so. Given this

tenet and the impact of microbes on host phenotypes, the target of selection at the so-called

host level will in many phenotypic cases be the holobiont—the extent to which is yet to be

determined. What we wish to emphasize here is that the holobiont does not imply group

level selection. Rather, variation in the hologenome yields variation in holobiont

phenotypes upon which various evolutionary (and ecological) processes at different levels

can act. The evolutionary response to selection can result in changes in the genetic

composition of the host genome, microbial genomes or both. We believe this concept

provides new perspectives.

As we have emphasized in different venues, it is also true that just as large parts of

the nuclear genome can evolve neutrally or be in conflict, so too can large parts of the

hologenome [1,2]. For example, "hologenomic drift can occur at all the different levels of the

holobiont from single genes of the microbes or the host to the holobiont itself" [2]. We would

be remiss to not be critical of our own inconsistent statements about the relative roles of

cooperation and conflict in hologenomic evolution. In *The Hologenome Concept*, some of us

stated that "evolution of animals and plants was driven primarily by natural selection for

cooperation between and with microorganisms" [2] while in other venues the concept

"places as much emphasis on cooperation as on competition" [25]. This latter statement is

more precisely aligned with the pluralistic nature of the holobiont, namely that "natural

selection...on holobiont phenotypes...can work to remove deleterious nuclear mutations or

microbes while spreading advantageous nuclear mutations or microbes" [1]. In fact, some of

us argued that conflicts of interests resulting from the nature of the transmission of

microbes to the next host could select for microbes that can manipulate the biology of their

host to improve their own transmission [26]. The holobiont is not a conglomerate that arises

solely from cooperation. Rather, it is a hierarchical level that can supersede the individual

host and encompass the organismal, genetic and phenotypic features of the host in

association with its microbial community, incorporating both competitive and cooperative

selective systems [27]. Hologenomes then exist as hierarchically nested levels of genomes, in

which all levels of selection are potentially in play.

In summary, we anticipate that many subdisciplines in biology will benefit from a

theoretical and experimental framework that broadly encompasses the ecology of

holobionts and evolution of hologenomes. The hologenome concept is a comprehensive

and relevant eco-evolutionary framework for which critical questions remain. For example,

can a response to selection on host traits be driven solely by changes in the genomes

and/or membership of a host-associated microbial community? How taxonomically

widespread is phylosymbiosis? What is the strength of selection on holobiotic interactions

required to maintain consistent association with environmentally-acquired microbes each

generation? How does selection operate on community phenotypes if H^2_C is variable

throughout the holobiont's lifetime due to the lability of microbial communities? Can the

lability of microbial communities explain inter-individual variations and help maintain a

higher phenotypic diversity within a holobiont population? Evolution of the hologenome

refers to the genetic basis of eco-evolutionary processes underlying community

phenotypes of the holobiont. We suggest that the null framework for the newly appreciated

complexities in the host-microbe consortia and their genomes is best reflected by the terms

holobiont and hologenome and their associated concepts.

 Figure Legend

Figure 1. Holobionts are entities comprised of the host and all of its symbiotic microbes, including those which affect the holobiont's phenotype and have coevolved with the host (blue), those which affect the holobiont's phenotype but have not coevolved with the host (red), and those which do not affect the holobiont's phenotype at all (gray). Microbes may be transmitted vertically or horizontally, acquired from the environment, and can be constant or inconstant in the host. Therefore, holobiont phenotypes can change in time and space as microbes come into and out of the holobiont. Microbes in the environment are not part of the holobiont (white). Hologenomes then encompass the genomes of the host and all of its microbes at any given time point, with individual genomes and genes falling into the same three functional categories of blue, red and gray. Holobionts and hologenomes are entities, whereas coevolution or host-symbiont interactions are processes.

199 **Acknowledgements**

200 We thank Jay Evans and Phil Pellett for feedback on the manuscript.

References

1   Bordenstein, S. R. & Theis, K. R. Host biology in light of the microbiome: Ten principles of
holobionts and hologenomes. *PLoS Biol* **13**, e1002226 (2015).
2   Rosenberg, E. & Zilber-Rosenberg, I. *The Hologenome Concept: Human, Animal and Plant*
Microbiota. (Springer, 2013).
3   Margulis, L. Symbiogenesis and symbionticism. In *Symbiosis as a Source of Evolutionary*
Innovation: Speciation and Morphogenesis   (eds L. Margulis & Rene Fester) 1-14 (MIT Press,
1991).
4   Mindell, D. P. Phylogenetic consequences of symbioses: Eukarya and Eubacteria are not
monophyletic taxa. *Biosystems* **27**, 53-62, doi:10.1016/0303-2647(92)90046-2 (1992).
5   Rohwer, F., Seguritan, V., Azam, F. & Knowlton, N. Diversity and distribution of coral-associated
bacteria. *Marine Ecology Progress Series* **243**, 1-10, doi:10.3354/meps243001 (2002).
6   Gordon, J., Knowlton, N., Relman, D. A., Rohwer, F. & Youle, M. Superorganisms and holobionts.
In *Microbe* Vol. 8   2 (American Society for Microbiology, Washington, DC, 2013).
7   Casadevall, A. & Pirofski, L. A. What is a host? Incorporating the microbiota into the damage-
response framework. *Infection and Immunity* **83**, 2-7, doi:10.1128/iai.02627-14 (2015).
8   Vandenkoornhuyse, P., Quaiser, A., Duhamel, M., Le Van, A. & Dufresne, A. The importance of
the microbiome of the plant holobiont. *New Phytologist* **206**, 1196-1206,
doi:10.1111/nph.13312 (2015).
9   Moran, N. A. & Sloan, D. B. The hologenome concept: Helpful or hollow? *PLoS Biol* **13**, e1002311
(2015).
10   Hester, E. R., Barott, K. L., Nulton, J., Vermeij, M. J. A. & Rohwer, F. L. Stable and sporadic
symbiotic communities of coral and algal holobionts. *The ISME Journal*,
doi:10.1038/ismej.2015.190 (In Press).
11   Chandler, J. A. & Turelli, M. Comment on "The hologenomic basis of speciation: Gut bacteria
cause hybrid lethality in the genus *Nasonia*". *Science* **345**, 2, doi:10.1126/science.1251997
(2014).
12   Fitzpatrick, B. M. Symbiote transmission and maintenance of extra-genomic associations.
Frontiers in Microbiology **5**, doi:10.3389/fmicb.2014.00046 (2014).
13   Brucker, R. M. & Bordenstein, S. R. The hologenomic basis of speciation: gut bacteria cause
hybrid lethality in the genus *Nasonia*. *Science* **341**, 667-669, doi:10.1126/science.1240659
(2013).
14   Fraune, S. & Bosch, T. C. G. Long-term maintenance of species-specific bacterial microbiota in
the basal metazoan Hydra. *Proceedings of the National Academy of Sciences* **104**, 13146-13151,
doi:10.1073/pnas.0703375104 (2007).
15   Brucker, R. M. & Bordenstein, S. R. The roles of host evolutionary relationships (Genus: *Nasonia*)
and development in structuring microbial communities. *Evolution* **66**, 349-362,
doi:10.1111/j.1558-5646.2011.01454.x (2012).
16   Brucker, R. M. & Bordenstein, S. R. Speciation by symbiosis. *Trends in Ecology & Evolution* **27**,
443-451, doi:10.1016/j.tree.2012.03.011 (2012).
17   Franzenburg, S. *et al.* Distinct antimicrobial peptide expression determines host species-specific
bacterial associations. *Proceedings of the National Academy of Sciences* **110**, E3730-E3738,
doi:10.1073/pnas.1304960110 (2013).
18   Whitham, T. G. *et al.* A framework for community and ecosystem genetics: from genes to
ecosystems. *Nature Reviews Genetics* **7**, 510-523, doi:10.1038/nrg1877 (2006).

19 Shuster, S. M., Lonsdorf, E. V., Wimp, G. M., Bailey, J. K. & Whitham, T. G. Community
heritability measures the evolutionary consequences of indirect genetic effects on community
structure. *Evolution* **60**, 991-1003, doi:10.1554/05-121.1 (2006).
20 van Opstal, E. J. & Bordenstein, S. R. Rethinking heritability of the microbiome. *Science* **349**,
1172-1173, doi:10.1126/science.aab3958 (2015).
21 Bailey, J. K., Wooley, S. C., Lindroth, R. L. & Whitham, T. G. Importance of species interactions to
community heritability: a genetic basis to trophic-level interactions. *Ecology Letters* **9**, 78-85,
doi:10.1111/j.1461-0248.2005.00844.x (2006).
22 Zilber-Rosenberg, I. & Rosenberg, E. Role of microorganims in the evolution of animals and
plants: the hologenome theory of evolution. *FEMS Microbiology Reviews* **32**, 723-735,
doi:10.1111/j.1574-6976.2008.00123.x (2008).
23 Rosenberg, E., Sharon, G., Atad, I. & Zilber-Rosenberg, I. The evolution of animals and plants via
symbiosis with microorganisms. *Environmental Microbiology Reports* **2**, 500-506,
doi:10.1111/j.1758-2229.2010.00177.x (2010).
24 Rosenberg, E. & Zilber-Rosenberg, I. Symbiosis and development: The hologenome concept.
Birth Defects Research Part C: Embryo Today: Reviews **93**, 56-66, doi:10.1002/bdrc.20196
(2011).
25 Rosenberg, E., Sharon, G. & Zilber-Rosenberg, I. The hologenome theory of evolution contains
Lamarckian aspects within a Darwinian framework. *Environmental Microbiology* **11**, 2959-2962,
doi:10.1111/j.1462-2920.2009.01995.x (2009).
26 Dheilly, N. M., Poulin, R. & Thomas, F. Biological warfare: Microorganisms as drivers of host-
parasite interactions. *Infection Genetics and Evolution* **34**, 251-259,
doi:10.1016/j.meegid.2015.05.027 (2015).
27 Lloyd, E. A. Holobionts as units of selection: Holobionts as interactors, reproducers, and
manifestors of adaptation. In *Landscapes of Collectivity in the Life Sciences Vienna Series in*
Theoretical Biology (eds S.B. Gissis, E. Lamm, & A. Shavit) (MIT Press, In Press).

Environmental metagenome

Hologenome

Host genome

Microbiome

Host and symbiont genes that alone and/or together affect a holobiont phenotype

Coevolved host and symbiont genes that affect a holobiont phenotype

Host genes and symbionts that do not affect a holobiont phenotype

Environmental microbes that are not part of the holobiont