Getting the Hologenome Concept Right:

An Eco-Evolutionary Framework for Hosts and Their Microbiomes

Kevin R. Theis[1,2#], Nolwenn M. Dheilly[3], Jonathan L. Klassen[4], Robert M. Brucker[5], John F. Baines[6,7], Thomas C.G. Bosch[8], John F. Cryan[9], Scott F. Gilbert[10], Charles J. Goodnight[11], Elisabeth A. Lloyd[12], Jan Sapp[13], Philippe Vandenkoornhuyse[14], Ilana Zilber-Rosenberg[15], Eugene Rosenberg[16], Seth R. Bordenstein[17#]

[1]Department of Immunology and Microbiology, Wayne State University, Detroit, MI 48201, USA.
[2]Perinatal Research Initiative, Wayne State University, Detroit, MI 48201, USA.
[3]School of Marine and Atmospheric Sciences, Stony Brook University, Stony Brook, NY 11794-5000, USA.
[4]University of Connecticut, Department of Molecular & Cell Biology, Storrs, CT 06269, USA.
[5]The Rowland Institute at Harvard, Harvard University, Cambridge, MA 02142, USA.
[6]Max Planck Institute for Evolutionary Biology, August-Thienemann-Strasse 2, D-24306 Plön, Germany.
[7]Institute for Experimental Medicine, Christian-Albrechts-University of Kiel, Arnold-Heller-Strasse 3, D-24105 Kiel, Germany.
[8]Zoological Institute, Christian-Albrechts-University, D-24118 Kiel, Germany.
[9]Department of Anatomy and Neuroscience, University College Cork, Cork, Ireland; Alimentary Pharmabiotic Centre, University College Cork, Cork, Ireland.
[10]Department of Biology, Swarthmore College, Swarthmore, PA 19081, USA.
[11]Department of Biology, University of Vermont, Burlington, VT 05405, USA.
[12]History and Philosophy of Science Department, Indiana University, Bloomington, IN 47405, USA.
[13]Department of Biology, York University, 4700 Keele Street, Toronto, Ontario, M3J 1P3, Canada.
[14]Université de Rennes 1, CNRS, UMR6553 EcoBio, F-35042 Rennes, France.
[15]Independent Scholar, Rachavat Ilan St., Givat Shmuel 51905, Israel.
[16]Department of Molecular Microbiology and Biotechnology, Tel-Aviv University, Tel Aviv, 69978, Israel.
[17]Departments of Biological Sciences and Pathology, Microbiology and Immunology, Vanderbilt University, Nashville, TN 37225, USA.

[#]Corresponding authors:

Kevin Theis (ktheis@med.wayne.edu), Seth Bordenstein (s.bordenstein@vanderbilt.edu)

Abstract word count: 150

Text word count: 2090

Abstract

Given the complexity of host-microbiota symbioses, scientists and philosophers are asking questions at new biological levels of hierarchical organization - What is a holobiont and hologenome? When should this vocabulary be applied? Are these concepts a null hypothesis for host-microbe systems or limited to a certain spectrum of symbiotic interactions such as host-microbial coevolution? Critical discourse is necessary in this nascent area, but productive discourse requires that skeptics and proponents use the same lexicon. For instance, critiquing the hologenome concept is not synonymous with critiquing coevolution, and arguing that an entity is not a primary unit of selection dismisses that the hologenome concept has always embraced multi-level selection. Holobionts and hologenomes are incontrovertible, multipartite entities that result from ecological, evolutionary and genetic processes at varying levels. They are not restricted to one special process but constitute a wider vocabulary and framework for host biology in light of the microbiome.

Main Text

Holobiont is a term used to describe an individual host and its microbial community,

including viruses and cellular microorganisms (1-6) (Figure 1). It is derived from the Greek word

holos that means whole or entire. Microbial symbionts can be constant or inconstant, vertically

or horizontally transmitted, and can interact with a host in a context-dependent manner as

harmful, harmless or helpful. In most cases, the net outcome of these interspecies relationships

varies with the presence of other symbionts. The term holobiont distinguishes itself by not only

recognizing hosts and their obligate symbionts, but also emphasizing the diversity of facultative

symbionts and their dynamic associations within a host. In contrast to binary host-microbial

interactions, the properties of complex microbial communities and their hosts are newly

appreciated and potentially universal. The host and microbial genomes of a holobiont are

collectively defined as its hologenome (1, 2), and the pluralistic attributes of a holobiont scale

directly to the hologenome (Figure 1). Microbial genomes can be stable or labile components of

the hologenome, vertically or horizontally transmitted, and the functional traits that they encode

are context dependent and may result in damage, benefit, or be of no consequence to the

holobiont (7). Having settled on these terms, one can look at holobionts and hologenomes as

incontrovertible realities of nature. Hologenome is fundamentally similar to the words genome

and chromosome in that they reflect different levels of biological information. The terms

holobiont and hologenome are therefore structural definitions, although their utility remains

subject to debate (8).

The first question then is why are these terms useful? They are useful because they

replace misnomers in the context of host-microbiota symbioses like superorganism (i.e., an

integrated social unit comprised of conspecifics), organ, and metagenome with a vocabulary that

aligns with recent advances demonstrating that host phenotypes are profoundly affected by their

complex microbial communities, in both cooperative and competitive ways (9-11). Holobionts

and their hologenomes are less entities that elucidate something *per se* than they are entities that

need elucidation.

The next question then is what is in need of elucidation? For any given symbiosis, host

genetic variation may affect susceptibility to colonization of diverse microbes or even promote it

in a highly specific way, microbial genetic variation may favor colonization while also affecting

competition with coinfecting microbes, and environmental variation may substantively influence

these dynamics and drive rapid microbial community changes. What is then in need of

elucidation is how common and influential these forces are across host-microbial systems.

Moreover, covariance between hosts and members of their microbiota is another important area

of future research. Covariance can be achieved via vertical inheritance or selective filtering from

the environment. The relative importance of these modes of holobiont assembly is not well

resolved, yet either way, covariance in genetic compartments of the hologenome can yield

variation in phenotypes upon which evolutionary processes can act. Finally, hologenomic

variation may arise not only by mutation and recombination in the host and microbiome, but also

by acquisition of new microbial strains from the environment, microbial amplification that

involves a change in microbial abundance, and horizontal gene transfer among microbes (2).

Another area in need of elucidation is whether variation in traits caused by different host-

microbiota assemblies drives a multigenerational response to selection. If there is a response to

selection, then did it occur at the host, microbe, or microbial community levels? Can shifts at the

microbial community level act akin to shifts in allele frequencies in host genomes? Preliminary

indications are that not only can this occur, but that we can capitalize on its occurrence by

artificially selecting (i.e., microbiome engineering) holobiont phenotypes in applied contexts (12,

13).

The hologenomic view of biology importantly does not prescribe host-centric or microbe-

centric attributes to changes in holobiont macrobe functions, but rather takes into account the

emergent interactions and outcomes of hosts and their microbiota. It is a relatively new view and

is therefore liable to be interpreted in ways that misrepresent its original conception. For

example, a recent paper expressed skepticism of the hologenome concept, yet did so by relying

on alternative definitions that incorrectly restricted hologenomes to only those situations when

holobionts are primary units of selection that arose by vertical inheritance and coevolution (8).

The result is a straw man argument. The hologenome concept requires evaluation as any new

idea does. However, to have a robust debate, skeptics and proponents must use the same

terminology and framework. Here we highlight errors in these recent narrow definitions of the

holobiont and hologenome, keeping them consistent with their original pluralistic definitions,

and attempt to stimulate understanding of the link between holobiont phenotype and genotype.

The first argument proposed against the hologenome concept is that if X did not coevolve

sensu stricto with Y, then the hologenome is not real (8, 14, 15). In this case, X and Y are

respectively a microbe/microbial community and a host. As emphasized above and in the

original literature, hologenome is a term that encompasses all of the genomes of the holobiont at

a given point in time. Thus, holobionts can be formed through neutral processes, selection at the

level of the host, symbiont or both (Figure 1). Although a component of it, coevolution is not the

sole feature of the hologenome and its associated concepts. By way of illustration, one would not

similarly say that if genes X and Y did not coevolve in a host, then they are not part of the same

genome. Evolution of genomes and hologenomes is not a monolithic process, nor is it simply

beanbag genetics. Genetic conflict, epistasis, selection, drift, etc. are all operational (1, 2). Thus,

objections to the hologenome concept based on a lack of coevolution misrepresent what

constitutes a hologenome and holobiont for that matter. To put it simply, coevolution is a

process; the hologenome is an entity that embraces the eco-evolutionary processes inherent in

much of macroscopic biology.

When referencing the original definitions of the hologenome, it was suggested that a non-

coevolutionary application of the word hologenome would make it "sufficiently general that it

can be interpreted in any number of ways" (8). This comment refers to the more generally

accepted definition of the hologenome as all of the genomes in the holobiont, all of which in turn

are evolving in that context (6). However, using the same logic, the word genome would be as

unhelpful to biology as the term hologenome because it would be an insufficiently general

definition of the types of evolutionary processes occurring within the genome. The main lesson

here is that coevolution, genetic conflict, selection, and drift at multiple levels (host genomes,

symbiont genomes, hologenomes) all occur. In arguing for a hologenomic status of macro-

organisms, we noted that interspecies interactions underlying holobiont phenotypes follow a

conceptual and theoretical continuum from genetic interactions or epistasis between genes in the

same genome (1, 16). Both types of interactions can be transient or stable under varying

conditions such as population structure and selection. Indeed, the (in)stability of a host and its

horizontally-transmitted microbes follows a theoretical continuum under the same math from the

(in)stability of interacting genes in the same genome that undergo recombination (16). Here we

note that vertical transmission versus horizontal transmission is a false dichotomy to draw

against the hologenome concept.

140   Prevalent misuse of coevolution in the microbiome literature is a legitimate concern and

141 was the impetus for some of us coining the word "phylosymbiosis" (17). It describes the

142 concordance between a host phylogeny (evolutionary relationships) and microbial community

143 dendrogram (ecological relationships) based on the degree of shared taxonomy and/or abundance

144 of members of the community (18-21). Phylosymbiosis does not *a priori* imply coevolution,

145 cospeciation, cocladogenesis, or codiversification because this latter vocabulary implies

146 concordant splitting of new species from a common ancestral one (19, 20, 22). Phylosymbiosis

147 avoids these assumptions because it "does not presume that microbial communities are stable or

148 even vertically transmitted from generation to generation" (19, 20). Rather, it refers to a pattern

149 in which changes in separate parts of the holobiont (host and microbiota) are related in a

150 concordant manner. It is also a stepping-stone from population genetics to community genetics

151 because when phylosymbiosis is observed under strictly controlled conditions, it tests whether

152 variation in holobiont assembly is primarily stochastic or deterministic (17, 19, 23). Stochastic

153 assembly means that each microbe has an equal opportunity of colonizing a host. Deterministic

154 assembly reflects ecological selection of a particular non-random microbial community and its

155 host, without reference to which partner, or potentially both, is doing the selecting, and it can be

156 affected by genetic variation in the host or microbial species. Controlled studies of microbial

157 community assembly across different species of *Nasonia* wasps and *Hydra* have yielded such

158 phylosymbiotic patterns (17, 18). When genetic variation in the interacting species affects

159 community assembly, it has been defined as broad sense "community heritability," or H^2_C (24,

160 25). Similar to population genetic heritability estimates of phenotypes that are abiotic, H^2_C

161 measures a "heritable basis to trophic-level interactions" (26). If there is a significant H^2_C, natural

selection can act on genetic variation affecting ecological community structure (23, 27),

including organization of the holobiont and its emergent phenotypes (25).

Discussion of evolutionary processes brings forth a second argument against the

hologenome concept, namely that holobionts and their hologenomes must be the "primary" unit

of selection (8). This strict claim leads biologists into error, as all of the literature emphasizes the

reality that multiple levels of selection can operate simultaneously. For example, selfish genetic

elements can be selected within a genome that is in turn selected for any number of phenotypes

that affect fitness—this is uncontroversial. While the holobiont is posited to be "a unit of

selection in evolution" (2, 28-30), it is naturally not proposed as the only or necessarily primary

unit of selection (1, 2). Primariness varies with what traits are targeted by natural selection.

As we have emphasized in different venues, it is also true that just as large parts of the

nuclear genome can evolve neutrally or be in conflict, so too can large parts of the hologenome

(1, 2). For example, "hologenomic drift can occur at all the different levels of the holobiont from

single genes of the microbes or the host to the holobiont itself" (2). We would be remiss to not be

critical of our own inconsistent statements about the relative roles of cooperation and conflict in

hologenomic evolution. In *The Hologenome Concept*, some of us stated that "evolution of

animals and plants was driven primarily by natural selection for cooperation between and with

microorganisms" (2) while in other venues the concept "places as much emphasis on cooperation

as on competition" (31). This latter statement is more precisely aligned with the pluralistic nature

of the holobiont, namely that "natural selection...on holobiont phenotypes...can work to remove

deleterious nuclear mutations or microbes while spreading advantageous nuclear mutations or

microbes" (1). In fact, some of us argued that conflicts of interests resulting from the nature of

the transmission of microbes to the next host could select for microbes that can manipulate the

biology of their host to improve their own transmission (32). The holobiont is not a conglomerate

that arises solely from cooperation. Rather, it is a hierarchical level that can supersede the

individual host that lives in association with its microbial community, incorporating both

competitive and cooperative selective systems (33). Hologenomes then exist as hierarchically

nested, although not necessarily integrated levels of genomes, in which all levels of selection are

in play.

In summary, we anticipate that many subdisciplines in biology will benefit from a

conceptual, theoretical, and experimental framework that broadly encompasses the ecology of

holobionts and evolution of hologenomes. The hologenome concept is a comprehensive and

relevant eco-evolutionary framework for which critical questions remain. For example, can a

response to selection on host traits be driven solely by changes in the genomes and/or

membership of a host-associated microbial community? How taxonomically widespread among

hosts is phylosymbiosis? How common is vertical inheritance of complex microbial

communities? What is the strength of selection required to maintain consistent association

between a host and environmentally-acquired microbes each generation? How does selection

operate on community phenotypes if H^2_C is variable due to the lability of microbial

communities? Evolution of the hologenome refers to the genetic basis of eco-evolutionary

processes underlying community phenotypes of the holobiont. This terminology and framework

for the newly appreciated complexities in the host-microbe consortia and their genomes is not

restricted to one special process but constitute an incontrovertible vocabulary and framework for

host biology in light of the microbiome.

Figure Legend

Figure 1. Holobionts are entities comprised of the host and all of its symbiotic microbes, including those which affect the holobiont's phenotype and have coevolved with the host (blue), those which affect the holobiont's phenotype but have not coevolved with the host (red), and those which do not affect the holobiont's phenotype at all (gray). Microbes may be transmitted vertically or horizontally, acquired from the environment, and can be constant or inconstant in the host. Therefore, holobiont phenotypes can change in time and space as microbes come into and out of the holobiont. Microbes in the environment are not part of the holobiont (white). Hologenomes then encompass the genomes of the host and all of its microbes at any given time point, with individual genomes and genes falling into the same three functional categories of blue, red and gray. Holobionts and hologenomes are entities, whereas coevolution or host-symbiont interactions are processes.

219 **Acknowledgements**

220   We thank Jay Evans and Phil Pellett for helpful feedback on the manuscript.

References

1. **Bordenstein SR, Theis KR.** 2015. Host biology in light of the microbiome: Ten principles of holobionts and hologenomes. PLoS Biol **13:**e1002226.
2. **Rosenberg E, Zilber-Rosenberg I.** 2013. The Hologenome Concept: Human, Animal and Plant Microbiota. Springer, Switzerland.
3. **Margulis L.** 1991. Symbiogenesis and symbionticism, p 1-14. *In* Margulis L, Fester R (ed), Symbiosis as a Source of Evolutionary Innovation: Speciation and Morphogenesis. MIT Press, Cambridge, MA.
4. **Mindell DP.** 1992. Phylogenetic consequences of symbioses: Eukarya and Eubacteria are not monophyletic taxa. Biosystems **27:**53-62.
5. **Rohwer F, Seguritan V, Azam F, Knowlton N.** 2002. Diversity and distribution of coral-associated bacteria. Marine Ecology Progress Series **243:**1-10.
6. **Gordon J, Knowlton N, Relman DA, Rohwer F, Youle M.** 2013. Superorganisms and holobionts, vol 8, p 2. American Society for Microbiology, Washington, DC.
7. **Casadevall A, Pirofski LA.** 2015. What is a host? Incorporating the microbiota into the damage-response framework. Infection and Immunity **83:**2-7.
8. **Moran NA, Sloan DB.** 2015. The hologenome concept: Helpful or hollow? PLoS Biol **13:**e1002311.
9. **Vandenkoornhuyse P, Quaiser A, Duhamel M, Le Van A, Dufresne A.** 2015. The importance of the microbiome of the plant holobiont. New Phytologist **206:**1196-1206.
10. **McFall-Ngai M, Hadfield MG, Bosch TCG, Carey HV, Domazet-Lošo T, Douglas AE, Dubilier N, Eberl G, Fukami T, Gilbert SF, Hentschel U, King N, Kjelleberg S, Knoll AH, Kremer N, Mazmanian SK, Metcalf JL, Nealson K, Pierce NE, Rawls JF, Reid A, Ruby EG, Rumpho M, Sanders JG, Tautz D, Wernegreen JJ.** 2013. Animals in a bacterial world, a new imperative for the life sciences. Proceedings of the National Academy of Sciences **110:**3229-3236.
11. **Bosch TCG, McFall-Ngai MJ.** 2011. Metaorganisms as the new frontier. Zoology **114:**185-190.
12. **Mueller UG, Sachs JL.** 2015. Engineering microbiomes to improve plant and animal health. Trends in Microbiology **23:**606-617.
13. **Voss JD, Leon JC, Dhurandhar NV, Robb FT.** 2015. Pawnobiome: manipulation of the hologenome within one host generation and beyond. Frontiers in Microbiology **6:**5.
14. **Hester ER, Barott KL, Nulton J, Vermeij MJA, Rohwer FL.** 2016. Stable and sporadic symbiotic communities of coral and algal holobionts. The ISME Journal **In Press**.
15. **Chandler JA, Turelli M.** 2014. Comment on "The hologenomic basis of speciation: Gut bacteria cause hybrid lethality in the genus *Nasonia*". Science **345:**2.
16. **Fitzpatrick BM.** 2014. Symbiote transmission and maintenance of extra-genomic associations. Frontiers in Microbiology **5**.
17. **Brucker RM, Bordenstein SR.** 2013. The hologenomic basis of speciation: gut bacteria cause hybrid lethality in the genus *Nasonia*. Science **341:**667-669.
18. **Fraune S, Bosch TCG.** 2007. Long-term maintenance of species-specific bacterial microbiota in the basal metazoan Hydra. Proceedings of the National Academy of Sciences **104:**13146-13151.

19.   **Brucker RM, Bordenstein SR.** 2012. The roles of host evolutionary relationships
(Genus: *Nasonia*) and development in structuring microbial communities. Evolution
66:349-362.
20.   **Brucker RM, Bordenstein SR.** 2012. Speciation by symbiosis. Trends in Ecology &
Evolution **27:**443-451.
21.   **Franzenburg S, Walter J, Kunzel S, Wang J, Baines JF, Bosch TCG, Fraune S.**
2013. Distinct antimicrobial peptide expression determines host species-specific bacterial
associations. Proceedings of the National Academy of Sciences of the United States of
America **110:**E3730-E3738.
22.   **de Vienne DM, Refregier G, Lopez-Villavicencio M, Tellier A, Hood ME, Giraud T.**
2013. Cospeciation vs host-shift speciation: methods for testing, evidence from natural
associations and relation to coevolution. New Phytologist **198:**347-385.
23.   **Whitham TG, Bailey JK, Schweitzer JA, Shuster SM, Bangert RK, Leroy CJ,**
Lonsdorf EV, Allan GJ, DiFazio SP, Potts BM, Fischer DG, Gehring CA, Lindroth
RL, Marks JC, Hart SC, Wimp GM, Wooley SC. 2006. A framework for community
and ecosystem genetics: from genes to ecosystems. Nature Reviews Genetics **7:**510-523.
24.   **Shuster SM, Lonsdorf EV, Wimp GM, Bailey JK, Whitham TG.** 2006. Community
heritability measures the evolutionary consequences of indirect genetic effects on
community structure. Evolution **60:**991-1003.
25.   **van Opstal EJ, Bordenstein SR.** 2015. Rethinking heritability of the microbiome.
Science **349:**1172-1173.
26.   **Bailey JK, Wooley SC, Lindroth RL, Whitham TG.** 2006. Importance of species
interactions to community heritability: a genetic basis to trophic-level interactions.
Ecology Letters **9:**78-85.
27.   **Goodnight CJ.** 2011. Evolution in metacommunities. Philosophical Transactions of the
Royal Society B-Biological Sciences **366:**1401-1409.
28.   **Zilber-Rosenberg I, Rosenberg E.** 2008. Role of microorganims in the evolution of
animals and plants: the hologenome theory of evolution. FEMS Microbiology Reviews
32:723-735.
29.   **Rosenberg E, Sharon G, Atad I, Zilber-Rosenberg I.** 2010. The evolution of animals
and plants via symbiosis with microorganisms. Environmental Microbiology Reports
2:500-506.
30.   **Rosenberg E, Zilber-Rosenberg I.** 2011. Symbiosis and development: The hologenome
concept. Birth Defects Research Part C: Embryo Today: Reviews **93:**56-66.
31.   **Rosenberg E, Sharon G, Zilber-Rosenberg I.** 2009. The hologenome theory of
evolution contains Lamarckian aspects within a Darwinian framework. Environmental
Microbiology **11:**2959-2962.
32.   **Dheilly NM, Poulin R, Thomas F.** 2015. Biological warfare: Microorganisms as drivers
of host-parasite interactions. Infection Genetics and Evolution **34:**251-259.
33.   **Lloyd EA.** In Press. Holobionts as units of selection: Holobionts as interactors,
reproducers, and manifestors of adaptation. *In* Gissis SB, Lamm E, Shavit A (ed),
Landscapes of Collectivity in the Life Sciences. MIT Press.

Environmental metagenome

Hologenome

Host genome

Microbiome

Host and symbiont genes that alone and/or together affect a holobiont phenotype

Coevolved host and symbiont genes that affect a holobiont phenotype

Host genes and symbionts that do not affect a holobiont phenotype

Environmental microbes that are not part of the holobiont