1 **Infection avoidance behaviour in adult fruit flies is sex-specific and**
2 **depends on prior exposure to a viral pathogen**
3

4 Pedro F. Vale[1,2,*], Michael D. Jardine[1]
5
6 [1] Institute of Evolutionary Biology, School of Biological Sciences, University of
7 Edinburgh. Edinburgh EH9-3FL
8
9 [2] Centre for Immunity, Infection and Evolution, University of Edinburgh.
10 Edinburgh EH9 3FL
11
12
* Email for correspondence : pedro.vale@ed.ac.uk

Abstract

Infection avoidance behaviours are the first line of defence against pathogenic
encounters. Behavioural plasticity in response to internal or external cues can
therefore generate heterogeneity in infection. We tested whether *Drosophila*
melanogaster exhibits infection avoidance behaviour during foraging, and
whether this behaviour is modified by prior exposure to Drosophila C Virus
(DCV) and by the risk of DCV encounter. We examined two measures of
infection avoidance: (1) the motivation to feed in the presence of an infection
risk and (2) the preference to feed on a clean food source over a potentially
infectious source. We found no clear evidence for preference of clean food
sources over potentially infectious ones. However, infection avoidance was
present in female fruit, which were less motivated to feed when presented
with a risk of encountering DCV, but this was only the case if they had been
previously exposed to this viral pathogen. We discuss the relevance of
plasticity in avoidance behaviours during ecologically relevant scenarios such
as foraging for host fitness and pathogen spread.

Key-words: Infection, avoidance behaviour, Drosophila, DCV, foraging

Background

Hosts vary considerably in their ability to acquire and transmit infection (Barron et al., 2015; Fellous et al., 2012; Paull et al., 2011; Susi et al., 2015; Vale and Little, 2009). Given the ubiquitous presence of pathogens and parasites in natural environments, mounting a timely and efficient immune response to all possible pathogenic challenges would be physiologically costly and ultimately ineffective. Individuals capable of reducing the probability of contacting parasites, infected conspecifics or infectious environments can therefore not only prevent the deleterious effects of infection, but also circumvent the undesirable energetic costs of immune responses, including immunopathology (Barron et al., 2015; Curtis, 2014). Avoiding infection is therefore the first line of non-immunological defence against infection(Parker et al., 2011), and it is known to occur across a broad range of host taxa, including humans (Curtis, 2014; Moore, 2013).

Like most traits, infection avoidance behaviours are likely to vary according to the context of infection, and pathogens are major drivers this context (Barron et al., 2015; Curtis, 2014; Lazzaro and Little, 2009; Moore, 2013; Vale et al., 2008; Wolinska and King, 2009). Pathogens may alter host responses in two ways. First, by altering the immuno-physiology of the host during infection, pathogens can alter host behaviour (Adamo, 2006; Adelman and Martin, 2009). Second, pathogens also modify the host external environment by increasing the likelihood of exposure to novel infections, and these external cues of infection risk are also known to influence host behavioural responses (Barron et al., 2015; Curtis, 2014). Understanding variation in infection avoidance behaviours therefore provides an important functional link between the neurological, behavioural and immunological processes that together govern the spread of disease (Adamo, 2006).

Insects are ideal systems to investigate the interplay between infection and behaviour (Adamo, 2006; Parker et al., 2010). The fruit fly *Drosophila* is especially amenable to these studies, as it is one of the best model systems for host-pathogen interactions (Neyen et al., 2014) and behavioural ecology and genetics (Dubnau, 2014; Sokolowski, 2001). One of the best-studied pathogenic interactions in *Drosophila* is the host response to systemic and

enteric infection with Drosophila C Virus (DCV) (Dostert et al., 2005; Ferreira

et al., 2014). DCV is a horizontally transmitted +ssRNA virus that naturally

infects the fly gut (Huszar and Imler, 2008), causing intestinal obstruction,

severe metabolic dysfunction and eventually death (Chtarbanova et al., 2014).

As a consequence of its pathology, female flies infected with DCV are also

known to exhibit behavioural modifications, such as reduced locomotion and

increased sleep (Vale and Jardine, 2015). The Drosophila-DCV interaction

therefore offers a powerful system to investigate the ecological consequences

that may arise from the physiological and behavioural effects of enteric viral

infections.

In the present study we used a combination of controlled experimental

infections and foraging choice assays, to test whether adult *D. melanogaster*

are able to avoid potentially infectious environments when foraging for food,

and if avoidance behaviour is modified in response to virus exposure history

and to different risks of acquiring DCV infection. We find evidence for

avoidance behaviours in the form of reduced motivation to feed according to

the risk of infection. However, these effects were only present in female flies,

indicating potentially important sexual dimorphism in infection avoidance,

and were only present when females were previously exposed to DCV.

Methods

Fly and virus stocks

All flies used were from a long-term laboratory stock of Wolbachia-

free *Drosophila melanogaster* Oregon R line, maintained on Lewis medium in

standard conditions: 25°C, with a 16:8h light:dark cycle. Fly stocks were

routinely kept on a 14-day cycle with non-overlapping generations under low

larval densities. The DCV culture used in this experiment was grown in

Schneider Drosophila Line 2 (DL2) as described in (Vale and Jardine, 2015).

Ten-fold serial dilutions of this culture (diluted in Ringers buffer solution)

were aliquoted and frozen at -80ºC for long-term storage before use.

Prior virus exposure

Flies used in the foraging choice assays were obtained by preparing 10

vials of Lewis medium and yeast containing ten mated females. Flies were

allowed to lay eggs for 48 hours resulting in age-matched progeny reared in

similar larval densities. Two to three days after eclosion, these progeny were

exposed to DCV via the oral route of infection (Vale and Jardine, 2015), in

order to test the effect of previous exposure to virus on avoidance behaviour

during foraging. Briefly, single-sex groups of 20 flies were placed in vials

containing agar previously sprayed with DCV ("exposed" to 50 µl of 10^8 viral

copies/ml) or the equivalent volume of Ringers buffer solution as a control

("not exposed"). This procedure produced 10 replicate vials of either healthy

or virus-exposed male or female flies. DCV exposure using this protocol

typically results in ~20% mortality (Fig. S1 and (Vale and Jardine, 2015)).

Foraging choice assays

Following 5 days of virus exposure, we set up independent foraging

choice assays in cages - cylindrical transparent plastic containers (12 cm in

diameter) containing two equally spaced plastic vials of standard Lewis fly

medium and yeast. For each combination of "exposed" and "not exposed" male

or female flies, we set up two sets of cages to simulate different risks of

infection: a "no risk" environment, with two clean vials (sprayed with sterile

Ringers solution), and a "high-risk" environment where one of the vials was

sprayed with DCV, as described above. Six replicate 20-fly groups were

allocated to the "high-risk" chambers and four replicates to the "no risk"

chambers, resulting in a total of 40 independent foraging choice cages. Flies

were added to the chamber from a neutrally placed hole in the lid, and the

number of flies that settled on each vial was recorded every 30 minutes for six

hours. Care was taken to randomise the position of the cages so that the

orientation of the light did not influence the choice of the flies in any

systematic way.

Statistical Analysis

To measure infection avoidance, we took two approaches. First, we

hypothesised that the motivation to feed would be lower in environments

where the risk of infection is higher (Curtis, 2014). We therefore compared

the motivation to feed between the "no risk" and "high-risk" cages, measured

as the proportion of flies inside a cage that chose to feed on any of the provided food sources. We also asked whether flies that chose to feed showed any evidence of avoiding potentially infectious food sources. For this analysis we focussed on the proportion of flies choosing the clean food source over the infectious food source in the "high risk" cages. In both analyses of 'motivation to feed' and 'infection avoidance', data on the proportion of flies choosing each food source within each replicate cage were analysed with a generalised linear model assuming binomial error and logit link function, and included fly 'sex', 'previous exposure' and 'infection risk' as fixed effects. 'Replicate cage' was included as a random effect, nested within treatments. We also analysed the average motivation to feed and infection avoidance across all time points, in a model including "time" as a random effect. Treatment specific contrasts were used to test the significance of pairwise comparisons. Analyses were carried out using JMP 12 (JMP).

Results

Once inside the cages, only a fraction of flies chose either of the food sources provided, and this motivation to feed increased over time for flies in all treatment groups (χ^2_1= 11.00, p=0.001; Fig. S2). The rate at which motivation increased differed between sexes ('Time × Sex' interaction, χ^2_1= 12.47, p=0.0004), and on average female flies showed greater motivation to feed than males (χ^2_1= 5.01, p=0.025), with 67% of female and 36% of male flies making a choice to feed on any of the provided substrates during the 6 hours of observation (Fig. 1).

Across the entire six-hour observation period, the motivation to feed differed between sexes, and depended both on their previous exposure and on their current risk of infection ('Sex' × 'risk of infection' × 'Previous exposure' interaction, χ^2_1= 21.82, p<0.0001). The proportion of males choosing any food substrate did not vary with previous exposure to DCV in either high-risk (χ^2_1= 2.21, p=0.137) or no-risk environments (χ^2_1= 0.09, p=0.764; Fig. 1).

In female flies however, previous exposure and current infection risk affected the motivation to feed on the provided food sources. When there was no risk of infection (Fig. 1, light grey bars) the motivation to feed was greater

in females that were previously exposed to DCV than in otherwise healthy,

non-exposed females (χ^2_1= 104.11, p<0.001). Among females that were

previously exposed to infection, we found that the presence of a risk of

acquiring infection resulted in lower foraging effort - with just over 50% of

flies making the choice to feed - compared to females in cages where there

was no risk of acquiring infection, where over 80% of flies made the choice to

feed (Fig. 1; χ^2_1= 168.48, p<0.001).

Once flies had made the choice to feed on one of the provided food

sources, the choice between a clean and a potentially infectious food source

was not affected by previous exposure to DCV ('previous exposure', χ^2_1=

0.513, p=0.47) in either male or females ('sex', χ^2_1= 0.595, p=0.44; Fig. 2).

Discussion

The ability to detect and discriminate between clean and potentially

infectious environments is vital to avoid the adverse consequences of

infection. In this study we tested if infection avoidance behaviour in

Drosophila melanogaster is modified by its previous exposure to a viral

pathogen and by the risk of infection with that same pathogen when

encountered during foraging.

The higher motivation to feed of some female flies when the risk of infection

was absent (Fig. 1) suggests flies were able to identify external cues of

infection risk. Identifying infection cues is a general prerequisite of avoidance

behaviours and occurs across a wide range of different taxa. For example,

lobsters are known to detect and avoid virus-infected conspecifics (Behringer

et al., 2006); fruit flies and nematodes are capable of avoiding pathogenic

bacteria (Babin et al., 2014; Meisel and Kim, 2014); gypsy moth larvae are able

to detect and avoid virus-contaminated foliage (Parker et al., 2010); sheep

have been found to prefer to graze in parasite-poor patches (Hutchings et al.,

2007); and it is has been argued that the disgust response in humans has

evolved because it decreases contact with potential infection (Curtis et al.,

2011). It is unclear how flies are able to detect food sources contaminated

with a viral pathogen. In *Drosophila* and *C. elegans* avoidance of pathogenic

bacteria is enabled by evolutionary conserved olfactory and chemosensory

pathways (Babin et al., 2014; Meisel and Kim, 2014), while avoidance of
parasitic wasps appears to be mainly enabled by the visual sensory system
(Kacsoh et al., 2013). While avoiding virus infected conspecifics is probably
driven by visual cues of infection (Behringer et al., 2006), it remains unclear
how virus-contaminated environments may be detected by *Drosophila*.

Infection avoidance, measured as a reduced motivation to feed, was
clearest when flies had been previously exposed to infection (Fig. 1). In
addition to responding to external cues of infection, internal physiological
cues therefore also modify avoidance behaviour. Behavioural modifications
due to infection are widely reported among animals (Barber and Dingemanse,
2010; Moore, 2013), and can be classified into (i) parasitic manipulation that
enhances parasite transmission (Moore, 2013) (ii) sickness behaviours that
benefit the host by conserving energetic resources during infection (Adelman
and Martin, 2009), or (iii) side-effects of pathogenicity that do not benefit the
host or the parasite (Barber and Dingemanse, 2010). Female flies infected
with DCV are known to experience increased lethargy and sleep (Vale and
Jardine, 2015), so these effects could also explain the reduced feeding activity
we detected in female flies that had been previously exposed to DCV. Another
potential explanation for reduced motivation to feed in previously exposed
flies is infection-induced anorexia (Ayres and Schneider, 2009), a commonly
described sickness behaviour (Adelman and Martin, 2009). However, it is
unlikely that a lower motivation feed is simply a symptom of a "sick" fly,
because it varied according risk of infection, and even reached 80% in
exposed flies when foraging in a 'no risk' environment (Fig. 1). This suggests
that flies are actively avoiding contact with the potentially contagious food
source by lowering their foraging effort.

The fact that only female flies demonstrated avoidance is an indication
that any potentially adaptive effects of avoiding infection may be related to
oviposition, which coincides with feeding. For flies previously exposed to DCV,
avoiding infection would not confer substantial direct benefits given the
physiological and behavioural costs of this infection (Arnold et al., 2013;
Chtarbanova et al., 2014; Vale and Jardine, 2015), but would however reduce
the exposure of future offspring to infection. While flies previously exposed to

DCV do not appear to immune primed following an initial viral exposure
(Longdon et al., 2013), our results point to a sort of behavioural priming,
where females previously exposed to infection avoid foraging in potentially
infectious environments.
246
Conclusions
Using a combination of experimental infections and behavioural
assays, we find evidence for avoidance behaviours in *Drosophila* in the form of
reduced motivation to feed, which was most pronounced when flies were
faced with an increased risk of encountering an infectious food source.
However, these effects were only present in female flies, indicating potentially
important sexual dimorphism in infection avoidance, and were only present
when females were previously exposed to DCV. Understanding how avoidance
behaviours may vary is therefore important for our understanding of how
disease will spread in natural populations (Barron et al., 2015), and more
broadly how pathogens might evolve in response to variation in host
responses to infection (Boots and Bowers, 1999; McLeod and Day, 2015).
259

260 **Competing interests**
261 The authors declare that they have no competing interests.
262

263 **Author contributions**
264 PFV conceived the study. PFV and MDJ designed the experiment. MDJ carried
265 out the experimental work. PFV analysed the data, wrote the manuscript and
266 provided all research consumables.
267

Acknowledgements
We thank H. Cowan and H. Borthwick for helpful technical assistance and the
members of the Vale, Obbard and Ross labs for helpful suggestions. This work
was supported by a strategic award from the Wellcome Trust for the Centre
for Immunity, Infection and Evolution (grant reference no. 095831), and by a
Society in Science – Branco Weiss fellowship, both awarded to P. Vale. P.Vale is
also supported by a Chancellor's Fellowship from the School of Biological
Sciences, University of Edinburgh.

Figure legends

Fig. 1. The motivation to feed, measured as the proportion of flies in the cage

that fed on any of the provided food sources. Single sex-groups of flies that

had either been previously exposed to DCV or to a sterile inoculum were

tested in a no-risk environment (choice between two clean vials; light grey) or

a high-risk environment (choice between a clean vial and a DCV-contaminated

vial; dark grey). Data are means ± SE.

Fig. 2. Infection avoidance, measured as the proportion of flies in the cage that

preferred to settle on the clean food source relative to the DCV-contaminated

food source. Single sex-groups of flies that had either been previously exposed

to DCV or to a sterile inoculum were tested in a high-risk environment (choice

between a clean vial and a DCV-contaminated vial). Data are means ± SE.

References

Adamo, S. A. (2006). Comparative psychoneuroimmunology: evidence from
the insects. *Behav. Cogn. Neurosci. Rev.* 5, 128–140.
doi:10.1177/1534582306289580.

Adelman, J. S., and Martin, L. B. (2009). Vertebrate sickness behaviors:
Adaptive and integrated neuroendocrine immune responses. *Integr.*
Comp. Biol. 49, 202–214. doi:10.1093/icb/icp028.

Arnold, P. A., Johnson, K. N., and White, C. R. (2013). Physiological and
metabolic consequences of viral infection in Drosophila melanogaster.
J. Exp. Biol. 216, 3350–3357. doi:10.1242/jeb.088138.

Ayres, J. S., and Schneider, D. S. (2009). The Role of Anorexia in Resistance and
Tolerance to Infections in Drosophila. *PLoS Biol* 7, e1000150.
doi:10.1371/journal.pbio.1000150.

Babin, A., Kolly, S., Schneider, F., Dolivo, V., Zini, M., and Kawecki, T. J. (2014).
Fruit flies learn to avoid odours associated with virulent infection. *Biol.*
Lett. 10, 20140048. doi:10.1098/rsbl.2014.0048.

Barber, I., and Dingemanse, N. J. (2010). Parasitism and the evolutionary
ecology of animal personality. *Philos. Trans. R. Soc. Lond. B Biol. Sci.*
365, 4077–4088. doi:10.1098/rstb.2010.0182.

Barron, D., Gervasi, S., Pruitt, J., and Martin, L. (2015). Behavioral competence:
how host behaviors can interact to influence parasite transmission
risk. *Curr. Opin. Behav. Sci.* 6, 35–40.
doi:10.1016/j.cobeha.2015.08.002.

Behringer, D. C., Butler, M. J., and Shields, J. D. (2006). Ecology: Avoidance of
disease by social lobsters. *Nature* 441, 421–421.
doi:10.1038/441421a.

Boots, M., and Bowers, R. G. (1999). Three mechanisms of host resistance to
microparasites-avoidance, recovery and tolerance-show different
evolutionary dynamics. *J. Theor. Biol.* 201, 13–23.
doi:10.1006/jtbi.1999.1009.

Chtarbanova, S., Lamiable, O., Lee, K.-Z., Galiana, D., Troxler, L., Meignin, C., et
al. (2014). Drosophila C virus systemic infection leads to intestinal
obstruction. *J. Virol.* doi:10.1128/JVI.02320-14.

Curtis, V. A. (2014). Infection-avoidance behaviour in humans and other
animals. *Trends Immunol.* 35, 457–464. doi:10.1016/j.it.2014.08.006.

Curtis, V., de Barra, M., and Aunger, R. (2011). Disgust as an adaptive system
for disease avoidance behaviour. *Philos. Trans. R. Soc. Lond. B. Biol. Sci.*
366, 389–401. doi:10.1098/rstb.2010.0117.

Dostert, C., Jouanguy, E., Irving, P., Troxler, L., Galiana-Arnoux, D., Hetru, C., et
al. (2005). The Jak-STAT signaling pathway is required but not

331   sufficient for the antiviral response of drosophila. *Nat. Immunol.* 6,
332   946–953. doi:10.1038/ni1237.

333   Dubnau, J. (2014). *Behavioral Genetics of the Fly (Drosophila Melanogaster)*.
334   Cambridge University Press.

335   Fellous, S., Duncan, A. B., Quillery, E., Vale, P. F., and Kaltz, O. (2012). Genetic
336   influence on disease spread following arrival of infected carriers. *Ecol.*
337   *Lett.* 15, 186–192. doi:10.1111/j.1461-0248.2011.01723.x.

338   Ferreira, Á. G., Naylor, H., Esteves, S. S., Pais, I. S., Martins, N. E., and Teixeira, L.
339   (2014). The Toll-Dorsal Pathway Is Required for Resistance to Viral
340   Oral Infection in Drosophila. *PLoS Pathog.* 10.
341   doi:10.1371/journal.ppat.1004507.

342   Huszar, T., and Imler, J. (2008). "Drosophila Viruses and the Study of Antiviral
343   Host-Defense," in *Advances in Virus Research* (Academic Press), 227–
344   265. Available at:
345   http://www.sciencedirect.com/science/article/pii/S0065352708004
346   065.

347   Hutchings, M. ., Knowler, K. ., McAnulty, R., and McEwan, J. . (2007). Genetically
348   resistant sheep avoid parasites to a greater extent than do susceptible
349   sheep. *Proc. R. Soc. B Biol. Sci.* 274, 1839–1844.
350   doi:10.1098/rspb.2007.0398.

351   JMP Cary, NC: SAS Institute Inc.

352   Kacsoh, B. Z., Lynch, Z. R., Mortimer, N. T., and Schlenke, T. A. (2013). Fruit
353   Flies Medicate Offspring After Seeing Parasites. *Science* 339, 947–950.
354   doi:10.1126/science.1229625.

355   Lazzaro, B. P., and Little, T. J. (2009). Immunity in a variable world. *Philos.*
356   *Trans. R. Soc. Lond. B. Biol. Sci.* 364, 15–26.
357   doi:10.1098/rstb.2008.0141.

358   Longdon, B., Cao, C., Martinez, J., and Jiggins, F. M. (2013). Previous Exposure
359   to an RNA Virus Does Not Protect against Subsequent Infection in
360   Drosophila melanogaster. *PLoS ONE* 8, e73833.
361   doi:10.1371/journal.pone.0073833.

362   McLeod, D. V., and Day, T. (2015). Pathogen evolution under host avoidance
363   plasticity. *Proc. Biol. Sci.* 282. doi:10.1098/rspb.2015.1656.

364   Meisel, J. D., and Kim, D. H. (2014). Behavioral avoidance of pathogenic
365   bacteria by Caenorhabditis elegans. *Trends Immunol.* 35, 465–470.
366   doi:10.1016/j.it.2014.08.008.

367   Moore, J. (2013). An overview of parasite-induced behavioral alterations – and
368   some lessons from bats. *J. Exp. Biol.* 216, 11–17.
369   doi:10.1242/jeb.074088.

Neyen, C., Bretscher, A. J., Binggeli, O., and Lemaitre, B. (2014). Methods to
study Drosophila immunity. *Methods San Diego Calif* 68, 116–128.
doi:10.1016/j.ymeth.2014.02.023.

Parker, B. J., Barribeau, S. M., Laughton, A. M., de Roode, J. C., and Gerardo, N.
M. (2011). Non-immunological defense in an evolutionary framework.
Trends Ecol. Evol. 26, 242–248. doi:10.1016/j.tree.2011.02.005.

Parker, B. J., Elderd, B. D., and Dwyer, G. (2010). Host behaviour and exposure
risk in an insect-pathogen interaction. *J. Anim. Ecol.* 79, 863–870.
doi:10.1111/j.1365-2656.2010.01690.x.

Paull, S. H., Song, S., McClure, K. M., Sackett, L. C., Kilpatrick, A. M., and Johnson,
P. T. (2011). From superspreaders to disease hotspots: linking
transmission across hosts and space. *Front. Ecol. Environ.* 10, 75–82.
doi:10.1890/110111.

Sokolowski, M. B. (2001). Drosophila: Genetics meets behaviour. *Nat. Rev.*
Genet. 2, 879–890. doi:10.1038/35098592.

Susi, H., Barrès, B., Vale, P. F., and Laine, A.-L. (2015). Co-infection alters
population dynamics of infectious disease. *Nat. Commun.* 6.
doi:10.1038/ncomms6975.

Vale, P. F., and Jardine, M. D. (2015). Sex-specific behavioural symptoms of
viral gut infection and Wolbachia in Drosophila melanogaster. *J. Insect*
Physiol. 82, 28–32. doi:10.1016/j.jinsphys.2015.08.005.

Vale, P. F., and Little, T. J. (2009). Measuring parasite fitness under genetic and
thermal variation. *Heredity* 103, 102–109.

Vale, P. F., Salvaudon, L., Kaltz, O., and Fellous, S. (2008). The role of the
environment in the evolutionary ecology of host parasite interactions.
Infect. Genet. Evol. 8, 302–305. doi:10.1016/j.meegid.2008.01.011.

Wolinska, J., and King, K. C. (2009). Environment can alter selection in host-
parasite interactions. *Trends Parasitol.* 25, 236–244.
doi:10.1016/j.pt.2009.02.004.