Infection avoidance behaviour: female fruit flies adjust foraging effort in

response to internal and external cues of viral infection

Pedro F. Vale[1,2,*], Michael D. Jardine[1]

[1] Institute of Evolutionary Biology, School of Biological Sciences, University of

Edinburgh. Edinburgh EH9-3FL

[2] Centre for Immunity, Infection and Evolution, University of Edinburgh.

Edinburgh EH9 3FL

* Email for correspondence : pedro.vale@ed.ac.uk

Infection avoidance behaviours are the first line of defence against pathogenic

encounters. Behavioural plasticity in response to internal or external cues can

therefore generate potentially significant heterogeneity in infection. We tested

whether *Drosophila melanogaster* exhibits infection avoidance behaviour

during foraging, and whether this behaviour is modified by prior exposure to

Drosophila C Virus (DCV) and by the risk of DCV encounter. We examined two

measures of infection avoidance: (1) the motivation to feed in the presence of

an infection risk and (2) the preference to feed on a clean food source over a

potentially infectious source. While we found no clear evidence for preference

of clean food sources over potentially infectious ones, female flies were less

motivated to feed when presented with a risk of encountering DCV, but only if

they had been previously exposed to the virus. We discuss the relevance of

behavioural plasticity during foraging for host fitness and pathogen spread.

Introduction

Hosts vary considerably in their ability to acquire and transmit infection [1–3], and much of this variation is caused by differences in the contact rate between susceptible individuals and sources of infection [4,5]. For example, viruses of *Drosophila* fruit flies are not only widely distributed, they also show very broad host range [6]. Given the high viral prevalence of pathogens in natural environments, mounting a timely and efficient immune response to all possible pathogenic challenges would be physiologically costly and ultimately ineffective. Hosts capable of reducing the probability of contacting parasites, infected conspecifics or infectious environments can therefore not only prevent the deleterious effects of infection, but also circumvent the undesirable energetic costs of immune responses, including immunopathology [4,7]. Avoiding infection is therefore the first line of non-immunological defence against infection [8], and is known to occur across a broad range of host taxa, including humans [7,9].

Like most traits, infection avoidance behaviours are likely to vary according to the context of infection, and pathogens are major drivers this context [4,7,9–11]. Pathogens may alter host responses in two ways. By altering the immunophysiology of the host during infection, pathogens can alter host behaviour [12,13]. Pathogens also modify the host external environment by increasing the likelihood of exposure to novel infections, and these external cues of infection risk are also known to influence host behavioural responses [4,7]. Understanding variation in infection avoidance behaviours therefore provides an important functional link between the neurological, behavioural and immunological processes that together govern the spread of disease [12].

Insects are ideal systems to investigate the interplay between infection

and behaviour [12,14]. The fruit fly *Drosophila* is especially amenable to these

studies, as it is one of the best developed model systems for host-pathogen

interactions [15] and behavioural ecology and genetics [16,17]. One of the most

studied pathogenic interactions in *Drosophila* is the host response to systemic

and enteric infection with Drosophila C Virus (DCV) [18,19]. DCV is a horizontally

transmitted +ssRNA virus that naturally infects the fly gut [19–21], causing

intestinal obstruction, severe metabolic dysfunction and eventually death [22,23].

As a consequence of its pathology, female flies infected with DCV are also

known to exhibit behavioural modifications, such as reduced locomotion and

increased sleep [24]. The Drosophila-DCV interaction therefore offers a powerful

system to investigate the ecological consequences that may arise from the

physiological and behavioural effects of enteric viral infections.

In the present study we used a combination of controlled experimental

infections and foraging choice assays, to test whether adult *D. melanogaster*

are able to avoid potentially infectious environments when foraging for food,

and if avoidance behaviour is modified in response to virus exposure history

and to different risks of acquiring DCV infection. We find evidence for

avoidance behaviours in the form of reduced motivation to feed according to

the risk of infection. However, these effects were only present in female flies

previously exposed to DCV, indicating potentially important sexual

dimorphism in infection avoidance.

Materials and methods

Fly and virus stocks

All flies used were from a long-term laboratory stock of Wolbachia-

free *Drosophila melanogaster* Oregon R line, maintained on Lewis medium in

standard conditions: 25°C, with a 12:12h light:dark cycle. Fly stocks were

routinely kept on a 14-day cycle with non-overlapping generations under low

larval densities. The DCV culture used in this experiment was grown in

Schneider Drosophila Line 2 (DL2) as described in [24]. Ten-fold serial dilutions

of this culture (diluted in Ringers buffer solution) were aliquoted and frozen

at -80°C for long-term storage before use.

Virus exposure

Flies used in the foraging choice assays were obtained by preparing 10 vials of

Lewis medium and yeast containing ten mated females. Flies were allowed to

lay eggs for 48 hours resulting in age-matched progeny reared in similar larval

densities. To test the effect of previous exposure to virus on avoidance

behaviour during foraging, We exposed these progeny to DCV via the oral

route of infection two to three days after eclosion. Oral DCV infection causes

small but significant reduction in fly survival[19] and in we have found that

orally infected flies experience changes affects fly mortality, fecundity, fecal

shedding (Vale, unpublished data), activity and sleep[24]. Briefly, single-sex

groups of 20 flies were placed in vials containing agar previously sprayed with

DCV ("exposed" to 50 μl of 10^8 viral copies/ml) or the equivalent volume of

Ringers buffer solution as a control ("not exposed"). This procedure produced

10 replicate vials of either healthy or virus-exposed male or female flies. The

viral dose used here was lower than previously reported methods[19], so we

first tested this dose was sufficient to result in viable DCV infections by

measuring changes in virus titres and fly survival in separate experiments

(Fig. 1). Fly survival was monitored on 5 replicate groups of 10 OreR female

flies per vial for 11 days following oral exposure. To measure changes in DCV

titre, twenty-five, 2-3 day-old female flies were individually housed in vials

previously sprayed with DCV as described above for 3 days. Five flies were

collected 1, 3, 6, 9 or 13 days after exposure and total RNA was extracted from

flies homogenised in Tri Reagent (Ambion), reverse-transcribed with M-MLV

reverse transcriptase (Promega) and random hexamer primers, and then

diluted 1:10 with nuclease free water. qRT-PCR was performed on an Applied

Biosystems StepOnePlus system using Fast SYBR Green Master Mix (Applied

Biosystems). We measured the relative fold change in DCV RNA relative to

rp49, an internal *Drosophila* control gene, calculated as $2^{-\Delta\Delta Ct}$ as described in [25].

Foraging choice assays

Following 3 days of virus exposure, we set up independent foraging

choice assays in cages - cylindrical transparent plastic containers (12 cm in

diameter) containing two equally spaced plastic vials of standard Lewis fly

medium supplemented with dry yeast. For each combination of "DCV

exposed" and "not exposed" male or female flies, we set up two sets of cages to

simulate different risks of infection: a "no risk" environment, with two clean

vials (sprayed with sterile Ringers solution), and a "high-risk" environment

where one of the vials was sprayed with DCV, as described above. Six replicate

20-fly groups were allocated to the "high-risk" chambers and four replicates to

the "no risk" chambers, resulting in a total of 40 independent foraging choice

cages. Flies were added to the chamber from a neutrally placed hole in the lid,

and the number of flies that settled on each vial was recorded every 30

minutes for five hours. Care was taken to randomise the position of the cages

so that the orientation of the light did not influence the choice of the flies in

any systematic way.

Statistical Analysis

To measure infection avoidance, we took two approaches. First, we

hypothesised that the motivation to feed would be lower in environments

where the risk of infection is higher [7]. We therefore compared the motivation

to feed between the "no risk" and "high-risk" cages, measured as the

proportion of flies inside each replicate cage that chose to feed on any of the

provided food sources. We also asked whether flies that chose to feed showed

any evidence of avoiding potentially infectious food sources. For this analysis

we focussed on the "high risk" cages and recorded the proportion of flies

choosing the clean food source over the infectious food source in each

replicate cage. In both analyses of 'motivation to feed' and 'infection

avoidance', data on the proportion of flies choosing each food source within

each replicate cage were analysed with a generalised linear model assuming

binomial error and logit link function, and included fly 'sex', 'previous

exposure' and 'infection risk' as fixed effects. 'Replicate cage' was included as a

random effect, nested within treatments. We also analysed the average

motivation to feed and infection avoidance across all time points, in a model

that inlcuded "time" as a random effect. Treatment specific contrasts were

used to test the significance of pairwise comparisons. Analyses were carried

out using JMP 12 [26].

Results and Discussion

The ability to detect and discriminate between clean and potentially

infectious environments is vital to avoid the adverse consequences of

infection. In this study we tested if infection avoidance behaviour in

Drosophila melanogaster is modified by its previous exposure to a viral

pathogen and by the risk of infection with that same pathogen when

encountered during foraging. Viral exposure prior to the behavioural assay

was achieved by placing flies in a DCV contaminated environment for 3 days,

allowing flies to acquire DCV infection orally. DCV acquired through the oral

route using this protocol continued to replicate within the fly, increasing by

10-100 fold by day 13 following oral exposure ($F_{4,19} = 8.78$, p=0.0003; Fig. 1A)

and resulted in up to 20% mortality within this period (Fig. 1B).

In the foraging choice assay, only a fraction of flies chose either of the

food sources provided, and this motivation to feed increased over time for

flies in all treatment groups (χ^2_1= 11.00, p=0.001; Fig. 2A). The rate at which

motivation increased differed between sexes ('Time × Sex' interaction, χ^2_1=

12.47, p=0.0004), and on average female flies showed greater motivation to

feed than males (χ^2_1= 5.01, p=0.025), with 67% of female and 36% of male

flies making a choice to feed on any of the provided substrates during the

observation period. Once flies had made the choice to feed on one of the

provided food sources, the choice between a clean and a potentially infectious

food source was not affected by previous exposure to DCV ('previous

exposure', χ^2_1= 0.513, p=0.47) in either male or female flies ('sex', χ^2_1= 0.595,

p=0.44).

183   Across the entire observation period, the motivation to feed differed

184 between sexes, and depended both on their previous exposure and on their

185 current risk of infection ('Sex' × 'risk of infection' × 'Previous exposure'

186 interaction, χ^2_1= 21.82, p<0.0001). The proportion of males choosing any food

187 substrate did not vary with previous exposure to DCV in either high-risk (χ^2_1=

188 2.21, p=0.137) or no-risk environments (χ^2_1= 0.09, p=0.764; Fig. 1). In female

189 flies however, previous exposure and current infection risk affected the

190 motivation to feed on the provided food sources. When there was no risk of

191 infection (Fig. 2B, light grey bars) the motivation to feed was greater in

192 females that were previously exposed to DCV than in otherwise healthy, non-

193 exposed females (χ^2_1= 104.11, p<0.001). Among females that were previously

194 exposed to infection, we found that the presence of a risk of acquiring

195 infection resulted in lower foraging effort - with just over 50% of flies making

196 the choice to feed - compared to females in cages where there was no risk of

197 acquiring infection, where over 80% of flies made the choice to feed (Fig. 2B;

198 χ^2_1= 168.48, p<0.001).

200   In addition to responding to external cues of infection (infection risk),

201 internal physiological cues (in this case, previous exposure to DCV) may also

202 modify avoidance behaviour. Behavioural modifications due to infection are

203 widely reported among animals [9,27], and can be classified into (i) parasitic

204 manipulation that enhances parasite transmission [9] (ii) sickness behaviours

205 that benefit the host by conserving energetic resources during infection [13], or

206 (iii) side-effects of pathogenicity that do not benefit the host or the parasite [27].

207 Female flies infected orally with DCV are known to experience increased

208 lethargy and sleep [24], so these effects could also explain the reduced feeding

activity we detected in female flies that had been previously exposed to DCV.

Another potential explanation for reduced motivation to feed in previously

exposed flies is infection-induced anorexia [28], a commonly described sickness

behaviour [13]. However, it is unlikely that a lower motivation feed is simply a

symptom of a "sick" fly, because it varied according to the risk of infection, and

even reached 80% in exposed flies when foraging in a 'no risk' environment

(Fig. 2). This suggests that flies are actively avoiding contact with the

potentially contagious food source by lowering their foraging effort.

The higher motivation to feed of some female flies when the risk of

infection was absent (Fig. 2) suggests flies were able to identify external cues

of infection risk. Identifying infection cues is a general prerequisite of

avoidance behaviours and occurs across a wide range of different taxa. For

example, lobsters are known to detect and avoid virus-infected conspecifics [29];

fruit flies and nematodes are capable of avoiding pathogenic bacteria [30,31];

gypsy moth larvae are able to detect and avoid virus-contaminated foliage [14];

sheep have been found to prefer to graze in parasite-poor patches [32]; and it is

has been argued that the disgust response in humans has evolved because it

decreases contact with potential infection [33]. It is unclear how flies are able to

identify food sources contaminated with a viral pathogen. In *Drosophila* and *C.*

elegans avoidance of pathogenic bacteria is enabled by evolutionary

conserved olfactory and chemosensory pathways [30,31], while avoidance of

parasitic wasps appears to be mainly enabled by the visual sensory system [34].

While avoiding virus infected conspecifics is probably driven by visual cues of

infection [29], it remains unclear how virus-contaminated environments may

trigger a lower motivation to feed in *Drosophila*.

The fact that only female flies demonstrated avoidance is an indication

that any potentially adaptive effects of avoiding infection may be related to

oviposition, which coincides with feeding. For flies previously exposed to DCV,

avoiding infection would not confer substantial direct benefits given the

physiological and behavioural costs of this infection [22-24], but would however

reduce the exposure of future offspring to infection. While flies previously

exposed to DCV do not appear to immune primed following an initial viral

exposure [35], our results point to a sort of behavioural priming, where females

previously exposed to infection avoid foraging in potentially infectious

environments.

In summary, using a combination of experimental infections and

behavioural assays, we find evidence for infection avoidance in *Drosophila* in

the form of reduced motivation to feed, which was most pronounced when

flies were faced with an increased risk of encountering an infectious food

source. However, these effects were only present in female flies, indicating

potentially important sexual dimorphism in infection avoidance.

Understanding how avoidance behaviours may vary is therefore important for

our understanding of how disease will spread in natural populations [4], and

more broadly how pathogens might evolve in response to variation in host

infection avoidance strategies [36,37].

Acknowledgements

We thank H. Cowan and H. Borthwick for preparing fly medium, K. Monteith

and S. Lewis for assistance with the qPCR, the members of the Vale, Obbard

and Ross labs for helpful suggestions. This work was supported by a strategic

award from the Wellcome Trust for the Centre for Immunity, Infection and

Evolution (grant reference no. 095831), and by a Society in Science – Branco

Weiss fellowship, both awarded to P. Vale. P.Vale is also supported by a

Chancellor's Fellowship from the School of Biological Sciences, University of

Edinburgh.

Competing interests

The authors declare that they have no competing interests.

Author contributions

PFV conceived the study. PFV and MDJ designed the experiment. MDJ and PFV

carried out the experimental work. PFV analysed the data, wrote the

manuscript and provided all research consumables.

References

1. Fellous, S., Duncan, A. B., Quillery, E., Vale, P. F. & Kaltz, O. Genetic

influence on disease spread following arrival of infected carriers. *Ecol.*

Lett. **15,** 186–192 (2012).

2. Vale, P. F. & Little, T. J. Measuring parasite fitness under genetic and

thermal variation. *Heredity* **103,** 102–109 (2009).

3. Susi, H., Barrès, B., Vale, P. F. & Laine, A.-L. Co-infection alters population

dynamics of infectious disease. *Nat. Commun.* **6,** (2015).

4. Barron, D., Gervasi, S., Pruitt, J. & Martin, L. Behavioral competence: how

host behaviors can interact to influence parasite transmission risk. *Curr.*

Opin. Behav. Sci. **6,** 35–40 (2015).

5. Paull, S. H. *et al.* From superspreaders to disease hotspots: linking

transmission across hosts and space. *Front. Ecol. Environ.* **10,** 75–82

(2012).

6. Webster, C. L. *et al.* The Discovery, Distribution, and Evolution of Viruses

Associated with Drosophila melanogaster. *PLOS Biol* **13,** e1002210

(2015).

7. Curtis, V. A. Infection-avoidance behaviour in humans and other animals.

Trends Immunol. **35,** 457–464 (2014).

8. Parker, B. J., Barribeau, S. M., Laughton, A. M., de Roode, J. C. & Gerardo, N.

M. Non-immunological defense in an evolutionary framework. *Trends Ecol.*

Evol. **26,** 242–248 (2011).

9. Moore, J. An overview of parasite-induced behavioral alterations – and

some lessons from bats. *J. Exp. Biol.* **216,** 11–17 (2013).

10. Wolinska, J. & King, K. C. Environment can alter selection in host-parasite

interactions. *Trends Parasitol.* **25,** 236–244 (2009).

11. Vale, P. F., Salvaudon, L., Kaltz, O. & Fellous, S. The role of the environment

in the evolutionary ecology of host parasite interactions. *Infect. Genet.*

Evol. **8,** 302–305 (2008).

12. Adamo, S. A. Comparative psychoneuroimmunology: evidence from the

insects. *Behav. Cogn. Neurosci. Rev.* **5,** 128–140 (2006).

13. Adelman, J. S. & Martin, L. B. Vertebrate sickness behaviors: Adaptive and

integrated neuroendocrine immune responses. *Integr. Comp. Biol.* **49,**

202–214 (2009).

14. Parker, B. J., Elderd, B. D. & Dwyer, G. Host behaviour and exposure risk in

an insect-pathogen interaction. *J. Anim. Ecol.* **79,** 863–870 (2010).

15. Neyen, C., Bretscher, A. J., Binggeli, O. & Lemaitre, B. Methods to study

Drosophila immunity. *Methods San Diego Calif* **68,** 116–128 (2014).

16. Dubnau, J. *Behavioral Genetics of the Fly (Drosophila Melanogaster)*.

(Cambridge University Press, 2014).

17. Sokolowski, M. B. Drosophila: Genetics meets behaviour. *Nat. Rev. Genet.* **2,**

879–890 (2001).

18. Dostert, C. *et al.* The Jak-STAT signaling pathway is required but not

sufficient for the antiviral response of drosophila. *Nat. Immunol.* **6,** 946–

953 (2005).

19. Ferreira, Á. G. *et al.* The Toll-Dorsal Pathway Is Required for Resistance to

Viral Oral Infection in Drosophila. *PLoS Pathog.* **10,** (2014).

20. Huszar, T. & Imler, J. in *Advances in Virus Research* **Volume 72,** 227–265

(Academic Press, 2008).

21. Kapun, M., Nolte, V., Flatt, T. & Schlötterer, C. Host Range and Specificity of

the Drosophila C Virus. *PLoS ONE* **5,** e12421 (2010).

22. Arnold, P. A., Johnson, K. N. & White, C. R. Physiological and metabolic

consequences of viral infection in Drosophila melanogaster. *J. Exp. Biol.*

216, 3350–3357 (2013).

23. Chtarbanova, S. *et al.* Drosophila C virus systemic infection leads to

intestinal obstruction. *J. Virol.* (2014). doi:10.1128/JVI.02320-14

24. Vale, P. F. & Jardine, M. D. Sex-specific behavioural symptoms of viral gut

infection and Wolbachia in Drosophila melanogaster. *J. Insect Physiol.* **82,**

28–32 (2015).

25. Livak, K. J. & Schmittgen, T. D. Analysis of relative gene expression data

using real-time quantitative PCR and the 2(-Delta Delta C(T)) Method.

Methods San Diego Calif **25,** 402–408 (2001).

26. *JMP*. (SAS Institute Inc.).

27. Barber, I. & Dingemanse, N. J. Parasitism and the evolutionary ecology of

animal personality. *Philos. Trans. R. Soc. Lond. B Biol. Sci.* **365,** 4077–4088

(2010).

28. Ayres, J. S. & Schneider, D. S. The Role of Anorexia in Resistance and

Tolerance to Infections in Drosophila. *PLoS Biol* **7,** e1000150 (2009).

29. Behringer, D. C., Butler, M. J. & Shields, J. D. Ecology: Avoidance of disease

by social lobsters. *Nature* **441,** 421–421 (2006).

30. Babin, A. *et al.* Fruit flies learn to avoid odours associated with virulent

infection. *Biol. Lett.* **10,** 20140048 (2014).

31. Meisel, J. D. & Kim, D. H. Behavioral avoidance of pathogenic bacteria by

Caenorhabditis elegans. *Trends Immunol.* **35,** 465–470 (2014).

32. Hutchings, M. ., Knowler, K. ., McAnulty, R. & McEwan, J. . Genetically

resistant sheep avoid parasites to a greater extent than do susceptible

sheep. *Proc. R. Soc. B Biol. Sci.* **274,** 1839–1844 (2007).

33. Curtis, V., de Barra, M. & Aunger, R. Disgust as an adaptive system for

disease avoidance behaviour. *Philos. Trans. R. Soc. Lond. B. Biol. Sci.* **366,**

389–401 (2011).

34. Kacsoh, B. Z., Lynch, Z. R., Mortimer, N. T. & Schlenke, T. A. Fruit Flies

Medicate Offspring After Seeing Parasites. *Science* **339,** 947–950 (2013).

35. Longdon, B., Cao, C., Martinez, J. & Jiggins, F. M. Previous Exposure to an

RNA Virus Does Not Protect against Subsequent Infection in Drosophila

melanogaster. *PLoS ONE* **8,** e73833 (2013).

36. Boots, M. & Bowers, R. G. Three mechanisms of host resistance to

microparasites-avoidance, recovery and tolerance-show different

evolutionary dynamics. *J. Theor. Biol.* **201,** 13–23 (1999).

37. McLeod, D. V. & Day, T. Pathogen evolution under host avoidance

plasticity. *Proc. Biol. Sci.* **282,** (2015).

Figure legends

369
Fig.1. Exposing flies to DCV by placing them in DCV-contaminated vials for

three days resulted in flies acquiring replicating virus as shown by the

increase in DCV titres over time (Fig. 1A). Grey points shown are individual

replicate female flies, black bars are mean titres. This orally acquired DCV

infection had a moderate effect on fly survival (full black line) compared to

uninfected control flies (dotted line) (Fig. 1B).

Fig. 2. Single sex-groups of flies that had been previously exposed either to

DCV or to a sterile inoculum were tested in a 'no-risk' environment (choice

between two clean vials; light grey) or a 'high-risk' environment (choice

between a clean vial and a DCV-contaminated vial; dark grey). The motivation

to feed, measured as the proportion of flies in the cage that fed on any of the

provided food sources, increased over time (Fig 2A). Fig 2B shows the average

of motivation to feed taken across the whole observation period for each

combination of fly sex, prior DCV exposure and current exposure risk ('no-

risk' environment (light grey) or a 'high-risk' environment (dark grey). Data

show means ± SE.