1
10   Programmed bet hedging in yeast responses to osmotic stress

17   Yoshikazu Hirate[1], Samuel Bottani[2] and Suzannah Rutherford[1]
18
19   1 Fred Hutchinson Cancer Research Center, Seattle, WA, 2 Laboratoire Matière et Systèmes
20   Complex, Paris Diderot, Paris, France

Abstract

Rapid reproduction contributes to evolutionary fitness but can be lethal under stress. Microbes are thought to limit risk in clonal populations by bet hedging; the stochastic expression of a low frequency of slow growing cells constitutively resistant to unpredictable environmental stresses including antibiotics. However fitness depends on rapid recovery and resumption of growth in potentially lethal environments whose severity and duration are also unpredictable.   Here we describe trade-offs between osmotic stress-responsive signaling, survival and proliferation in 50 ecologically distinct strains of budding yeast. By contrast with prior examples, programmed bet hedging responses were heritable, stress-specific and varied continuously in our population. During rapid growth strong osmotic stress signaling promoted survival. Weak signaling predicted lower viability, intense rebound signaling, and robust recovery. Older cultures survived and adapted to unprecedented stress with fitness depending on reproducible, strain-specific proportions of cells with divergent strategies. The most 'cautious' cells survive extreme stress without dividing; the most 'reckless' cells attempt to divide too soon and fail, killing both mother and daughter.   Heritable proportions of cautious and reckless cells generate a tunable, rapidly diversifying template for microbial bet hedging that resembles natural variation and would evolve in different patterns of environmental stress.

In order to understand the evolutionary trajectories of populations and species we need to understand the effects of natural genetic variation on mechanisms of development and expression of phenotypic variation. The mapping between genetic variation and the spectrum of attributes and behaviors upon which selection acts defines population-level properties such as the capacity to evolve (evolvability), the capacity to withstand genetic and environmental perturbation (robustness or canalization), and the optimization, within a given genotype, of phenotypic responses across different environments (reaction norms)[1–3]. Here we describe strain-specific differences in hyperosmotic stress responsive signaling and associated behaviors in a synthetic population of budding yeast.

The high osmolarity glycerol (HOG) signaling pathway is central to an elaborate stress response that reduces cellular damage and death in unpredictably changing osmotic environments where the balance between external solutes and free water pressure in the cell can change suddenly[4]. A main function of the HOG pathway is the production and accumulation of intracellular glycerol, which restores water balance and, as demonstrated by a large body of work from many labs, is essential for survival, adaptation and proliferation in hyperosmotic stress[4–9].   In the wild, yeast and other microorganisms must balance immediate survival against evolutionary fitness. Multiplicative fitness favors clonal populations that respond as rapidly as possible to improved conditions with earlier cell cycle reentry and proliferation[10]. On the other hand, individual survival requires that cells carefully sense the amplitude and direction of environmental change to more safely reenter the cell cycle after stress[7]. The HOG pathway consists of at least two highly-conserved, multi-component osmotic stress sensors linked to a parallel series of at least 15 kinases and accessory proteins that ultimately alter the activity of nearly 10% of the yeast genome[4,6]. The sheer numbers of genes involved in HOG signaling, their conservation, and their elaborate circuitry suggest that a nuanced response to osmotic stress has been crucial and strongly selected throughout evolutionary history. As the hyperosmotic stress response of budding yeast has well-characterized and accessible signaling and phenotypic traits that can be measured in the lab and are almost certainly under strong selection in nature, this system is ideal for characterizing the mapping between signaling and behavior in a diverse population[4,6,7].

We measured osmotic stress signaling, survival and adaptation in both exponentially growing and nearly quiescent cultures of diploid yeast. As glycerol-3-phosphate dehydrogenase (*GPD1*) is rate-limiting for glycerol production[11], the synthesis and accumulation of green fluorescent protein (GFP) integrated into the gene for *GPD1* was a proxy for HOG pathway activity. A synthetic population of diverse yeast genotypes was made by crossing *GPD1*::GFP in the genetic background of a standard laboratory strain (BY4742 Mat alpha) to a panel of wild and industrial genetic backgrounds –e.g. fifty diverse haploids of the opposite mating type extracted from globally diverse, sequence-validated *Saccharomyces cerivisiae* diploid strains deposited to the collection of the Royal Netherlands Academy of Arts and Sciences over the past 100 years (CBS; Table 1 and Table S1).

HOG signaling and behavior in young cultures

The behavior of single cells before and after their exposure to osmotic stress was followed by time-lapse video microscopy of monolayer cultures in custom microfluidics devices[12]. When cells in exponential growth were exposed to sudden hyperosmotic stress, cell volume decreased, cell division and budding immediately stopped, and daughter cells retracted[13].   After a lag period proportional to the severity of the stress GFP fluorescence driven by the *GPD1* promoter began to accumulate in the cytoplasm of surviving cells.   Cells that did not accumulate *GPD1*::GFP to high levels did not survive or adapt, developed large vacuoles, and began to die, remaining in view as shrunken cell ghosts. As GFP accumulated to saturation levels in the surviving cells, they adapted to the higher osmotic pressure, resumed cell division, budded and began to divide with a longer half time, producing daughter cells with similarly high fluorescence[13].

Viability per culture and *GPD1*::GFP accumulation per cell were measured using flow cytometry of statistically large numbers of cells from all 50 strains (~10,000 cells / sample; Figure S1). The rate and extent of mean *GPD1*::GFP accumulation in exponentially growing cultures exposed to hyperosmotic media depended on the severity of the stress and the genetic background of each strain (Figure 1A).   Prior to the osmotic stress mean *GPD1*::GFP fluorescence and viability were uncorrelated.   After 2 hours in moderate 0.75 M KCl viability decreased and became steeply correlated with accumulated *GPD1*::GFP (Figure1B and C).   As expected, natural variation in the strength of HOG signaling was directly responsible for variation among the strains in osmotic stress survival.

Negative feedback drives a robust recovery

The initially strong positive correlation between variation in *GPD1*::GFP accumulation and variation in viability reversed as cells adapted and began to divide (Figure 1C; 4 hours). This distinguished two phases of the response, an early phase (0 – 2 hours) when viability decreased markedly and acute HOG signaling promoted osmotic stress survival and a later phase (2 – 4 hours) when viability recovered but became negatively correlated with HOG signaling and *GPD1*::GFP accumulation. The switch from positive to negative correlations might have indicated that stronger HOG signaling, initially beneficial, suddenly caused lower viability.   However we think it likely that negative feedback increased signaling in the surviving cells of the less viable strains. Negative feedback controls, occurring at many levels and timescales, are present in essentially all of the varied mechanisms that act in concert to increase intracellular glycerol and restore water balance. For example (1) unequal water pressures activate osmotic stress sensors, glycerol channels and other pressure-sensitive components whose activities control and depend on water balance (e.g. see Figure 5 in Hohman 2002[4,6]), (2) GPD1p indirectly controls and is controlled by osmotic stress-sensitive kinases that respond to upward and downward changes in water balance[14], and (3) nuclear Hog-1 MAP kinase increases the transcription of phosphatases that restore its own cytoplasmic localization and basal activity[15–17].

Consistent with acting negative regulation, we found a strong and highly significant
correlation between early mortality (0 – 2 hour decreases in viability) and later accumulations of
GPD1::GFP (2 – 4 hours; Table 2).   We reasoned that cells and strains that adapt quickly
experience lower and less sustained effects of osmotic stress (e.g. water loss) with more rapidly
attenuated HOG pathway activity and lower *GPD1*::GFP accumulation.   Conversely, surviving
cells of strains that were slower to adapt and less viable would experience higher and more
sustained osmotic stress (and likely other stresses). Prolonged osmotic stress would sustain
HOG signaling and maintain *GPD1* transcription – also activated by general stress responses[18] –
further promoting *GPD1*::GFP accumulation (e.g. negative feedback regulation of viability at the
level of general stress responses). Indeed, even as *GPD1*::GFP and viability became negatively
correlated, their *rates of change* remained positively correlated (2 – 4 hours; Table 2) prompting
a parsimonious interpretation that osmotic stress signaling promotes adaptation and viability
during both the initial and recovery phases of the response (Figure 1C, insets).
After 4 hours all strains had adapted to a new steady state in 0.75M KCl and later
viability remained largely unchanged (Figure 1C inset, lower right). Interestingly, initial decreases
in steady-state viability (0 – 2 hour mortality) were almost perfectly restored by 4 hours (Figure
1D).   Notably, by 6 hours early mortality and recovery were 98% correlated ($R^2 = 0.9852$,
P<0.0001; not shown).   The biological robustness of adaptation and complete recovery of
steady state viability further support the idea that negative feedback restores viability through
continued activation of stress responses. Indeed, continued accumulation of *GPD1* and glycerol
– directly responsible for restoration of water balance and reduction of osmotic stress –suggests
that the accumulation of glycerol integrates the cumulative activities of many facets of the
osmotic stress response (e.g. a plausible biological mechanism for "integral feedback"[19]). The
presence of integral feedback virtually assures perfect adaptation[17,19]. However, despite their
resilience, strains that were relatively slower to adapt would be ultimately less fit than rapidly
adapting strains due to their higher death rate, slower recovery, and lower steady-state viabilities
before and after hyperosmotic shifts.
142

Extreme stress resistance of older cultures
By contrast with exponential cultures, when the aging yeast cultures (post-diauxic) were exposed
to hyperosmotic media they survived and adapted after long periods in unprecedented conditions
(Movies 1 and 2). As aging cultures deplete available glucose in their media they undergo a
metabolic change called the diauxic shift[20]. During post-diauxic growth stress response proteins
accumulate, cell division slows and then stops, and cells enter quiescence[21]. Post-diauxic
cultures survived up to 5 weeks in 3 M KCl (41/50 strains). They could not adapt and did not
grow in 3 M KCl, but recovered rapidly and grew when plated on fresh isotonic media ('static
viability'; Figure S2). When we tested their limits of adaptation in increasing concentrations of KCl
all but one strain could grow on 2.6 M KCl media and, remarkably, three strains grew on media

containing 2.9 M KCl (Table 3). We are unaware of previous reports of such extreme osmotic

stress survival or adaption limits for budding yeast of any growth stage or genotype.

Heterogeneity of older cultures

By contrast with cultures in exponential growth, in post diauxic growth the genetically identical

cells within each strain and culture were surprisingly heterogeneous in size, shape and signaling

behaviors (compare Figures S1 and S3).   Neither total $GPD1$::GFP fluorescence nor rates of

change in fluorescence was strongly correlated with viability.   After several hours in 2.5 M KCl

$GPD1$::GFP increased sharply in one group of cells as they began to divide. More surprising,

other cells also induced $GPD1$::GFP to high levels, started to divide and then popped, killing both

the mother and daughter (Movie 3). Another group of cells had slower signaling and cell division

while the most 'cautious' cells failed to signal or divide but remained in a cellular state of static

viability without dividing.

166   We used a Gaussian mixture model to assign the cells in each sample to four

Gaussian distributions described by eight parameters – means and covariances – numbered

according to increasing levels of fluorescence (G_0-G_3 distributions; Figure S3).   Only

$GPD1$::GFP pre-accumulated into cells of the G_3 distribution during post-diauxic growth *prior* to

the osmotic challenge and therefore unrelated to osmotic stress signaling, predicted survival at

any time.   The expression of a low frequency of cells in each strain expressing stress resistance

in rich media and normal growth conditions could be analogous to previously-defined stochastic

bet hedging in yeast and bacteria[22,23].   The amount of $GPD1$::GFP accumulated in G_{3_0} cells

predicted early but not later viability, and this relationship was better fit by 2^{nd} order quadratic

rather than linear functions of $GPD1$::GFP, suggesting early survival was higher in strains with

intermediate G_3 level signaling (more variation explained and lower mean square errors; Table

S3).   Despite the fine-scaled characterization of osmotic stress signaling behaviors of different

groups of cells in each strain, none of the distributions learned by the Gaussian mixture model –

neither pre-accumulated G_{3_0}, total $GPD1$::GFP fluorescence, nor stress-induced $GPD1$::GFP in

any distribution – embodied features of the osmotic stress signaling that were important for later

survival.

Continuous variation in signaling behavior

In order to map osmotic stress-responsive signaling onto survival more directly we next

quantified osmotic stress signaling behaviors of cells in each strain directly, as they unfolded

over time, unbiased by Gaussian assumptions or approximations. First, normalized levels of

induced $GPD1$::GFP were binned (as in a histogram), creating a 'sample vector' of cell numbers

in each of 100 successive intervals of fluorescence intensity. Next, sample vectors for increasing

time points in each strain were linked to create time line vectors (700-mers). For comparison of

averaged strain behaviors the time line vectors were clustered using a mean distance matrix,

constructed from 17,000 randomly chosen permutations of the data over replicate samples and

times (mean clusters; see methods).   Finally, the fraction of permutations in which each strain
grouped with more than half of the other strains in its mean cluster was used to rank that strain's
behavior relative to the other strains in its group (clustering statistics; Table 4).
The unique signaling trajectories of most strains were highly reproducible (Figure S3).
Based on their shared and strain-specific (heritable) signaling behaviors the 50 strains rapidly
converged onto two large groups made up of six mean clusters (Figure 2). Each strain could be
further ordered within and between mean clusters and clades based on their clustering statistics,
with their rank order describing increasingly rapid accumulations of *GPD1*::GFP and 'reckless'
signaling (Figure S4). Tellingly, both mean cluster and rank now predicted viability over time
(Figure S5 and Table 4; respectively) thereby confirming the biological relevance of 'cautious'
versus 'reckless' osmotic stress signaling, validating the clustering method and supporting the
role of natural osmotic stress signaling differences between strains in shaping variation in fitness
during osmotic stress.
205
Evidence for bet hedging
As cautious and reckless behaviors were found both within and between strains, we wondered
whether bet hedging, the expression of alternate, conditionally-adaptive phenotypes within a
clone of genetically identical organisms[10,24–27], could explain the observed variation in osmotic
stress signaling and survival.   Indeed, rank predicted survival in several different osmotic
environments. Higher-ranked strains with more aggressive osmotic stress signaling strategies
were favored in milder conditions, but with increasing time in extreme osmotic stress more
cautious strains and behaviors became more fit (Table 4). For example, W178 at rank 50 was
most viable in moderate 0.75 M KCl, but the optimum rank shifted to 25 after 20 hours in 2.5 M
KCl, 20 after 72 hours, and 9.6 after 168 hours (1 week).   Viability after 168 hours decreased
most markedly among the most reckless strains (Figure S5).
Demonstration of differential relative fitness of strains with more cautious behavior
under increasing osmotic stress constitutes strong empirical evidence for bet hedging[25]. To
confirm that the relative fitness of 'cautiousness' was really correlated to the amount of stress, we
tested the idea that the yeast experienced increasing stress over time in severe osmotic
conditions. We again incubated cultures in 2.5 M KCl for 168 hours, but they were first exposed
to a mild pre-stress (2 hours in 0.5 M KCl) to pre-induce osmotic stress proteins and make them
more resistant to subsequent stress. If the optimum rank depended solely on time independent of
the degree of stress experienced by different strains, then it should be unaffected by the short
pre-stress. However optimum rank was shifted toward more reckless behaviors (rank 9.6 to rank
18; P<0.0001) after the pre-stress and viability increased by ~10% as expected if the pre-treated
cells experienced lower osmotic stress.   After 168 hours in 2.5 M KCl the most reckless cells in
the highest-ranking strains began to selectively disappear.   For example, replicate cultures of
strain W242 (rank 49) that had lower viability also had fewer cells with high accumulations of
GPD1::GFP, smaller G3 distributions, and correspondingly larger distributions with lower mean

GPD1::GFP (Figure 3A).   The loss of cells with the highest accumulations of *GPD1*::GFP could

indicate that *GPD1*::GFP levels simply decrease over time.   However, G3 distributions were

stable over most time points and in most strains.   More likely, after 168 hours in 2.5 M KCl the

most aggressive cells in the highest-ranking strains attempt to divide and fail (e.g. Movie 3). This

suggests that rapid signaling and adaptation, a fitness advantage in milder conditions, becomes

a liability in severe or prolonged osmotic stress. On the other hand static viability – the survival of

non-dividing cultures in 3 M KCl (Figure 3B) – would usually have dramatically reduced

evolutionary fitness, but allows more cautious cells and strains to survive severe stress.

Evolution of bet hedging

Heritable osmotic stress signaling behavior predicted survival in increasing severity and duration

of osmotic stress. Since cautious and reckless strains reliably expressed a range of cells with

different behaviors and fitness depending on the environment we wondered if a simple, 2-state

bet hedging model including a heritable control of the relative proportions of cautious and

reckless cell types could account for the observed variation in osmotic stress signaling and

explain the complex relationship between rank and viability. If more aggressive osmotic stress

signaling with rapid recovery and resumption of growth in potentially lethal environments were

the default, ancestral behaviors, then cautious signaling and behavior could have arisen as a

later, adaptive response to the unpredictable severity and duration of potentially lethal osmotic

environments. We therefore assumed that the heritable probability of daughters with cautious

signaling and behavior (P) was the derived, bet-hedging trait and asked whether it could evolve.

252   We considered changes between three very general osmotic stress environments that

discriminate cautious versus reckless behaviors: (E0) a permissive environment in which both

cautious and reckless cells grow equally well, (E1) a restrictive environment approximating

moderate osmotic stress reckless cells divide and cautious cells survive without dividing, and

(E2) a killing osmotic stress where reckless cells die and cautious cells survive without cell

division.   After several generations including each of 9 possible environmental shifts between

the three environments, we calculated the relative fitness (cell numbers) of each strategy

(probability (P) of cautious cells where $0 \leq P \leq 1$). Most environmental shifts favor optimum

strategies of either all cautious (P = 1) or all reckless cell types (P = 0; Figure 3D). Strictly

intermediate strategies ($0 < P < 1$) and bet hedging prevail only when the osmotic environment

shifted from moderate to more severe, with the optimum P depending on the number of

generations in the first environment (E1 -> E2). Shorter lag periods – corresponding to less

severe osmotic conditions – and more cell divisions in E1 initially favor lower P and a higher

proportion of reckless cells.   Longer lag periods – corresponding to more severe conditions and

fewer cell divisions – favor higher P and a higher proportion of cautious cells.   Indeed,

worsening osmotic environments are common in nature (for example, during fermentation or

drying). A worsening environment was experienced with increasing time by the cells and strains

in our experiments; lower-ranked strains with more cautious signaling behaviors, longer lag

periods and fewer attempted cell divisions were increasingly fit over time and with increasing

severity of osmotic stress (Table 4).

272   While the 2-state model of bet hedging behaviors may be overly simplistic relative to up

to four learned distributions of cell types that best fit the data (Figure S3), it provides a conceptual

framework for understanding the observed variation in osmotic stress signaling strategies and

generates testable hypotheses for further studies.   Bet hedging in microorganisms has been

previously thought to arise almost exclusively through stochastic switching with a low probability

of alternate phenotypes that are independent of the environmental challenge[10,22,23,28].   By

contrast, the variation in osmotic stress signaling and behavior we observe is heritable, occurs in

a programmed response of cells to the osmotic stress and, since strains have a low to high

frequency of cells displaying cautious behavior, is one of few known examples in a

microorganism similar to the rapidly diversifying, within-clutch bet hedging strategies of plants

and animals[10,28].   If the heritable probabilities of cautious cells and behavior we observed were

sculpted by selection, we reasoned that milder and/or slowly worsening osmotic environments

would generate more reckless strains (e.g. W242, rank 49 isolated from rotting banana; W282,

rank 48 isolated from fermenting fruit) and more severe or rapidly deteriorating osmotic

environments would generate more cautious strains and behavior (e.g. W455, rank 1 from

molasses; W217, rank 4 from sugar cane syrup).   Indeed, even though our synthetic population

is represented by lab/wild heterozygotes rather than inbred strains, we found that the general

environment from which each strain was isolated weakly predicted its aggressiveness (but not

rank; mean percent of cells above threshold, $P = 0.030$ by ANOVA), plausibly suggesting

adaptation to growth in rotting fruit or wine fermentation favors more aggressive osmotic stress

behavior than growth in more hyperosmotic sugar cane syrups, molasses or olive wastes.

293   Classical evolutionary models assign fitness directly to genotypes, mutations, and

mean trait values without consideration of the genotype-to-phenotype map; molecular models

provide detailed mechanistic understanding of development but rarely consider the effects of

natural genetic variation.   The osmotic stress response is extensively characterized in a few

strains and genotypes but until now osmotic stress signaling and behavior in a population had

not been documented. Labyrinthine developmental mechanisms – themselves controlled by

genetic variation – translate genotypes into phenotypes with a variable fidelity that allows for the

possibility of phenotypic heterogeneity and the evolution of bi-stable states[29].   Our view of

osmotic stress signaling and response on a backdrop of natural variation enabled identification of

negative feedback controlling a robust recovery of steady-state viability in exponential growth.

In post diauxic cultures continuous, heritable variation in the distribution of cautious to reckless

osmotic stress signaling is a risk spreading strategy.   Microorganisms in nature including yeast

spend a high fraction of their time in post-diauxic or quiescent phases[21]. Our simple, 2-state

model demonstrates how cells and strains can balance constraints between survival and

evolutionary fitness through programmed bet hedging responses that are adapted to different

patterns of environmental stress.

Materials and methods

Strain acquisition and deposition

Wild and industrial diploid strains of *Saccharomyces cerevisiae* were obtained from the fungal

diversity collection of Centraalbureau voor Schimmelcultures (CBS), an institute of the Royal

Netherlands Academy of Arts and Sciences in Utrecht, Netherlands

(http://www.cbs.knaw.nl/index.php/collection).   Strains modified for this report are listed in

Tables 1 and S1.   They have been deposited to the Yeast Genetic Resources Lab of the

National BioResource Project in Osaka, Japan (http://yeast.lab.nig.ac.jp/nig/index_en.html).

Creation of haploid *MATa* library of wild and industrial genotypes

For unique identification of each strain and prevention of homothalism, the *HO* locus of each wild

or industrial diploid strain was deleted and replaced with the KanMX4 marker gene and unique

barcode sequences[30,31]. The KanMX4 gene was PCR-amplified for transformation of each strain

with primers containing unique sets of barcode sequences[32] (Table S1). Kanamycin-resistant

transformants were sporulated in 1% potassium acetate sporulation medium and *MATa* haploids

were selected by testing for schmoo formation in 96-well plates containing alpha factor. Mating

type was confirmed by test mating with a dominantly marked *MATa* tester strain on replica plates.

The *URA3* gene of the sporulated wild haploids (*MATa*) was deleted using standard gene

deletion method.   In brief, pJL164 plasmid DNA was transformed and cells were plated onto

YPD plate grown for overnight at 30°C.   Cells were replica plated on 5-FOA and grown for 2

days at 30°C. The *ho* and *ura3* deletions and barcode sequences were verified by PCR and

resulting wild haploids were in many cases validated by their colony morphologies.   Forty-nine

wild strains and a laboratory strain met these criteria (see Tables 1 and S1 for strain details).

Construction of synthetic population of *GPD1*::GFP wild/lab diploids

The common laboratory strain BY stably integrated *GFP* reporter strain was created by replacing

the *GPD1* gene with *GFP* gene using a deletion cassette containing a *URA3* marker for selection

of transformants on SC-URA plates[32,33].   Wild *MATa* haploids were mated with *MATalpha* GFP

reporter strains.   Briefly, *MATa* and *MATalpha* cells were mixed on SC-URA plates for 2 hours

and streaked onto selective SC-URA+G418 plates. To test the BY4742 background for low

function alleles that could critically compromise osmotic stress function, a homozygous

laboratory strain was made by crossing BY4742a to cells of the opposite mating type to create a

50th strain in our population (BY4741alpha; Table 2).

Ploidy analysis

Polyploids had higher GFP and tolerance to osmotic stress.   To screen for triploids or

tetraploids obtained after mating, wild/lab yeast were inoculated into 2 ml YPD with appropriate

antibiotics and cultured for 24 h at 21°C on rotator after which 40 ul of the cultured cells were

then added to fresh 2 ml YPD and cultured for 4 h at 21°C on rotator.   800 ul of cells were fixed

with 1.7 ml cold ethanol and stored at -20^{o}C at least 20 minutes, centrifuged at 2300 rpm for 5

min, washed with 100 mM Tris-HCl (pH 7.8) twice, and suspended in 500 ul of 100 mM Tris-HCl

(pH 7.8).   To digest ribonucleic acids 10 ul of 10 mg/ml RNase A was added to cell suspension

and they were incubated for at least 60 min at 37^{o}C.   Digested cells were then centrifuged at

2300 rpm for 5 min and suspended in 500 ul of FACS buffer (0.5 M Tris-HCl, pH 7.5, MgCl,

NaCl.).   To dissociate cell clumps cells, they were sonicated for 10 sec at setting 3 (Sonifier Cell

Disrupter, Model W185).   Cells were stained with 30 ug/ml Propidium Iodide (PI).   After leaving

them for 20 min in the dark, cells were examined for their ploidy with FACS Calibur (Becton

Dickinson).

Plating assays

Cells cultured in YPD with G418 for 4 d at 21 ^{o}C on rotator were diluted to OD$_{600}$ 0.1 with

exhausted YPD and sonicated for 5 sec at setting 2.5.   5 ul of cells were plated on YPD with or

without KCl and incubated at 21 ^{o}C for 2 months. To determine adaptation limits at which strains

could resume growth in concentrations up to 3 M KCl, we inoculated PD cells in a 96-well plate,

agar media containing increasing concentrations of KCl ranging from 2 to 3 M.   After 2-months

incubation at 21^{o}C, we documented adaptation limit, the highest concentration of KCl in which

colonies could grow.

Microfluidics

We employed custom made microfluidics devices with two fluid inputs as described previously[12].

Experiments were run at ambient room temperature.   Yeast cells were observed using Nikon

TS100 and recorded using a digital camera (Photometrics CoolSnap HQ2) operated by Metavue

(Molecular Dynamics).   Analysis of acquired images was performed using Image J software.

Flow cytometry

After osmotic stress challenge 4 ml of PBS was added to each tube.   Cells were isolated by

centrifugation and resuspended in 1 ml PBS, transferred to FACS tubes, sonicated, and stained

with 3 ug/ml propidium iodide (PI) to monitor viability.   After 20 min GFP fluorescence and

viability were quantified in a FACS Calibur flow cytometer (Becton Dickinson).   Data were

initially analyzed using magnetic windows in FlowJo software to eliminate cell fragments and

dead (PI-positive) cells.

Flow cytometry data analysis

Raw cell data were initially processed and trimmed to identify cells and the fraction of viable cells

using magnetic windows in FloJO data analysis software.   Cell data for each sample including

forward scatter and GFP fluorescence data were then extracted into an executable SQL

database for analysis of single cell data.   We performed multi-normal fits to extract quantitative

values of GPD1::GFP for the different cellular conditions of a strain at a given time point. The

exact number of distributions to be fit is a parameter that is not determined by the model and was

determined from the data. We found that fitting both *GPD1*::GFP and forward scatter worked

slightly better than *GPD1*::GFP in fitting the data, and that 4 distributions provided more stable

fits than 3 distributions.< Figure S3- fit distributions G0-G4). Multi-normal fits were performed with

the sklearn.mixture.GMM Gaussian Mixture Model algorithm of Python scikit package

(http://scikit-learn.org/stable/index.html). The exact number of distributions to be fit is a

parameter that is not determined by the model, not determined from the data.   We found that

fitting both *GPD1*::GFP and forward scatter worked slightly better than *GPD1*::GFP in fitting the

data, and that 4 distributions was better (smaller ??) than 3 distributions.< Figure S3- fit

distributions G0-G4).

Clustering

We used a clustering algorithm of *GDP1*::GFP accumulations to characterize the signaling

behaviors of the strains over time. In brief, the *GPD1*::GFP distribution of each strain at every

time point was converted into a vector of cell density (numbers of cells) in a small section (100

sections) of a normalized density plot of cells at each time point.   Two-dimensional clustering on

the Kullbak-Leibler distance function was used to follow the behavior of each strain over time.

Strains and timepoints were replicated between 4 and 15 times. To control for variation in

sampling and clustering outcomes samples were randomly drawn for each strain and time point

with equal probability for 17000 permutations.

407   Strains were ranked from most static signaling to most aggressive, providing a

continuous relationship between signaling, survival and adaptation with the number of

permutations in which strains clustered with a majority of other strains in their clade was used to

rank strains from the most static to the most aggressive.   Computational sorting of time-series

distributions resolved 6 clades differentiated for rates of GFP accumulation, adaptation and

survival.   We used a clustering algorithm of *GDP1*::GFP and forward scatter accumulations to

characterize the signaling behaviors of the strains over time. We found that clustering based on

both the combined *GPD1*::GFP and forward scatter   data worked slightly better than only

GPD1::GFP. In brief, for an experiment we consider for each cell the   *GPD1*::GFP and forward

scatter data. This 2D distribution of each strain at every time point was converted first into a 2D

100x100 grid with binned values, then into a normalized linear 1000 elements state vector

characteristic for the *GPD1*::GFP and forward scatter data of cells at each time point. The

vectors for all the *n* time points were then successively appended to form a nx1000 time line

vector representing the time course of the combined GPD1::GFP and forward scatter data. The

vectors were then used to compute a distance matrix between strains at each time point using a

Kullback-Leibler divergence based distance function. We used hierarchical clustering with

Ward"s method using the fastcluster implementation in Python[34](http://www.jstatsoft.org/v53/i09/).

We found stronger cluster when performing the same analysis using the combined GPD1::GFP

and forward scattering for each cell in the flow cytometry, and unless stated otherwise the

reported results are in this case. Data from experimental replicates were representatively

sampled for each time point to generate a random average sampling of time line vectors for each

strain.

Bet hedging model

The bet hedging 'strategy' P was defined as the heritable probability of cautious cells for $0 \leq P \leq$

1. Relative fitness was measured for representative strategies (0, 0.1,0.2,…1.0) after 10

generations in each environment. Nine possible 2-state environmental shifts between three

general osmotic stress environments were considered: permissive (E0; all cells grow equally

well), restrictive (E1; reckless cells divide, cautious cells survive without dividing), and killing (E2;

reckless cells die, cautious cells survive without cell division). For simplicity, the natural attrition

of older cells and rates of cell division were assumed to be equal for all cell types.

Results were independent of the number of generations in the first environment except as shown

when E1 was the first environment.

References

1. Kirschner, M. & Gerhart, J. Evolvability. *Proc. Natl. Acad. Sci. U. S. A.* **95,** 8420–8427 (1998).
2. West-Eberhard, M. J. *Developmental plasticity and evolution. Nature* **424,** (2003).
3. Rutherford, S. L. From genotype to phenotype: buffering mechanisms and the storage of genetic information. *Bioessays* **22,** 1095–105 (2000).
4. Hohmann, S. Osmotic Stress Signaling and Osmoadaptation in Yeasts. *Microbiol. Mol. Biol. Rev.* **66,** 300–372 (2002).
5. Hohmann, S., Krantz, M. & Nordlander, B. *Yeast Osmoregulation. Methods Enzymol.* **428,** (Elsevier Masson SAS, 2007).
6. Saito, H. & Posas, F. Response to hyperosmotic stress. *Genetics* **192,** 289–318 (2012).
7. Clotet, J. & Posas, F. *Control of Cell Cycle in Response to Osmostress: Lessons from Yeast. Methods Enzymol.* **428,** (Elsevier Masson SAS, 2007).
8. Nadal, E. de, Alepuz, P. & Posas, F. Dealing with osmostress through MAP kinase activation. *EMBO Rep.* **3,** 735–40 (2002).
9. Babazadeh, R., Furukawa, T., Hohmann, S. & Furukawa, K. Rewiring yeast osmostress signalling through the MAPK network reveals essential and non-essential roles of Hog1 in osmoadaptation. *Sci. Rep.* **4,** 4697 (2014).
10. Ratcliff, W. C., Hawthorne, P. & Libby, E. Courting disaster: How diversification rate affects fitness under risk. *Evolution* 1–10 (2014). doi:10.1111/evo.12568
11. Remize, F., Barnavon, L. & Dequin, S. Glycerol export and glycerol-3-phosphate dehydrogenase, but not glycerol phosphatase, are rate limiting for glycerol production in Saccharomyces cerevisiae. *Metab. Eng.* **3,** 301–312 (2001).
12. Cookson, S., Ostroff, N., Pang, W. L., Volfson, D. & Hasty, J. Monitoring dynamics of single-cell gene expression over multiple cell cycles. *Mol. Syst. Biol.* **1,** 2005.0024 (2005).
13. Miermont, A. *et al.* Severe osmotic compression triggers a slowdown of intracellular signaling, which can be explained by molecular crowding. *Proc. Natl. Acad. Sci. U. S. A.* **110,** 5725–30 (2013).
14. Lee, Y. J., Jeschke, G. R., Roelants, F. M., Thorner, J. & Turk, B. E. Reciprocal Phosphorylation of Yeast Glycerol-3-Phosphate Dehydrogenases in Adaptation to Distinct Types of Stress. *Mol. Cell. Biol.* **32,** 4705–4717 (2012).
15. Jacoby, T. *et al.* Two Protein-tyrosine Phosphatases Inactivate the Osmotic Stress Response Pathway in Yeast by Targeting the Mitogen-activated Protein Kinase, Hog1. *J. Biol. Chem.* **272,** 17749–17755 (1997).
16. Wurgler-Murphy, S. M., Maeda, T., Witten, E. a & Saito, H. Regulation of the Saccharomyces cerevisiae HOG1 mitogen-activated protein kinase by the PTP2 and PTP3 protein tyrosine phosphatases. *Mol. Cell. Biol.* **17,** 1289–1297 (1997).
17. Muzzey, D., Gómez-Uribe, C. a., Mettetal, J. T. & van Oudenaarden, A. A Systems-Level Analysis of Perfect Adaptation in Yeast Osmoregulation. *Cell* **138,** 160–171 (2009).
18. Boy-Marcotte, E., Perrot, M., Bussereau, F., Boucherie, H. & Jacquet, M. Msn2p and Msn4p control a large number of genes induced at the diauxic transition which are repressed by cyclic AMP in Saccharomyces cerevisiae. *J. Bacteriol.* **180,** 1044–52 (1998).
19. Yi, T. M., Huang, Y., Simon, M. I. & Doyle, J. Robust perfect adaptation in bacterial chemotaxis through integral feedback control. *Proc. Natl. Acad. Sci. U. S. A.* **97,** 4649–4653 (2000).
20. Galdieri, L., Mehrotra, S., Yu, S. & Vancura, A. Transcriptional Regulation in Yeast during Diauxic Shift and Stationary Phase. *Omi. A J. Integr. Biol.* **14,** 629–638 (2010).
21. Gray, J. V *et al.* 'Sleeping beauty': quiescence in Saccharomyces cerevisiae. *Microbiol. Mol. Biol. Rev.* **68,** 187–206 (2004).
22. Levy, S. F., Ziv, N. & Siegal, M. L. Bet hedging in yeast by heterogeneous, age-correlated

492   expression of a stress protectant. *PLoS Biol.* **10,** (2012).

493   23.   De Jong, I. G., Haccou, P. & Kuipers, O. P. Bet hedging or not? A guide to proper
494   classification of microbial survival strategies. *BioEssays* **33,** 215–223 (2011).

495   24.   Philippi, T. & Seger, J. Hedging one's evolutionary bets, revisited. *Trends Ecol. Evol.*
496   *(Personal Ed.* **4,** 41–44 (1989).

497   25.   Simons, A. M. Modes of response to environmental change and the elusive empirical
498   evidence for bet hedging. *Proc. Biol. Sci.* **278,** 1601–1609 (2011).

499   26.   King, O. D. & Masel, J. The evolution of bet-hedging adaptations to rare scenarios. *Theor.*
500   *Popul. Biol.* **72,** 560–575 (2007).

501   27.   Meyers, L. A. & Bull, J. J. Fighting change with change: Adaptive variation in an uncertain
502   world. *Trends Ecol. Evol.* **17,** 551–557 (2002).

503   28.   Ratcliff, W. C. & Denison, R. F. Individual-Level Bet Hedging in the Bacterium
504   Sinorhizobium meliloti. *Curr. Biol.* **20,** 1740–1744 (2010).

505   29.   Rutherford, S. L. Between genotype and phenotype: protein chaperones and evolvability.
506   *Nat. Rev. Genet.* **4,** 263–74 (2003).

507   30.   Giaever, G. *et al.* Functional profiling of the Saccharomyces cerevisiae genome. *Nature*
508   **418,** 387–391 (2002).

509   31.   Shoemaker, D. D., Lashkari, D. a, Morris, D., Mittmann, M. & Davis, R. W. Quantitative
510   phenotypic analysis of yeast deletion mutants using a highly parallel molecular bar-coding
511   strategy. *Nat. Genet.* **14,** 450–456 (1996).

512   32.   Wach, A., Brachat, A., Pöhlmann, R. & Philippsen, P. New heterologous modules for
513   classical or PCR-based gene disruptions in Saccharomyces cerevisiae. *Yeast* **10,** 1793–
514   1808 (1994).

515   33.   Gietz, R. D. & Woods, R. a. Transformation of yeast by lithium acetate/single-stranded
516   carrier DNA/polyethylene glycol method. *Methods Enzymol.* **350,** 87–96 (2002).

517   34.   Müllner, D. fastcluster : Fast Hierarchical , Agglomerative Clustering Routines for R and
518   Python. *J. Stat. Softw.* **53,** 1–18 (2013).

519   35.   Mortimer, R. K. & Johnston, J. R. Genealogy of principal strains of the yeast genetic stock
520   center. *Genetics* **113,** 35–43 (1986).

521   36.   Brachmann, C. B. *et al.* Designer deletion strains derived from Saccharomyces cerevisiae
522   S288C: A useful set of strains and plasmids for PCR-mediated gene disruption and other
523   applications. *Yeast* **14,** 115–132 (1998).

Supplementary information

Python script for bet hedging model

Flow cytometry database

3 movies

Acknowledgements

The wild strain collection was made in the Rutherford lab thanks to the help of numerous teams

of summer undergraduates. We thank Wyming Lee Pang and Jeff Hasty for help with

microfluidics and Stéphane Douady for advice on scaling single cell data. This research was

supported by a Scholar Award from the Damon Runyon Cancer Research Foundation (SR), NIH

R01 GM068873 (SR) and the Lady Tata Foundation (YH).

Author information

The current address for YH is Center for Experimental Animals, Tokyo Medical and Dental

University, 1-5-45 Yushima, Bunkyo-ku, Tokyo 113-8510, Japan.   None of the authors have

competing interests.

Tables

Table 1. Strains and aliases used in this study. See Table S1 for details about each of the 49

wild haploid strain derivatives (WHXXX). For brevity, figures are labeled with the wild parent

strain number (WXXX; see Table S1 for details).

Strain	Genotype	Source
WHXXX	*MATa ura3D0 hoD::barcode::KanMX4*	This study; haploid Mata isolates of wild strains. See Table S1 for details.
BY41	*MATa his3D1 leu2D0 met15D0 ura3D0 flo8-1 hoD::barcode::KanMX4*	Mata derivative of BY4742.
BY4742	*MATalpha his3D1 leu2D0 lys2D0 ura3D0 flo8-1*	This study; BY4741 and BY4742 backgrounds derive from a wild diploid isolated in Nerced, California in 1938 on figs (EM93[35]; S228C[36]). They are distinguished primarily by the many generations it has been under laboratory selection.
G01	*MATalpha his3D1 leu2D0 lys2D0 ura3D0 flo8-1 gpd1D::GFP::URA3*	This study; BY4742 background. For monitoring GPD1.
WXXX.BY16	*MATa/MATalpha LYS2/lys2D0 ura3D0/URA3 FLO8/flo8-1 hoD::barcode::KanMX4/ho*	This study; 49 wild/lab plus 1 BY41 control. See Table S1 for details.
WXXX.G01	*MATa/MATalpha HIS3/his3D1 LEU2/leu2D0 LYS2/lys2D0 ura3D0/ura3D0 FLO8/flo8-1 hoD::barcode::KanMX4/ho GPD1/gpd1D::GFP::URA3*	This study. 49 wild/lab plus 1 lab/lab diploids for GPD1 quantification. See Table S1 for details.

Table 2. Negative feedback between rates of change in *GPD1*::GFP accumulation and viability.   To control for potential deviations from normality, non-parametric pairwise correlations between rates of change in *GPD1*::GFP accumulation and viability within (upper 3 rows) and between 2 hours time intervals (below).   Changes occurring in earlier intervals, which were potentially causal for later changes, are listed first. Results were qualitatively similar for Pearson's correlations with all significant P-values remaining <0.0001.   Significant comparisons are indicated in bold.

Interval	Variable	Interval	Variable	Spearman's r	Prob>|r|
0 – 2 hours	**Δ*GPD1*::GFP**	**0 - 2 hours**	**Δviability**	**0.7725**	**<.0001**
2 – 4 hours	**Δ*GPD1*::GFP**	**2 - 4 hours**	**Δviability**	**0.7217**	**<.0001**
4 - 6 hours	Δ*GPD1*::GFP	4 - 6 hours	Δviability	-0.1992	0.4282
0 – 2 hours	**Δviability**	**2 - 4 hours**	**Δ*GPD1*::GFP**	**-0.7411**	**<.0001**
0 – 2 hours	**Δviability**	**2 - 4 hours**	**Δviability**	**-0.9503**	**<.0001**
0 – 2 hours	**Δ*GPD1*::GFP**	**2 - 4 hours**	**Δ*GPD*1::GFP**	**-0.7697**	**<.0001**
0 – 2 hours	**Δ*GPD1*::GFP**	**2 - 4 hours**	**Δviability**	**-0.7696**	**<.0001**
1 - 2 hours	Δviability	4 - 6 hours	Δviability	-0.2178	0.3854
1 - 2 hours	Δviability	4 - 6 hours	Δ*GPD1*::GFP	-0.2549	0.3073
1 - 2 hours	Δ*GPD1*::GFP	4 - 6 hours	Δviability	-0.0464	0.8548
1 - 2 hours	Δ*GPD1*::GFP	4 - 6 hours	Δ*GPD1*::GFP	-0.4572	0.0565
2 - 4 hours	Δviability	4 - 6 hours	Δviability	0.0733	0.7726
2 - 4 hours	Δviability	4 - 6 hours	Δ*GPD1*::GFP	0.3333	0.1765
2 - 4 hours	Δ*GPD1*::GFP	4 - 6 hours	Δviability	0.1207	0.6332
2 - 4 hours	Δ*GPD1*::GFP	4 - 6 hours	Δ*GPD1*::GFP	0.3602	0.1421

Table 3. Growth of post diauxic cells at unprecedented limits of adaptation.   Shown are concentrations of agar media on which post diauxic strains could grow and form colonies.

[KCl] M	Wild/lab (G*PD1*) diploids*
2.0	W455
2.6	W027, W035, W167, W202, W203, W242, W285, W454
2.7	W033, W041, W042, W134, W136, W150, W166, W178, W195, W215, W217, W219, W235, W248, W282, W291, W292, W294, BC41
2.8	W037, W044, W050, W153, W157, W163, W164, W179, W189, W206, W238, W244, W245, W249, W255, W276, W301, W340
2.9	W173, W211, W343

Table 4.   Osmotic stress signaling behavior (rank) predicted early and late viability of post diauxic cultures in osmotic stress. Least squares predictions of early and late viability by linear and 2^{nd} order quadratic fits of fluorescence pre-accumulated into the G3 Gaussian at time 0 (G3_0) and ranked signaling behavior.   The Bonferroni cutoff at the 0.05 level, based on 4 tests per data set, was 0.0125.   Significant fits with lowest root mean squared errors and highest fraction of variation explained (R^2) shown in bold, predicted values for optimum (x) and value at optimum (y) for non-significant fits are marked NS.

	G3_0 fluorescence (AU)		Signaling (rank)	
	quadratic	linear	quadratic	linear
0 hours, 0M KCl				
probability > F	**0.0006**	0.0016	0.9629	0.8454
R_square	**0.2715**	0.1891	0.0016	0.0008
root_mean_square_error	**1.3935**	1.4548	1.6313	1.6149
max_viability at optimum (%)	**98.8**		NS 98.2	
optimum	**2626.6**		NS 31.9	
20 hours, 2.5M KCl				
probability > F	**< 0.0001**	< 0.0001	**0.0055**	0.6023
R_square	**0.4286**	0.3138	**0.1987**	0.0057
root_mean_square_error	**7.0949**	7.6940	**8.4022**	9.2615
max_viability at optimum (%)	**86.7**		**86.3**	
optimum	**2662.5**		**24.4**	
48 hours, 2.5M KCl				
probability > F	**< 0.0001**	0.0003	**0.0013**	0.9159
R_square	**0.3148**	0.2371	**0.2464**	0.0002
root_mean_square_error	**7.6281**	8.1323	**8.1679**	9.3096
max_viability at optimum (%)	**82.2**		**82.9**	
optimum	**2622.3**		**25.7**	
72 hours, 2.5M KCl				
probability > F	**0.0010**	0.0033	**0.0062**	0.8464
R_square	**0.2556**	0.1660	**0.1943**	0.0008
root_mean_square_error	**9.9877**	10.4608	**10.3903**	11.4501
max_viability at optimum (%)	**73.5**		**74.3**	
optimum	**2586.8**		**25.1**	
96 hours, 2.5M KCl				
probability > F	**0.0060**	0.0018	**0.0065**	0.0954
R_square	**0.1956**	0.1862	**0.1927**	0.0569
root_mean_square_error	**8.4186**	8.3789	**8.4338**	9.0201
max_viability at optimum (%)	**71.7**		**71.0**	
optimum	**3389.8**		**21.3**	
120 hours, 2.5M KCl				
probability > F	0.0791	0.0413	**0.0004**	0.0037
R_square	0.1023	0.0839	**0.2859**	0.1625
root_mean_square_error	8.5666	8.5636	**7.6405**	8.1881
max_viability at optimum (%)	NS 67.7		**74.7**	
optimum	NS 1679		**18.1**	

144 hours, 2.5M KCl				
probability > F	0.3473	0.1675	**0.0047**	0.0501
R_square	0.0440	0.0393	**0.2038**	0.0776
root_mean_square_error	9.6837	9.6059	**8.8377**	9.4123
max_viability at optimum (%)	NS 61.9		**68.3**	
optimum	NS 1481.3		**20.4**	
168 hours, 2.5M KCl				
probability > F	0.1785	0.3433	**<0.0001**	< 0.0001
R_square	0.0707	0.0187	**0.5733**	0.4911
root_mean_square_error	12.3795	12.5879	**8.3885**	9.0652
max_viability at optimum (%)	NS 56.2		**68.4**	
optimum	NS 2077		**9.7**	
24 hours, 3M KCl				
probability > F	0.0107	0.0292	**< 0.0001**	< 0.0001
R_square	0.1757	0.0952	**0.5254**	0.2980
root_mean_square_error	11.8963	12.3332	**9.0267**	10.8635
max_viability at optimum (%)	71.7		**65.0**	
optimum rank	2029.2		**18.1**	
48 hours, 3M KCl				
probability > F	0.0294	0.0140	**< 0.0001**	0.0003
R_square	0.1394	0.1193	**0.5382**	0.2402
root_mean_square_error	12.8903	12.9029	**9.4420**	11.9849
max_viability at optimum (%)	NS 57.1		**62.3**	
optimum	NS 2867.3		**19.7**	
72 hours, 3M KCl				
probability > F	0.0519	0.0395	**< 0.0001**	0.0002
R_square	0.1183	0.0853	**0.5459**	0.2460
root_mean_square_error	12.4241	12.5218	**8.9158**	11.3687
max_viability at optimum (%)	NS 51.4		**56.9**	
optimum	NS 2651		**19.7**	
Adaptation limit				
probability > F	0.5435	0.2699	**0.0002**	0.0086
R_square	0.0262	0.0258	**0.3099**	0.1379
root_mean_square_error	0.1344	0.1330	**0.1132**	0.1251
max_concentration optimum (M)	NS 2.8		**2.8**	
optimum	NS -95.6		**31.2**	

575 **Table S1. Haploid derivatives of wild strains.**   The source for all wild strains in this study
576 was the strain collection of the Royal Netherlands Academy of Arts and Sciences over the past
577 100 years (Table 1 and Table S1).   This resource has been deposited at the Yeast Genetic
578 Resources Lab of the National BioResource Project in Osaka, Japan.
579

Alias	MAT	Genotype	Comments
WH027	a	MATa ura3D0 hoD::barcode::KanMX4	Original CBS #: CBS 459; isolated in 1938 in Castellina, Italy from grape must; barcode #29(Uptag sequence: GGCCCGCACACAATTAGGAA, Downtag sequence: GCGCCGCATTAACTAAACTA)
WH030	a	MATa ura3D0 hoD::barcode::KanMX4	Original CBS #: CBS 1508; isolated in 1927 from starter for sorghum brandy; YH note: mating defective, hard to make wild/lab diploids; barcode #16(Uptag sequence: GTCCGAACTATCAACACGTA, Downtag sequence: GCGCACGAGAAACCTCTTAA)
WH033	a	MATa ura3D0 hoD::barcode::KanMX4	Original CBS #: CBS 405 ; isolated in 1925 in West Africa from catabo for billi wine, from Osbeckia grandiflora; barcode #187(Uptag sequence: CCGTGTACTGAATTACGATC, Downtag sequence: CCATCTTTGGTAATGTGAGG)
WH035	a	MATa ura3D0 hoD::barcode::KanMX4	Original CBS #: CBS 5822; isolated in 1967 from malt wine; barcode #30(Uptag sequence: GGTCTATGCAAACACCCGAA, Downtag sequence: GCCGTCTTGACAACCTTATA)
WH037	a	MATa ura3D0 hoD::barcode::KanMX4	Original CBS #: CBS 1395; isolated in 1922 from an unknown source ; barcode #235(Uptag sequence: GGCTAAGGGACAACACCTCA, Downtag sequence: GCCCGGCACATAGAAGTAAC)
WH041	a	MATa ura3D0 hoD::barcode::KanMX4	Original CBS #: CBS 5635; isolated in 1958 in South Africa from grape must; barcode #2(Uptag sequence: CCATGATGTAAACGATCCGA, Downtag sequence: TATATGGCAGCAGATCGCCG)
WH042	a	MATa ura3D0 hoD::barcode::KanMX4	Original CBS #: CBS 3081; isolated in 1958 in Spain from alpechin; barcode #12(Uptag sequence: GTGCGAACCAACGTACTACA, Downtag sequence: GCAGGAACACCACAGGGTTA)
WH044	a	MATa ura3D0 hoD::barcode::KanMX4	Original CBS #: CBS 422; isolated in 1926 in Odessa, Ukraine from beer; barcode #135(Uptag sequence: CCCGCGATTGTAATGAATAG, Downtag sequence: CATACTACGTGGGACAGTTG)
WH050	a	MATa ura3D0 hoD::barcode::KanMX4	Original CBS #: CBS 5112; isolated in 1962 in Spain from grape must; barcode #49(Uptag sequence: CTTACTGATAGCGTAGAGGT, Downtag sequence: GTGGTCTGCAAACCCAACAA)
WH134	a	MATa ura3D0 hoD::barcode::KanMX4	Original CBS #: CBS 2964; isolated in 1947 in Copenhagen, Denmark from distiller's rum yeast; barcode #18(Uptag sequence: GCCCTGATAACAAGGTGTAA, Downtag sequence: GCGCCTATTACACAAACGTA)

WH136	a	MATa ura3D0 hoD::barcode::KanMX4	Original CBS #: CBS 2811; isolated in 1957 from wine; wine yeast; barcode #20(Uptag sequence: GTGAGCGAAACACCGCGTAA, Downtag sequence: GGTAATACGCAACTCCTCTA)
WH150	a	MATa ura3D0 hoD::barcode::KanMX4	Original CBS #: CBS 2962; isolated in 1947 in Copenhagen, Denmark from ; distiller's rum yeast; barcode #15(Uptag sequence: GCCGTAGCCACAAGAGTTAA, Downtag sequence: GCGGCCACTTACACAAATTA)
WH153	a	MATa ura3D0 hoD::barcode::KanMX4	Original CBS #: CBS 6458; isolated in 1972 location not reported; barcode #37(Uptag sequence: GGGACCGCCAAAGCTATCAA, Downtag sequence: GTGAACAATAACGGCCTTGA)
WH157	a	MATa ura3D0 hoD::barcode::KanMX4	Original CBS #: CBS 6506; isolated in 1973 in UK brewery; killer yeast; barcode #53(Uptag sequence: CTGAGCGTAGGATATTCCGT, Downtag sequence: GCCGGTCGCAAACTCATAAA)
WH163	a	MATa ura3D0 hoD::barcode::KanMX4	Original CBS #: CBS 6914; isolated in 1977 in Spain from white wine; barcode #51(Uptag sequence: CTACGTCGGCTCATAGTCGT, Downtag sequence: GCTCTCGGCCAAGGAAACAA)
WH164	a	MATa ura3D0 hoD::barcode::KanMX4	Original CBS #: CBS 6978; isolated in 1984 from wine; wine yeast; barcode #59(Uptag sequence: CACTCGGATTCAGTTCTAGT, Downtag sequence: GGCCTTGCCAAACAGTCAAA)
WH166	a	MATa ura3D0 hoD::barcode::KanMX4	Original CBS #: CBS 7072; isolated in 1980 from distillate; barcode #62(Uptag sequence: CCTAGTTCGAGATTGCGAGT, Downtag sequence: GTGGTCGCCCAAGCAACAAA)
WH167	a	MATa ura3D0 hoD::barcode::KanMX4	Original CBS #: CBS 7173; isolated in 1985 from catabo for billi wine; wine yeast; barcode #43(Uptag sequence: CAGTATGCTAGATTCCGGGT, Downtag sequence: GTCCTCGCAAGAAAGGCCAA)
WH173	a	MATa ura3D0 hoD::barcode::KanMX4	Original CBS #: CBS 8615; isolated in 1998 in Italy from grape must; dry yeast for wine making; barcode #61(Uptag sequence: CCTGTAGTACGAGTATGAGT, Downtag sequence: GGTCTGCCCAAAGTCACAAA)
WH178	a	MATa ura3D0 hoD::barcode::KanMX4	Original CBS #: CBS 1192; isolated in 1928 from wine; wine yeast; barcode #155(Uptag sequence: CGCACACGATTAAGGTCCAG, Downtag sequence: CACTGTTGGTAAGGTCTATG)
WH179	a	MATa ura3D0 hoD::barcode::KanMX4	Original CBS #: CBS 1193; isolated in 1928 from wine; wine yeast; barcode #70(Uptag sequence: CAATAGGGTGTGACAGTTCT, Downtag sequence: CTACTTCGCGTGAGCTGGTT)
WH189	a	MATa ura3D0 hoD::barcode::KanMX4	Original CBS #: CBS 1241; isolated in 1930 from an unknown source ; barcode #212(Uptag sequence: CCACTTAGTTCAATAGGCGC, Downtag sequence: CCGAGTATTACATTCTCACG)

WH195	a	MATa ura3D0 hoD::barcode::KanMX4	Original CBS #: CBS 1256; isolated in 1937 from port wine; barcode #123(Uptag sequence: CGTGGAGCAGTTCGTATAAT, Downtag sequence: CTCGACGCTGGACGTTATGT)
WH202	a	MATa ura3D0 hoD::barcode::KanMX4	Original CBS #: CBS 7372; isolated in 1988; killer yeast, K2Rd (Young & Yagiu), K2R2 (Wickner); barcode #119(Uptag sequence: CAACGTAGAGTGAGGTACAT, Downtag sequence: CACTTAGCTTAGACTCGTGT)
WH203	a	MATa ura3D0 hoD::barcode::KanMX4	Original CBS #: CBS 7438; isolated in 1989 from wine; wine yeast; barcode #65(Uptag sequence: CTTTCGGACGTATGTGCAGT, Downtag sequence: CCTTGATGATAGAGGGCTTT)
WH206	a	MATa ura3D0 hoD::barcode::KanMX4	Original CBS #: CBS 7833; isolated in 1994 in Missouri, USA from lung of a man with immune deficiency syndrome; virulent strain; barcode #82(Uptag sequence: CATACAAAGAGAGGTGTCCT, Downtag sequence: CCCTTGCGATTGGTGCAGTT)
WH211	a	MATa ura3D0 hoD::barcode::KanMX4	Original CBS #: CBS 7838; isolated in 1994 in USA from patient; barcode #144(Uptag sequence: CGATACAAGTAAGTTGCGAG, Downtag sequence: CCTCTTACGAGATAGCGGTG)
WH215	a	MATa ura3D0 hoD::barcode::KanMX4	Original CBS #: CBS 7962; isolated in 1984 in Sao Paulo, Brazil from fermenting concentrated syrup from sugar cane; barcode #94(Uptag sequence: CCCGATTGAGGCATGGTTAT, Downtag sequence: CGCTTCGAGTATGGGATATT)
WH217	a	MATa ura3D0 hoD::barcode::KanMX4	Original CBS #: CBS 7964; isolated in 1995 in Sao Paulo, Brazil from fermenting concentrated syrup from sugar cane; barcode #92(Uptag sequence: CGCGGAGTATAGAGCTTTAT, Downtag sequence: CAATCGCTCGGAGGCGTATT)
WH219	a	MATa ura3D0 hoD::barcode::KanMX4	Original CBS #: CBS 439 ; isolated in 1933 in Lager Schartel, Germany from Silvaner grapes; barcode #93(Uptag sequence: CGACCCTGATGATCCTTTAT, Downtag sequence: CTACGGGCTCGATGCCTATT)
WH235	a	MATa ura3D0 hoD::barcode::KanMX4	Original CBS #: CBS 5952; isolated in 1968; barcode #102(Uptag sequence: GGCTACGATACATCTTCATC, Downtag sequence: CATTTGTAACCAGTTCGCTC)
WH238	a	MATa ura3D0 hoD::barcode::KanMX4	Original CBS #: CBS 6223; isolated in 1969 in Chile from grape juice; radiation resistant; barcode #104(Uptag sequence: CTATGTGCGGTAAGACGTAT, Downtag sequence: CGGCGTAGATTGTTAGCATT)
WH242	a	MATa ura3D0 hoD::barcode::KanMX4	Original CBS #: CBS 6333; isolated in 1942 in Costa Rica from rotting banana; Strain name NRRL Y-1350 (synonymous designation; NRRL YB-210, NRRL-210, NRRL-B210; Mortimer and Johnston (1986), Genetics 113: 35); barcode #56(Uptag sequence: CCTGTAGATTGACGTGTAGT, Downtag sequence: GCCCTCGTGACAAATCGAAA)

WH244	a	MATa ura3D0 hoD::barcode::KanMX4	Original CBS #: CBS 8266; isolated in 1996 in from rotting fig; Strain name X2180, derived from S288C by self-diploidization (Mortimer and Johnston (1986), Genetics 113: 35). Did not survive freeze-drying.; barcode #122(Uptag sequence: CAGAGGGCACTGTTCTTAAT, Downtag sequence: CCCTGCTGTAGAGGTTATGT)
WH245	a	MATa ura3D0 hoD::barcode::KanMX4	Original CBS #: CBS 3000; isolated in 1956 in Pakistan from palm wine; wine yeast; barcode #138(Uptag sequence: CACATCGTTTAACACTGGAG, Downtag sequence: CTAGGAGGTTACAGTCATTG)
WH248	a	MATa ura3D0 hoD::barcode::KanMX4	Original CBS #: CBS 8049; isolated in 1981 from fish food; feed for fish and crustaceans; barcode #108(Uptag sequence: CGACCCGATGTAGTAGATAT, Downtag sequence: CCGCCGGATGTGATATAATT)
WH249	a	MATa ura3D0 hoD::barcode::KanMX4	Original CBS #: CBS 6069; isolated in 1981; hybrid strain (Y55-2 x JJ101); barcode #87(Uptag sequence: CACTGTGACCGAGGGATACT, Downtag sequence: CGCGCTATTATACTCGACTT)
WH255	a	MATa ura3D0 hoD::barcode::KanMX4	Original CBS #: CBS 3090; isolated in 1958 from white grape must; barcode #72(Uptag sequence: CACTGTGGACGATACGGTCT, Downtag sequence: CTGTACGTGCGATACTCGTT)
WH276	a	MATa ura3D0 hoD::barcode::KanMX4	Original CBS #: CBS 1394; isolated in 1924 from pressed yeast; distillery yeast; barcode #176(Uptag sequence: CCACCGATGTAATTTGAGTC, Downtag sequence: CACTCTGCGTTAATGTTGGG)
WH282	a	MATa ura3D0 hoD::barcode::KanMX4	Original CBS #: CBS 1460; isolated in 1927 in Indonesia from fermenting fruit; barcode #115(Uptag sequence: CATACTTAGGGATCAGGGAT, Downtag sequence: CCTTGTCTGAGAGCCGTTGT)
WH285	a	MATa ura3D0 hoD::barcode::KanMX4	Original CBS #: CBS 1479; isolated in 1928 from wine; wine yeast; barcode #240(Uptag sequence: GCGGCCAATAGTAAACTTCA, Downtag sequence: GCCGCCGTGATAAGAAACAC)
WH291	a	MATa ura3D0 hoD::barcode::KanMX4	Original CBS #: CBS 1576; isolated in 1931 in Sulawesi, Indonesia from sap of Arenga palm; barcode #117(Uptag sequence: CCTGAGGACTTATTCACGAT, Downtag sequence: CATTGGATTAGACCGTGTGT)
WH292	a	MATa ura3D0 hoD::barcode::KanMX4	Original CBS #: CBS 1582; isolated in 1948 in Portugal from wine; barcode #118(Uptag sequence: CCGATTAGAGGTTGACAGAT, Downtag sequence: CACTGACTTCGAGGTCGTGT)
WH294	a	MATa ura3D0 hoD::barcode::KanMX4	Original CBS #: CBS 1585; isolated in 1934 from sake-moto; sake yeast; barcode #180(Uptag sequence: CATTAAGGCGCACGTTTATC, Downtag sequence: CTATCCTAGAGATTTGAGGG)
WH301	a	MATa ura3D0 hoD::barcode::KanMX4	Original CBS #: CBS 1594; isolated in 1936 from juice of aren palm; barcode #182(Uptag sequence: CACGTTTGCGAATAGGTATC, Downtag sequence: CAGATACTATTAAGTGCCGG)

WH340	a	MATa ura3D0 hoD::barcode::KanMX4	Original CBS #: CBS 2805; isolated in 1954 from wine; wine yeast, particularly suitable for fruit wines; barcode #233(Uptag sequence: GCCGGGCTTAAATTGAATCA, Downtag sequence: GCTCCGACTGAAGAACTAAC)
WH343	a	MATa ura3D0 hoD::barcode::KanMX4	Original CBS #: CBS 2808; isolated in 1954 from grapes (Blauer Portugieser); wine yeast, suitable for fruit wines, yields more than 18% of alcohol; barcode #224(Uptag sequence: CCCGTGAATATAAGTGAAGC, Downtag sequence: CCTGGATTTGAAGCGTATAG)
WH454	a	MATa ura3D0 hoD::barcode::KanMX4	Original CBS #: CBS 6412; isolated in 1952 from sake; sake yeast; barcode #192(Uptag sequence: CCTTAGGGATAATGAGTTGC, Downtag sequence: CCAGTGTTCTAACGTGCAGG)
WH455	a	MATa ura3D0 hoD::barcode::KanMX4	Original CBS #: CBS 440 ; isolated in 1934 in Taiwan from molasses; barcode #249(Uptag sequence: GCCCAGGCTAAATGTTAAGA, Downtag sequence: GAAGTACGCTCAAGACCGAC)
BC4741 (BY41)	a	MATa his3D1 leu2D0 met15D0 ura3D0 flo8-1 hoD::barcode::KanMX4	Original CBS #: Lab strain, BY4741; isolated in 1938 in Merced, CA, USA from rotting fig; derived from S288C, of which strain 88% of the gene pool is contributed by strain EM93 (Mortimer and Johnston (1986), Genetics 113: 35). Barcode #266(Uptag sequence: GGCCTAACTCAACAGACGGA, Downtag sequence: GCGCTCGACTAAGAGAAACC)

582 **Table S2. Clustering statistics used to rank signaling behavior.** Statistics showing the

583 fraction of 17,000 permutations in which strains ranked with 50% of other strains in each mean

584 cluster.   These data were used to rank total signaling behaviors from most cautious (1) to most

585 reckless (50) based on the fraction of time each strain was associated with its mean cluster

586 (characteristic of that cluster).   See Figure 2.

Rank	Cluster	Strain	MC0	MC1	MC3	MC2	MC5	MC4
1	MC0	W455	0.8484	0.1516	0	0	0	0
2	MC0	W167	0.8365	0.1635	0	0	0	0
3	MC1	W219	0.6375	0.3625	0	0	0	0
4	MC1	W217	0.5536	0.4460	0.0004	0	0	0
5	MC1	W027	0.5214	0.4755	0.0028	0.0002	0	0
6	MC1	W042	0.4850	0.4892	0.0191	0.0044	0.0023	0.0001
7	MC1	W235	0.1976	0.5532	0.2266	0.0226	0	0.0002
8	MC1	W340	0.1700	0.5803	0.2354	0.0142	0	0
9	MC1	W454	0.3355	0.6610	0.0034	0	0	0
10	MC1	W134	0.1686	0.7376	0.0821	0.0117	0	0
11	MC1	W276	0.1617	0.7643	0.0658	0.0083	0	0
12	MC1	W294	0.1692	0.7834	0.0440	0.0034	0	0
13	MC1	W157	0.1195	0.7851	0.0866	0.0088	0	0
14	MC1	W202	0.1671	0.7855	0.0446	0.0027	0	0
15	MC1	W238	0.1337	0.8048	0.0580	0.0036	0	0
16	MC1	W035	0.1293	0.8086	0.0572	0.0050	0	0
17	MC1	W248	0.1494	0.8331	0.0172	0.0003	0	0
18	MC3	W130	0.0003	0.0671	0.7020	0.2307	0	0
19	MC3	W136	0	0.0092	0.6572	0.3334	0.0002	0
20	MC3	W203	0.0025	0.1348	0.6410	0.2214	0.0002	0
21	MC3	W285	0.0003	0.0806	0.6245	0.2924	0.0022	0
22	MC3	W163	0.0011	0.1474	0.6228	0.2285	0.0002	0
23	MC3	W206	0	0.0036	0.6203	0.3747	0.0014	0
24	MC3	BY41	0	0.0123	0.6102	0.3768	0.0008	0
25	MC3	W041	0.0002	0.0233	0.5625	0.4090	0.0050	0
26	MC3	W343	0.0012	0.0432	0.5522	0.3910	0.0123	0.0002
27	MC3	W292	0.0201	0.1962	0.4883	0.2893	0.0062	0
28	MC3	W189	0.0001	0.0205	0.4757	0.4715	0.0319	0.0003
29	MC3	W211	0	0	0.3698	0.5572	0.0696	0.0033
30	MC2	W245	0	0	0.2579	0.6898	0.0523	0
31	MC2	W291	0	0	0.2211	0.6786	0.0994	0.0009
32	MC2	W164	0	0	0.1517	0.6601	0.1852	0.0031

33	MC2	W249	0	0	0.3349	0.6417	0.0233	0.0001
34	MC2	W166	0	0	0.3372	0.6407	0.0220	0.0001
35	MC2	W179	0	0.0002	0.3569	0.6212	0.0217	0
36	MC2	W244	0	0	0.1478	0.6160	0.2225	0.0137
37	MC2	W173	0	0.0003	0.3801	0.6005	0.0188	0.0002
38	MC2	W215	0	0.0012	0.3162	0.6000	0.0815	0.0011
39	MC2	W037	0	0	0.0714	0.5308	0.3863	0.0115
40	MC2	W255	0	0.0011	0.4771	0.5141	0.0077	0
41	MC2	W050	0	0	0.0567	0.4672	0.4417	0.0343
42	MC5	W044	0	0	0.0021	0.0728	0.8501	0.0750
43	MC5	W195	0	0	0.0019	0.0744	0.8501	0.0736
44	MC5	W301	0	0	0.0029	0.0882	0.8203	0.0887
45	MC5	W150	0.0001	0.0004	0.0006	0.0414	0.7978	0.1598
46	MC5	W033	0	0	0	0.0279	0.7196	0.2525
47	MC5	W153	0	0	0	0.0181	0.6545	0.3274
48	MC4	W282	0	0	0	0	0.0596	0.9404
49	MC4	W242	0	0	0	0.0002	0.1477	0.8521
50	MC4	W178	0	0	0	0.0017	0.1697	0.8286

590 **Table S3**

591 **Plate key**

592 **Left**

	1	2	3	4	5	6
A	W027.BY01	W044.BY01	W153.BY01	W173.BY01	W203.BY01	W235.BY01
B	W033.BY01	W050.BY01	W157.BY01	W178.BY01	W206.BY01	W238.BY01
C	W035.BY01	W130.BY01	W163.BY01	W179.BY01	W211.BY01	W242.BY01
D	W037.BY01	W134.BY01	W164.BY01	W189.BY01	W215.BY01	W244.BY01
E	W041.BY01	W136.BY01	W166.BY01	W195.BY01	W217.BY01	W245.BY01
F	W042.BY01	W150.BY01	W167.BY01	W202.BY01	W219.BY01	W248.BY01

594 **Right**

	1	2	3
A	W249.BY01	W292.BY01	W455.BY01
B	W255.BY01	W294.BY01	
C	W276.BY01	W301.BY01	
D	W282.BY01	W340.BY01	
E	W285.BY01	W343.BY01	
F	W291.BY01	W454.BY01	

Figure legends

Figure 1. Rate of osmotic stress signaling in exponential cultures with negative feedback

predicts early survival and robust recovery in moderate osmotic stress.

A. Time course of mean accumulated *GPD1*::GFP fluorescence in exponential cultures

exposed to 0.75 and 1.5 M KCl. Each point represents an independent replicate

measurement; curves connect strain means at each time. In the absence of stress, all

strains had high steady-state viability (propidium iodide dye exclusion; range 96.3 –

98.7%; mean 97.6%) and relatively low mean GFP fluorescence indicating low

background activity of HOG pathway signaling through the GPD1 promoter and low

GPD1::GFP accumulation (range 12.7 – 34.8 AU; mean 18.8 AU).

B. Pie charts show relative changes after 2 hours in 0.75M KCl of mean viability (shaded

area), mortality (white area) and *GPD1*::GFP accumulation (percent opacity). Strains

ordered by viability at 2 hours, which was proportional to viability of non-disrupted

controls having two intact copies of the GPD1 gene (R^2 = 0.7085; P<0.0001;).

C. Relationship between mean *GPD1*::GFP accumulation and viability in mid-exponential

cultures exposed to 0.75 M KCl for 0, 2, 4, and 6 hours (h). Each data point represents

the average of at least three replicates per strain (~10,000 cells/ sample). The ellipses

indicate correlations between viability and fluorescence at alpha = 0.95. The inserts show

relationships between changes in *GPD1*::GFP and viability over each time interval.

D. Robust recovery of steady-state viability after 4 hours in 0.75 M KCl driven by negative

feedback (see text; recovery = (0.7670) early mortality + 3.4936; R^2 = 0.9351; P<

0.0001).

E. Plausible integrating negative control via multiple feedback mechanisms resulting in

accumulation of intracellular glycerol. Integrating feedback control would assure perfect

adaptation of stress responses, water balance and steady state viability. Note that the

persistence of dead cells over the course of the experiment precludes 100% recovery.

Figure 2. Signaling behaviors predict the survival and fitness of older cultures in severe

osmotic stress.

A. Strains classified by mean cluster (MC0 – MC5) and ranked top (1) to bottom (50)

according to changes in GPD1::GFP accumulation over time (see Table S2 and methods).

Each time point shows representative distributions of GPD1::GFP accumulation (green)

and relative survival red (99.7% viability) to blue (11.7% viability). Cells above the 89[th]

percentile (top 11%) are shown in black. Prior to osmotic challenge steady-state viabilities

were uniformly high (range 93.0 – 99.6%; mean 98.2%). Rank-ordered mean clusters are

topographically equivalent to a sequential ordering.

B. B. Relative viability of post-diauxic cultures (WXXX.BY01 controls) incubated in 3 M KCl

before plating on iso-osmolar media. Cultures were re-ordered according to the ranked

signaling behavior given in Figure 2A. The same experiment as originally plated is shown

in Figure S1. Strains color coded as in Figure 1C for comparison of exponential and post

diauxic cultures.

Figure 3. Extreme cautious and reckless cell types observed in post diauxic cultures

produce observed variation in cautious to reckless signaling strategies in a simple bet

hedging model.

a. Cells with the most aggressive signaling began to die after long periods in severe stress

leaving increasing fraction of cells with lower *GPD1*::GPF accumulations.   Shown are

distributions of accumulated *GPD1*::GFP and viability in replicate cultures of W242 (rank

49) after 168 hours in 2.5 M KCl.   Mean (x), standard deviation (std), and weight (w; the

fraction of cells in each distribution) are given.   Sum (red) shows the cumulative fit of the

4 learned Gaussians.

b. Static viability of post diauxic cells of strain W027 exposed to 3 M in microfluidic

chambers.   Individual cell behaviors mirror population behaviors measured by flow

cytometry – e.g longer lag periods and increased accumulations of GPD1::GFP with

increasing osmotic stress. Colored traces indicate accumulated fluorescence in

representative cells in 1.5 (green), 2.0 (blue), 2.5 (red) and 3.0 M KCl (yellow).   Arrows

indicate average time to the first cell division +/- standard deviations.

c. Generation of intermediate optima in simple 2-state bet hedging model with heritable

production of cautious (static) and reckless cell types.   Bet hedging strategy P was

defined as the probability of cautious cells for $0 \leq P \leq 1$. Relative fitness was measured

for all strategies after 10 generations in each environment. All nine possible

environmental shifts are shown. Nine possible 2-state environmental shifts between three

general osmotic stress environments were considered: permissive (E0; all cells grow

equally well), restrictive (E1; reckless cells divide, cautious cells survive without dividing),

and killing (E2; reckless cells die, cautious cells survive without cell division). Only in

shifts from moderate to more severe (E1 -> E2) do bet hedging, intermediate strategies (0

< P < 1) prevail.   The optimum strategy P depends on the number of generations in the

first moderate stress environment. Results were independent of the number of

generations in the first environment except as shown when E1 was the first environment.

Figure S1. Monophasic signaling behavior of exponential cultures described well by

mean *GPD1*::GFP accumulations at each time point.

Mean (x), standard deviation (std), and weight (w; the fraction of cells in each distribution) are

given.   Sum (red) shows the cumulative fit of the 4 learned Gaussians.   The representative

strains shown are color-coded as in Figure 1B.

Figure S2. Static viability and survival of post diauxic cultures in extreme osmotic stress.

Strains were incubated for up to 5 weeks in 3 M KCl plated on iso-osmolar media.   A plate key

is given in Table S3.   The same data are shown sorted by rank in Figure 2B.

Figure S3a–c. Reproducibility of cell distributions in strains with cautious to reckless

signaling.   Learned distributions of GPD1::GFP accumulation for post diauxic strains exposed

to 2.5 M KCl for the times shown.   Mean (x), standard deviation (std), and weight (w; the

fraction of cells in each distribution) are given.   Sum (red) shows the cumulative fit of the 4

learned Gaussians.   The representative strains shown are color-coded as in Figure 1B.

Figure S4. Rank predicts increasingly aggressive osmotic stress signaling.   Rank

predicts the relative proportion of cells above a threshold set at the top 11% of accumulation of

GPD1::GFP in post diauxic cultures exposed to 2.5 M KCl for the times shown.   Mean clusters

are indicated with increasingly lighter grey scale in order of signaling (MC0, MC1, MC3, MC2,

MC5, MC4).

Figure S5. Mean clusters were differentiated for predicts viability of post diauxic cultures.

Mean cluster predicted viability.   Shown are average viabilities each mean cluster were

analyzed by ANOVA or where appropriate by Welch's ANOVA at each time.   Asterisks indicate

significance at the ≤0.05 level.   Decreasing mean viabilities at each time are indicated

(horizontal lines).

695