Plant diversity accurately predicts insect diversity in two tropical landscapes: BioRxiv preprint

Zhang, K.,*†[1] Lin, S.L., ‡[1] Ji, Y.Q.,* Yang, C.X.,* Wang, X.Y.,*† Yang, C.Y.,*

Wang, H.,§ Jiang, H.S.,‡ Harrison, R.D.¶**$ and Yu, D.W.*††

* State Key Laboratory of Genetic Resources and Evolution, Kunming Institute of Zoology, Chinese Academy

of Sciences, Kunming, Yunnan 650223, China

† Kunming College of Life Science, University of the Chinese Academy of Sciences, Kunming 650204, China

‡ School of Life Sciences, South China Normal University, Guangzhou, Guangdong 510631, China

§ Hainan Yinggeling National Nature Reserve, Baisha, Hainan 572800, China

¶ World Agroforestry Center, East and Central Asia Regional Office, Kunming 650201, China

** Center for Mountain Ecosystem Studies (CMES), Kunming Institute of Botany, Chinese Academy of

Sciences, Kunming 650201, China

$ Current address: World Agroforestry Centre, East & Southern Africa Region, 13 Elm Road, Woodlands,

Lusaka, Zambia.

†† School of Biological Sciences, University of East Anglia, Norwich Research Park, Norwich, Norfolk NR47TJ,

UK

Key-words: Arthropoda, biodiversity, biomonitoring, China, host specificity, insect-plant interactions,

species richness, surrogate species

co-corresponding authors. DWY +86 871 68125438 (ofc), 86 871 65110887 (fax),

dougwyu@gmail.com; RDH +86 18687136498 (ofc), 86 871 65223377 (fax), r.harrison@cgiar.org

Send proofs to DWY.

Running title: Plants accurately predict insects

[1] Shared first-authorship.

Abstract

Plant diversity surely determines arthropod diversity, but only moderate correlations between arthropod and plant species richness had been observed until Basset *et al.* (2012, Science 338: 1481-1484) finally undertook an unprecedentedly comprehensive sampling of a tropical forest and demonstrated that plant species richness could indeed accurately predict arthropod species richness. We now require a high-throughput pipeline to operationalize this result so that we can (1) test competing explanations for tropical arthropod megadiversity, (2) improve estimates of global eukaryotic species diversity, and (3) use plant and arthropod communities as efficient proxies for each other, thus improving the efficiency of conservation planning and of detecting forest degradation and recovery. We therefore applied metabarcoding to Malaise-trap samples across two tropical landscapes in China. We demonstrate that plant species richness can accurately predict arthropod (mostly insect) species richness and that plant and insect community compositions are highly correlated, even in landscapes that are large, heterogeneous, and anthropogenically modified. Finally, we review how metabarcoding makes feasible highly replicated tests of the major competing explanations for tropical megadiversity.

Introduction

The relationship between plant diversity and insect diversity is fundamental to ecology because (1) it underpins global species estimates of arthropods based on plant diversity (Condon *et al.* 2008; Hamilton *et al.* 2013; Stork *et al.* 2015); (2) it improves our understanding of the drivers of arthropod diversity and assembly structure (Novotny *et al.* 2006; Lewinsohn & Roslin 2008; Pellissier *et al.* 2013); and (3) a strongly predictive relationship could open the way to using plant community metrics as surrogates for arthropod communities (and vice versa), thus improving the efficiency of efforts to conserve biodiversity, and ecosystem functions and services (Castagneyrol & Jactel 2012). In particular, arthropod species richness and community composition (i.e. species identities and frequencies) could serve as a sensitive method for detecting and quantifying the degree of forest degradation and recovery (Ji *et al.* 2013; Edwards *et al.* 2014), which is especially needed for the monitoring and verification of contracts to pay local populations and governments to protect and restore standing forest, also known as PES (Payments for Environmental Services) and REDD+ schemes (Reduction in Emissions from Deforestation and Degradation) (Bustamante *et al.* 2015).

   A priori, plant diversity must surely predict insect diversity (Lewinsohn & Roslin 2008; Haddad *et al.* 2009), because insects depend directly (via herbivory, pollination, and housing) and indirectly (via consumption of herbivores) on plant species, and insect herbivores show dietary specialization to subsets of plant taxa (Novotny & Basset 2005). In addition, plant and insect coevolutionary interactions have driven the evolution of the vast diversity of plant and insect species today (Thompson 1994; Cruaud *et al.* 2012; Edger *et al.* 2015). (N.B. In practice, studies of terrestrial arthropod diversity tend to focus on insects, because insects

make up the majority of described arthropods and can be easier to sample. This study will also

follow this practice.)

Not surprisingly, many papers have reported significant correlations between arthropod (mostly insect) and plant beta and alpha diversities (reviews in Lewinsohn & Roslin 2008; Castagneyrol & Jactel 2012; Pellissier *et al.* 2013). In particular, work by Novotny *et al.* (2002, 2006, 2007) has strongly suggested that the primary driver of high species richness among herbivorous insects in tropical forests is simply the greater number of plant species in the tropics, rather than either higher levels of host specificity and beta diversity or more insect species per area of foliage. In short, the local number of insect species should increase nearly linearly with the local number of plant species, and with a slope greater than one, since each plant species is associated with multiple herbivore and predator species (Castagneyrol & Jactel 2012).

However, correlations between arthropod (mostly insect) and plant species-richness have shown only moderate fit. Castagneyrol and Jactel's (2012) comprehensive meta-analysis reported mean correlations of only 0.39 and 0.51 for studies in single habitats and across multiple habitats, respectively, and a regression slope < 1, even for studies that focused on herbivores and pollinators.

Four possible reasons for this apparent lack of explanatory power are (1) geographic variation in the ratio of herbivores to plants and of non-herbivores to herbivores, due to coevolutionary and ecological interactions amongst plants, herbivores and their predators (Hamilton *et al.* 2013); (2) variation across plant species in their geographic ranges, which is positively correlated with total insect richness (Condon *et al.* 2008); (3) correlations and linear regressions being inappropriate models; and (4) incomplete taxon sampling (Lewinsohn & Roslin 2008). The last explanation is straightforward to test at least in principle. For instance,

although Pellissier *et al.* (2013) successfully demonstrated a correlation between phylogenetic

beta diversities of plant and butterfly communities, they also found that plant phylogenetic

90   alpha diversity did not explain butterfly phylogenetic alpha diversity. One reason was that

some of the local plant taxa were not consumed by Lepidoptera and therefore contributed to

variance in plant alpha diversity but not to explanatory power. Presumably, those plant

species are consumed by other insect clades, and a taxonomically more complete sample

might have uncovered a positive relationship between plant and insect alpha diversity.

95   Thus, in a groundbreaking study involving 102 investigators and 129 494 arthropod

specimens collected in twelve 0.04-ha quadrats of tropical forest (0.48 ha total), Basset *et al.*

(2012) demonstrated that local tree species richness could predict the local species richness of

both herbivore and non-herbivore arthropod taxa exceptionally well. For each of their

eighteen taxon datasets (corresponding to ordinal or sub-ordinal guilds), Basset *et al.* (2012)

100   used tree-species data from the 0.48 ha of sampling effort to extrapolate total arthropod

species richness for the entire 6000-ha reserve. Overall, they found that what they called their

"plant models," which were parameterized species-accumulation curves that predicted the

accumulation of arthropod species from the accumulation of tree species, were consistently

able to predict "to a precision of 1%" independently derived best estimates of total arthropod

105   species richness for the entire 6000-ha reserve.

In summary, Basset *et al.* (2012) showed that, given comprehensive taxon sampling and a

more sophisticated statistical approach than correlations, plant species richness could indeed

accurately predict arthropod species richness. However, due to their unprecedentedly huge

sampling and taxonomic effort, Basset *et al.*'s (2012) protocol is effectively unrepeatable (and

110   itself was unavoidably limited to a tiny area [0.48 ha]), but it would be highly desirable to be

able to repeat this protocol efficiently in large numbers and over large spatial scales, i.e. to

'operationalize' the approach. At larger spatial scales (i.e. within and across landscapes), additional determinants of community structure can start to contribute, such as variation in environmental conditions and variation in regional species pools (Castagneyrol & Jactel

115   2012). It is at these larger spatial scales that plant community data would be most valuable in management for acting as a surrogate for arthropod diversity (and vice versa).

Metabarcoding is emerging as a promising way of advancing biodiversity research (Taberlet *et al.* 2012; Cristescu 2014). In metabarcoding, bulk samples of eukaryotes or environmental DNA are extracted, amplified, and sequenced for one or more taxonomically

120   informative genes (DNA 'barcodes') (Taberlet *et al.* 2012; Yu *et al.* 2012; Ji *et al.* 2013; Cristescu 2014). Most importantly, despite false negatives (species failing to be detected) and false positives (falsely present species) being found in metabarcoding, due to primer bias (Yu *et al.* 2012; Clarke *et al.* 2014; Deagle *et al.* 2014; Piñol *et al.* 2015) and other errors in the metabarcoding pipeline (sequence errors and chimeras from PCR, library prep, and/or

125   pyrosequencing, and species lumping and splitting in OTU clustering and taxonomic assignment), species richness and composition estimates from metabarcoded arthropod samples have been shown to correlate well with estimates calculated from standard morphological identification, even when the focal taxa are different (Yu *et al.* 2012; Ji *et al.* 2013; Edwards *et al.* 2014).

130   We therefore used metabarcoding to scale up Basset *et al.*'s (2012) approach, and we asked if plant diversity can predict insect diversity at landscape scales. Specifically, 1) does plant species richness predict insect species richness (alpha diversity); 2) does plant community composition predict insect community composition (beta diversity); and 3) is the predictive power of the plant model consistent across insect orders and over different seasons?

135   We report here that plant models parameterized with metabarcoding data produce

landscape-scale estimates of total insect species richness that are very close to independent

non-parametric estimates of insect species richness, and we also find a high degree of

correlation between insect and plant community compositions in two widely separated

tropical landscapes. As a result, we conclude that, armed with high-throughput methods, it

140   should indeed be possible to operationalize Basset *et al.*'s (2012) important result that plant

diversity can accurately predict arthropod diversity.

One potential benefit is that modern remote-sensing technologies, which show increasing

promise at efficient assessment of vegetation (Féret & Asner 2014; Baldeck *et al.* 2015),

might now also make possible the efficient management of a large proportion of animal

145   biodiversity. Another benefit, and perhaps the most important one, is that it should now be

possible to conduct highly replicated tests of the major competing explanations for tropical

megadiversity (Lewinsohn & Roslin 2008), and we explain this in detail in the Discussion.

Materials and Methods

Study sites

150   We conducted our study in two montane landscapes in tropical southern China (Fig. 1), which

differ in the nature of environmental heterogeneity they encompass and provide contrasting

case studies of the relationship between plant and insect diversity at landscape scales.

    Yinggeling Nature Reserve is located in central Hainan province (UTM/WGS84: 49N

328731 E, 2102468 N), a land-bridge island, and is the largest nature reserve in Hainan with

155   an area of > 500 km^2. The elevation ranges from 180 m to 1812 m, and the annual mean

temperature correspondingly ranges between 24°C to 20°C. Mean annual rainfall is 1800–

2700 mm. The principal vegetation types are tropical montane rainforest and tropical montane

evergreen broadleaf forest (Lin *et al.* 2013). Over 64% of the vegetation in the reserve is in a

near-pristine state, although many of the carnivores have been extirpated (Lau *et al.* 2010).

160   Mengsong (UTM/WGS84: 47N 656355 E, 2377646 N) is a sub-catchment of the upper

Mekong River, with an area of ~100 km^2. The elevation ranges from 800 m to 2000 m.

Mengsong has a subtropical climate influenced by the Indian monsoon. The annual mean

temperature is 18°C (at 1600 m asl). Mean annual rainfall varies between 1600–1800 mm, 80%

of which falls in May–October. Mengsong has a > 200-year history of occupation by

165   indigenous farmers, who formerly practiced swidden agriculture (Xu *et al.* 2009). Hence,

today the landscape is a mosaic of mature forest with a history of selective cutting, forest that

has naturally regenerated from clearance, and currently open land, such as terrace tea fields

and grasslands. The principal primary vegetation types are seasonal montane rain forest in

valleys, which grades into tropical montane evergreen broadleaf forest on upper slopes and

170   ridges (Zhu *et al.* 2005). Part of Mengsong was included in Bulong Nature Reserve

established in 2009. As with Yinggeling, many of the larger vertebrates have been extirpated

(Sreekar *et al.* 2015).

Biodiversity sampling

Yinggeling. - Twenty-nine 50×50 m plots were set up in Yinggeling in May 2009 (10 plots)

175   and September 2011 (19 plots) (Fig. 1). The plot locations were selected from a satellite

image to incorporate as much of the substantial topographic variation found within the nature

reserve as logistically possible. However, plot locations were not strictly randomly chosen.

Plots established in 2009 were clustered, so for our study, only one plot was selected

randomly from each cluster to minimize pseudo-replication. In total, 21 plots in Yinggeling

180   were included. All trees ≥ 5 cm DBH ('diameter at breast height,' which is set at 1.3 m from

the soil surface) in each plot were surveyed. Species name, DBH, height and crown width

were recorded. Field identifications were conducted by local experts, and trees that were not identified to the species level were excluded from further analyses.

Insects were sampled in the wet season (September to November 2011) using a Malaise

185   trap located at the center of each plot for an average of 16 days (range: 12-25) depending on the weather, which affects capture efficiency. Malaise traps are widely used in sampling insect diversity, and are designed to capture flying insects that escape upwards (mainly Diptera and Hymenoptera), but are generally not suitable for Coleoptera, which drop when they hit a barrier (review in Russo *et al.* 2011). The collecting bottles on the Malaise traps were filled

190   with 99.9% ethanol. Upon collection, the contents of each bottle were sieved to remove ethanol and placed in a storage bottle with fresh 99.9% ethanol. Between samples, the sieve and other equipment were rinsed with water and ethanol-flamed to prevent DNA cross-contamination.

Mengsong. - Twenty-eight 100×100 m plots were set up from April 2010 to May 2011,

195   based on a stratified random sampling design described in Paudel *et al.* (2015) (Fig. 1). Plots covered a gradient from heavily disturbed shrubland and grassland (*n* = 6), through regenerating forest (*n* = 12) to mature forest (*n* = 10). Each plot consisted of nine 10-m radius subplots arranged on a square grid with 50 m spacing (Beckschäfer *et al.* 2014). All trees, bamboos, and lianas with ≥ 10 cm DBH were recorded within a 10-m radius of the subplot

200   center, and all trees, bamboos, and lianas with 2-10 cm DBH were recorded within a 5-m radius of the subplot center. Species name, DBH, distance and angle to the subplot center were recorded. All herbs, ferns, and woody seedlings with < 2 cm DBH were surveyed within 1-m radius of the subplot center using a Braun-Blanquet coverage estimator (total coverage for each species was estimated visually and recorded using cover-abundance scale within six

205   cover classes). Vouchers of every species in each plot were collected, and field identifications

were confirmed (or adjusted) based on comparison to herbarium material at the

Xishuangbanna Tropical Botanical Garden (HITBC). The vouchers were later deposited at the

Kunming Institute of Botany.

    Insects were collected with Malaise traps in five subplots (four corners and the middle

210   subplot) over six days at the end of the wet season (late September-December 2010, hereafter

wet season) and at the end of the dry season (April-June 2011, hereafter dry season). The

collection and laboratory processing protocol were same as for Yinggeling. Subplot samples

were pooled within each plot for further analyses.

DNA extraction, PCR amplification, pyrosequencing, and bioinformatic analysis

215   Samples were prepared by using one leg from all specimens equal to or larger than a large fly

(~5 mm length) and whole bodies of everything smaller, added with 4 ml Qiagen ATL buffer

(Hilden, Germany) (20 mg/ml proteinase $k = 9 : 1$) per 1 g of sample, homogenized with

sterile 0.25-inch ceramic spheres in a FastPrep-24® system (MP Biomedicals, Santa Ana, CA,

USA) set on 5 m/s for 1 min at room temperature, and incubated overnight at 56 °C. The

220   genomic DNA was extracted with the Qiagen DNeasy Blood & Tissue Kit from 10% of the

lysed solution, with ≤ 900 μL per spin column, and quality-checked using the Nanodrop 2000

spectrophotometer (Thermo Fisher Scientific, Wilmington, DE, USA). DNA was PCR

amplified for the standard mtCOI barcode region using the degenerate primers, *Fol-degen-for*

5′-TCNACNAAYCAYAARRAYATYGG-3′ and *Fol-degen-rev* 5′-

225   TANACYTCNGGRTGNCCRAARAAYCA-3′ (Yu *et al.* 2012). The standard Roche A-

adaptor and a unique 10 bp MID (Multiplex IDentifier) tag for each sample were attached to

the forward primer. PCRs were performed in 20 μL reaction volumes containing 2 μL of 10 ×

buffer, 1.5 mM MgCl$_2$, 0.2 mM dNTPs, 0.4 μM each primer, 0.6 U HotStart Taq DNA

polymerase (TaKaRa Biosystems, Ohtsu, Japan), and ~60 ng of genomic DNA. We used a

230 touchdown thermocycling profile of 95 °C for 2 min; 11 cycles of 95 °C for 15 s; 51 °C for 30

s; 72 °C for 3 min, decreasing the annealing temperature by 1 degree every cycle; then 17

cycles of 95 °C for 15 s, 41 °C for 30 s, 72 °C for 3 min and a final extension of 72 °C for 10

min. We used non-proofreading Taq and fewer, longer cycles to reduce chimera production

(Lenz & Becker 2008; Yu *et al.* 2012). DNA from each sample was amplified in three

235 independent reactions and pooled to reduce amplification stochasticity. A negative control

was included for each sample during PCR runs to detect contamination. For pyrosequencing,

PCR products were gel-purified by using a Qiagen QIAquick PCR purification kit, quantified

using the Quant-iT PicoGreen dsDNA Assay kit (Invitrogen, Grand Island, New York, USA),

pooled and A-amplicon-sequenced on a Roche GS FLX at the Kunming Institute of Zoology.

240 The 21 Yinggeling samples were sequenced on two 1/8 regions (one 1/8 region shared with

other samples). The 28 Mengsong samples were sequenced on one whole run (four 1/4

regions, late September-December 2010: wet season) and two 1/4 regions (April-June 2011:

dry season), respectively.

We followed an experimentally validated bioinformatic pipeline (Yu *et al.* 2012; Ji *et al.*

245 2013) to denoise, deconvolute, and cluster the reads into 97%-similarity Operational

Taxonomic Units (OTUs). *Quality control*: Header sequences and low-quality reads were

removed from the raw output in the QIIME 1.5.0 environment (split_libraries.py: -l 100 -L

700 -H 9 -M 2 -b 10) (Caporaso *et al.* 2010b). We removed any sequences < 100 bp.

Denoising, deconvoluting and chimera removal: PyNAST 1.1 (Caporaso *et al.* 2010a) was

250 used to align reads against a high-quality, aligned data set of Arthropoda sequences (Yu *et al.*

2012), and sequences that failed to align at ≥ 60% similarity were removed. The remaining

sequences were clustered at 99% similarity with USEARCH 5.2.236 (Edgar 2010), a

consensus sequence was chosen for each cluster, and the UCHIME function was used to

perform *de novo* chimera detection and removal. A clustering step is required for chimera

255   detection because chimeric reads are expected to be rare and thus belong to small clusters only.

The final denoising step used MACSE 0.8b2 (Ranwez *et al.* 2011), which aligns at the amino

acid level to high-quality reference sequences and uses any stop codons in COI to infer

frameshift mutations caused by homopolymers. *OTU-picking and Taxonomic assignment*:

Sequences were clustered at 97% similarity using CROP 1.33 (Hao *et al.* 2011). Each cluster

260   of sequences represents a set of COI reads that are more similar to each other than to any

other cluster, and is called an operational taxonomic unit (OTU), which should approximate

or somewhat underestimate biological species. OTUs were assigned taxonomies using SAP

1.0.12 (Munch *et al.* 2008), keeping only taxonomic levels for which the posterior probability

was $\geq$ 80%. OTUs containing only one read (which tend to be PCR or sequencing errors and

265   are uncertain presences [Yu *et al*. 2012; Ficetola *et al.* 2015]) or assigned to non-Arthropoda

taxa were discarded.

Statistical analysis

Analyses were mostly performed using R 3.2.2 (R Core Team 2015) and packages *BAT* 1.3.1

(Cardoso *et al.* 2015) and *vegan* 2.3-0 (Oksanen *et al.* 2015). We converted metabarcoding

270   read numbers to presence/absence data before analyses, because read numbers are unlikely to

reflect biomass or abundance (Yu *et al.* 2012). We first analyzed all Insecta-assigned OTUs

together and then separately analyzed each Insecta order with $\geq$ 50 OTUs, including

Coleoptera, Diptera, Hemiptera, Hymenoptera, Lepidoptera, and Psocoptera (the last for

Mengsong only). More than 90% of OTUs were identified to order rank. We did not conduct

275   analysis at family level, since less than half of the OTUs were identified to this rank. We

tentatively included Arachnida from Mengsong (n = 84 and 83 OTUs for wet and dry season,

respectively) in our analyses of species richness, although they are a by-catch of Malaise traps

and hence may have a more stochastic pattern.

Species richness

280 To test whether plant species richness can predict insect species richness estimated from

metabarcoded Malaise-trap samples (i.e. Insecta OTU richness), we first calculated Pearson's

correlations (*cor.test* function) to allow comparison with the wider literature (Castagneyrol &

Jactel 2012). We then applied the "plant model" approach of Basset *et al.* (2012), as follows:

First, the mean number of arthropod OTUs and of plant species found with each additional

285 vegetation sampling plot (i.e. rarefaction curves) (*specaccum* in *vegan*) were calculated. To

control for the order in which plots are added, we randomly subsampled the data without

replacement (method = "random", permutations = 9999).

Second, we used CurveExpert 1.4 (Hyams 2009; default settings, except maximum

iterations = 1000) to fit functions to the relationship between the mean number of arthropod

290 OTUs and the mean number of plant species found with each additional sampling plot.

Following Basset *et al.* (2012), we used AICc to choose the best function from three

candidates: linear, power, and Weibull functions, and called the best function the "plant

model." For comparison, we also chose the best function from a broader selection of 25

candidates, including ones used in other studies (Dengler 2009). These alternative candidates

295 included linear (including quadratic fit and 3^{rd} degree polynomial fit), exponential, power,

growth, sigmoidal and rational functions (Hyams 2009). We selected the top three functions

based on AICc. Since statistical models offering a good fit to the data do not necessarily result

in a robust species richness estimates (Basset *et al.* 2012), we again fitted the top three

functions to a random subset of data (20 out of 28 plots for Mengsong, 15 out of 21 plots for

300 Yinggeling) to check for robustness. Then we predicted the arthropod OTU richness at 297

tree species (the number of species in all the Yinggeling survey plots) or at 807 vascular plant

species (the number of species in all the Mengsong survey plots) with these newly

parameterized models, and we compared these predicted richnesses against the observed

arthropod OTU richnesses in our metabarcode datasets. The best function was the one with

305   the smallest absolute difference (similar to the "lowest error of extrapolation" in Dengler

2009). The results from three and 25 candidate functions proved similar, and we thus focused

on the results from the first approach, using three candidate functions.

Third, we extrapolated the best function (i.e. the "plant model") to the total plant species

richness in the landscape to generate the plant model's prediction of total arthropod species

310   richness. In Yinggeling, there are 603 tree species known from the total reserve (Zhang *et al.*

2013). In Mengsong, no information on total vascular plant species richness is available, so

we used non-parametric estimators to extrapolate from the plot data to total vascular plant

species richness (*alpha.accum* in *BAT*).

Fourth, we used non-parametric estimators to independently estimate total arthropod OTU

315   richness in the landscape directly from the arthropod dataset (*alpha.accum* in *BAT*), and we

compared this extrapolation with the prediction from the plant model ('same-site prediction').

Specifically, we calculated the explained variance (R^2) when fit to a y = x model, and also

calculated correlations (*cor.test*, method = "pearson") for insect orders.

Note that there exists no 'true' biodiversity dataset to test against. Basset *et al.* (2012) used

320   both statistical (best-fitting function with lowest error of extrapolation) and biological

arguments (relevant surveys in the world with large sampling efforts) to get their best

estimates of arthropod diversity. As no comparable surveys with metabarcoding techniques

are available, we necessarily used non-parametric estimators, choosing those (Jackknife1,

Jackknife2 and Chao) that have been shown to perform better than other estimators (Walther

325   & Moore 2005; Hortal *et al.* 2006). Non-parametric estimators use the species

abundance/occurrence relationships (e.g. the number of species occurring in only one or two

sites throughout the samples) to estimate the total number of species (Hortal *et al.* 2006). We

further applied a correction factor (*P*) for these non-parametric estimators to improve

performance under conditions of low sampling effort, which is usually the case in arthropod

330   surveys (Lopez *et al.* 2012). In Mengsong, the above approach was firstly applied to the

whole landscape, and then separately to forests (mature and regenerating forest) and open

lands. We also included only tree data to build the plant model in the Mengsong forests.

Finally, to evaluate the generality of our plant models, we used Yinggeling's plant model to

try to predict Mengsong insect diversity, and used Mengsong's plant model to try to predict

335   Yinggeling insect diversity ('cross-site prediction'). Yinggeling and Mengsong are good

candidates for such a test, as they are in the same zoogeographic region (Holt 2013) but are

far from each other (~1000 km). However, Yinggeling and Mengsong have different

landscape histories, and their vegetation was sampled differently. To maximize comparability,

we used only the Mengsong plots (*n* = 16) located within the forest of Bulong Nature Reserve

340   (~60 km^2) (Fig. 1) and only included trees ≥ 5 cm DBH in each plot.

Community composition

To test whether plant species compositions can predict insect species compositions, we could

use Mantel tests, Procrustes analysis, or co-correspondence analysis, with each approach

offering advantages and drawbacks (reviewed in Gioria *et al.* 2011). We elected to use

345   Procrustes analysis, because it is generally more powerful than Mantel tests and is more

widely used than co-correspondence analysis, facilitating comparison with other studies.

Procrustes analysis superimposes one ordination on top of another by minimizing the sum of

the squared distances between points from the first to the second ordination. The probability

of the fit is calculated by comparing the observed sum of squared distances against those from

350   a null distribution obtained by repeated Procrustes fitting of permuted data (Oksanen *et al.*

2015). We used a non-metric multidimensional scaling (NMDS) ordination (*metaMDS* in

vegan, distance = "jaccard") of community composition data as the input data matrices for the

Procrustes analyses (*protest* in *vegan*, symmetric = TRUE). Because the Procrustes analysis

requires an identical number of axes in both ordinations, we constrained the number of axes to

355   two (k = 2) for Yinggeling and four (k = 4) for Mengsong across all analyses. Initial

exploratory analyses found that two/four axes were optimal for most groups (low stress and

consistent results). Stress values ranged from 0.08 – 0.24. We also used these approaches to

compare variation in community compositions among insect orders and between the two

seasons in Mengsong.

360   **Results**

Species richness

The 21 Yinggeling samples produced 40 261 sequence reads, and the 28 Mengsong samples

produced 519 865 and 253 025 reads for wet season and dry season, respectively. After

bioinformatic processing, we obtained 1 995 Insecta OTUs from Yinggeling, and we obtained

365   2 946 Insecta OTUs from Mengsong, including 2 073 in the wet season and 2 215 in the dry

season samples. None of the PCR negative controls detected sample contamination.

All the best-fit 'plant models' were Weibull functions, except for Hymenoptera in the dry

season in Mengsong, which was a power function. All the 'plant models' exhibited very close

fits to the non-parametric estimates of total OTU richness for insects as a whole (Insecta) and

370   for individual orders (Coleoptera, Diptera, Hemiptera, Hymenoptera, Lepidoptera and

Psocoptera) in both Yinggeling (Figs. 2 and S1, all Pearson's r > 0.98) and Mengsong (Figs. 2

and S1, all Pearson's r > 0.99). Similar results were obtained when we analyzed forests and

open land separately in Mengsong (Figs. S2 and S3), and similar results were obtained when

we used only trees to build the plant models in Mengsong forests (Fig. S4). In contrast, and

375   consistent with the results compiled by Castagneyrol & Jactel (2012), simple Pearson

correlations between insect OTU richness and plant species richness at the survey plot level

were low (Yinggeling: all r ≤ 0.14; Mengsong: all r ≤ 0.29 for both wet and dry seasons).

    The cross-site plant-model predictions (Yinggeling plant model predicting Mengsong

Insecta OTU richness and vice versa) lay within an error of 2X for all Insecta and for three

380   orders (Coleoptera, Hemiptera, Hymenoptera), but not for Diptera and Lepidoptera (Fig. 3).

Given the observed scatter, all correlations were, not surprisingly, very low (Mengsong's wet-

season plant model predicting Yinggeling: all Pearson's r < 0.1; Yinggeling's plant model

predicting Mengsong's wet season: all Pearson's r ≤ 0.1. Similar results were obtained when

we used Mengsong's dry-season data, Fig. S5. In all analyses, we excluded the Insecta points

385   to avoid double counting.)

Community composition

Community compositions in Insecta and plants were highly correlated in both Yinggeling and

Mengsong (Fig. 4, Table 1). Correlations were reduced somewhat but were still high when we

considered insect orders separately, likely reflecting the smaller sample size at this taxonomic

390   level (Figs. S6 and S7, Table 1). High correlations were maintained even when we limited our

analyses to only forests in Mengsong (Table S1).

    In Mengsong, 61% of Insecta OTUs recorded in the wet season were also recorded in the

dry season. Interestingly, community compositions remained highly correlated between these

two seasons for all Insecta and for individual insect orders, except Lepidoptera and Psocoptera

395   (Table 1), showing that despite turnover across seasons, the different insect species

compositions contain the same 'ecological information,' meaning that they consistently reveal the persistent compositional differences between the different vegetation plots, which themselves reflect differences in microhabitats, food sources and histories.

Finally, community compositions were highly correlated between most pairs of insect
400   orders in both Yinggeling and Mengsong, with the exceptions of Lepidoptera and Hemiptera in Yinggeling and Lepidoptera and Psocoptera in the wet season of Mengsong (Table 1). This suggests that different insect orders also contain similar ecological information about habitat differences. Again, these results were upheld even when analyzing only forests in Mengsong (Table S1).

405   **Discussion**

Our study has demonstrated (1) a close fit between estimates of total insect species richness that have been derived from plant models and from non-parametric estimators (alpha diversity, Figs. 2 and S1) and (2) a high degree of correlation between insect communities and plant communities (beta diversity, Fig. 4). We replicated our results in two landscapes (Yinggeling
410   and Mengsong), in two seasons in one of the landscapes (wet and dry in Mengsong), and across multiple insect orders (Figs. S6 and S7). Furthermore, we have extended the plant-model approach from tropical America to tropical Asia, from a homogeneous forest of 60 km^2 to two heterogeneous, anthropogenically modified landscapes (~100-500 km^2), and from a labor-intensive dataset of morphologically identified specimens to an efficiently processed
415   dataset of metabarcoded samples.

Our results thus strongly support Basset *et al.*'s (2012) finding that plant species richness can be an <u>accurate</u> predictor of insect species richness in tropical forest, and we show that plant and insect species compositions are highly correlated. Also, given that the Mengsong

plant model was able to predict Arachnida species richness (Figs. 2 and S2), we find some

420 support for the broader conclusion that plant diversity can be an accurate predictor of

arthropod diversity. Of course, it will be necessary to carry out taxonomically more

comprehensive sampling to be able to support the last conclusion strongly.

We also found that plant models from one landscape could predict the species richnesses of

Coleoptera, Hemiptera, Hymenoptera, and all Insecta in another landscape to within an error

425 of 2X (cross-site predictions, Figs. 3 and S5). One clear reason for the error in the cross-site

predictions is that our sampling effort and design were different across sites, due to

unavoidable logistical constraints. For instance, in Mengsong, each quadrat was sampled with

five Malaise traps, but in Yinggeling, only one trap was used.

Regardless, it seems likely that different landscapes will have differently parameterized

430 plant models, and we argue below that this provides an opportunity for testing theories of

tropical megadiversity. Note also that for biodiversity management and conservation, precise

estimates of insect species richness are not necessary. It is already useful to know that plant

species diversity can indeed be used as a close surrogate for insect diversity and *vice versa*.

We propose that metabarcoded insect samples should be tested as a sensitive and perhaps

435 early indicator of forest degradation.

Malaise traps and metabarcoding

When resources are limited, which they always are, a feasible way to carry out arthropod

diversity surveys at large scales is to combine mass trapping (here, Malaise traps) with a high-

throughput taxonomic method (here, metabarcoding). Naturally, this places limitations on the

440 informational content of the resulting datasets. Any given trap type can collect only a portion

of total arthropod biodiversity (Russo *et al.* 2011), and the downstream processes of DNA

extraction, PCR amplification, high-throughput sequencing, and bioinformatic processing will

result in false negatives ('dropout species') and false positives ('artefactual species' created by PCR-induced sequence chimeras, and PCR, sequencing, and clustering errors) (Bohmann *et al.* 2014).

PCR primers and software pipelines have been developed to minimize these errors (here, Yu *et al.* 2012) but more important is to understand how to interpret metabarcoding outputs judiciously. Multiple studies (Yu *et al.* 2012; Ji *et al.* 2013; Yang *et al.* 2014) have shown using both mock and real biodiversity samples that, despite false negatives and positives, metabarcoding datasets are nonetheless reliable for estimating *community-level* metrics of alpha and beta diversity. In other words, the degree to which arthropod samples (and thus locations) differ in species richness and composition can be quantified with metabarcoding, which is precisely the requirement of our study. We were thus able to recapitulate Basset *et al.* (2012) in finding that Weibull-function plant models accurately predict insect communities.

However, we cannot directly compare the parameter values of our plant models with Basset *et al.*'s (2012) models for two reasons. Most importantly, we used only Malaise traps, which are designed to capture flying insects that escape upwards, whereas many beetles and nonflying species are missed. Basset *et al.*'s (2012) collections were more comprehensive. Less importantly, our species concept is based on COI sequence similarity, which will differ somewhat (but not hugely) from morphological concepts in the Arthropoda (e.g. Schmidt *et al.* 2015). In any event, the use of DNA barcodes as a major input to species delimitation is now mainstream (Ratnasingham & Hebert 2013; Riedel *et al.* 2013; Tang *et al.* 2014), and barcodes are advantageous because they more efficiently reveal cryptic species (Condon *et al.* 2008).

465   *Explaining tropical herbivore megadiversity*

Lewinsohn and Roslin (2008) partitioned the causes of tropical herbivore megadiversity into four components: (A) more host plant species in the tropics combined with some level of host specificity, (B) more arthropod species per tropical plant species, (C) higher host specificity of tropical herbivores, and (D) higher rates of species turnover (beta diversity) within the same

470   host species in the tropics. Studies by Novotny et al. (2002, 2006, 2007) in Papua New Guinea, with temperate-zone contrasts in Central Europe, have supported component A (more host plant species) over the other three components, whereas a compilation of feeding experiments by Dyer et al. (2007) has supported component C: higher host specificity in tropical species. Two important observations made by Dyer et al. (2007) are that broad host range in the

475   temperate-zone is more obvious when more host plant species are surveyed and that different host plant species in the tropics show higher levels of insect community differentiation than do different temperate-zone host plants, suggesting higher host specificity in tropical insects.

Given our results here and elsewhere that metabarcoding can deliver reliable metrics of arthropod communities (Yu et al. 2012; Ji et al. 2013; Edwards et al. 2014; Yang et al. 2014),

480   we suggest that metabarcoding can now be used to carry out the large numbers of surveys needed to test the four competing (and perhaps coexisting) explanations of Lewinsohn and Roslin (2008). Components A (more tropical plant species) and B (more tropical arthropod species per plant species) can be differentiated by parameterizing plant models along a tropical to temperate gradient. A is self-evidently true, but if B is important then we should

485   observe a steeper slope of the plant model in the tropics. Following Dyer et al. (2007), components C (tropical insects having narrower host ranges) and D (more rapid spatial turnover in tropical insects) can be tested and differentiated by the relative contributions to beta diversity of changing host plant species and spanning geographic distance, in tropical and

490 temperate habitats. Although in many parts of the world, DNA-barcode databases are not yet sufficiently comprehensive to be able to identify most insects to species level, it should be possible to use a combination of sequence matching (Ratnasingham & Hebert 2007) and phylogenetic placement (Matsen et al. 2010; Berger et al. 2011) to identify most specimens to at least family level, allowing differentiation of herbivores from non-herbivores in the near future.

Acknowledgments

495 We are very grateful to the three reviewers and to the editor for comments and suggestions. We thank Mei Long for field assistance, Zhaoli Ding for sequencing, Philip Beckschäfer for providing the Mengsong GIS layer, and Mengsong village, Bulong Nature Reserve and Hainan Yinggeling National Nature Reserve for field support. Funding for fieldwork in Mengsong was provided by GIZ/BMZ (Project Nos. 08.7860.3-001.00, 13.1432.7-001.00) on behalf of the Government of the Federal Republic of Germany. DWY was supported by Yunnan Province (20080A001), the Chinese Academy of Sciences (0902281081, KSCX2-YW-Z-1027), the National Natural Science Foundation of China (31170498), the Ministry of Science and Technology of China (2012FY110800), the University of East Anglia and its GRACE computing cluster, and the State Key Laboratory of Genetic Resources and Evolution at the Kunming Institute of Zoology.

References

Baldeck CA, Asner GP, Martin RE *et al.* (2015) Operational tree species mapping in a diverse tropical forest with airborne imaging spectroscopy. *Plos ONE*, **10**, e0118403.

Basset Y, Cizek L, Cuénoud P *et al.* (2012) Arthropod diversity in a tropical forest. *Science*, **338**, 1481–1484.

Beckschäfer, P., Mundhenk, P., Kleinn, C., Ji, Y., Yu, D.W. & Harrison, R.D. (2013) Enhanced structural complexity index: an improved index for describing forest structural complexity. *Open Journal of Forestry*, **3**, 23–29.

Berger SA, Krompass D, Stamatakis A (2011) Performance, accuracy, and web server for evolutionary placement of short sequence reads under maximum likelihood. *Systematic Biology*, **60**, 291–302.

Bohmann K, Evans A, Gilbert MTP *et al.* (2014) Environmental DNA for wildlife biology and biodiversity monitoring. *Trends in Ecology & Evolution*, **29**, 358–367.

Bustamante MMC, Roitman I, Aide TM *et al.* (2015) Towards an integrated monitoring framework to assess the effects of tropical forest degradation and recovery on carbon stocks and biodiversity. *Global Change Biology*. doi: 451 10.1111/gcb.13087.

Caporaso JG, Bittinger K, Bushman FD *et al.* (2010a) PyNAST: A flexible tool for aligning sequences to a template alignment. *Bioinformatics*, **26**, 266–267.

Caporaso JG, Kuczynski J, Stombaugh J *et al.* (2010b) QIIME allows analysis of high-throughput community sequencing data. *Nature methods*, **7**, 335–6.

Cardoso P, Rigal F, Carvalho JC (2015) BAT - Biodiversity Assessment Tools, an R package for the measurement and estimation of alpha and beta taxon, phylogenetic and functional diversity. *Methods in Ecology and Evolution*, **6**, 232–236.

Castagneyrol B, Jactel H (2012) Unraveling plant-animal diversity relationships: A meta-regression analysis. *Ecology*, **93**, 2115–2124.

Clarke LJ, Soubrier J, Weyrich LS, Cooper A (2014) Environmental metabarcodes for insects: in silico PCR reveals potential for taxonomic bias. *Molecular Ecology Resources*, **14**, 1160–1170.

Condon MA, Scheffer SJ, Lewis ML, Swensen SM (2008) Hidden neotropical diversity: greater than the sum of its parts. *Science*, **320**, 928–931.

Cristescu ME (2014) From barcoding single individuals to metabarcoding biological communities: towards an integrative approach to the study of global biodiversity. *Trends in Ecology & Evolution*, **29**, 566–571.

Cruaud A, Ronsted N, Chantarasuwan B *et al.* (2012) An Extreme case of plant-insect codiversification: Figs and fig-pollinating wasps. *Systematic Biology*, **61**, 1029–1047.

Deagle BE, Jarman SN, Coissac E, Taberlet P (2014) DNA metabarcoding and the cytochrome c oxidase subunit I marker: not a perfect match. *Biology Letters*, **10**, 20140562.

Dengler J (2009) Which function describes the species-area relationship best? A review and empirical evaluation. *Journal of Biogeography*, **36**, 728–744.

Dyer LA, Singer MS, Lill JT *et al.* (2007) Host specificity of Lepidoptera in tropical and temperate forests. *Nature*, **448**, 696–699.

Edgar RC (2010) Search and clustering orders of magnitude faster than BLAST. *Bioinformatics*, **26**, 2460–2461.

Edger PP, Heidel-fischer HM, Bekaert M *et al.* (2015) The butterfly plant arms-race escalated
by gene and genome duplications. *Proceedings of the National Academy of Sciences of*
the United States of America, **112**, 8362–8366.
Edwards DP, Magrach A, Woodcock P *et al.* (2014) Selective-logging and oil palm: multi-
taxon impacts, biodiversity indicators, and trade-offs for conservation planning.
Ecological Applications, **24**, 2029–2049.
Féret JB, Asner GP (2014) Mapping tropical forest canopy diversity using high-fidelity
imaging spectroscopy. *Ecological Applications*, **24**, 1289–1296.
Ficetola GF, Pansu J, Bonin A *et al.* (2015) Replication levels, false presences and the
estimation of the presence/absence from eDNA metabarcoding data. *Molecular Ecology*
Resources, **15**, 543–556.
Gioria M, Bacaro G, Feehan J (2011) Evaluating and interpreting cross-taxon congruence:
Potential pitfalls and solutions. *Acta Oecologica*, **37**, 187–194.
Haddad NM, Crutsinger GM, Gross K *et al.* (2009) Plant species loss decreases arthropod
diversity and shifts trophic structure. *Ecology Letters*, **12**, 1029–1039.
Hamilton AJ, Novotný V, Waters EK *et al.* (2013) Estimating global arthropod species
richness: refining probabilistic models using probability bounds analysis. *Oecologia*, **171**,
357–65.
Hao X, Jiang R, Chen T (2011) Clustering 16S rRNA for OTU prediction: A method of
unsupervised Bayesian clustering. *Bioinformatics*, **27**, 611–618.
Holt BG (2013) An update of Wallace's zoogeographic regions of the world. *Science*, **339**,
74–78.
Hortal J, Borges P a V, Gaspar C (2006) Evaluating the performance of species richness
estimators: Sensitivity to sample grain size. *Journal of Animal Ecology*, **75**, 274–287.
Hyams D (2009) CurveExpert: A curve fitting system for Windows. URL
http://www.curveexpert.net (accessed 15 Feb 2016).
Ji Y, Ashton L, Pedley SM *et al.* (2013) Reliable, verifiable and efficient monitoring of
biodiversity via metabarcoding. *Ecology Letters*, **16**, 1245–1257.
Lau MW-N, Fellows JR, Chan BPL (2010) Carnivores (Mammalia: Carnivora) in South
China: a status review with notes on the commercial trade. *Mammal Review*, **40**, 247–
292.
Lenz TL, Becker S (2008) Simple approach to reduce PCR artefact formation leads to reliable
genotyping of MHC and other highly polymorphic loci--implications for evolutionary
analysis. *Gene*, **427**, 117–123.
Lewinsohn TM, Roslin T (2008) Four ways towards tropical herbivore megadiversity.
Ecology Letters, **11**, 398–416.
Lin J, Ng S-C, Zhang M *et al.* (2013) Vegetation of Yinggeling Nature Reserve. In:
Biodiversity and conservation of Hainan Yinggeling Nature Reserve (eds Jiang H, Chan
BPL, Zhou Y, *et al.*), pp. 214–259. China Forestry Press, Beijing, China.
Lopez LCS, De Aguiar Fracasso MP, Mesquita DO, Palma ART, Riul P (2012) The
relationship between percentage of singletons and sampling effort: A new approach to
reduce the bias of richness estimates. *Ecological Indicators*, **14**, 164–169.
Matsen FA, Kodner RB, Armbrust EV (2010) pplacer: linear time maximum-likelihood and
Bayesian phylogenetic placement of sequences onto a fixed reference tree. *BMC*
Bioinformatics, **11**, 538.
Munch K, Boomsma W, Huelsenbeck JP, Willerslev E, Nielsen R (2008) Statistical
assignment of DNA sequences using Bayesian phylogenetics. *Systematic biology*, **57**,
750–7.

Novotny V, Basset Y (2005) Host specificity of insect herbivores in tropical forests.
Proceedings of the Royal Society. B, **272**, 1083–1090.
Novotny V, Basset Y, Miller SE *et al.* (2002) Low host specificity of herbivorous insects in a
tropical forest. *Nature*, **416**, 841–844.
Novotny V, Drozd P, Miller SE *et al.* (2006) Why are there so many species of herbivorous
insects in tropical rainforests? *Science*, **313**, 1115–8.
Novotny V, Miller SE, Hulcr J *et al.* (2007) Low beta diversity of herbivorous insects in
tropical forests. *Nature*, **448**, 692–697.
Oksanen J, Blanchet FG, Kindt R *et al.* (2015) vegan: Community Ecology Package. URL
http://CRAN.R-project.org/package=vegan.
Paudel E, Dossa GGO, Xu J, Harrison RD (2015) Litterfall and nutrient return along a
disturbance gradient in a tropical montane forest. *Forest Ecology and Management*, **353**,
97–106.
Pellissier L, Ndiribe C, Dubuis A *et al.* (2013) Turnover of plant lineages shapes herbivore
phylogenetic beta diversity along ecological gradients. *Ecology Letters*, **16**, 600–608.
Piñol J, Mir G, Gomez-Polo P, Agustí N (2015) Universal and blocking primer mismatches
limit the use of high-throughput DNA sequencing for the quantitative metabarcoding of
arthropods. *Molecular Ecology Resources*, **15**, 819–830.
R Core Team (2015) R: a language and environment for statistical computing. URL
https://www.R-project.org/.
Ranwez V, Harispe S, Delsuc F, Douzery EJP (2011) MACSE: Multiple Alignment of
Coding SEquences accounting for frameshifts and stop codons. *PLoS ONE*, **6**, e22594.
Ratnasingham S, Hebert PDN (2007) BOLD: The Barcode of Life Data System (http://www.
barcodinglife.org). *Molecular Ecology Notes*, **7**, 355–364.
Ratnasingham S, Hebert PDN (2013) A DNA-based registry for all animal species: The
Barcode Index Number (BIN) system. *PLoS ONE*, **8**, e66213.
Riedel A, Sagata K, Suhardjono YR, Tänzler R, Balke M (2013) Integrative taxonomy on the
fast track - towards more sustainability in biodiversity research. *Frontiers in Zoology*, **10**,
15.
Russo L, Stehouwer R, Heberling JM, Shea K (2011) The composite insect trap: an
innovative combination trap for biologically diverse sampling. *PLoS ONE*, **6**, e21079.
Schmidt S, Schmid-Egger C, Morinière J, Haszprunar G, Hebert PDN (2015) DNA barcoding
largely supports 250 years of classical taxonomy: identifications for Central European
bees (Hymenoptera, Apoidea partim). *Molecular Ecology Resources*, **15**, 985–1000.
Sreekar R, Zhang K, Xu J, Harrison RD (2015) Yet another empty forest: Consider the
conservation value of a recently established tropical nature reserve. *PLoS ONE*, **10**,
e0117920.
Stork NE, McBroom J, Gely C, Hamilton AJ (2015) New approaches narrow global species
estimates for beetles, insects, and terrestrial arthropods. *Proceedings of the National*
Academy of Sciences, **112**, 7519–7523.
Taberlet P, Coissac E, Pompanon F, Brochmann C, Willerslev E (2012) Towards next-
generation biodiversity assessment using DNA metabarcoding. *Molecular Ecology*, **21**,
2045–2050.
Tang CQ, Humphreys AM, Fontaneto D, Barraclough TG (2014) Effects of phylogenetic
reconstruction method on the robustness of species delimitation using single-locus data.
Methods in Ecology and Evolution, **5**, 1086–1094.
Thompson JN (1994). The coevolutionary process. University of Chicago Press, Chicago,
USA.

Walther BA, Moore JL (2005) The concepts of bias, precision and accuracy, and their use in
testing the performance of species richness estimators, with a literature review of
estimator performance. *Ecography*, **28**, 815–829.
Xu J, Lebel L, Sturgeon J (2009) Functional links between biodiversity, livelihoods, and
culture in a Hani swidden landscape in southwest China. *Ecology and Society*, **14**, 20.
Yang C, Wang X, Miller J a. *et al.* (2014) Using metabarcoding to ask if easily collected soil
and leaf-litter samples can be used as a general biodiversity indicator. *Ecological*
Indicators, **46**, 379–389.
Yu DW, Ji Y, Emerson BC *et al.* (2012) Biodiversity soup: metabarcoding of arthropods for
rapid biodiversity assessment and biomonitoring. *Methods in Ecology and Evolution*, **3**,
613–623.
Zhang R, Xing F, Ng S-C *et al.* (2013) Spermatophyte species diversity of Yinggeling Nature
Reserve. In: Biodiversity and conservation of Hainan Yinggeling Nature Reserve (eds
Jiang H, Chan BPL, Zhou Y, et al.), pp. 68–171. China Forestry Press, Beijing, China.
Zhu H, Shi JP, Zhao CJ (2005) Species composition, physiognomy and plant diversity of the
tropical montane evergreen broad-leaved forest in Southern Yunnan. *Biodiversity and*
Conservation, **14**, 2855–2870.
664

Data Accessibility

DNA sequences: Genbank's Short Read Archive (Accession numbers: SRP065001 and SRP065147) and Dryad doi: http://dx.doi.org/10.5061/dryad.37b53.

Bioinformatic script: Dryad doi: http://dx.doi.org/10.5061/dryad.37b53.

All input data sets to R and R script: Dryad doi: http://dx.doi.org/10.5061/dryad.37b53.

Conflicts of interest

DWY is a co-founder of a UK company, NatureMetrics, which provides DNA metabarcoding services to the private and public sector. All other authors declare no conflicts of interest.

Author Contributions

KZ, DWY, RDH, and HJ designed the study. KZ, RDH, SL, CXY, CYY and HW collected

data, led by RDH and SL. KZ, YJ, CXY, XW performed the molecular experiments. KZ and

YJ performed the bioinformatic analyses. KZ performed the statistical analyses and wrote the

first draft of the manuscript. DWY and RDH contributed substantially to revisions.

Table 1 Procrustes correlations among plant, Insecta, and individual insect order communities in (a) Yinggeling and (b) Mengsong (9999 permutations), with the input NMDS (non-metric multidimensional scaling) ordinations calculated from binary Jaccard dissimilarities (k = 2 axes used in Yinggeling, k = 4 in Mengsong). Correlations with plants are bolded. In Mengsong, where insects were sampled in two seasons, wet vs. dry-season Procrustes correlations are presented on the diagonal and are underlined, and the proportions of wet-season Operational Taxonomic Units (OTUs) that were also collected in the dry season are reported below as percentages.

(a) Yinggeling

	Coleoptera	Diptera	Hemiptera	Hymenoptera	Lepidoptera	**Trees (*n* = 297)**
Insecta (*n* = 1995)						**0.76*****
Coleoptera (*n* = 239)	0.54**	0.47*	0.63***	0.43*	**0.44***
Diptera (*n* = 848)		0.64***	0.57**	0.53**	**0.73*****
Hemiptera (*n* = 205)			0.45*	0.39	**0.60****
Hymenoptera (*n* = 163)				0.55**	**0.56***
Lepidoptera (*n* = 263)					**0.65*****

689   (b) Mengsong

		Insecta	Coleoptera	Diptera	Hemiptera	Hymenoptera	Lepidoptera	Psocoptera	Vascular plants (*n* = 807)
Insecta	wet (*n* = 2073)	0.75***							**0.78***
	dry (*n* = 2215)	60.6%							**0.75***
Coleoptera	wet (*n* = 375)		0.45*	0.60***	0.58***	0.55***	0.43*	0.52**	**0.53***
	dry (*n* = 603)		38.0%	0.75***	0.70***	0.63***	0.51***	0.60***	**0.63***
Diptera	wet (*n* = 414)			0.52***	0.67***	0.71***	0.63***	0.52***	**0.71***
	dry (*n* = 413)			70.9%	0.78***	0.61***	0.52***	0.50**	**0.65***
Hemiptera	wet (*n* = 435)				0.67***	0.58***	0.44*	0.50**	**0.70***
	dry (*n* = 370)				71.1%	0.61***	0.44*	0.57***	**0.77***
Hymenoptera	wet (*n* = 409)					0.68***	0.50**	0.57***	**0.64***
	dry (*n* = 360)					70.0%	0.51***	0.61***	**0.72***
Lepidoptera	wet (*n* = 78)						0.33	0.40	**0.46**
	dry (*n* = 95)						61.1%	0.47**	0.38
Psocoptera	wet (*n* = 71)							0.42	**0.55***
	dry (*n* = 63)							85.7%	**0.52***

690   * *p* < 0.05, ** *p* < 0.01, *** *p* < 0.001, non-significant values shown in gray, after adjustment for multiple tests (*p.adjust*, method = "fdr")

Figure 1 Inventory plots in Yinggeling ($n = 21$) and Mengsong ($n = 28$). Colors stand for different elevation categories (orange = < 600 m; blue = 600-800 m; purple = ≥ 800 m) in Yinggeling, and for different habitat types (red = mature forest; green = regenerating forest; yellow = open land) in Mengsong. The Mengsong area left of the dashed line is included in Bulong Nature Reserve.

Figure 4 Procrustes superimposition plots between plant and Insecta communities (9999 permutations), with the input of non-metric multidimensional scaling (NMDS) ordinations calculated from binary Jaccard dissimilarities (k = 2 axes used in Yinggeling, k = 4 in Mengsong). All Procrustes correlation coefficients (r) are significantly different from zero at $p < 0.001$ (Table 1). Each point is a census site; solid points indicate plant data, and hollow points insect data. Colors as in Figure 1.

Figure 3 Cross-site predictions. Scatterplot of plant-model estimates versus non-parametric estimates of total OTU richness. Symbols as in Figure 2. Plant models from one landscape were used to predict non-parametric estimates in the other landscape. Shown here are the Mengsong wet-season results. Mengsong dry-season results are similar and shown in Figure S5. The shaded area encompasses a two-fold difference between the two estimates (y = 0.5x to y = 2x). The plant-model functions were Weibull for all taxa in Yinggeling and Mengsong, except for Lepidoptera in Mengsong, which was a linear function.

Figure 2 Same-site predictions. Scatterplot of plant-model estimates versus non-parametric estimates of total OTU (Operational Taxonomic Unit) richness, for Arachnida (for Mengsong only), Insecta and insect orders that contained ≥ 50 OTUs. To quantify the goodness-of-fit of these two estimates, explained variances (R^2) for insect orders were calculated from a y = x model (dashed line). Circles, squares and triangles stand for P-corrected versions of the Jackknife1, Jackknife2 and Chao estimators, respectively. Different colors stand for different taxa; only taxa absent from Yinggeling are labeled in the Mengsong figures. The plant-model functions were Weibull for all taxa in Yinggeling and Mengsong, except for Hymenoptera (dry season) in Mengsong, which was a power function. Note breaks in the axes.

Plant diversity accurately predicts insect diversity in two tropical landscapes: Supporting Information

Kai Zhang, * † [1] Siliang Lin, ‡ [1] Yinqiu Ji, * Chenxue Yang, * Xiaoyang Wang, * † Chunyan Yang, * Hesheng Wang, § Haisheng Jiang, ‡ Rhett D. Harrison ¶ ** and Douglas W. Yu * ††

* State Key Laboratory of Genetic Resources and Evolution, Kunming Institute of Zoology, Chinese Academy of Sciences, Kunming, Yunnan 650223, China
† Kunming College of Life Science, University of the Chinese Academy of Sciences, Kunming 650204, China
‡ School of Life Sciences, South China Normal University, Guangzhou, Guangdong 510631, China
§ Hainan Yinggeling National Nature Reserve, Baisha, Hainan 572800, China
¶ World Agroforestry Center, East and Central Asia Regional Office, Kunming 650201, China
** Center for Mountain Ecosystem Studies (CMES), Kunming Institute of Botany, Chinese Academy of Sciences, Kunming 650201, China
†† School of Biological Sciences, University of East Anglia, Norwich Research Park, Norwich, Norfolk NR47TJ, UK

co-corresponding authors. DWY +86 871 68125438 (ofc), 86 871 65110887 (fax), dougwyu@gmail.com; RDH +86 18687136498 (ofc), 86 871 65223377 (fax), r.harrison@cgiar.org

[1] Shared first-authorship.

This PDF file includes:
Figs. S1 to S7
Table S1
NOTE: R Markdown output document (*Plants accurately predict insects_R Markdown output.html*) is not include here and presented independently.

Figure S1 Same-site prediction. Scatterplot of plant-model (25 candidate functions) estimates versus non-parametric estimates of total OTU richness. Symbols as in Figure 2. Note breaks in the axes.

Figure S2 Same-site predictions. Scatterplot of plant-model estimates versus non-parametric estimates of total OTU richness in the (a) forests and (b) open land in Mengsong. Symbols as in Figure 2. The plant-model functions for (a) forests were Weibull for all taxa except Hymenoptera (dry season), which was a power function. The plant-model functions for (b) open land were power for all taxa except Coleoptera (wet season), which was a linear function. Note breaks in the axes.

Figure S3 Same-site predictions. Scatterplot of plant-model (25 candidate functions) estimates versus non-parametric estimates of total OTU richness in the (a) forests and (b) open land in Mengsong. Symbols as in Figure 2. Note breaks in the axes.

Figure S4 Same-site predictions. Scatterplot of plant-model estimates versus non-parametric estimates of total OTU richness in the forests in Mengsong. Only tree data were included to build the plant model. Symbols as in Figure 2. The plant-model functions were Weibull for all taxa except Hymenoptera (dry season), which was a linear function. Note breaks in the axes.

Figure S5 Cross-site predictions. Scatterplot of plant-model estimates versus non-parametric estimates of total OTU richness. Symbols as in Figure 3. Shown here are the Mengsong dry-season results. The plant-model functions were Weibull for all taxa in both Yinggeling and Mengsong.

Figure S6 Procrustes superimposition plots between plant and insect order communities in Yinggeling. All Procrustes correlation coefficients (r) are significantly different from zero at $p < 0.05$ (Table 1). Symbols as in Figure 4.

a

Figure S7 Procrustes superimposition plots between plant and insect order communities in (a) wet and (b) dry seasons of Mengsong. All Procrustes correlation coefficients (r) are significantly different from zero at $p < 0.01$ for wet season and at $p < 0.001$ for dry season except for Lepidoptera (Table 1). Symbols as in Figure 4.

Table S1 Procrustes correlations among plant, Insecta, and insect order communities in the forests of Mengsong ($n = 22$) (9999 permutations), with the input of non-metric multidimensional scaling (NMDS) ordinations calculated from binary Jaccard dissimilarities (k = 4). The percentage calculations, and the bold and underlined statistics as in Table 1.

		Insecta	Coleoptera	Diptera	Hemiptera	Hymenoptera	Lepidoptera	Psocoptera	Vascular plants ($n = 747$)
Insecta	wet ($n = 1780$)	<u>0.68***</u>							**0.69***
	dry ($n = 1913$)	55.9%							**0.70***
Coleoptera	wet ($n = 323$)		<u>0.55**</u>	0.63***	0.59***	0.60***	0.47	0.50*	**0.52**
	dry ($n = 510$)		34.9%	0.76***	0.72***	0.71***	0.56***	0.67***	**0.68***
Diptera	wet ($n = 363$)			<u>0.50*</u>	0.65***	0.65***	0.58***	0.60***	**0.62***
	dry ($n = 381$)			64.8%	0.71***	0.68***	0.53**	0.62***	**0.65***
Hemiptera	wet ($n = 360$)				<u>0.53*</u>	0.58***	0.48*	0.52**	**0.60***
	dry ($n = 293$)				65.5%	0.54**	0.53**	0.65***	**0.73***
Hymenoptera	wet ($n = 360$)					<u>0.64***</u>	0.52**	0.65***	**0.58***
	dry ($n = 308$)					66.9%	0.52**	0.60***	**0.66***
Lepidoptera	wet ($n = 70$)						<u>0.46</u>	0.45	**0.48*
	dry ($n = 84$)						56.0%	0.50*	0.36
Psocoptera	wet ($n = 60$)							<u>0.44</u>	**0.52**
	dry ($n = 59$)							78.0%	**0.62***

* $p < 0.05$, ** $p < 0.01$, *** $p < 0.001$, non-significant values shown in gray, after adjustment for multiple tests (*p.adjust*, method = "fdr")