Plant diversity accurately predicts insect diversity in two tropical landscapes

Zhang, K.,*†[1] Lin, S.L., ‡[1] Ji, Y.Q.,* Yang, C.X.,* Wang, X.Y.,*† Yang, C.Y.,*

Wang, H.,§ Jiang, H.S.,‡ Harrison, R.D.¶** and Yu, D.W.*††

* State Key Laboratory of Genetic Resources and Evolution, Kunming Institute of Zoology, Chinese Academy

of Sciences, Kunming, Yunnan 650223, China

† Kunming College of Life Science, University of the Chinese Academy of Sciences, Kunming 650204, China

‡ School of Life Sciences, South China Normal University, Guangzhou, Guangdong 510631, China

§ Hainan Yinggeling National Nature Reserve, Baisha, Hainan 572800, China

¶ World Agroforestry Center, East and Central Asia Regional Office, Kunming 650201, China

** Center for Mountain Ecosystem Studies (CMES), Kunming Institute of Botany, Chinese Academy of

Sciences, Kunming 650201, China

†† School of Biological Sciences, University of East Anglia, Norwich Research Park, Norwich, Norfolk NR47TJ,

UK

Key-words: Arthropoda, biodiversity, biomonitoring, China, host specificity, insect-plant interactions,

species richness, surrogate species

co-corresponding authors. DWY +86 871 68125438 (ofc), 86 871 65110887 (fax),

dougwyu@gmail.com; RDH +86 18687136498 (ofc), 86 871 65223377 (fax), r.harrison@cgiar.org

Send proofs to DWY.

Running title: Plants accurately predict insects

[1] Shared first-authorship.

Abstract

Plant diversity surely determines arthropod diversity, but only moderate correlations between arthropod and plant species richness had been observed until Basset *et al.* (2012, Science 338: 1481-1484) finally undertook an unprecedentedly comprehensive sampling of a tropical forest and demonstrated that plant species richness could indeed accurately predict arthropod species richness. We now require a high-throughput pipeline to operationalize this result so that we can (1) test competing explanations for tropical arthropod megadiversity, (2) improve estimates of global eukaryotic species diversity, and (3) use plant and arthropod communities as efficient proxies for each other, thus improving the efficiency of conservation planning and of detecting forest degradation and recovery. We therefore applied metabarcoding to Malaise-trap samples across two tropical landscapes in China. We demonstrate that plant species richness can accurately predict arthropod (mostly insect) species richness and that plant and insect community compositions are highly correlated, even in landscapes that are large, heterogeneous, and anthropogenically modified. Finally, we review how metabarcoding makes feasible highly replicated tests of the major competing explanations for tropical megadiversity.

Introduction

The relationship between plant diversity and insect diversity is fundamental to ecology

because (1) it underpins global species estimates of arthropods based on plant diversity

(Condon *et al.* 2008; Hamilton *et al.* 2013; Stork *et al.* 2015); (2) it improves our

understanding of the drivers of arthropod diversity and assembly structure (Novotny *et al.*

2006; Lewinsohn & Roslin 2008; Pellissier *et al.* 2013); and (3) a strongly predictive

relationship could open the way to using plant community metrics as surrogates for arthropod

communities (and vice versa), thus improving the efficiency of efforts to conserve

biodiversity, and ecosystem functions and services (Castagneyrol & Jactel 2012). In

particular, arthropod species richness and community composition could serve as a sensitive

method for detecting and quantifying the degree of forest degradation and recovery (Ji *et al.*

2013; Edwards *et al.* 2014), which is especially needed for the monitoring and verification of

contracts to pay local populations and governments to protect and restore standing forest, also

known as PES (Payments for Environmental Services) and REDD+ schemes (Reduction in

Emissions from Deforestation and Degradation) (Bustamante *et al.* 2015).

A priori, plant diversity must surely predict insect diversity (Lewinsohn & Roslin 2008;

Haddad *et al.* 2009), because insects depend directly (via herbivory, pollination, and housing)

and indirectly (via consumption of herbivores) on plant species, and insect herbivores show

dietary specialization to subsets of plant taxa (Novotny & Basset 2005). In addition, plant and

insect coevolutionary interactions have driven the vast diversity of plant and insect species

today (Thompson 1994; Cruaud *et al.* 2012; Edger *et al.* 2015). (N.B. In practice, studies of

terrestrial arthropod diversity tend to focus on insects because insects make up the majority of

described arthropods and can be easier to sample. This study will also follow this practice.)

Not surprisingly, many papers have reported significant correlations between arthropod

(mostly insect) and plant beta and alpha diversities (reviews in Lewinsohn & Roslin 2008;

Castagneyrol & Jactel 2012; Pellissier *et al.* 2013). In particular, work by Novotny *et al.*

(2002, 2006, 2007) has strongly suggested that the primary driver of high species richness

among herbivorous insects in tropical forests is simply the greater number of plant species in

the tropics, rather than either higher levels of host specificity and beta diversity or more insect

species per area of foliage. In short, the local number of insect species should increase nearly

linearly with the local number of plant species, and with a slope greater than one, since each

plant species is associated with multiple herbivore and predator species (Castagneyrol &

Jactel 2012).

However, correlations between arthropod (mostly insect) and plant species-richness have

shown only moderate fit. Castagneyrol and Jactel's (2012) comprehensive meta-analysis

reported mean correlations of only 0.39 and 0.51 for studies in single habitats and across

multiple habitats, respectively, and a regression slope < 1, even for studies that focused on

herbivores and pollinators.

Four possible reasons for this apparent lack of explanatory power are (1) geographic

variation in the ratio of herbivores to plants and of non-herbivores to herbivores, due to

coevolutionary and ecological interactions amongst plants, herbivores and their predators

(Hamilton *et al.* 2013); (2) variation across plant species in their geographic ranges, which is

positively correlated with total insect richness (Condon *et al.* 2008); (3) correlations and

linear regressions being inappropriate models; and (4) incomplete taxon sampling (Lewinsohn

& Roslin 2008). The last explanation is straightforward to test. For instance, although

Pellissier *et al.* (2013) successfully demonstrated a correlation between phylogenetic beta

diversities of plant and butterfly communities, they also found that plant phylogenetic alpha

diversity did not explain butterfly phylogenetic alpha diversity. One reason was that some of

the local plant taxa were not consumed by Lepidoptera and therefore contributed to variance

in plant alpha diversity but not to explanatory power. Presumably, those plant species are

consumed by other insect clades, and a taxonomically more complete sample might have

uncovered a positive relationship between plant and insect alpha diversity.

Thus, in a groundbreaking study involving 102 investigators and 129 494 arthropod

specimens collected in twelve 0.04-ha quadrats of tropical forest (0.48 ha total), Basset *et al.*

(2012) demonstrated that local tree species richness could predict the local species richness of

both herbivore and non-herbivore arthropod taxa exceptionally well. For each of their

eighteen taxon datasets (corresponding to ordinal or sub-ordinal guilds), Basset *et al.* (2012)

used tree-species data from the 0.48 ha of sampling effort to extrapolate total arthropod

species richness for the entire 6000-ha reserve. Overall, they found that what they called their

"plant models," which were parameterized species-accumulation curves that predicted the

accumulation of arthropod species from the accumulation of tree species, were consistently

able to predict "to a precision of 1%" independently derived best estimates of total arthropod

species richness for the entire 6000-ha reserve.

In summary, Basset *et al.* (2012) showed that, given comprehensive taxon sampling and a

more sophisticated statistical approach than correlations, plant species richness could indeed

accurately predict arthropod species richness. However, due to their unprecedentedly huge

sampling and taxonomic effort, Basset *et al.*'s (2012) protocol is effectively unrepeatable (and

itself was unavoidably limited to a tiny area [0.48 ha]), but it would be highly desirable to be

able to repeat this protocol efficiently in large numbers and over large spatial scales, i.e. to

'operationalize' the approach. At larger spatial scales (i.e. within and across landscapes),

additional determinants of community structure can start to contribute, such as variation in

environmental conditions and variation in regional species pools (Castagneyrol & Jactel

2012). It is at these larger spatial scales that plant community data would be most valuable in

management for acting as a surrogate for arthropod diversity (and vice versa).

Metabarcoding is emerging as a promising way of advancing biodiversity research

(Taberlet *et al.* 2012; Cristescu 2014). In metabarcoding, bulk samples of eukaryotes or

environmental DNA are extracted, amplified, and sequenced for one or more taxonomically

informative genes (DNA 'barcodes') (Taberlet *et al.* 2012; Yu *et al.* 2012; Ji *et al.* 2013;

Cristescu 2014). Most importantly, despite false negatives (species failing to be detected) and

false positives (falsely present species) being found in metabarcoding, due to primer bias (Yu

et al. 2012; Clarke *et al.* 2014; Deagle *et al.* 2014; Piñol *et al.* 2015) and other errors in the

metabarcoding pipeline (sequence errors and chimeras from PCR, library prep, and/or

pyrosequencing, and species lumping and splitting in OTU clustering and taxonomic

assignment), species richness and composition estimates from metabarcoded arthropod

samples have been shown to correlate well with estimates calculated from standard

morphological identification, even when the focal taxa are different (Yu *et al.* 2012; Ji *et al.*

2013; Edwards *et al.* 2014).

We therefore used metabarcoding to scale up the Basset *et al.*'s (2012) approach, and we

asked if plant diversity can predict insect diversity at landscape scales. Specifically, 1) does

plant species richness predict insect species richness (alpha diversity); 2) does plant

community composition predict insect community composition (beta diversity); and 3) is the

predictive power of the plant model consistent across insect orders and over different

seasons?

We report here that plant models parameterized with metabarcoding data produce

landscape-scale estimates of total insect species richness that are very close to independent

non-parametric estimates of insect species richness (alpha diversity), and we also find a high

degree of correlation between insect and plant community compositions (beta diversity) in

two widely separated tropical landscapes. As a result, we conclude that, armed with high-

throughput methods, it should indeed be possible to operationalize Basset *et al.*'s (2012)

important result that plant diversity can accurately predict arthropod diversity.

One potential benefit is that modern remote-sensing technologies, which show increasing

promise at efficient assessment of plant community composition (Asner *et al.* 2014; Baldeck

et al. 2015), might now also make possible the efficient management of a large proportion of

animal biodiversity. Another benefit, and perhaps the most important one, is that it should

now be possible to conduct highly replicated tests of the major competing explanations for

tropical megadiversity (Lewinsohn & Roslin 2008), and we explain this in detail in the

Discussion.

Materials and Methods
146

Study sites
147

We conducted our study in two montane landscapes in tropical southern China (Fig. 1), which

differ in the nature of environmental heterogeneity they encompass and provide contrasting

case studies of the relationship between plant and insect diversity at landscape scales.

Yinggeling Nature Reserve is located in central Hainan province (UTM/WGS84: 49N

328731 E, 2102468 N), a land-bridge island, and is the largest nature reserve in Hainan with

an area of > 500 km^2. The elevation ranges from 180 m to 1812 m, and the annual mean

temperature correspondingly ranges between 24°C to 20°C. Mean annual rainfall is 1800–

2700 mm. The principal vegetation types are tropical montane rainforest and tropical montane

evergreen broadleaf forest (Lin *et al.* 2013). Over 64% of the vegetation in the reserve is in a

near-pristine state, although many of the carnivores have been extirpated (Lau *et al.* 2010).

Mengsong (UTM/WGS84: 47N 656355 E, 2377646 N) is a sub-catchment of the upper

Mekong River, with an area of ~100 km^2. The elevation ranges from 800 m to 2000 m.

Mengsong has a subtropical climate influenced by the Indian monsoon. The annual mean

temperature is 18°C (at 1600 m asl). Mean annual rainfall varies between 1600–1800 mm, 80%

of which falls in May–October. Mengsong has a > 200-year of occupation by indigenous

farmers, who formerly practiced swidden agriculture (Xu *et al.* 2009). Hence, today the

landscape is a mosaic of mature forest with a history of selective cutting, forest that has

naturally regenerated from clearance, and currently open land, such as terrace tea fields and

grasslands. The principal primary vegetation types are seasonal montane rain forest in valleys,

which grades into tropical montane evergreen broadleaf forest on upper slopes and ridges

(Zhu *et al.* 2005). Part of Mengsong was included in Bulong Nature Reserve established in

2009. As with Yinggeling, many of the larger vertebrates have been extirpated (Sreekar *et al.*

2015).

Biodiversity sampling

Yinggeling. - Twenty-nine 50×50 m plots were set up in Yinggeling in May 2009 (10 plots)

and September 2011 (19 plots) (Fig. 1). The plot locations were selected from a satellite

image to incorporate as much of the substantial topographic variation found within the nature

reserve as logistically possible. However, plot locations were not strictly randomly chosen.

Plots established in 2009 were clustered, so for our study, only one plot was selected

randomly from each cluster to minimize pseudo-replication. In total, 21 plots in Yinggeling

were included. All trees ≥ 5 cm DBH ('diameter at breast height,' which is set at 1.3 m from

the soil surface) in each plot were surveyed. Species name, DBH, height and crown width

were recorded. Field identifications were conducted by local experts.

Insects were sampled in the wet season (September to November 2011) using a Malaise

trap located at the center of each plot for an average of 16 days (range: 12-25) depending on

the weather, which affects capture efficiency. The collecting bottles on the Malaise traps were

filled with 99.9% ethanol. Upon collection, the contents of each bottle were sieved to remove

ethanol and placed in a storage bottle with fresh 99.9% ethanol. Between samples, the sieve

and other equipment were rinsed with water and ethanol-flamed to prevent DNA cross-

contamination.

Mengsong. - Twenty-eight 100×100 m plots were set up from April 2010 to May 2011,

based on a stratified random sampling design described in Paudel *et al.* (2015) (Fig. 1). Plots

covered a gradient from heavily disturbed shrubland and grassland (*n* = 6), through

regenerating forest (*n* = 12) to mature forest (*n* = 10). Each plot consisted of nine 10-m radius

subplots arranged on a square grid with 50 m spacing (Beckschäfer *et al.* 2014). All trees,

bamboos, and lianas with ≥ 10 cm DBH were recorded within a 10-m radius of the subplot

center, and all trees, bamboos, and lianas with 2-10 cm DBH were recorded within a 5-m

radius of the subplot center. Species name, DBH, distance and angle to the subplot center

were recorded. All herbs, ferns, and woody seedlings with < 2 cm DBH were surveyed within

1-m radius of the subplot center using a Braun-Blanquet coverage estimator (total coverage

for each species was estimated visually and recorded using cover-abundance scale within six

cover classes). Vouchers of every species in each plot were collected, and field identifications

were confirmed (or adjusted) based on comparison to herbarium material at the

Xishuangbanna Tropical Botanical Garden (HITBC). The vouchers were later deposited at the

Kunming Institute of Botany.

Insects were collected with Malaise traps in five subplots (four corners and the middle

subplot) over six days at the end of the wet season (November-December 2010, hereafter wet

season) and at the end of the dry season (May-June 2011, hereafter dry season). The

collection and laboratory processing protocol were same as for Yinggeling. Subplot samples

were pooled within each plot for further analyses.

DNA extraction, PCR amplification, pyrosequencing, and bioinformatic analysis

Samples were prepared by using one leg from all specimens equal to or larger than a large fly

(~5 mm length) and whole bodies of everything smaller, added with 4 ml Qiagen ATL buffer

(Hilden, Germany) (20 mg/ml proteinase $k = 9 : 1$) per 1 g of sample, homogenized with

sterile 0.25-inch ceramic spheres in a FastPrep-24[®] system (MP Biomedicals, Santa Ana, CA,

USA) set on 5 m/s for 1 min at room temperature, and incubated overnight at 56 °C. The

genomic DNA was extracted with the Qiagen DNeasy Blood & Tissue Kit, with ≤ 900 μL per

spin column, and quality-checked using the Nanodrop 2000 spectrophotometer (Thermo

Fisher Scientific, Wilmington, DE, USA). DNA was PCR amplified for the standard mtCOI

barcode region using the degenerate primers, *Fol-degen-for* 5′-

TCNACNAAYCAYAARRAYATYGG-3′ and *Fol-degen-rev* 5′-

TANACYTCNGGRTGNCCRAARAAYCA-3′ (Yu *et al.* 2012). The standard Roche A-

adaptor and a unique 10 bp MID (Multiplex IDentifier) tag for each sample were attached to

the forward primer. PCRs were performed in 20 μL reaction volumes containing 2 μL of 10 ×

buffer, 1.5 mM MgCl$_2$, 0.2 mM dNTPs, 0.4 μM each primer, 0.6 U HotStart Taq DNA

polymerase (TaKaRa Biosystems, Ohtsu, Japan), and ~60 ng of genomic DNA. We used a

touchdown thermocycling profile of 95 °C for 2 min; 11 cycles of 95 °C for 15 s; 51 °C for 30

s; 72 °C for 3 min, decreasing the annealing temperature by 1 degree every cycle; then 17

cycles of 95 °C for 15 s, 41 °C for 30 s, 72 °C for 3 min and a final extension of 72 °C for 10

min. We used non-proofreading Taq and fewer, longer cycles to reduce chimera production

(Lenz & Becker 2008; Yu *et al.* 2012). DNA from each sample was amplified in three

independent reactions and pooled to reduce amplification stochasticity. A negative control

was included for each sample during PCR runs to detect contamination. For pyrosequencing,

PCR products were gel-purified by using a Qiagen QIAquick PCR purification kit, quantified

using the Quant-iT PicoGreen dsDNA Assay kit (Invitrogen, Grand Island, New York, USA),

pooled and A-amplicon-sequenced on a Roche GS FLX at the Kunming Institute of Zoology.

The 21 Yinggeling samples were sequenced on two 1/8 regions (one 1/8 region shared with

other samples). The 28 Mengsong samples were sequenced on one whole run (four 1/4

regions, November-December 2010: wet season) and two 1/4 regions (May-June 2011: dry

season), respectively.

We followed an experimentally validated bioinformatic pipeline (Yu *et al.* 2012; Ji *et al.*

2013) to denoise, deconvolute, and cluster the reads into 97%-similarity Operational

Taxonomic Units (OTUs). *Quality control*: Header sequences and low-quality reads were

removed from the raw output in the QIIME 1.5.0 environment (split_libraries.py: -l 100 -L

700 -H 9 -M 2 -b 10) (Caporaso *et al.* 2010b). We removed any sequences < 100 bp.

Denoising, deconvoluting and chimera removal: PyNAST 1.1 (Caporaso *et al.* 2010a) was

used to align reads against a high-quality, aligned data set of Arthropoda sequences (Yu *et al.*

2012), and sequences that failed to align at ≥ 60% similarity were removed. The remaining

sequences were clustered at 99% similarity with USEARCH 5.2.236 (Edgar 2010), a

consensus sequence was chosen for each cluster, and the UCHIME function was used to

perform *de novo* chimera detection and removal. A clustering step is required for chimera

detection because chimeric reads are expected to be rare and thus belong to small clusters

only. The final denoising step used MACSE 0.8b2 (Ranwez *et al.* 2011), which aligns at the

amino acid level to high-quality reference sequences and uses any stop codons in COI to infer

frameshift mutations caused by homopolymers. *OTU-picking and Taxonomic assignment*:

Sequences were chain-clustered at 97% similarity using CROP 1.33 (Hao *et al.* 2011). Each

cluster of sequences represents a set of COI reads that are more similar to each other than to

any other cluster, and is called an operational taxonomic unit (OTU), which should

approximate or somewhat underestimate biological species. OTUs were assigned taxonomies

using SAP 1.0.12 (Munch *et al.* 2008), keeping only taxonomic levels for which the posterior

probability was ≥ 80%. OTUs containing only one read (which tend to be PCR or sequencing

errors and are uncertain presences [Ficetola *et al.* 2015]) or assigned to non-Arthropoda taxa

were discarded.

Statistical analysis

Analyses were mostly performed using R 3.2.2 (R Core Team 2015) and packages *BAT* 1.3.1

(Cardoso *et al.* 2015) and *vegan* 2.3-0 (Oksanen *et al.* 2015). We converted metabarcoding

read numbers to presence/absence data before analyses, because read numbers are unlikely to

reflect biomass or abundance (Yu *et al.* 2012). We first analyzed all Insecta-assigned OTUs

together and then separately analyzed each Insecta order with ≥ 50 OTUs, including

Coleoptera, Diptera, Hemiptera, Hymenoptera, Lepidoptera, and Psocoptera (the last for

Mengsong only). More than 90% of OTUs were identified to order rank. We did not conduct

analysis at family level, since less than half of the OTUs were identified to this rank. We

tentatively included Arachnida from Mengsong (n = 84 and 83 OTUs for wet and dry season,

respectively) in our analyses of species richness, although they are a by-catch of Malaise traps

and hence may have a more stochastic pattern.

Species richness

To test whether plant species richness can predict insect species richness estimated from

metabarcoded Malaise-trap samples (i.e. Insecta OTU richness), we first calculated Pearson's

correlations (*cor.test* function) to allow comparison with the wider literature (Castagneyrol &

Jactel 2012). We then applied the "plant model" approach of Basset *et al.* (2012), as follows:

First, the mean number of arthropod OTUs and of plant species found with each additional

vegetation sampling plot (i.e. rarefaction curves) (*specaccum* in *vegan*) were calculated. To

control for the order in which plots are added, we randomly subsampled the data without

replacement (method = "random", permutations = 9999).

Second, we used CurveExpert 1.4 (Hyams 2009; default settings, except maximum

iterations = 1000) to fit functions to the relationship between the mean number of arthropod

OTUs and the mean number of plant species found with each additional sampling plot.

Following Basset *et al.* (2012), we used AICc to choose the best function from three

candidates: linear, power and Weibull functions, and called the best function the "plant

model." For comparison, we also chose the best function from a broader selection of 25

candidates, including ones used in other studies (Dengler 2009). These alternative candidates

included linear (including quadratic fit and 3[rd] degree polynomial fit), exponential, power,

growth, sigmoidal and rational functions (Hyams 2009). We selected the top three functions

based on AICc. Since statistical models offering a good fit to the data do not necessarily result

in a robust species richness estimates (Basset *et al.* 2012), we again fitted the top three

functions to a random subset of data (20 out of 28 plots for Mengsong, 15 out of 21 plots for

Yinggeling) to check for robustness. Then we predicted the arthropod OTU richness at 297

tree species (the number of species in all the Yinggeling survey plots) or at 807 vascular plant

species (the number of species in all the Mengsong survey plots) with these newly

parameterized models, and we compared these predicted richnesses against the observed

arthropod OTU richnesses in our metabarcode datasets. The best function was the one with

the smallest absolute difference (similar to the "lowest error of extrapolation" in Dengler

2009). The results from three and 25 candidate functions proved similar, and we thus focused

on the results from the first approach, using three candidate functions.

Third, we extrapolated the best function (i.e. the "plant model") to the total plant species

richness in the landscape to generate the plant model's prediction of total arthropod species

richness. In Yinggeling, there are 603 tree species known from the total reserve (Lin *et al.*

2013). In Mengsong, no information on total vascular plant species richness is available, so

we used non-parametric estimators to extrapolate from the plot data to total vascular plant

species richness (*alpha.accum* in *BAT*).

Fourth, we used non-parametric estimators to independently estimate total arthropod OTU

richness in the landscape directly from the arthropod dataset (*alpha.accum* in *BAT*), and we

compared this extrapolation with the prediction from the plant model ('same-site prediction').

Specifically, we calculated the explained variance (R^2) when fit to a y = x model, and also

calculated correlations (*cor.test*, method = "pearson") for insect orders.

Note that there exists no 'true' biodiversity dataset to test against. Basset *et al.* (2012) used

both statistical (best-fitting function with lowest error of extrapolation) and biological

arguments (relevant surveys in the world with large sampling efforts) to get their best

estimates of arthropod diversity. As no comparable surveys with metabarcoding techniques

are available, we necessarily used non-parametric estimators, choosing those (Jackknife1,

Jackknife2 and Chao) that have been shown to perform better than other estimators (Walther

& Moore 2005; Hortal *et al.* 2006). Non-parametric estimators use the species

abundance/occurrence relationships (e.g. the number of species occurring in only one or two

sites throughout the samples) to estimate the total number of species (Hortal *et al.* 2006). We

further applied a correction factor (P) for these non-parametric estimators to improve

performance under conditions of low sampling effort, which is usually the case in arthropod

surveys (Lopez *et al.* 2012). In Mengsong, the above approach was firstly applied to the

whole landscape, and then separately to forests (mature and regenerating forest) and open

lands. We also included only tree data to build the plant model in the Mengsong forests.

Finally, to evaluate the generality of our plant models, we used Yinggeling's plant model to

try to predict Mengsong insect diversity, and used Mengsong's plant model to try to predict

Yinggeling insect diversity ('cross-site prediction'). Yinggeling and Mengsong are good

candidates for such a test, as they are in the same zoogeographic region (Holt 2013) but are

far from each other (~1000 km). However, Yinggeling and Mengsong have different

landscape histories, and their vegetation had been sampled differently. To maximize

comparability, we used only the Mengsong plots (n = 16) located within the forest of Bulong

Nature Reserve (~60 km^2) (Fig. 1) and only included trees $\geq$ 5 cm DBH in each plot.

Community composition

To test whether plant species compositions can predict insect species compositions, we could

use Mantel tests, Procrustes analysis, or co-correspondence analysis, with each approach

offering advantages and drawbacks (reviewed in Gioria *et al.* 2011). We elected to use

Procrustes analysis, because it is generally more powerful than Mantel tests and is more

widely used than co-correspondence analysis, facilitating comparison with other studies.

Procrustes analysis superimposes one ordination on top of another by minimizing the sum of

the squared distances between points from the first to the second ordination. The probability

of the fit is calculated by comparing the observed sum of squared distances against those from

a null distribution obtained by repeated Procrustes fitting of permuted data (Oksanen *et al.*

2015). We used a non-metric multidimensional scaling (NMDS) ordination (*metaMDS* in

vegan, distance = "jaccard") of community composition data as the input data matrices for the

Procrustes analyses (*protest* in *vegan*, symmetric = TRUE). Because the Procrustes analysis

requires an identical number of axes in both ordinations, we constrained the number of axes to

two (k = 2) for Yinggeling and four (k = 4) for Mengsong across all analyses. Initial

exploratory analyses found that two/four axes were optimal for most groups (low stress and

consistent results). Stress values ranged from $0.08 - 0.24$. We also used these approaches to

compare variation in community compositions among insect orders and between the two

seasons in Mengsong.

Results

Species richness

The 21 Yinggeling samples produced 40 261 sequence reads, and the 28 Mengsong samples

produced 519 865 and 253 025 reads for wet season and dry season, respectively. After

bioinformatic processing, we obtained 1 995 Insecta OTUs from Yinggeling, and we obtained

2 946 Insecta OTUs from Mengsong, including 2 073 in the wet season and 2 215 in the dry

season samples. None of the PCR negative controls detected sample contamination.

All the 'plant models' exhibited very close fits to the non-parametric estimates of total

OTU richness for insects as a whole (Insecta) and for individual orders (Coleoptera, Diptera,

Hemiptera, Hymenoptera, Lepidoptera and Psocoptera) in both Yinggeling (Figs. 2 and S1, all

Pearson's r > 0.98) and Mengsong (Figs. 2 and S1, all Pearson's r > 0.99). Similar results

were obtained when we analyzed forests and open land separately in Mengsong (Figs. S2 and

S3), and similar results were obtained when we used only trees to build the plant models in

Mengsong forests (Fig. S4). In contrast, and consistent with the results compiled by

Castagneyrol & Jactel (2012), simple Pearson correlations between insect OTU richness and

plant species richness at the survey plot level were low (Yinggeling: all r ≤ 0.14; Mengsong:

all r ≤ 0.29 for both wet and dry seasons).

The cross-site plant-model predictions (Yinggeling plant model predicting Mengsong

Insecta OTU richness and vice versa) lay within an error of 2X for all Insecta and for three

orders (Coleoptera, Hemiptera, Hymenoptera), but not for Diptera and Lepidoptera (Fig. 3).

Given the observed scatter, all correlations were, not surprisingly, very low (Mengsong's wet-

season plant model predicting Yinggeling: all Pearson's r < 0.1; Yinggeling's plant model

predicting Mengsong's wet season: all Pearson's r ≤ 0.1. Similar results were obtained when

we used Mengsong's dry-season data, Fig. S5. In all analyses, we excluded the Insecta points

to avoid double counting.)

Community composition

Community compositions in Insecta and plants were highly correlated in both Yinggeling and

Mengsong (Fig. 4, Table 1). Correlations were reduced somewhat but were still high when we

considered insect orders separately, likely reflecting the smaller sample size at this taxonomic

level (Figs. S6 and S7, Table 1). High correlations were maintained even when we limited our

analyses to only forests in Mengsong (Table S1).

In Mengsong, 61% of Insecta OTUs recorded in the wet season were also recorded in the

dry season. Interestingly, community compositions remained highly correlated between these

two seasons for all Insecta and for individual insect orders, except Lepidoptera and

Psocoptera (Table 1), showing that despite turnover across seasons, the different insect

species compositions contain the same 'ecological information,' meaning that they

consistently reveal the persistent compositional differences between the different vegetation

plots, which themselves reflect differences in microhabitats, food sources and histories.

Finally, community compositions were highly correlated between most pairs of insect

orders in both Yinggeling and Mengsong, with the exceptions of Lepidoptera and Hemiptera

in Yinggeling and Lepidoptera and Psocoptera in the wet season of Mengsong (Table 1). This

suggests that different insect orders also contain similar ecological information about habitat

differences. Again, these results were upheld even when analyzing only forests in Mengsong

(Table S1).

Discussion

Our study has demonstrated (1) a close fit between estimates of total insect species richness

that have been derived from plant models and from non-parametric estimators (Figs. 2 and

S1), and (2) a high degree of correlation between insect communities and plant communities

(Fig. 4). Moreover, we replicated our results in two landscapes (Yinggeling and Mengsong),

in two seasons in one of these landscapes (wet and dry in Mengsong), and across multiple

insect orders (Figs. S6 and S7).

Furthermore, we have extended the plant-model approach from tropical America to

tropical Asia, from a homogeneous forest of 60 km^2 to two heterogeneous, anthropogenically

modified landscapes (~100-500 km^2), and from a labor-intensive dataset of morphologically

identified specimens to an efficiently processed dataset of metabarcoded samples. We even

found that plant models from one landscape could predict the species richnesses of Coleoptera,

Hemiptera, Hymenoptera, and all Insecta in another landscape to within an error of 2X (cross-

site predictions, Figs. 3 and S5). However, cross-site predictions were quantitatively

inaccurate, suggesting that a general plant model (at least for our sampling protocol) does not

exist.

Our results thus strongly support Basset *et al.*'s (2012) finding that plant species richness

can be an <u>accurate</u> predictor of insect species richness in tropical forest, and we show that

plant and insect species compositions are highly correlated. Also, given that the Mengsong

plant model was able to predict Arachnida species richness (Figs. 2 and S2), we find some

support for the broader conclusion that plant diversity can be an accurate predictor of

arthropod diversity. Of course, it will be necessary to carry out taxonomically more

comprehensive sampling to be able to support the last conclusion strongly.

Malaise traps and metabarcoding

When resources are limited, which they always are, a feasible way to carry out arthropod

diversity surveys at large scales is to combine mass trapping (here, Malaise traps) with a high-

throughput taxonomic method (here, metabarcoding). Naturally, this places limitations on the

informational content of the resulting datasets. Any given trap type can collect only a portion

of total arthropod biodiversity, and the downstream processes of DNA extraction, PCR

amplification, high-throughput sequencing, and bioinformatic processing will result in false

negatives ('dropout species') and false positives ('artefactual species' created by PCR-

induced sequence chimeras, and PCR, sequencing, and clustering errors) (Bohmann *et al.*

2014).

PCR primers and software pipelines have been developed to minimize these errors (here,

Yu *et al.* 2012), but more important is to understand how to interpret metabarcoding outputs

judiciously. Multiple studies (Yu *et al.* 2012; Ji *et al.* 2013; Yang *et al.* 2014) have shown

using both mock and real biodiversity samples that, despite false negatives and positives,

metabarcoding datasets are nonetheless reliable for estimating *community-level* metrics of

alpha and beta diversity. In other words, the degree to which arthropod samples (and thus

locations) differ in species richness and composition can be quantified with metabarcoding,

which is precisely the requirement of our study. We were thus able to recapitulate Basset *et al.*

(2012) in finding that Weibull-function plant models accurately predict insect communities.

However, we cannot directly compare the parameter values of our plant models with

Basset *et al.*'s (2012) models for two reasons. Most importantly, we used only Malaise traps,

which are designed to capture flying insects that escape upwards (many beetles drop when

they hit a barrier and are thus not captured, and nonflying species are missed), whereas Basset

et al.'s (2012) collections were more comprehensive. Less importantly, our species concept is

based on COI sequence similarity, which will differ somewhat (but not hugely) from

morphological concepts in the Arthropoda (e.g. Schmidt *et al.* 2015). In any event, the use of

DNA barcodes as a major input to species delimitation is now mainstream (Ratnasingham &

Hebert 2013; Riedel *et al.* 2013; Tang *et al.* 2014), and barcodes are advantageous because

they more efficiently reveal cryptic species (Condon *et al.* 2008).

Explaining tropical herbivore megadiversity

Lewinsohn and Roslin (2008) partitioned the causes of tropical herbivore megadiversity into

four components: (A) more host plant species in the tropics combined with some level of host

specificity, (B) more arthropod species per tropical plant species, (C) higher host specificity

of tropical herbivores, and (D) higher rates of species turnover (beta diversity) within the

same host species in the tropics. Studies by Novotny *et al.* (2002, 2006, 2007) in Papua New

Guinea, with temperate-zone contrasts in Central Europe, have supported component A (more

host plant species) over the other three components, whereas a compilation of feeding

experiments by Dyer *et al.* (2007) has supported component C: higher host specificity in

tropical species. Two important observations made by Dyer *et al.* (2007) are that broad host

range in temperate-zone is more obvious when more hostplant species are surveyed, and that

different hostplant species in the tropics show higher levels of insect community

differentiation than do different temperate-zone hostplants, suggesting higher host specificity

in tropical insects.

Given our results here and elsewhere that metabarcoding can deliver reliable metrics of

arthropod communities (Yu *et al.* 2012; Ji *et al.* 2013; Edwards *et al.* 2014; Yang *et al.* 2014),

we suggest that metabarcoding can now be used to carry out the large numbers of surveys

needed to test the four competing (and perhaps coexisting) explanations of Lewinsohn and

Roslin (2008). Components A (more tropical plant species) and B (more tropical arthropod

species per plant species) can be differentiated by parameterizing plant models along a

tropical to temperate gradient. A is self-evidently true, but if B is important then we should

observe a steeper slope of the plant model in the tropics. Following Dyer *et al.* (2007),

components C (tropical insects having narrower host ranges) and D (more rapid spatial

turnover in tropical insects) can be tested and differentiated by the relative contributions to

beta diversity of changing hostplant species and spanning geographic distance, in tropical and

temperate habitats. Although in many parts of the world, DNA-barcode databases are not yet

sufficiently comprehensive to be able to identify most insects to species level, it should be

possible to use a combination of sequence matching (Ratnasingham & Hebert 2007) and

phylogenetic placement (Matsen *et al.* 2010; Berger *et al.* 2011) to identify most specimens to

at least family level, allowing differentiation of herbivores from non-herbivores in the near

future.

Acknowledgments

We are very grateful to the three reviewers and to the editor for comments and suggestions. We thank Mei Long for field assistance, Zhaoli Ding for sequencing, Philip Beckschäfer for providing Mengsong GIS layer, and Mengsong village, Bulong Nature Reserve and Hainan Yinggeling National Nature Reserve for field support. Funding for fieldwork in Mengsong was provided by GIZ/BMZ (Project Nos. 08.7860.3-001.00, 13.1432.7-001.00) on behalf of the Government of the Federal Republic of Germany. DWY was supported by Yunnan Province (20080A001), the Chinese Academy of Sciences (0902281081, KSCX2-YW-Z-1027), the National Natural Science Foundation of China (31170498), the Ministry of Science and Technology of China (2012FY110800), the University of East Anglia and its GRACE computing cluster, and the State Key Laboratory of Genetic Resources and Evolution at the Kunming Institute of Zoology.

References

Asner GP, Anderson CB, Martin RE *et al.* (2014) Landscape-scale changes in forest structure and functional traits along an Andes-to-Amazon elevation gradient. *Biogeosciences*, **11**, 843–856.

Baldeck CA, Asner GP, Martin RE *et al.* (2015) Operational tree species mapping in a diverse tropical forest with airborne imaging spectroscopy. *Plos One*, **10**, e0118403.

Basset Y, Cizek L, Cuénoud P *et al.* (2012) Arthropod diversity in a tropical forest. *Science*, **338**, 1481–1484.

Beckschäfer P, Fehrmann L, Harrison RD, Xu J, Kleinn C (2014) Mapping Leaf Area Index in subtropical upland ecosystems using RapidEye imagery and the randomForest algorithm. *iForest*, **7**, 1–11.

Berger SA, Krompass D, Stamatakis A (2011) Performance, accuracy, and web server for evolutionary placement of short sequence reads under maximum likelihood. *Systematic Biology*, **60**, 291–302.

Bohmann K, Evans A, Gilbert MTP *et al.* (2014) Environmental DNA for wildlife biology and biodiversity monitoring. *Trends in Ecology & Evolution*, **29**, 358–367.

Bustamante MMC, Roitman I, Aide TM *et al.* (2015) Towards an integrated monitoring framework to assess the effects of tropical forest degradation and recovery on carbon stocks and biodiversity. *Global Change Biology*. doi: 451 10.1111/gcb.13087.

Caporaso JG, Bittinger K, Bushman FD *et al.* (2010a) PyNAST: A flexible tool for aligning sequences to a template alignment. *Bioinformatics*, **26**, 266–267.

Caporaso JG, Kuczynski J, Stombaugh J *et al.* (2010b) QIIME allows analysis of high-throughput community sequencing data. *Nature methods*, **7**, 335–6.

Cardoso P, Rigal F, Carvalho JC (2015) BAT - Biodiversity Assessment Tools, an R package for the measurement and estimation of alpha and beta taxon, phylogenetic and functional diversity. *Methods in Ecology and Evolution*, **6**, 232–236.

Castagneyrol B, Jactel H (2012) Unraveling plant-animal diversity relationships: A meta-regression analysis. *Ecology*, **93**, 2115–2124.

Clarke LJ, Soubrier J, Weyrich LS, Cooper A (2014) Environmental metabarcodes for insects: in silico PCR reveals potential for taxonomic bias. *Molecular Ecology Resources*, **14**, 1160–1170.

Condon MA, Scheffer SJ, Lewis ML, Swensen SM (2008) Hidden neotropical diversity: greater than the sum of its parts. *Science*, **320**, 928–931.

Cristescu ME (2014) From barcoding single individuals to metabarcoding biological communities: towards an integrative approach to the study of global biodiversity. *Trends in Ecology & Evolution*, **29**, 566–571.

Cruaud A, Ronsted N, Chantarasuwan B *et al.* (2012) An Extreme case of plant-insect codiversification: Figs and fig-pollinating wasps. *Systematic Biology*, **61**, 1029–1047.

Deagle BE, Jarman SN, Coissac E, Taberlet P (2014) DNA metabarcoding and the cytochrome c oxidase subunit I marker: not a perfect match. *Biology Letters*, **10**, 20140562.

Dengler J (2009) Which function describes the species-area relationship best? A review and empirical evaluation. *Journal of Biogeography*, **36**, 728–744.

Dyer LA, Singer MS, Lill JT *et al.* (2007) Host specificity of Lepidoptera in tropical and temperate forests. *Nature*, **448**, 696–699.

Edgar RC (2010) Search and clustering orders of magnitude faster than BLAST.
Bioinformatics, **26**, 2460–2461.
Edger PP, Heidel-fischer HM, Bekaert M *et al.* (2015) The butterfly plant arms-race escalated
by gene and genome duplications. *Proceedings of the National Academy of Sciences of*
the United States of America, **112**, 8362–8366.
Edwards DP, Magrach A, Woodcock P *et al.* (2014) Selective-logging and oil palm: multi-
taxon impacts, biodiversity indicators, and trade-offs for conservation planning.
Ecological Applications, **24**, 2029–2049.
Ficetola GF, Pansu J, Bonin A *et al.* (2015) Replication levels, false presences and the
estimation of the presence/absence from eDNA metabarcoding data. *Molecular Ecology*
Resources, **15**, 543–556.
Gioria M, Bacaro G, Feehan J (2011) Evaluating and interpreting cross-taxon congruence:
Potential pitfalls and solutions. *Acta Oecologica*, **37**, 187–194.
Haddad NM, Crutsinger GM, Gross K *et al.* (2009) Plant species loss decreases arthropod
diversity and shifts trophic structure. *Ecology Letters*, **12**, 1029–1039.
Hamilton AJ, Novotný V, Waters EK *et al.* (2013) Estimating global arthropod species
richness: refining probabilistic models using probability bounds analysis. *Oecologia*, **171**,
357–65.
Hao X, Jiang R, Chen T (2011) Clustering 16S rRNA for OTU prediction: A method of
unsupervised Bayesian clustering. *Bioinformatics*, **27**, 611–618.
Holt BG (2013) An update of Wallace's zoogeographic regions of the world. *Science*, **339**,
74–78.
Hortal J, Borges P a V, Gaspar C (2006) Evaluating the performance of species richness
estimators: Sensitivity to sample grain size. *Journal of Animal Ecology*, **75**, 274–287.
Hyams D (2009) CurveExpert: A curve fitting system for Windows. URL
http://www.curveexpert.net (accessed 15 Feb 2016).
Ji Y, Ashton L, Pedley SM *et al.* (2013) Reliable, verifiable and efficient monitoring of
biodiversity via metabarcoding. *Ecology Letters*, **16**, 1245–1257.
Lau MW-N, Fellows JR, Chan BPL (2010) Carnivores (Mammalia: Carnivora) in South
China: a status review with notes on the commercial trade. *Mammal Review*, **40**, 247–
292.
Lenz TL, Becker S (2008) Simple approach to reduce PCR artefact formation leads to reliable
genotyping of MHC and other highly polymorphic loci--implications for evolutionary
analysis. *Gene*, **427**, 117–123.
Lewinsohn TM, Roslin T (2008) Four ways towards tropical herbivore megadiversity.
Ecology Letters, **11**, 398–416.
Lin J, Ng S-C, Zhang M *et al.* (2013) Vegetation of Yinggeling Nature Reserve. In:
Biodiversity and conservation of Hainan Yinggeling Nature Reserve (eds Jiang H, Chan
BPL, Zhou Y, *et al.*), pp. 214–259. China Forestry Press, Beijing, China.
Lopez LCS, De Aguiar Fracasso MP, Mesquita DO, Palma ART, Riul P (2012) The
relationship between percentage of singletons and sampling effort: A new approach to
reduce the bias of richness estimates. *Ecological Indicators*, **14**, 164–169.
Matsen FA, Kodner RB, Armbrust EV (2010) pplacer: linear time maximum-likelihood and
Bayesian phylogenetic placement of sequences onto a fixed reference tree. *BMC*
Bioinformatics, **11**, 538.
Munch K, Boomsma W, Huelsenbeck JP, Willerslev E, Nielsen R (2008) Statistical
assignment of DNA sequences using Bayesian phylogenetics. *Systematic biology*, **57**,
750–7.

Novotny V, Basset Y (2005) Host specificity of insect herbivores in tropical forests. *Proceedings of the Royal Society. B*, **272**, 1083–1090.

Novotny V, Basset Y, Miller SE *et al.* (2002) Low host specificity of herbivorous insects in a tropical forest. *Nature*, **416**, 841–844.

Novotny V, Drozd P, Miller SE *et al.* (2006) Why are there so many species of herbivorous insects in tropical rainforests? *Science*, **313**, 1115–8.

Novotny V, Miller SE, Hulcr J *et al.* (2007) Low beta diversity of herbivorous insects in tropical forests. *Nature*, **448**, 692–697.

Oksanen J, Blanchet FG, Kindt R *et al.* (2015) vegan: Community Ecology Package. URL http://CRAN.R-project.org/package=vegan.

Paudel E, Dossa GGO, Xu J, Harrison RD (2015) Litterfall and nutrient return along a disturbance gradient in a tropical montane forest. *Forest Ecology and Management*, **353**, 97–106.

Pellissier L, Ndiribe C, Dubuis A *et al.* (2013) Turnover of plant lineages shapes herbivore phylogenetic beta diversity along ecological gradients. *Ecology Letters*, **16**, 600–608.

Piñol J, Mir G, Gomez-Polo P, Agustí N (2015) Universal and blocking primer mismatches limit the use of high-throughput DNA sequencing for the quantitative metabarcoding of arthropods. *Molecular Ecology Resources*, **15**, 819–830.

R Core Team (2015) R: a language and environment for statistical computing. URL https://www.R-project.org/.

Ranwez V, Harispe S, Delsuc F, Douzery EJP (2011) MACSE: Multiple Alignment of Coding SEquences accounting for frameshifts and stop codons. *PLoS ONE*, **6**, e22594.

Ratnasingham S, Hebert PDN (2007) BOLD: The Barcode of Life Data System (http://www.barcodinglife.org). *Molecular Ecology Notes*, **7**, 355–364.

Ratnasingham S, Hebert PDN (2013) A DNA-based registry for all animal species: The Barcode Index Number (BIN) system. *PLoS One*, **8**, e66213.

Riedel A, Sagata K, Suhardjono YR, Tänzler R, Balke M (2013) Integrative taxonomy on the fast track - towards more sustainability in biodiversity research. *Frontiers in Zoology*, **10**, 15.

Schmidt S, Schmid-Egger C, Morinière J, Haszprunar G, Hebert PDN (2015) DNA barcoding largely supports 250 years of classical taxonomy: identifications for Central European bees (Hymenoptera, Apoidea partim). *Molecular Ecology Resources*, **15**, 985–1000.

Sreekar R, Zhang K, Xu J, Harrison RD (2015) Yet another empty forest: Consider the conservation value of a recently established tropical nature reserve. *PLoS One*, **10**, e0117920.

Stork NE, McBroom J, Gely C, Hamilton AJ (2015) New approaches narrow global species estimates for beetles, insects, and terrestrial arthropods. *Proceedings of the National Academy of Sciences*, **112**, 7519–7523.

Taberlet P, Coissac E, Pompanon F, Brochmann C, Willerslev E (2012) Towards next-generation biodiversity assessment using DNA metabarcoding. *Molecular Ecology*, **21**, 2045–2050.

Tang CQ, Humphreys AM, Fontaneto D, Barraclough TG (2014) Effects of phylogenetic reconstruction method on the robustness of species delimitation using single-locus data. *Methods in Ecology and Evolution*, **5**, 1086–1094.

Thompson JN (1994). The coevolutionary process. University of Chicago Press, Chicago, USA.

632   Walther BA, Moore JL (2005) The concepts of bias, precision and accuracy, and their use in
633   testing the performance of species richness estimators, with a literature review of
634   estimator performance. *Ecography*, **28**, 815–829.
635   Xu J, Lebel L, Sturgeon J (2009) Functional links between biodiversity, livelihoods, and
636   culture in a Hani swidden landscape in southwest China. *Ecology and Society*, **14**, 20.
637   Yang C, Wang X, Miller J a. *et al.* (2014) Using metabarcoding to ask if easily collected soil
638   and leaf-litter samples can be used as a general biodiversity indicator. *Ecological
639   Indicators*, **46**, 379–389.
640   Yu DW, Ji Y, Emerson BC *et al.* (2012) Biodiversity soup: metabarcoding of arthropods for
641   rapid biodiversity assessment and biomonitoring. *Methods in Ecology and Evolution*, **3**,
642   613–623.
643   Zhu H, Shi JP, Zhao CJ (2005) Species composition, physiognomy and plant diversity of the
644   tropical montane evergreen broad-leaved forest in Southern Yunnan. *Biodiversity and
645   Conservation*, **14**, 2855–2870.

Data Accessibility

647   DNA sequences: Genbank's Short Read Archive (Accession numbers: SRP065001 and

648   SRP065147) and Dryad doi: http://dx.doi.org/10.5061/dryad.37b53.

649   Bioinformatic script: Dryad doi: http://dx.doi.org/10.5061/dryad.37b53.

650   All input data sets to R and R script: Dryad doi: http://dx.doi.org/10.5061/dryad.37b53.

Author Contributions

KZ, DWY, RDH, and HJ designed the study. KZ, RDH, SL, CXY, CYY and HW collected

data, led by RDH and SL. KZ, YJ, CXY, XW performed the molecular experiments. KZ and

YJ performed the bioinformatic analyses. KZ performed the statistical analyses and wrote the

first draft of the manuscript. DWY and RDH contributed substantially to revisions.

659 **Table 1** Procrustes correlations among plant, Insecta, and individual insect order communities in (a) Yinggeling and (b) Mengsong (9999
660 permutations), with the input NMDS (non-metric multidimensional scaling) ordinations calculated from binary Jaccard dissimilarities (k = 2
661 axes used in Yinggeling, k = 4 in Mengsong). Correlations with plants are bolded. In Mengsong, where insects were sampled in two seasons, wet
662 vs. dry-season Procrustes correlations are presented on the diagonal and are underlined, and the proportions of wet-season Operational
663 Taxonomic Units (OTUs) that were also collected in the dry season are reported below as percentages.
664
665 (a) Yinggeling

	Coleoptera	Diptera	Hemiptera	Hymenoptera	Lepidoptera	**Trees (*n* = 297)**
Insecta (*n* = 1995)						**0.76*** **
Coleoptera (*n* = 239)		0.54**	0.47*	0.63***	0.43*	**0.44***
Diptera (*n* = 848)			0.64***	0.57**	0.53**	**0.73*** **
Hemiptera (*n* = 205)				0.45*	0.39	**0.60*** *
Hymenoptera (*n* = 163)					0.55**	**0.56***
Lepidoptera (*n* = 263)						**0.65*** **

667   (b) Mengsong

		Insecta	Coleoptera	Diptera	Hemiptera	Hymenoptera	Lepidoptera	Psocoptera	Vascular plants (n = 807)
Insecta	wet (n = 2073)	0.75***							**0.78***
	dry (n = 2215)	60.6%							**0.75***
Coleoptera	wet (n = 375)		0.45*	0.60***	0.58***	0.55***	0.43*	0.52**	**0.53***
	dry (n = 603)		38.0%	0.75***	0.70***	0.63***	0.51***	0.60***	**0.63***
Diptera	wet (n = 414)			0.52***	0.67***	0.71***	0.63***	0.52***	**0.71***
	dry (n = 413)			70.9%	0.78***	0.61***	0.52***	0.50**	**0.65***
Hemiptera	wet (n = 435)				0.67***	0.58***	0.44*	0.50**	**0.70***
	dry (n = 370)				71.1%	0.61***	0.44*	0.57***	**0.77***
Hymenoptera	wet (n = 409)					0.68***	0.50**	0.57***	**0.64***
	dry (n = 360)					70.0%	0.51***	0.61***	**0.72***
Lepidoptera	wet (n = 78)						0.33	0.40	**0.46**
	dry (n = 95)						61.1%	0.47**	0.38
Psocoptera	wet (n = 71)							0.42	**0.55***
	dry (n = 63)							85.7%	**0.52***

668   * *p* < 0.05, ** *p* < 0.01, *** *p* < 0.001, non-significant values shown in gray, after adjustment for multiple tests (*p.adjust*, method = "fdr")

Figure 1 Inventory plots in Yinggeling ($n = 21$) and Mengsong ($n = 28$). Colors stand for different elevation categories (orange = < 600 m; blue = 600-800 m; purple = ≥ 800 m) in Yinggeling, and for different habitat types (red = mature forest; green = regenerating forest; yellow = open land) in Mengsong. The Mengsong area left of the dashed line is included in Bulong Nature Reserve.

Figure 2 Same-site predictions. Scatterplot of plant-model estimates versus non-parametric estimates of total OTU (Operational Taxonomic Unit) richness, for Arachnida (for Mengsong only), Insecta and insect orders that contained ≥ 50 OTUs. To quantify the goodness-of-fit of these two estimates, explained variances (R^2) for insect orders were calculated from a y = x model (dashed line). Circles, squares and triangles stand for P-corrected versions of the Jackknife1, Jackknife2 and Chao estimators, respectively. Different colors stand for different taxa; only taxa absent from Yinggeling are labeled in the Mengsong figures. The plant-model functions were Weibull for all taxa in Yinggeling and Mengsong, except for Hymenoptera (dry season) in Mengsong, which was a power function. Note breaks in the axes.

Figure 3 Cross-site predictions. Scatterplot of plant-model estimates versus non-parametric estimates of total OTU richness. Symbols as in Figure 2. Plant models from one landscape were used to predict non-parametric estimates in the other landscape. Shown here are the Mengsong wet-season results. Mengsong dry-season results are similar and shown in Figure S5. The shaded area encompasses a two-fold difference between the two estimates (y = 0.5x to y = 2x). The plant-model functions were Weibull for all taxa in Yinggeling and Mengsong, except for Lepidoptera in Mengsong, which was a linear function.

Figure 4 Procrustes superimposition plots between plant and Insecta communities (9999 permutations), with the input of non-metric multidimensional scaling (NMDS) ordinations calculated from binary Jaccard dissimilarities (k = 2 axes used in Yinggeling, k = 4 in Mengsong). All Procrustes correlation coefficients (r) are significantly different from zero at $p < 0.001$ (Table 1). Each point is a census site; solid points indicate plant data, and hollow points insect data. Colors as in Figure 1.

Plant diversity accurately predicts insect diversity in two tropical landscapes: Supporting Information

Kai Zhang, * † [1] Siliang Lin, ‡ [1] Yinqiu Ji, * Chenxue Yang, * Xiaoyang Wang, * † Chunyan Yang, * Hesheng Wang, § Haisheng Jiang, ‡ Rhett D. Harrison ¶ ** and Douglas W. Yu * ††

* State Key Laboratory of Genetic Resources and Evolution, Kunming Institute of Zoology, Chinese Academy of Sciences, Kunming, Yunnan 650223, China
† Kunming College of Life Science, University of the Chinese Academy of Sciences, Kunming 650204, China
‡ School of Life Sciences, South China Normal University, Guangzhou, Guangdong 510631, China
§ Hainan Yinggeling National Nature Reserve, Baisha, Hainan 572800, China
¶ World Agroforestry Center, East and Central Asia Regional Office, Kunming 650201, China
** Center for Mountain Ecosystem Studies (CMES), Kunming Institute of Botany, Chinese Academy of Sciences, Kunming 650201, China
†† School of Biological Sciences, University of East Anglia, Norwich Research Park, Norwich, Norfolk NR47TJ, UK

co-corresponding authors. DWY +86 871 68125438 (ofc), 86 871 65110887 (fax), dougwyu@gmail.com; RDH +86 18687136498 (ofc), 86 871 65223377 (fax), r.harrison@cgiar.org

[1] Shared first-authorship.

This PDF file includes:
Figs. S1 to S7
Table S1
NOTE: R Markdown output document (*Plants accurately predict insects_R Markdown output.html*) is not include here and presented independently.

Figure S1 Same-site prediction. Scatterplot of plant-model (25 candidate functions) estimates versus non-parametric estimates of total OTU richness. Symbols as in Figure 2. Note breaks in the axes.

Figure S2 Same-site predictions. Scatterplot of plant-model estimates versus non-parametric estimates of total OTU richness in the (a) forests and (b) open land in Mengsong. Symbols as in Figure 2. The plant-model functions for (a) forests were Weibull for all taxa except Hymenoptera (dry season), which was a power function. The plant-model functions for (b) open land were power for all taxa except Coleoptera (wet season), which was a linear function. Note breaks in the axes.

Figure S3 Same-site predictions. Scatterplot of plant-model (25 candidate functions) estimates versus non-parametric estimates of total OTU richness in the (a) forests and (b) open land in Mengsong. Symbols as in Figure 2. Note breaks in the axes.

Figure S4 Same-site predictions. Scatterplot of plant-model estimates versus non-parametric estimates of total OTU richness in the forests in Mengsong. Only tree data were included to build the plant model. Symbols as in Figure 2. The plant-model functions were Weibull for all taxa except Hymenoptera (dry season), which was a linear function. Note breaks in the axes.

Figure S5 Cross-site predictions. Scatterplot of plant-model estimates versus non-parametric estimates of total OTU richness. Symbols as in Figure 3. Shown here are the Mengsong dry-season results. The plant-model functions were Weibull for all taxa in both Yinggeling and Mengsong.

Figure S6 Procrustes superimposition plots between plant and insect order communities in Yinggeling. All Procrustes correlation coefficients (r) are significantly different from zero at $p < 0.05$ (Table 1). Symbols as in Figure 4.

Figure S7 Procrustes superimposition plots between plant and insect order communities in (a) wet and (b) dry seasons of Mengsong. All Procrustes correlation coefficients (r) are significantly different from zero at $p < 0.01$ for wet season and at $p < 0.001$ for dry season except for Lepidoptera (Table 1). Symbols as in Figure 4.

Table S1 Procrustes correlations among plant, Insecta, and insect order communities in the forests of Mengsong ($n = 22$) (9999 permutations), with the input of non-metric multidimensional scaling (NMDS) ordinations calculated from binary Jaccard dissimilarities (k = 4). The percentage calculations, and the bold and underlined statistics as in Table 1.

		Insecta	Coleoptera	Diptera	Hemiptera	Hymenoptera	Lepidoptera	Psocoptera	Vascular plants ($n = 747$)
Insecta	wet ($n = 1780$)	0.68***							**0.69***
	dry ($n = 1913$)	55.9%							**0.70***
Coleoptera	wet ($n = 323$)		0.55**	0.63***	0.59***	0.60***	0.47	0.50*	**0.52**
	dry ($n = 510$)		34.9%	0.76***	0.72***	0.71***	0.56***	0.67***	**0.68***
Diptera	wet ($n = 363$)			0.50*	0.65***	0.65***	0.58***	0.60***	**0.62***
	dry ($n = 381$)			64.8%	0.71***	0.68***	0.53**	0.62***	**0.65***
Hemiptera	wet ($n = 360$)				0.53*	0.58***	0.48*	0.52**	**0.60***
	dry ($n = 293$)				65.5%	0.54**	0.53**	0.65***	**0.73***
Hymenoptera	wet ($n = 360$)					0.64***	0.52**	0.65***	**0.58***
	dry ($n = 308$)					66.9%	0.52**	0.60***	**0.66***
Lepidoptera	wet ($n = 70$)						0.46	0.45	**0.48*
	dry ($n = 84$)						56.0%	0.50*	0.36
Psocoptera	wet ($n = 60$)							0.44	**0.52**
	dry ($n = 59$)							78.0%	**0.62***

* $p < 0.05$, ** $p < 0.01$, *** $p < 0.001$, non-significant values shown in gray, after adjustment for multiple tests (*p.adjust*, method = "fdr")