Macroevolutionary trade-offs in plant-feeding insects

Daniel A. Peterson,[1,2] Nate B. Hardy,[3] and Benjamin B. Normark[1]

[1]Department of Biology and Graduate Program in Organismic and Evolutionary Biology, University of

Massachusetts, Amherst, Massachusetts

[2]E-mail: dapeters@cns.umass.edu

[3]Department of Entomology and Plant Pathology, Auburn University, Auburn, Alabama

Summary

Most plant-feeding insects are ecological specialists restricted to one or a few closely related

host-plant species (Forister et al. 2015). A long-standing hypothesis asserts that natural selection favors

host specialization because trade-offs between performance on alternative host species limit the fitness

of generalists, yet empirical evidence for such trade-offs is scarce (Futuyma and Moreno 1988; Forister

et al. 2012). Here we show that trade-offs between adaptations to alternative hosts occur over both

long- and short-term macroevolutionary timescales, but positive associations between host-use traits

are also abundant. Host-use records of 1604 caterpillar (Lepidoptera) species revealed negative

associations between adaptations to two diverse groups of host-plant taxa over 150 million years

(Misof et al. 2014) of caterpillar evolutionary history, but a different division between use of

angiosperm and pine hosts among closely related caterpillars. In contrast, host-use records of 955 true

bug (Hemiptera) species suggested uniformly positive associations between adaptations to the same

host taxa both over the 300-million-year (Misof et al. 2014) evolutionary history of true bugs and

among closely related species. The lack of consistent patterns across insect orders and timescales

suggests that host-use trade-offs are historically contingent rather than universal constraints, reflecting

the diversity of mechanisms driving host-specialization in plant-feeding insects.

Main Text

The ubiquity of ecological specialists (Forister et al. 2012, 2015) is often attributed to

evolutionary constraints on generalism (Futuyma and Moreno 1988; Futuyma et al. 1995). Adaptations

to one environment may come at the cost of performance in other environments (Futuyma and Moreno

1988; Forister et al. 2012), and such trade-offs are crucial elements of most theoretical models of the

evolution of specialization (Ravigné et al. 2009). In many biological systems, trade-offs are implied by

negative correlations between traits (Shoval et al. 2012), but performance measures on alternative host

species are most often uncorrelated or positively correlated in wild and experimentally selected

populations of plant-feeding insects (Futuyma 2008; Forister et al. 2012). In fact, although plant-

feeding insects are a model system for the study of ecological specialization (Futuyma and Moreno

1988; Forister et al. 2012), the role of trade-offs in the evolution of insect host-specialization remains

generally unsupported.

However, trade-offs that drive specialization in plant-feeding insects may be difficult to detect

within species (Joshi and Thompson 1995). Host-use traits are phylogenetically conserved in many

insect groups (Futuyma and Agrawal 2009), and genetic variation for use of novel hosts is often absent

within a single population (Futuyma et al. 1995). These constraints likely stem from the fact that

resource use is a complex trait involving multiple interacting genetic loci (Remold 2012), so the

evolutionary trajectories that lead to high fitness on alternative resources may be mutually exclusive.

For example, the evolution of improved performance on one host can select for a preference for that

host, driving a positive feedback loop of further specialization (Fry 1996). When epistatic interactions

are strong, the greatest variation in host-use strategies will evolve between reproductively isolated

lineages over long time scales (Satterwhite and Cooper 2015), with trade-off outcomes determined by

each lineage's evolutionary history (Rodriguez-Verdugo et al. 2014). Analagous macroevolutionary

trade-offs have been described in plants; the presence of alternative defensive strategies are negatively

correlated over plant evolutionary history (Campbell and Kessler 2013). However, it remains unknown

whether this functional diversification across hosts has created macroevolutionary trade-offs for plant-

feeding insects; performance on alternative hosts may instead be positively associated if generalist

adaptations have been prevalent during the evolution of insect herbivores (Peterson et al. 2015).

We investigated trade-offs between adaptations to diverse host taxa over long- and short-term

macroevolutionary timescales in two orders of plant-feeding insects: Lepidoptera (caterpillars) and

Hemiptera (true bugs). Assuming that trade-offs should produce negative correlations between recorded

presence on alternative host taxa for individual insect species (Peterson et al. 2015), we quantified the

macroevolutionary relationships between use of the most common host-plant orders for North

American insects (Figure 1). Long- and short-term effects were partitioned by estimating separate

correlation parameters for phylogenetic and residual effects (Figure 2) in a phylogenetic mixed model.

We assembled the pairwise correlations between use of all focal hosts to produce network graphs of

host-use relationships, revealing overall patterns of host-use evolution in each insect order. Specifically,

we investigated two potential trade-off scenarios: segregation of hosts into discrete clusters and

isolation of individual host orders.

We found that the network of long-term evolutionary relationships between lepidopteran use of

the focal host orders was significantly structured ($P < 0.01$), revealing two large clusters of host taxa

(Figure 3a). Clustering is a common feature of ecological networks (Olesen et al. 2007) that implies

functional relationships between organisms which often correlate with phylogenetic distances (Krasnov

et al. 2012; Hadfield et al. 2014). However, cluster membership in our results was surprising: the

gymnosperm Pinales (pines) and monocot Poales (grasses) were each affiliated with a different set of

eudicot orders. Instead of host-insect coevolution, long-term evolutionary trade-offs experienced by

lepidopteran lineages appear to reflect a broad-scale ecological division between plant hosts. The

relevant ecological traits that differ between the two host clusters are not obvious, but may correspond

to the habitat types where those hosts are found (Futuyma 1976).

Short-term correlations between lepidopteran use of the focal host taxa also showed significant

network structure ($P < 0.01$) but use of all angiosperm hosts formed a single cluster of mostly positive

associations (Figure 3b). The disappearance over short evolutionary timescales of the trade-off between

angiosperm hosts suggests that individual generalist species have overcome the constraints of historical

contingency rather than merely maximizing host breadth within their lineage's niche. On the other

hand, we do not account for differences in fecundity between specialist and generalist insects on

particular hosts; it may be that generalists usually have lower fitness – i.e. they are jacks of all trades

but masters of none (Futuyma and Moreno 1988). However, costs of generalism have been difficult to

document (Forister et al. 2012; Gompert et al. 2015), so the positive residual correlations we observed

may instead represent evolutionary breakthroughs made possible by novel mechanisms of phenotypic

plasticity or other generalist adaptations (Barrett and Heil 2012).

Nevertheless, although we observed few constraints to short-term evolution of lepidopteran

host-use within the focal angiosperm taxa, we did find evidence for a short-term evolutionary trade-off

between use of angiosperms and Pinales (pines). Lepidopteran presence on pines and many of the focal

angiosperm taxa were positively correlated over long timescales, yet individual lepidopteran species

found on pines were less likely to use all other focal host taxa (mean pairwise correlation -0.23, $P <$

0.01). This pattern suggests that although pines and some angiosperms are used by insects with similar

evolutionary backgrounds, a trade-off between evolutionarily labile traits tends to prevent the

simultaneous use of those host taxa by individual caterpillar species. Such trade-offs between labile but

mutually exclusive host-use traits may be particularly significant because they can promote rapid

speciation (Nosil et al. 2002), driving the adaptive radiations that have produced much of the plant-

feeding insect diversity on earth (Farrell 1998; Janz et al. 2006).

In contrast to the observed patterns in the Lepidoptera, hemipteran host-use showed no evidence

of trade-offs over long or short evolutionary timescales. Associations between hemipteran presence on

all focal host orders were uniformly positive (Figure 3c,d) and network structure showed significant

support for a single host-use cluster encompassing all focal hosts for both phylogenetic ($P < 0.01$) and

residual correlations ($P < 0.01$). These results suggest that generalist adaptations that increase fitness

across multiple hosts have been more important for Hemiptera than specialist adaptations to particular

hosts (Peterson et al. 2015). Moreover, hemipteran generalism appears completely unrestrained by host

taxonomy even over long time scales, leading to the evolution of both super-generalist species and

clades (Normark and Johnson 2011). The trade-offs observed in lepidopteran host-use are therefore not

universal to plant-feeding insects; they depend on evolutionary context.

There are many differences between Lepidoptera and Hemiptera, but their fundamentally

distinct relationships with host plants may be particularly important to understanding why the role of

trade-offs appears to be very different in the two groups. Hemiptera are sucking insects, while

Lepidoptera are generally leaf-chewers (Forister et al. 2015). These two feeding modes elicit different

modes of plant defensive responses (Ali and Agrawal 2012), and sap-sucking may be particularly

amenable to generalist adaptations that circumvent host defenses (Barrett and Heil 2012). In contrast,

Lepidoptera often rely on specialized enzymes to detoxify defensive chemicals, which may constrain

the evolution of generalism, although super-generalist Lepidoptera do exist, possibly powered by

phenotypic plasticity in enzyme expression (Li et al. 2002).

Overall, we found strong evidence that both labile traits and long-term, historically contingent

evolutionary processes produce important host-use constraints in plant-feeding insects. Nevertheless,

the relatively few, broad-scale trade-offs found here fail to explain the observation that a vast majority

of plant-feeding insects are restricted to a single host family (Forister et al. 2015). It is clear that trade-

offs can vary based on genetic context (Rodriguez-Verdugo et al. 2014), so we cannot rule out the

hypothesis that species-specific, idiosyncratic trade-offs drive ecological specialization. Nevertheless,

although individual species-specific trade-offs have been documented (Nosil et al. 2002), our results

suggest that most microevolutionary trade-offs do not persist over macroevolutionary timescales.

Trade-offs play an intuitive and possibly inescapable role in constraining performance across multiple

tasks (Shoval et al. 2012), yet host-specialization in plant-feeding insects is not necessarily driven by

host-specific limitations on survival and growth. Alternative factors, such as neural constraints

(Bernays 2001), mate-finding (Hawthorne and Via 2002), or genetic drift (Gompert et al. 2015) may

ultimately shape the evolution of each species' ecological niche.

Figure Legends

Figure 1 | M**apping host-use traits on insect phylogenies**. For each host-plant order, colored blocks

indicate which insect species have been observed on that host. Insect species with no hosts shown were

observed only on non-focal hosts or had no host-use information associated with their locality records

(Hemiptera only). Insect families are indicated around the phylogenies as follows: **a**, Lepidoptera – 1:

Noctuoidea, 2: Nymphalidae, 3: Lycaenidae, 4: Hesperiidae, 5: Pyralidae, 6: Sphingidae, 7:

Saturniidae, 8: Geometridae, 9: Tortricidae, 10: Gracillariidae. **b**, Hemiptera – 11: Cicadellidae, 12:

Membracidae, 13: Cicadidae, 14: Miridae, 15: Tingidae, 16: Pentatomidae, 17: Scutelleridae, 18:

Coreidae, 19: Rhopalidae, 20: Lygaeidae, 21: Delphacidae, 22: Fulgoridae, 23: Flatidae, 24: Aphididae,

25: Diaspididae, 26: Coccidae, 27: Pseudococcidae, 28: Psylloidea, 29: Aleyrodidae.

Figure 2 | H**ypothetical scenarios of evolutionary correlation between presence on two hosts**. **a**,

Negative phylogenetic correlation. **b**, Positive phylogenetic correlation. **c**, Negative residual

correlation. **d**, Positive residual correlation. **e**, Negative phylogenetic and positive residual correlations.

f, Positive phylogenetic and negative residual correlations. In each example, filled red squares on the

left indicate which species in the phylogeny are present on host 1, and filled blue squares on the right

indicate which species are present on host 2.

Figure 3 | **Network graphs of inferred pairwise host-use correlations**. **a**, Lepidoptera – phylogenetic

correlations. **b**, Lepidoptera – residual correlations. **c**, Hemiptera – phylogenetic correlations. **d**,

Hemiptera – residual correlations. Each vertex represents a host order, with vertex area proportional to

the number of insects that were observed on that host. Positive interactions between presence a pair of

hosts are represented by solid, black lines and negative correlations by dashed, red lines, with line

thickness proportional to the magnitude of the correlation. Network spatial structure was determined

using the Kamada-Kawai algorithm (Kamada and Kawai 1989), a force-directed layout method in

which "repulsion" between vertices was proportional to the inverse of one plus the correlation values

between the respective hosts. Vertices are labelled with the following abbreviations – Ast: Asterales,

Car: Caryophyllales, Eri: Ericales, Fab: Fabales, Fag: Fagales, Lam: Lamiales, Mal: Malpighiales, Pin:

Pinales, Poa: Poales, Ros: Rosales, Sap: Sapindales. Vertices are colored by taxonomic group –

Eudicots: blue, Monocots: yellow, Conifers: green. Statistically significant modules ($P < 0.05$) are

indicated by grey bubbles. Individual host orders with mean correlations of significantly higher

magnitude than expected ($P < 0.05$) are indicated by bold vertex outlines (black for positive means, red

for negative means).

Supplementary Information is available in the online version of the paper.

Acknowledgements We thank D. Moen, L. Doubleday, and S. Noda for comments that improved the

manuscript. This work was supported by the National Science Foundation (EF-1115191 and DEB-

1258001)

Author Contributions All authors were involved in the study design and wrote the manuscript. D.A.P.

and N.B.H. analysed the data.

Author Information The authors declare no competing financial interests. Correspondence and

requests for materials should be addressed to D.A.P. (dapeters@cns.umass.edu)

Materials and Methods

Data. Lepidopteran host-use data were downloaded from the HOSTS database (nhm.ac.uk/hosts) (Robinson et al. 2015), a collection of published records of worldwide caterpillar host-plants. Hemipteran host-use data were downloaded from the Tri-Trophic Thematic Collection Network database (tcn.amnh.org), a compilation of insect collection records from academic and governmental museums in the United States. For both datasets, we restricted our analysis to records from North America (all localities labelled USA, Canada, Mexico or Nearctic). All plant taxonomic names were standardized with the Taxonomic Name Resolution Service (Boyle et al. 2013) and insect taxonomic names with the python package TaxonNamesResolver and the following reference databases: Aphid Species File (Favret 2015), Integrated Taxonomic Information System (itis.gov), and Catalogue of Life (catalogueoflife.org).

We created binary presence/absence matrices of lepidopteran and hemipteran species by host plant taxon, with insects considered present on all hosts for which they had at least one host-use record. To focus computational resources on host taxa with enough statistical power to detect evolutionary host-use interactions, we restricted our analyses to focal host orders used by at least 100 insect species in one insect order (~10% of the total focal insect species per order). Eleven host orders met this cut-off, and each of these met the cut-off for both Hemiptera and Lepidoptera: Asterales, Caryophyllales, Ericales, Fabales, Fagales, Lamiales, Malpighiales, Pinales, Poales, Rosales, and Sapindales. Interactions with these focal orders accounted for 77% of total interactions in the Lepidoptera dataset and 57% of total interactions in the Hemiptera dataset.

We characterized shared ancestry among the North American lepidopteran and hemipteran species in our host-use dataset by estimating time-scaled phylogenies using a phyloinformatic approach (see Supplemental Materials for details). Phylogenetic data were not available for all species in the host-use dataset, but there was an overlap of host-use and phylogenetic data for 1604 lepidopteran species and 955 hemipteran species. Phylogenies and host-use matrices for these species will be

available on Dryad (datadryad.org) upon publication.

Statistical Analysis. We used a phylogenetic meta-analysis approach to test for evolutionary

correlations between insect use of the focal host-plant orders (Hadfield and Nakagawa 2010). Treating

each pairwise comparison between host taxa individually, we set up a phylogenetic mixed model for

each insect order using the package MCMCglmm (Hadfield 2010) in the R statistical framework (R

Core Team 2015). We estimated the evolutionary interaction between the pair of host traits using the

"`random=~us(trait):Insect`" and "`rcov=~us(trait):units`" syntax to capture both

phylogenetic and residual effects. Prior parameter distributions were specified as "`prior<-`

`list(R=list(V=diag(2),nu=2), G=list(G1=list(V=diag(2),nu=2)))`". All

chains were run for 10 million iterations with a burn-in of 1 million iterations, and we ran ten MCMC

chains for each model to test for convergence. Gelman-Rubin convergence analysis of the ten chains

for each model produced potential scale reduction factors under 1.05 in every case (99% were under

1.01), suggesting that all chains successfully converged (Gelman and Rubin 1992).

215   We were interested in evaluating two potential trade-off scenarios: isolation of individual host-

use traits, and divisions of host-use traits into clusters (characterized by positive interactions within

clusters and negative interactions between them). To characterize isolation of individual host-use traits,

we calculated the mean correlation of the ten pairwise comparisons involving each host-use trait

(separately for phylogenetic and residual correlations). To characterize associations among groups of

host-use traits, we found the network structure that maximized the structure score for the entire network

of host-use traits, defined as the sum of all pairwise correlations between host-use traits in the same

cluster minus the sum of all pairwise correlations between host-use traits in different clusters (See

Supplementary Materials for details). We tested the statistical significance of the resulting values by

producing a series of 100 null datasets that simulated independent Brownian motion of a continuous

character for performance on each host order along the insect phylogenies, plus additional normally

distributed residual variation. We converted the resulting continuous host performance values to a

binary host presence/absence character by assuming that only the insect species with the highest

performance values for each host taxon were present on that host, with the threshold set by matching

the number of species using that host in the empirical data. We then calculated all pairwise correlations

between use of the focal hosts, mean correlations per host-use trait, and whole-network structure as we

did for the empirical data. Empirical individual host-use trait mean correlations were considered

statistically significant when their absolute values exceeded the maximum absolute values of any

individual mean in 95% of null datasets. Empirical network structures was considered statistically

significant when their scores exceeded the network structure scores of 95% the null datasets.

Literature Cited

Ali, J. G., and A. A. Agrawal. 2012. Specialist versus generalist insect herbivores and plant defense.
Trends Plant Sci. 17:293–302.

Barrett, L. G., and M. Heil. 2012. Unifying concepts and mechanisms in the specificity of plant-enemy
interactions. Trends Plant Sci. 17:282–292.

Bernays, E. A. 2001. Neural limitations in phytophagous insects: implications for diet breadth and
evolution of host affiliation. Annu. Rev. Entomol. 46:703–727.

Boyle, B., N. Hopkins, Z. Lu, J. A. Raygoza Garay, D. Mozzherin, T. Rees, N. Matasci, M. L. Narro,
W. H. Piel, S. J. McKay, S. Lowry, C. Freeland, R. K. Peet, and B. J. Enquist. 2013. The taxonomic
name resolution service: an online tool for automated standardization of plant names. BMC
Bioinformatics 14:16.

Campbell, S. A., and A. Kessler. 2013. Plant mating system transitions drive the macroevolution of
defense strategies. Proc. Natl. Acad. Sci. 110:3973–3978.

Farrell, B. D. 1998. "Inordinate fondness" explained: why are there so many beetles? Science 281:555–
559.

Favret, C. 2015. Aphid species file. http://aphid.speciesfile.org

Forister, M. L., L. A. Dyer, M. Singer, J. O. Stireman, and J. T. Lill. 2012. Revisiting the evolution of
ecological interactions. Ecology 93:981–991.

Forister, M. L., V. Novotny, A. K. Panorska, L. Baje, Y. Basset, P. T. Butterill, L. Cizek, P. D. Coley, F.
Dem, I. R. Diniz, P. Drozd, M. Fox, A. E. Glassmire, R. Hazen, J. Hrcek, J. P. Jahner, O. Kaman, T. J.

Kozubowski, T. a Kursar, O. T. Lewis, J. Lill, R. J. Marquis, S. E. Miller, H. C. Morais, M. Murakami,
H. Nickel, N. a Pardikes, R. E. Ricklefs, M. S. Singer, A. M. Smilanich, J. O. Stireman, S. Villamarín-
Cortez, S. Vodka, M. Volf, D. L. Wagner, T. Walla, G. D. Weiblen, and L. a Dyer. 2015. The global
distribution of diet breadth in insect herbivores. Proc. Natl. Acad. Sci. U. S. A. 112:442–7.

Fry, J. 1996. The evolution of host specialization: are trade-offs overrated? Am. Nat. 148:S84–107.

Futuyma, D. J. 1976. Food plant specialization and environmental predictability in Lepidoptera. Am.
Nat. 110:285–292.

Futuyma, D. J. 2008. Sympatric speciation: norm or exception? Pp. 136–147 *in* Specialization,
Speciation, and Radiation: The Evolutionary Biology of Herbivorous Insects.

Futuyma, D. J., and A. A. Agrawal. 2009. Macroevolution and the biological diversity of plants and
herbivores. Proc. Natl. Acad. Sci. U. S. A. 106:18054–18061.

Futuyma, D. J., M. Keese, and D. J. Funk. 1995. Genetic constraints on macroevolution: the evolution
of host affiliation in the leaf beetle genus *Ophraella*. Evolution 49:797–809.

Futuyma, D. J., and G. Moreno. 1988. The evolution of ecological specialization. Annu. Rev. Ecol.
Syst. 19:207–233.

Gelman, A., and D. B. Rubin. 1992. Inference from iterative simulation using multiple sequences. Stat.
Sci. 7:457–472.

Gompert, Z., J. P. Jahner, C. F. Scholl, J. S. Wilson, L. K. Lucas, V. Soria-Carrasco, J. A. Fordyce, C. C.
Nice, C. A. Buerkle, and M. L. Forister. 2015. The evolution of novel host use is unlikely to be
constrained by tradeoffs or a lack of genetic variation. Mol. Ecol. 24:2777–2793.

Hadfield, J. D. 2010. MCMC methods for multi-response generalized linear mixed models: The
MCMCglmm R package. J. Stat. Softw. 33:1–22.

Hadfield, J. D., B. R. Krasnov, R. Poulin, and S. Nakagawa. 2014. A tale of two phylogenies:
comparative analyses of ecological interactions. Am. Nat. 183:174–87.

Hadfield, J. D., and S. Nakagawa. 2010. General quantitative genetic methods for comparative biology:
Phylogenies, taxonomies and multi-trait models for continuous and categorical characters. J. Evol.
Biol. 23:494–508.

Hawthorne, D. J., and S. Via. 2002. The genetic architecture of ecological specialization: correlated
gene effects on host use and habitat choice in pea aphids. Am. Nat. 159:S76–S88.

Janz, N., S. Nylin, and N. Wahlberg. 2006. Diversity begets diversity: host expansions and the
diversification of plant-feeding insects. BMC Evol. Biol. 6:4.

Joshi, A., and J. N. Thompson. 1995. Trade-offs and the evolution of host specialization. Evol. Ecol.
9:82–92.

Kamada, T., and S. Kawai. 1989. An algorithm for drawing general undirected graphs. Inf. Process.

Lett. 31:7–15.

Krasnov, B. R., M. A. Fortuna, D. Mouillot, I. S. Khokhlova, G. I. Shenbrot, and R. Poulin. 2012.
Phylogenetic signal in module composition and species connectivity in compartmentalized host-
parasite Networks. Am. Nat. 179:501–511.

Li, X., M. A. Schuler, and M. R. Berenbaum. 2002. Jasmonate and salicylate induce expression of
herbivore cytochrome P450 genes. Nature 419:712–715.

Misof, B., S. Liu, K. Meusemann, R. S. Peters, A. Donath, C. Mayer, P. B. Frandsen, J. Ware, T. Flouri,
R. G. Beutel, O. Niehuis, M. Petersen, F. Izquierdo-Carrasco, T. Wappler, J. Rust, A. J. Aberer, U.
Aspock, H. Aspock, D. Bartel, A. Blanke, S. Berger, A. Bohm, T. R. Buckley, B. Calcott, J. Chen, F.
Friedrich, M. Fukui, M. Fujita, C. Greve, P. Grobe, S. Gu, Y. Huang, L. S. Jermiin, A. Y. Kawahara, L.
Krogmann, M. Kubiak, R. Lanfear, H. Letsch, Y. Li, Z. Li, J. Li, H. Lu, R. Machida, Y. Mashimo, P.
Kapli, D. D. McKenna, G. Meng, Y. Nakagaki, J. L. Navarrete-Heredia, M. Ott, Y. Ou, G. Pass, L.
Podsiadlowski, H. Pohl, B. M. von Reumont, K. Schutte, K. Sekiya, S. Shimizu, A. Slipinski, A.
Stamatakis, W. Song, X. Su, N. U. Szucsich, M. Tan, X. Tan, M. Tang, J. Tang, G. Timelthaler, S.
Tomizuka, M. Trautwein, X. Tong, T. Uchifune, M. G. Walzl, B. M. Wiegmann, J. Wilbrandt, B.
Wipfler, T. K. F. Wong, Q. Wu, G. Wu, Y. Xie, S. Yang, Q. Yang, D. K. Yeates, K. Yoshizawa, Q.
Zhang, R. Zhang, W. Zhang, Y. Zhang, J. Zhao, C. Zhou, L. Zhou, T. Ziesmann, S. Zou, X. Xu, H.
Yang, J. Wang, K. M. Kjer, and X. Zhou. 2014. Phylogenomics resolves the timing and pattern of insect
evolution. Science 346:763–767.

Normark, B. B., and N. A. Johnson. 2011. Niche explosion. Genetica 139:551–564.

Nosil, P., B. J. Crespi, and C. P. Sandoval. 2002. Host-plant adaptation drives the parallel evolution of
reproductive isolation. Nature 417:440–3.

Olesen, J. M., J. Bascompte, Y. L. Dupont, and P. Jordano. 2007. The modularity of pollination
networks. Proc. Natl. Acad. Sci. U. S. A. 104:19891–6.

Peterson, D. A., N. B. Hardy, G. E. Morse, I. C. Stocks, A. Okusu, and B. B. Normark. 2015.
Phylogenetic analysis reveals positive correlations between adaptations to diverse hosts in a group of
pathogen-like herbivores. Evolution 69:2785–2792.

R Core Team. 2015. R: A language and environment for statistical computing. R Foundation for
Statistical Computing, Vienna, Austria.

Ravigné, V., U. Dieckmann, and I. Olivieri. 2009. Live where you thrive: joint evolution of habitat
choice and local adaptation facilitates specialization and promotes diversity. Am. Nat. 174:E141–E169.

Remold, S. 2012. Understanding specialism when the jack of all trades can be the master of all. Proc.
R. Soc. B Biol. Sci. 279:4861–4869.

Robinson, G. S., P. R. Ackery, I. J. Kitching, G. W. Beccaloni, and L. M. Hernández. 2015. HOSTS - a
database of the world's Lepidopteran hostplants.

Rodriguez-Verdugo, A., D. Carrillo-Cisneros, A. Gonzalez-Gonzalez, B. S. Gaut, and A. F. Bennett.

2014. Different tradeoffs result from alternate genetic adaptations to a common environment. Proc.
Natl. Acad. Sci. 111:12121–12126.

Satterwhite, R. S., and T. F. Cooper. 2015. Constraints on adaptation of *Escherichia coli* to mixed-
resource environments increase over time. Evolution 69:2067–2078.

Shoval, O., H. Sheftel, G. Shinar, Y. Hart, O. Ramote, A. Mayo, E. Dekel, K. Kavanagh, and U. Alon.
2012. Evolutionary trade-offs, Pareto optimality, and the geometry of phenotype space. Science
336:1157–1160.

a

b

Asterales
Caryophyllales
Ericales
Fabales
Fagales
Lamiales
Malpighiales
Pinales
Poales
Rosales
Sapindales