Short title: A new tomato resource for high resolution QTL mapping

Corresponding authors:

Neelima R. Sinha

Department of Plant Biology

1002 Life Sciences

One Shields Ave.

Phone: (530) 754-8441

Fax: (530) 752-5410

e-mail: nrsinha@ucdavis.edu

Julin N. Maloof

Department of Plant Biology

1002 Life Sciences

One Shields Ave.

Phone: (530) 752-8077

Fax: (530) 752-5410

e-mail: jnmaloof@ucdavis.edu

Research area: Genes, Development and Evolution

<u>Title: A new advanced backcross tomato population enables high resolution leaf</u>

<u>QTL mapping and gene identification</u>

Fulop, Daniel*; Ranjan, Aashish[#a]*; Ofner, Itai[§]; Covington, Michael F.; Chitwood, Daniel

H.[#b]; West, Donelly; Ichihashi, Yasunori[#c]; Headland, Lauren[#d]; Zamir, Daniel[§]; Maloof,

Julin N.^; Sinha, Neelima R.^.

Department of Plant Biology, University of California at Davis, Davis, California, United

States of America

[§] Institute of Plant Sciences and Genetics in Agriculture, The Hebrew University of

Jerusalem, Rehovot, Israel

* These authors contributed equally to this work

^ Corresponding authors

Summary: Determining the genes responsible for species differences is central to

biology. Finding such genes is typically done using experimental populations derived

from crossing closely-related species and correlating phenotypes to genotypes in the

progeny of these crosses. In this work we present a powerful new experimental

population for tomato, derived from a cross between a modern cultivated tomato variety

(M82) and the wild species *Solanum pennelli*. This Backcrossed Inbred Line (BIL)

population was derived through repeated backcrossing to the M82 parent. We

demonstrate that compared to other populations available for tomato, this BIL

population has substantially smaller segments of *S. pennelli* DNA, that can enable the

identification of single or few candidates genes for traits of interest. Using the BILs we

determined the genomic regions controlling dissection of the leaves into leaflets, leaflet

shape, and time to flower. Moreover, we demonstrated the benefit of populations with

multiple introgressions per line, such as the BILs, for finding putative epistatic

interactions among loci. Finally, we present newly developed statistical inference and

visualization tools, and demonstrate how state of the art regression techniques can

assist in analyzing populations of this type.

Footnotes:

Funding information: This work was supported through a National Science Foundation grant (IOS-0820854) awarded to NRS and JNM. DHC was a fellow of the Life Sciences Research Foundation funded through the Gordon and Betty Moore Foundation.

Present Address:

[#a] : National Institute of Plant Genome Research, New Delhi, India.

[#b] : Donald Danforth Plant Science Center, St. Louis, Missouri, United States of America.

[#c] : RIKEN Center for Sustainable Resource Science, Yokohama, Kanagawa, Japan.

[#d] : University Campus Suffolk, Suffolk, United Kingdom

Corresponding author emails: nrsinha@ucdavis.edu and jnmaloof@ucdavis.edu

Abstract

Quantitative Trait Locus (QTL) mapping is a powerful technique for dissecting the

genetic basis of traits and species differences. Established tomato mapping populations

between domesticated tomato and its more distant interfertile relatives typically follow a

near isogenic line (NIL) design, such as the *Solanum pennellii* Introgression Line (IL)

population, with a single wild introgression per line in an otherwise domesticated genetic

background. Here we report on a new advanced backcross QTL mapping resource for

tomato, derived from a cross between the M82 tomato cultivar and *S. pennelli*. This so-

called Backcrossed Inbred Line (BIL) population is comprised of a mix of BC_2 and BC_3

lines, with domesticated tomato as the recurrent parent. The BIL population is

complementary to the existing *S. pennellii* IL population, with which it shares parents.

Using the BILs we mapped traits for leaf complexity, leaflet shape, and flowering time.

We demonstrate the utility of the BILs for fine-mapping QTL, particularly QTL initially

mapped in the ILs, by fine-mapping several QTL to single or few candidate genes.

Moreover, we confirm the value of a backcrossed population with multiple introgressions

per line, such as the BILs, for epistatic QTL mapping. Our work was further enabled by

the development of our own statistical inference and visualization tools, namely a

heterogeneous Hidden Markov Model for genotyping the lines, and by using state of the

art sparse regression techniques for QTL mapping.

Introduction

Leaves are the primary sites for capturing light energy, for gas exchange, and for

synthesizing metabolic compounds through photosynthesis. Leaf morphology can

significantly affect growth and overall performance of the plant through the interception

of environmental signals (Nicotra et al., 2011), and is constrained by developmental,

environmental, and phylogenetic contexts. Therefore, understanding the genetic and

gene regulatory mechanisms underlying leaf development is of fundamental importance

for increasing photosynthetic efficiency and optimizing crop performance. The basic

genetic mechanisms underlying leaf development, such as leaf initiation (Waites et al.,

1998; Timmermans et al., 1999; Tsiantis et al., 1999), establishment of leaf polarity

(Kerstetter et al., 2001; McConnell et al., 2001), and leaf differentiation (Nath et al.,

2003), have been investigated in detail in species with simple leaves (Tsukaya, 2013).

In contrast to the simple leaves of Antirrhinum, maize, and Arabidopsis, leaf architecture

in many plants is complex with correspondingly protracted development and leaflet

specification. Tomato is a classic example of a species with a complex or compound

leaf. Tomato leaf lamina are subdivided into two or three orders of leaflets produced by

repeated generation of auxin maxima (Koenig et al., 2009) that pattern leaflets along the

marginal blastozone (Kim et al., 2003a; Kim et al., 2003b), which is competent to

respond to the activity of class I KNOX genes necessary for leaflet initiation (Bharathan

et al., 2002; Efroni et al., 2010). Most of our current understanding of simple and

compound leaf development derives from mutant analyses, expression studies, and

gene expression modulation in transgenic systems (Goliber et al., 1999; Tsiantis and

Hay, 2003; Kimura et al., 2008). Although these studies have provided a basic

understanding of leaf development, the genetic regulation of leaf development at the

level of entire genomes and incorporating epistatic interactions among many different

loci has only just begun to be deciphered.

Quantitative genetics, through the identification of QTL (Quantitative Trait Loci),

provides an effective means to identify genetic loci regulating complex traits, such as

leaf morphology, by capitalizing on existing phenotypic diversity and genetic variation.

Considering the relationship of leaf traits with water use efficiency and crop yield, as

well as with developmental mechanisms common to all lateral organs in plants,

identification of genetic loci regulating leaf traits is of critical importance. Surprisingly,

only a few systematic studies have focused on investigating the quantitative basis of

leaf traits to identify genetic loci underlying yield, physiological, or morphological traits

(Jiang et al., 2000; Pérez-Pérez et al., 2002; Langlade et al., 2005; Tian et al., 2011; Ku

et al., 2012; Chitwood et al., 2014b). Genome-wide association on a nested association

mapping population identified *liguleless*, a gene regulating upright leaf angles in maize,

as a factor underlying yield (Tian et al., 2011). However, this maize study and one

recent study in *Mimulus* suggest that leaf shape is generally highly heritable and

polygenic, with many additive effects, yet without much evidence for epistasis (Ferris et

al., 2015). Nonetheless, there is a need for a comprehensive examination of the role of epistasis in regulating leaf shape and complexity in other species, particularly using statistical techniques that facilitate the inference of epistatic QTL.

Leaf traits, including leaf shape and complexity, changed repeatedly during plant evolution (Bharathan et al., 2002; Kim et al., 2003a; Koenig and Sinha, 2010; Townsley and Sinha, 2012). Elucidation of the genetic basis of morphological changes that occurred during the evolution and divergence of lineages has been a major challenge in biology (Hoekstra and Coyne, 2007; Carroll, 2008). Phenotypic differences between populations and species can be ascribed to QTL inferred from experimental crosses, and thus analysis of QTL can address significant questions about the process of evolutionary divergence. A classic example of the use of QTL in an evolutionary context is the identification of genetic loci that are crucial to the evolution of modern maize from teosinte (Doebley and Wang, 1997; Westerbergh and Doebley, 2002).

The tomato clade offers an excellent system to investigate the natural variation in many plant traits, as its species have similar genetic constitution but divergent morphological and anatomical features and disparate ecologies (Stevens and Rick, 1986; Moyle, 2008; Ranjan et al., 2012). The evolution of many traits of modern domesticated tomato have been deciphered through QTL studies. Since tomato is one of the most extensively used vegetable crops in the world, most studies have focused on investigating variation in fruit size, shape and sugar content (Frary et al., 2000; Fridman et al., 2004; Xiao et al., 2008). Tomato is also well-established as a model system for the study of compound leaf development in Angiosperms, and the tomato clade offers a unique opportunity for investigating natural variation in leaf complexity and other leaf traits. *Solanum pennellii*, one of the most distant interfertile relatives of domesticated tomato, *Solanum lycopersicum*, possesses distinct leaf size, complexity, and morphology. In addition, *S. pennellii* shows striking differences in fruit characteristics, drought resistance, disease resistance and water use efficiency as compared to *S. lycopersicum* (Moyle, 2008; Chitwood et al., 2013; Koenig et al., 2013). All aspects of leaf complexity are reduced in *S. pennellii* relative to domesticated tomato (Holtan and Hake, 2003). *S. pennellii* leaves are rounder and less serrated, and leaf

initiation is delayed while leaf development is accelerated relative to tomato (Ichihashi et al., 2014). In addition, the leaves of *S. pennellii* are more densely covered with epidermal hairs, and show a characteristic high accumulation of anthocyanin at the base of trichomes that gives the appearance of purple spots on these leaves. In spite of these dramatic differences in leaf form, only a few studies have been undertaken to identify genetic loci, and their network of downstream and epistatic interactions, that regulate the various leaf traits at the entire genome level.

Solanum lycopersicum and *Solanum pennellii* are interfertile, enabling genetic dissection of trait differences such as those seen for leaf complexity. The interfertility of these two species was exploited to develop a genetic resource consisting of 76 introgressions lines (IL) (Eshed and Zamir, 1994; Liu and Zamir). Most of these introgression lines contain a single introgression from the wild species in an otherwise domesticated background. These ILs, which introgress the complete *S. pennellii* genome in the cultivated tomato (M82) background, have been extensively phenotyped for numerous traits, such as morphology, yield, fruit quality, fruit primary and secondary metabolites for identification of QTL (Rousseaux et al., 2005; Schauer et al., 2006; Semel et al., 2006; Stevens et al., 2007; Schauer et al., 2008; Steinhauser et al., 2011). The striking differences in leaf morphology between *Solanum lycopersicum* and *Solanum pennellii* were capitalized on to identify QTL associated with leaf dissection (Holtan and Hake, 2003). The IL population was also used to investigate the quantitative basis of adaptive evolution of leaf traits, indicating that leaf traits are an important component of climatic niche adaptation in wild tomatoes(Chitwood et al., 2012a; Muir et al., 2014). The diverse leaf morphologies of wild tomato species (Chitwood et al., 2012b) have been compared and contextualized within the shape range of the tomato IL lines and parents (Chitwood et al., 2014a).

A few genes underlying QTL in the tomato IL population have been fine mapped and cloned. High-resolution mapping applied to the *S. pennellii* ILs led to the map-based cloning of the first two QTL: the fruit weight QTL, *fw2.2*, and the sugar yield QTL, *Brix9-2-5* (Frary et al., 2000; Fridman et al., 2000). Fine genetic mapping using the same introgression population identified the *RXopJ4* locus, a 190-kb segment on

chromosome 6 of *S. pennellii*, that confers resistance to bacterial spot disease

(Sharlach et al., 2012). Advances in next-generation sequencing and bioinformatic

approaches allowed ultra high-density genotyping of the IL population, thus increasing

the resolution of QTL mapping within this population (Chitwood et al., 2013).

Recently we used the IL population to identify more than a thousand QTL relating

not only to leaf morphological features such as leaf shape, size, complexity and

serration, but also to cellular features, such as epidermal cell morphology, and stomatal

density and patterning (Chitwood et al., 2013). Though numerous QTL for various traits

have been identified using the IL population, there has been limited success in fine

mapping or narrowing the QTL interval to enable the identification of strong candidate

genes that could then be tested for causality, in part because of the large size of most of

the introgressions. Further, since almost all of the ILs contain a single introgression on a

specific chromosome, the epistatic interactions regulating a trait cannot be explored

using this population. Therefore, additional introgression lines, where the large

introgression regions of the ILs are subdivided into smaller regions, would facilitate the

identification of strong candidate genes. Moreover, lines with multiple introgressions

would be greatly enable the inference of the genetic interactions regulating a trait.

Here we report on a new genetic mapping resource, an advanced backcross

experimental population with ~500 Backcrossed Inbred Lines (BILs). This population

was generated by repeated backcrossing (2 or 3 times) of a F1 hybrid of *S.*

lycopersicum cv. M82 and *S. pennellii* and its progeny to M82, followed by selfing. For

this work we sampled progeny after four generations of selfing (Fig. 1). Ultrahigh-

density genotyping using reduced representation libraries revealed that the BILs provide

a much higher resolution of introgressions in terms of size compared to the ILs. In

addition, most of these BILs have more than one introgression, which allows the

investigation of genetic interactions underlying traits. We identified QTL regulating

various leaf traits and flowering time, and also characterized potential epistatic

interactions regulating these traits. Furthermore, the BIL population was also

investigated for its potential to fine map the genes likely responsible for the purple spot

(Punctate) phenotype of *S. pennellii* leaves, as well as to fine map leaf traits previously

mapped in the ILs. Similarly, several recent studies have used a subset of the BILs with

introgressions in a region of interest to further fine-map a QTL found in the ILs (Ning et

al., 2015; Fan et al., 2016; Müller et al., 2016). The precise genotypes of these lines,

their utility in fine mapping traits, the greater phenotypic spectrum seen in the lines, and

their utility for mapping single locus and epistatic QTL suggests that this population will

be an extremely useful resource for basic and applied research in tomato. In this study

we also strove to overcome some of the limitations of QTL mapping, such as large QTL

intervals encompassing many genes and methodological obstacles to revealing the

genetic architecture of traits, by developing our own statistical inference and

visualization tools for high precision genotyping and by using state of the art sparse

regression techniques to reveal the genetic architecture of traits through the inference of

epistatic interactions without preconditioning on main effect QTL.

Results and Discussion

High precision genotyping by means of a novel HMM

The Backcrossed Inbred Lines (BILs) were genotyped at high resolution using

Restriction Enzyme Sequence Comparative ANalysis (RESCAN), a reduced

representation genomic sequencing method (Monson-Miller et al., 2012; Seymour et al.,

2012). A polymorphism database was created for sequence variants that differentiate

the two parents, the domesticated tomato variety *S. lycopersicum* cv. M82 and its wild

relative, *S. pennellii* LA0716. The database consisted of 346,645 polymorphisms

between M82 and *S. pennellii* (295,571 SNPs and 51,074 inserted or deleted

nucleotides), which was used to identify parental SNPs and INDELs for each BIL across

the entire genome.

Because of the generally low and sparse sequence coverage of RESCAN, we

used a novel heterogeneous Hidden Markov Model (HMM) to define the introgressed

regions for each BIL. The HMM proved indispensable for accurately genotyping about

15% of the lines, which were problematic due to a variety of issues (Fig. 2). Moreover,

the HMM also enabled us to infer sharper recombination boundaries

(github.com/mfcovington/bil-paper). The HMM inferred narrow transition boundaries in

spite of noise and/or low coverage in some of the lines (Fig. 2).

We were able to establish the genotype of 493 out of a total 545 BILs with

confidence. However, 52 lines were found to be genotypically identical to the M82

parent, and two out of the remaining 441 were identical to two other lines. This could be

due to seed or pollen contamination, and/or random effects during backcrossing and

selfing. We were not able to establish the genotype of 52 other lines either due to lack

of germination or excessive noise in the raw genotype data. The 439 uniquely

genotyped BILs covered the entire tomato genome except for a small gap of ~400Kb on

chromosome 1. We detected heterozygous introgressions on almost all chromosomes

(Figs. 2C, 3A, and 4; Supplemental Fig. S1). However most of these heterozygous

introgressions were covered by another homozygous introgression in the same region

in a different BIL. Nevertheless, although the BIL lines were not fully inbred after 4

generations of selfing, overall we found little residual heterozygosity (Figs. 3A and 4).

The comprehensive genotype of the 439 BILs along with the chromosome-wise

distribution of introgressions for the population is presented in Supplemental Datasets

S1 and S2. In addition, the plots of the genotyped BILs are presented in an online

resource for this paper (github.com/mfcovington/bil-paper).

The number of introgressions for each BIL varied from 1 to 12 with a mean of

3.125 introgressions per BIL (Fig. 3B). Approximately 80% of the lines had from 1 to 4

introgressions (Fig. 3B). As opposed to the single introgressions per line of the IL

population, the presence of multiple introgressions in a single BIL enables the study of

the genetic interactions regulating complex traits. The BIL population provides a

synergistic complement to the IL population (Eshed and Zamir, 1994), an additional

resource for linking QTL identified in the ILs to genes as multiple BILs possess smaller

overlapping introgressions encompassed within the larger introgression regions of the IL

population. The mean and median numbers of genes per introgression are

approximately 677 and 485 for the ILs, and 479 and 293 for the BILs. For example, the

introgression in IL4-3 has been identified as having the largest contribution to tomato

leaf traits in terms of the overall number of QTL mapped to this region (Holtan and

Hake, 2003; Chitwood et al., 2013). We identified at least 50 BILs with breakpoints

overlapping the introgressed region of IL4-3, allowing fine-scale dissection of QTL in

this region. We also explored the BIL population for the number of introgressions on

specific chromosomes (Table 1). A relatively small number of BILs (54) have an

introgression on chromosome 1, the largest chromosome by physical distance, whereas

a surprisingly high number of BILs (236) have introgressions on chromosome 11 (Table

1; Figs. 3 and 4). Chromosome 5 also harbors a large number of introgressions (171).

The BILs showed remarkable variation in terms of the percentage of *S. pennellii*

genome in each line (Fig. 3C), ranging from <1% to >10%. The introgression size

ranged from 16.7 Kb on chromosome 10 in BIL-025, containing only 4 genes, to the

largest introgressions on chromosome 12 in BIL-083 and BIL-444, which span almost

the entire chromosome and contain more than 2400 genes. Taken together, the

remarkable variation in the number of introgressions, size of introgressions, and the

level of genome contribution makes the BILs a highly useful resource for fine mapping

of complex traits and identification of the underlying genes.

The precise boundary information for the genotyped BILs was used to define

non-contiguous "bins": regions defined by the pattern of overlap among introgressions in

the population and within which no line changes genotype (Chitwood et al., 2013; Liu

and Zamir). A total of 1049 unique bins (87 per chromosome on average) were detected

across the entire genome with the maximum number of bins on chromosomes 5 and 11,

with 156 and 128 bins respectively (Table 1; Fig. 3D; Supplemental Dataset S3). Thus,

the BILs provide an increase of nearly an order of magnitude in the number of bins as

compared to the 112 bins defined in the IL population (Chitwood et al., 2013). We used

our precision genotyping and introgression boundary calls to define the gene content of

bins with near certainty. A list of annotated genes within the BIL bins is provided as

Supplemental Dataset S4. Interestingly almost half of the bins defined for BILs (455 out

of total 1049 bins) had 10 or fewer genes, which will greatly facilitate the identification of

candidate genes in future studies. Conversely, only 9 out of 126 IL bins (at the sub-bin

level) have 10 or fewer genes. For the ILs, the summary statistics for the number of

genes per bin (at the sub-bin level) are: range = 1 - 1890, mean = 262.06, and median =

150. In contrast, for the BILs these values are: range = 1 - 984, mean = 23.15, and

median = 7.

Analysis of leaf trait variation in the population

As described above, the parents of this population show striking differences in

leaf morphology and complexity. Not surprisingly, there is substantial variation for these

same traits segregating in the BILs. In order to account for replication and pseudo-

replication of trait measurements, linear mixed models (LMM) were used to estimate

model-fitted BIL trait means for QTL mapping. As expected due to the back-crossed

population design, the distribution of many of the traits is roughly centered around the

recurrent parent, *S. lycopersicum* cv. M82, yet often present a skew towards the wild

parent, *S. pennellii* (Supplemental Figs. S4-S29). This is most evident in traits such as

flowering time and roundness, for which the only BILs significantly different from M82

are in the *S. pennellii* direction (Supplemental Figs. S13 and S11).

Overall, we found positive correlation among leaf complexity traits (i.e., the

number of different types of leaflets and the total number of leaflets). There is stronger

correlation between intercalary and secondary leaflet counts, than between either of

these and primary complexity. Likewise, intercalary and secondary leaflets are also

more strongly correlated than primary leaflets to the total number of leaflets

(Supplemental Fig. S30). We found no correlation between leaf complexity and leaflet

shape traits. Aspect ratio and roundness are very strongly negatively correlated, which

is to be expected. Circularity and solidity are strongly correlated. In the context of

tomato leaflet shape, dentation and lobing of the leaflet margin similarly affects

circularity and solidity. As in our previous study of tomato leaflet shape in the ILs

(Chitwood et al., 2013), we used an outline morphometrics technique, elliptical fourier

descriptors decomposed into their principal components (EFD-PC), to examine leaflet

shape in the BILs (Supplemental Figs. S14-S29). However, the EFD-PC traits showed

low heritability and repeatability in the BILs, perhaps owing to the lower replication in

this dataset (Supplemental Table S1), so we excluded these traits from further

discussion.

QTL regulating leaf traits

We used the precisely defined BIL bins to identify QTL regulating various leaf
developmental traits, such as complexity, aspect ratio, circularity, and solidity. Two
methods, marginal regression (Chitwood et al., 2013) and SparseNet (Mazumder et al.,
2011), were used to identify BINs associated with different traits. Hundreds of bins were
significantly associated with complexity and leaf shape traits by marginal regression at a
q-value of < 0.01 (Supplemental Dataset S5; Supplemental Figs. S31-S45).
Corroborating what we found in the ILs, where ILs 4-3 and 5-4 harbored the largest
number of bins associated with leaf traits (Chitwood et al., 2013), numerous BIL bins in
chromosomes 4 and 5 were associated with leaf complexity and leaflet shape traits.
To increase mapping resolution beyond what is possible with marginal regression
we used the regularized regression method SparseNet, which performs variable
selection and shrinkage of the regression coefficients (i.e., inferred association between
bins and traits). SparseNet reduced the number of trait associated bins by around an
order of magnitude (Fig. 5; Supplemental Figs. S46-S66; Supplemental Dataset S6).
Only 28, 36, 26 and 7 bins were found to be associated with primary, intercalary,
secondary, and total complexity, respectively. This method also identified a remarkably
reduced number of bins and sharp QTL peaks for leaf shape traits. The bins associated
with various leaf traits were distributed on multiple chromosomes across the tomato
genome, emphasizing the polygenic regulation of leaf developmental traits.

Epistatic interactions for the leaf traits

Our previous work showed that tomato leaf complexity and leaflet shape is
polygenic (Chitwood et al., 2013). This is also evident in the current study, where we
detected numerous loci that contribute to leaf traits distributed over the entire tomato
genome. Although functional genetic studies have reported evidence for epistasis in
tomato leaf development (Koltai and Bird, 2000; Ori et al., 2007; Naz et al., 2013), the
genome-wide pattern of epistasis regulating tomato leaf traits has yet to be investigated.

The variable selection capability of SparseNet enabled us to search the full

epistatic QTL model space without preconditioning on QTL with a main (i.e., single

locus) effect. The 1049 BIL bins have 549,676 possible two-locus combinations, which

define the set of potential epistatic QTL. We observed 306,291 (or roughly 56%) of said

two-locus combinations in the BIL lines. Thus, the search space for the epistatic QTL

model comprised 307,340 variables, 1049 single locus plus 306,291 two-locus

variables. We inferred 25, 3, 33 and 23 epistatic interactions regulating primary,

secondary, intercalary and total leaf complexity (Fig. 5 A and B; Supplemental Figs.

S46-S47; Supplemental Dataset S6). We further uncovered 42, 45, 53 and 61 possible

epistatic interactions regulating aspect ratio, circularity, roundness and solidity of

leaflets (Fig. 5 C and D; Supplemental Figs. S48-S49; Supplemental Dataset S6).

Although we did not do so, inclusion of a per-line effect in the regression model would

help guard against false positive epistatic QTL in a backcrossed population such as the

BILs with uneven number of lines covering different genomic regions. Similarly, higher

replication per line would also lower the chance for false positives.

Candidate genes underlying the detected QTL

In order to define candidate genes underlying the identified single locus and

epistatic QTL, we used a previously established set of literature-curated leaf

developmental genes (Ichihashi et al., 2014). We reasoned that any causal gene would

have amino acid and/or *cis*-driven mRNA abundance differences segregating in the

population. To identify genes with *cis*-driven mRNA abundance differences we filtered

the literature-curated leaf developmental genes to retain those exhibiting *cis* eQTL using

transcriptome profiling and expression-QTL analysis of the IL population (Chitwood et

al., 2013; Ranjan et al. in revision). Out of 384 leaf developmental genes, 62 were

detected with an underlying eQTL. Using these 62 genes, we identified roughly 10

candidate genes for each of the leaf complexity traits, and 10-20 genes for each leaf

shape phenotype. *GROWTH-REGULATING FACTOR, ASYMMETRIC LEAVES 2,*

AINTEGUMENTA, ARGONAUTE, LEUNIG, LONELY GUY, STM, BEL1, TCP, YUCCA

and *IAA* were some of the salient candidate genes for both leaf complexity and shape

traits (Supplemental Dataset S7). We used permutation analyses to determine the

significance of enrichment of leaf developmental genes among the QTL identified.

Significant enrichment of the literature-curated leaf developmental gene set was

observed for leaf shape QTL, such as aspect ratio, circularity, solidity and roundness,

but the enrichment was not significant for leaf complexity traits (Supplemental Table

S2). One possible explanation for this is that since the literature-curated leaf gene

reference list was primarily based on Arabidopsis (simple) leaf development, it is likely

biased for leaf shape and not representative of leaf complexity in tomato.

We also detected candidate genes underlying the epistatic QTL. One possible

interacting pair of genes underlying an epistatic QTL for primary complexity is *GRF2* on

chromosome 2 and *ANT* on chromosome 4 (Fig. 5A; Supplemental Dataset S8).

Surprisingly, our transcriptomic study of tomato, *S. pennellii*, and *S. habrochaites*

compound leaf development found *GRF2* and *ANT* to be highly-connected hub genes at

the core of the gene co-expression network (Ichihashi et al., 2014). That these

independent analyses using different methodologies identified the same genes supports

their importance for compound leaf development. This finding also underscores the

importance of the regulation of cell proliferation for compound leaf development.

Another epistatic QTL for primary complexity with candidate genes for both loci

was that between chromosomes 7 and 12, which contains *LSH10* and *TCP4* in one

locus and *IAA CARBOXYMETHYLTRAFERASE1* in the other. *BEL1*, *ARGONAUTE 5*

and *GRF4* were other strong candidate genes for epistatic QTL, but without obvious

interacting candidates from the reference gene set on their paired interval. It should be

noted that some of the epistatic QTL may in fact be broad-peaked single locus QTL, as

the inferred pairs of interacting bins are on the same chromosome and possibly too

close to have been separated by recombination in a few generations of crossing.

Therefore, for the purpose of candidate gene identification for epistatic QTL we

removed QTL whose bins are less than 20 cM apart. Following this filtering, we focused

on the top 10 epistatic QTL by effect magnitude (Supplemental Dataset S8).

The above analysis only considered the candidate genes showing gene

expression differences between the two parents and does not account for amino acid

changes that could underlie the leaf shape and complexity differences between the two

parents. Therefore, we used PROVEAN (Choi et al., 2012) to predict amino acid

changes of functional significance between the two parents for the curated leaf

developmental genes. This added 42 genes to the candidate genes reference list.

Redoing the candidate gene search with an updated gene list, including both eQTL and

PROVEAN data, increased the number of candidate genes for each trait but at the cost

of significance in the permutation analysis (Supplemental Table S2). Noteworthy

PROVEAN candidate gene additions were *PIN-FORMED 1*, *GIBBERELLIN OXIDASE*s,

CLAVATA 1-like, *IAA4*, *YUUCA8*, and *GRF5*.

Lastly, in order to explore if gene regulatory interactions underlie some of the

inferred epistatic QTL we assessed whether IL *trans*-eQTL were enriched among the

set of paired regions that make up the BIL epistatic QTL. We did this with two sets of

trans-eQTL derived from eQTL clustering (Ranjan et al. in revision). The above set of

literature-curated leaf development genes were highly represented in two eQTL

clusters, one enriched for leaf development GO (gene ontology) terms and another

enriched for photosynthesis GO terms. Therefore, we tested for IL *trans*-eQTL

enrichment among BIL epistatic QTL by permutation using the leaf development cluster

(set 1), as well as this cluster plus the photosynthesis cluster (set 2). Both sets yielded

highly significant results with p-values of 0.0028 and $< 1.0E\text{-}04$, respectively for set 1

and 2 (Supplemental Table S3).

***Punctate:* a fine-mapping test case**

To experimentally test the usefulness of the BILs in resolving genetic loci

associated with a trait, we conducted a case study for the *Punctate* phenotype, which

enhances the accumulation of anthocyanins in trichome bases creating a "purple spot"

phenotype. Among ILs, IL10-3 shows the *Punctate* phenotype in leaves, a phenotype

observed in *S. pennellii* but not M82. Notably, the Tomato Genetics Resource Center

lists the *Punctate* locus as residing on chromosome 8 (see accessions LA0812,

LA0998, and LA3089, tgrc.ucdavis.edu), suggesting a previously unknown locus on

chromosome 10 is responsible for the trait variation segregating in the ILs and BILs.

The *S. pennellii* introgression in IL10-3 spans 2.65 Mbp and 385 genes (Chitwood et al.,

2013), making candidate gene identification difficult. To determine the extent to which

the BILs could improve the mapping resolution, we screened the BILs and identified 54

lines with the phenotype. Of the 54 BILs exhibiting the "purple spot" phenotype, 46

possessed an introgression on chromosome 10 that overlapped the IL10-3

introgression. The remaining 8 BILs did not show the *Punctate* phenotype consistently.

Among BILs with a chromosome 10 introgression and the "purple spot" phenotype BIL-

228 had the smallest introgression, covering 310 Kbp and containing only 51 genes,

reducing the number of candidate genes from 385. By examining BILs with

chromosome 10 introgressions yet without the *Punctate* phenotype, we further

narrowed the associated genomic region to a 114 Kbp segment with only 18 genes (Fig.

6). Thus, the BILs enabled us to dramatically reduce the region and the number of

candidate genes for the "purple spot" phenotype by 96% and 95% respectively.

Among the 18 genes in the *Punctate* fine-mapped region, there were four

tandemly-duplicated genes encoding MYB transcription factors. Overexpression of an

Arabidopsis homolog of these MYB genes (MYB114, AT1G66380), part of an

independently-derived tandem duplication of MYB genes that regulates anthocyanin

biosynthesis, results in a *Punctate*-like phenotype with accumulation of anthocyanins in

trichome bases (Gonzalez et al., 2008). Remarkably, a similar tandem duplication of

four MYB homologs in grape (*Vitis vinifera*) (*VvMybA1, VvMybA2, VvMybA3*) explains

62% of the variation of anthocyanin accumulation in berries in a Syrah x Grenache F1

QTL study (Fournier-Level et al., 2009). This reinforces the involvement of MYB

transcription factors in the regulation of the *Punctate* phenotype in *S. pennellii* leaves,

and the importance of tandemly duplicated homologs found in lineages as diverse as

Arabidopsis and *Vitis vinifera* in regulating anthocyanin accumulation in a tissue-specific

manner. Similarly, BILs could facilitate the identification and cloning of genes regulating

various other developmental and metabolic traits.

Flowering time QTL and candidate genes

We also used the BIL population to infer QTL for flowering time and identify their

candidate genes. Given the well-characterized pathways and genes for flowering time,

this provides a good opportunity to further test the utility of the BIL population for

candidate identification. Marginal regression identified 69 significant bins associated

with flowering time in tomato. Interestingly, 49 of those were contiguous bins on

chromosome 5 (Supplemental Fig. S39). The remaining bins were localized on

chromosomes 8, 9 and 12. Three narrow QTL, one each on chromosomes 5, 8 and 12,

were found to be associated with flowering time using the more stringent SparseNet

method (Fig. 6B, ring A). *SELF-PRUNING 5G (SP5G)*, a *FLOWERING LOCUS T*

homolog, is present in the QTL on chromosome 5 and is the strongest candidate gene,

by virtue of its function, that we identified for flowering time in the BILs (Carmel-Goren et

al., 2003). The QTL on chromosome 12 also has some important developmental genes

within it, such as *AP2, ULT* and *YABBY*, that are plausible candidates for the regulation

of flowering time in tomato.

The QTL on chromosome 5 is the major flowering time QTL in the BILs by effect

size and is coincident with a large effect QTL in the ILs (Fig. 7A, ring C). The overlap

between the BIL and IL QTL on chromosome 5 defines a 297-Kbp region containing 37

genes, among which *SP5G* is the only flowering time gene (Fig. 6B, inset E;

Supplemental Table S4; Jones et al., 2007). Thus, the *SP5G* example not only

highlights the utility of the BILs for fine-mapping, but also the advantage of using the

BILs in conjunction with the ILs for QTL confirmation and fine-mapping.

QTL confirmation and fine-mapping

Beyond the above two test cases, we explored the benefit of combining results

from the IL and BIL populations for the purposes of QTL confirmation and fine-mapping

since we measured the same traits in both populations. As mentioned above, the IL and

BIL populations share the same parents, but have different crossing schemes and

broadly disparate numbers of lines. The ILs have a near isogenic line (NIL) design,

whereas the BILs have a mixed BC_2 / BC_3 advanced backcross design.

One advantage of utilizing both of these populations is that QTL can be validated

in an independent experiment using a different set of lines. In our case, we found broad

congruence between the IL and BIL QTL mapping results (Fig. 7; Supplemental Figs.

S67-S72; Supplemental Dataset S9). Apart from the *Punctate* example several IL QTL

were significantly narrowed in the BILs, such as QTL for total leaf complexity on

chromosomes 8 and 9 (Fig. 7B), circularity QTL on chromosomes 8 and 11 (Fig. 7C),

and several QTL for primary, secondary, and intercalary complexity (Supplemental Figs.

S67-S69). Likewise, several recent studies have used a subset of the BILs with

introgressions in a region of interest to further fine-map a QTL found in the ILs (Ning et

al., 2015; Fan et al., 2016; Müller et al., 2016).

Furthermore, the use of these two populations is quite complementary. The ILs

allow for higher replication of trait measurements due to the lower number of lines. NIL

populations, such as the ILs, have also been shown to enable the detection of smaller

effect QTL (Keurentjes et al., 2007). In contrast, the BILs can greatly facilitate higher

resolution mapping (i.e., narrower QTL intervals) due to their generally narrower bins.

Moreover, the BILs enable the mapping of epistatic QTL, which is not possible with the

ILs. Our ability to infer epistatic QTL was further empowered by our high precision

genotyping HMM and use of the regularized regression method SparseNet.

Conclusions

The BILs provide a valuable new resource for the tomato research community,

one that is complementary to the long-established IL population yet also extends the

research toolkit considerably. We demonstrated the utility of this new population for

corroborating and fine-mapping IL QTL. Furthermore, we established the capacity of the

BIL population for epistatic QTL mapping and for revealing the genetic architectures of

tomato traits. The present study also highlights the dramatic gains that can be reaped

from the use of advanced statistical techniques, such as our novel heterogeneous

genotyping HMM and the application of regularized regression for QTL mapping.

Materials and methods

Plant materials, growth conditions, experimental design, and phenotyping

Back-crossed Introgression Line (BIL) seeds were generously provided by Itai

Ofner and Dani Zamir (Hebrew University, Rehovot, Israel). These BILs were generated

by crossing the wild species *S. pennellii* LA0716 with the tomato cultivar M82 and

backcrossing the resulting hybrid to M82 two to three times. Four rounds of consecutive

selfing gave rise to the BIL population assayed in this study, which consisted of

approximately 545 lines.

Seeds were washed in 50% bleach for ~2 minutes, rinsed, and placed on water-

soaked paper towels in Phytatrays (Sigma) to germinate. Seeds were stratified in

darkness at room temperature for three days before moving to a 16:8 light cycle in

growth chambers for four days. Seedlings were then transplanted into 5 x 10 subdivided

trays (11" x 22" in dimension) with Sunshine Mix soil (Sun Gro) and grown in a

greenhouse. Approximately three weeks after transplantation, trays were moved to a

lath house where seedlings were hardened by vigorous top watering and allowing the

soil to dry between waterings.

In early May 2011, hardened seedlings were transplanted into a field in Davis,

CA (USA). The hypocotyl of the seedlings was buried in soil. Seedlings were initially

sprinkler-watered, but later transitioned to ditch irrigation. Three randomized blocks

were planted adjacent to each other, each of which contained all 545 BIL genotypes

once and each of the *S. lycopersicum* cv. M82 and *S. pennellii* LA0716 parents five

times.

Phenotyping for the *Punctate* locus was performed on seedlings (when the

phenotype is most penetrant) at the 2-3 leaf stage when the plants were still in the

greenhouse. Seedlings were scored for the punctate phenotype as a discrete,

presence/absence trait. Most genotypes were scored as absent or present in all three

replicates, but a few were not, and this information was noted and used during map-

based cloning of the trait.

Leaf complexity was measured in the field by pairs of researchers each counting

leaflets on four leaves per plant, a pair of leaves per person. Primary, intercalary,

secondary, and total leaflet counts were noted. Leaflet shape was measured as

previously described (Chitwood et al., 2013). Briefly, for each plant, the terminal and two

distal lateral leaflets for five leaves per plant were placed into plastic Ziploc bags and

transported back to lab. The three leaflets for each of five leaves per plant were

dissected and arranged under non-reflective glass. A copy stand (Adorama, 36" Deluxe

Copy Stand) with mounted camera (Olympus SP-500 UZ) controlled remotely using

Cam2Com software (Sabsik) was used to photograph images. ImageJ (Abramoff et al.,

2004) was used to isolate and threshold individual leaflets, which were saved as

individual files. Leaflet outlines were then batch measured in ImageJ for area, aspect

ratio, circularity, roundness and solidity values. Leaflet outlines were then processed

using the software package SHAPE (Iwata and Ukai, 2002) where outlines were

converted to chain code, normalized Elliptical Fourier Descriptors (EFDs) calculated

relative to the orientation of the proximal-distal axis of the leaflet, and a Principal

Component Analysis (PCA) performed on coefficients from the first 20 harmonics.

DNA library preparation and sequencing

After phenotyping, developing leaves and shoot apices were collected for DNA

isolation and library preparation for genotyping. DNA was isolated using the DNeasy

Plant Mini Kit (QIAGEN) and reduced representation libraries were prepared using

restriction enzyme based RESCAN method (Monson-Miller et al., 2012). RESCAN

library preparation involved restriction digestion of genomic DNA with the restriction

enzyme NlaIII followed by adapter ligation and PCR enrichment. The PCR-enriched

libraries were sequenced at the Vincent J. Coates Genomics Sequencing Laboratory at

UC Berkeley on the Illumina HiSeq2000 platform (Illumina, Inc., San Diego, CA) to

generate 100 bp single-end reads.

Read processing and mapping

Reads were pre-processed, quality filtered and barcode sorted using the

Fastx_toolkit as described in Chitwood et al., 2013. To remove reads originating from

repeat-rich genomic regions, RESCAN sequencing reads were initially mapped to the

Sol Genomic Network's tomato repeat database using BWA (BWA parameters: -e 15 -

i10 -k 1 -l 25 -n 0.05) (we created the fasta file for this from the gff3 file available at

ftp://ftp.sgn.cornell.edu/genomes/Solanum_lycopersicum/

annotation/ITAG2.3_release/ITAG2.3_repeats.gff3). Reads not mapped to the repeat

database were extracted using bam2fastq program

(http://www.hudsonalpha.org/gsl/software/bam2fastq.php). Subsequently, these repeat-

filtered reads were mapped to the Heinz reference genome using the same BWA

parameters. Samtools (with the '–bq 1' option) was used to retain the reads that

mapped uniquely to the reference genome.

Polymorphism identification

We used SNPTools (https://github.com/mfcovington/SNPtools) v0.2.4 to identify

SNPs and INDELs between *Solanum lycopersicum* cv. M82 and *S. pennellii*. SNPTools

is a variant-detection, genotype-scoring, and visualization tool composed of Perl

modules, Perl scripts, and R scripts. The use of an earlier version of SNPTools, v0.1.5,

has been described elsewhere (Devisetty et al., 2014).

In brief, the SNPTools polymorphism identification pipeline consists of the

following steps. Putative Heinz:cv M82 and Heinz:*S. pennellii* polymorphisms were

identified using a pileup-comparison-based approach. Positions where the M82 and *S.*

pennellii alleles differ were retained and the Heinz-specific polymorphisms were

discarded. SNPTools also has a polymorphism noise reduction step. The cv M82 and *S.*

pennellii sequence alignments were genotyped at each putative polymorphism. For a

true cv M82:*S. pennellii* polymorphism, the large majority of reads (>90%) should match

the appropriate allele for both alignments. Putative polymorphisms that contradict this

expectation were discarded.

We performed an additional, non-SNPTools, polymorphism-filtering step to

remove outliers in regards to allele ratios and coverage within the BIL population. The

sequence alignment files from all of the BILs were merged together and genotyped

using the post-noise-reduction polymorphism database.

Polymorphisms were kept if the *S. pennellii* allele ratio was between the 25th and

75th percentile (relative to the current chromosome) and the non-parental allele ratio

was less than 0.025. Since a small number of positions had excessive coverage relative

the rest of the genome, we kept data only for polymorphisms with a maximum coverage

of 2,000 reads.

Detailed descriptions of the polymorphism identification pipeline, including the

code that was executed and genotype plots at different steps in the pipeline, are

available at: https://github.com/mfcovington/bil-paper/blob/master/docs/polymorphism-

identification.md.

Heterogeneous Hidden Markov Model (HMM)

M82, heterozygous, and *S. pennellii* genotype blocks in the BILs were inferred by

means of a Hidden Markov Model (HMM) implemented using the StochHMM C library

plus our own C++ code for the heterogeneous transition matrix (Lott and Korf, 2014).

The HMM may be eventually released as standalone software. Our HMM uses a

heterogeneous transition matrix in which the transition probability between states (i.e.,

M82, heterozygous, and *S. pennellii*) depends on the genetic distance between

adjacent SNPs. With the transition matrix of our HMM we in essence modeled

recombination between pairs of SNPs. Given the small number of crosses and the short

distance between SNPs, we made the simplifying assumption that recombination

between any two adjacent SNPs in one line of the population occurred only once (i.e., in

a single generation) or not at all. We tested recombination probability equations derived

from classic genetic map functions and found them to be adequate, but nevertheless

unstable in a number of lines with high noise and/or low sequence coverage. Hence, we

explored alternative recombination probability equations with simple parameterizations

that would approximate Haldane's and Kosambi's equations at large genetic distances

yet be more stable at short distances (i.e., less prone to switch states due local noise).

The cells of the transition matrix are calculated according to the following equation for

the probability of changing states:

1
$$\frac{1- e^{-d^2/2\sigma^2}}{2} \text{ (1)}$$

2   and its reciprocal for the probability of not changing states:

3
$$\frac{e^{-d^2/2\sigma^2}}{2} \text{ (2)},$$

where d is genetic distance in cM between a pair of adjacent SNPs and σ is a free

parameter (Supplemental Fig. S73). We empirically chose a value of 0.5 for σ for the

present study. We plan to make the σ parameter tunable, by means of the Baum-Welch

algorithm, in the training phase of the HMM in future versions of our method. Equation 1

has a low probability of transition at small genetic distances because at such distances

it has an opposite curvature to Haldane's and Kosambi's recombination probability

equations.

The prior probabilities for the hidden states are 0.88 : 0.10 : 0.02 for M82 : *S.*

pennellii : heterozygous, which are roughly the expected proportion of the states in the

BILs given their crossing scheme. The input SNP genotype probabilities were derived

by proportionally scaling the SNP likelihoods from bcftools so that they sum to one,

which implicitly assumes a uniform prior. Following previous work on genotyping HMMs

(Andolfatto et al., 2011), an emission error rate of 1% was assumed, which captures

several potential genotype error sources. The posterior decoding algorithm was used to

infer posterior probabilities for each SNP sampled in a given BIL. Following convention,

a posterior probability of 0.95 was used as the threshold for inferring a change in state

(i.e., genotype).

In order to estimate the posterior probability of ancestry of each genomic region

(i.e., which parent it came from), our HMM used the genotype probabilities at the

parentally ascertained and preselected SNPs and a heterogeneous transition model, in

which the transition probability between hidden states (i.e., genotypes) is a function of

the genetic distance between SNPs (Fig. 2). We did so in order to achieve high

resolution and precision of genotype boundary calls. Common work-arounds for dealing

with variable distances between SNPs within a homogeneous HMM are to bin SNPs

into equidistant windows (e.g., 1 Kb in size) that are assumed to be homogenous in

genotype, or to simply ignore the differences in distance and treat SNPs as if

equidistant. Both of these workarounds would have degraded the resolution and

precision of the inferred genotype boundaries.

BIL genotyping, plotting, and definition of bins

The BILs were genotyped using the Perl script 'extract+genotype_pileups.pl' from

SNPtools v0.2.4 (https://github.com/mfcovington/SNPtools/). BIL sequence alignments

were interrogated at the position of each cv M82 : *S. pennellii* polymorphism to create a

set of genotype files per BIL. Recorded within a genotype file is the number of reads

matching each allele for every polymorphism with read coverage on a chromosome.

To detect and fine-tune genotype bins (i.e., regions of a single parental genotype

within a BIL) and the boundaries between them, we used a set of Perl scripts from

detect-boundaries v0.6.1 (https://github.com/mfcovington/detect-boundaries/). The

'filter-snps.pl' script identifies bins and boundaries on a per BIL scale. The 'merge-

boundaries.pl' script takes the bin and boundary data for all of the individual BILs and

merges them to identify every sub-bin within the population and calculate related

boundary and bin statistics. The 'fine-tune-boundaries.pl' script provides an automated

approach for human verification of each boundary between bins of opposite parental

genotype for a BIL. This command-line tool displays color-coded genotype data

together with the currently-defined bin boundaries. Using shortcut keys, the operator

can quickly and easily approve or fine-tune a boundary (at which point, the next

boundary is instantly displayed for approval).

BIL genotype plots were created with the R script 'plot-genotype-and-bin.R',

which uses a BIL's genotype and bin boundaries files.

Phenotypic estimates used for mapping

BIL phenotypic estimates for plotting and QTL mapping were derived from

Gaussian linear mixed effect models fitted using the lme4 R package (Bates 2015). For

all traits a block random effect was used. The model used for flowering time had no

additional random effects. For leaf complexity traits and asymmetric leaflet EFD

principal components an additional plant level random effect was used to accommodate

pseudoreplicate measurements. For shape attribute traits and symmetric leaflet EFD

principal components the models included a leaflet type random effect nested within the

plant level effect. For asymmetric leaflet EFD principal components, the terminal leaflet

data was discarded (as these should be symmetrical or near so), and the absolute

values for the left and right lateral leaflets were treated equally as they are mirror

images of each other.

Model-fitted trait mean estimates were used for BIL trait plots (Supplemental

Figs. S4-S29) and for marginal regression QTL mapping. For QTL mapping by

SparseNet (see below) we generated predictions from the LMM fits such that

pseudoreplication was eliminated (i.e., generated a single value per plant) and then the

pseudoreplicate averaged residuals were added to the predictions.

Heritability and Repeatability estimates

Broad sense heritability and repeatability estimates for the leaf traits were

inferred by frequentist and bayesian random effect regressions using the R packages

lme4 and MCMCglmm (Hadfield, 2010; Bates et al., 2015), respectively. The

experimental design random effects (e.g., block, plant, etc.) used in this case were the

same as for the phenotypic estimates.

QTL mapping

<u>Marginal regression</u>

Each individual BIL was factored for the presence or absence of *S. pennellii*

introgressed DNA at a particular bin, and this process was repeated for each bin. For

marginal regression, a linear model was fitted between each trait as a function of the

presence for each bin. The multiple test adjustment for the significance values for each

bin with respect to a given trait was performed using the Benjamini–Hochberg method.

The threshold for adjusted significance value (q-value) was set at 0.001 to detect bins

significantly associated with a trait.

<u>SparseNet</u>

We mapped QTL by regressing bin genotypes on phenotypic traits using the

regularized regression method SparseNet (Mazumder et al., 2011), which uses the MC+

penalty family (an L0 to L1 family of penalties) to impose shrinkage and sparsity on the

number and magnitude of regression coefficients. For each trait we performed two QTL

mapping analyses with SparseNet: A) one in which we only considered non-epistatic or

monogenic (single bin) QTL (which we denote as the Additive Model), B) and a second

analysis in which we considered single locus QTL and all potential epistatic two-locus

QTL whose bin combination were observed in the BILs (which we denote as the

Epistatic Model). The Additive Model had 1,049 possible effects, i.e., equal to the

number of BINs. The Epistatic Model had 307,340 possible effects, 1,049 single loci

plus the 306,291 two-locus combinations observed in the population; out of the 549,676

possible two BIN combinations among the 1,049 BINs, 243,385 were not observed in

the BILs.

By using a penalized method such as SparseNet we were able to map each trait on all

bins and all bins plus all observed two bin combinations simultaneously. In order to

choose the tuning parameter values for SparseNet, for each trait we averaged the alpha

and gamma values from ten 6-fold cross-validations; the averaging was done to achieve

reproducible tuning of the model. Briefly, alpha controls the severity of the penalty, and

gamma controls the mixture between L0 and L1 penalties. Following cross-validation

the SparseNet model was fit to all the data of a given trait using the chosen values for

alpha and gamma. Because methods such as LASSO and SparseNet select a single

coefficient among a set of highly correlated coefficients, we took the set of BINs with a

correlation of 0.90 to a SparseNet selected BIN as a sort of empirical QTL interval within

which to search for candidate genes.

PROVEAN analysis

PROVEAN (PRotein Variation Effect ANalyzer) is a command-line tool to predict

the functional impact an amino acid substitution or indel has on a protein (Choi et al.,

2012). PROVEAN-Assembly-Line v0.3.2 (https://github.com/mfcovington/PROVEAN-

Assembly-Line) was used to run a set of Perl scripts to automate high-throughput bulk

PROVEAN analysis.

The 'find-aa-substitutions.pl' script was used to generate the amino acid

substitution files required to run PROVEAN. The input files for this script are SNP files,

a genomic FASTA reference file (S_lycopersicum_chromosomes.2.40.fa), and the

corresponding GFF3 annotation file (ITAG2.3_gene_models.gff3).

The 'run-provean.pl' script was used to run the PROVEAN tool in parallel for

every missense mutation, every nonsense mutation, and every mutation that caused the

loss of a STOP codon. The input files for this script are the amino acid substitution files

generated in the previous step and a CDS FASTA reference file

(ITAG2.3_cds_SHORTNAMES.fasta).

The 'filter-provean-results.pl' was used to filter PROVEAN results for a single

gene, a file containing a list of genes, or all genes within a genomic range.

Candidate gene Identification

A set of literature-curated leaf developmental genes was used as a reference for

the finding candidate genes within QTL bins for different leaf traits (Ichihashi et al.,

2014). We selected only those genes from the set that were either differentially

expressed between the two parents based on expression-QTL analysis of the IL

population or that showed amino acid changes between the two parents that were

predicted to be functionally significant by PROVEAN. The QTL bins were then scanned

for the presence of these genes. Following, we performed permutation analyses in the

R statistical environment to check whether bins associated with leaf traits were enriched

for the selected literature-curated leaf developmental genes. The data were permuted

10,000 times, and the results were considered significant if $< 5\%$ of the permuted

datasets equaled or exceeded the real data set.

IL eQTL enrichment among BIL epistatic QTL

We assessed whether IL *trans*-eQTL were enriched among the set of paired

regions that make up the BIL epistatic QTL using two eQTL clusters. One cluster was

enriched for leaf development GO terms and the other for photosynthesis GO terms.

Together these two clusters contained the majority of the literature-curated leaf

development genes used for the candidate gene search (Ichihashi et al., 2014; Ranjan

et al. in revision). The IRanges R/Bioconductor package (Lawrence et al., 2013) was

used to find the overlaps between the IL *trans*-eQTL and the BIL epistatic QTL. We

tested for IL *trans*-eQTL enrichment among BIL epistatic QTL by permutation using the

leaf development cluster (set 1), as well as this cluster plus the photosynthesis cluster

(set 2). The data were permuted 10,000 times, and the results were considered

significant if $< 5\%$ of the permuted datasets equaled or exceeded the real data set.

QTL confirmation and fine-mapping

The IRanges R/Bioconductor package (Lawrence et al., 2013) was used to find

overlaps between the IL phenotypic QTL and the BIL single locus QTL, and the results

were tabulated and plotted.

<u>**Sequence submission**</u>

The quality filtered, barcode-sorted and trimmed short read dataset, was deposited to

the NCBI Short Read Archive under accessions SRR##### - SRR##### (Bioproject

accession SRP#####)

<u>**Supplemental datasets and weblinks**</u>

Supplemental Weblink 1. Recombination boundary plots:

github.com/mfcovington/bil-paper/data/plots-with-boundaries/

Supplemental Weblink 2. Recombination boundary files:

github.com/mfcovington/bil-paper/data/boundaries/

Supplemental Dataset 1. Introgressions by BIL. Number and location of

introgressions organized by BILs.

Supplemental Dataset 2. Introgressions by chromosome. Number and location of introgressions organized by chromosome.

Supplemental Dataset 3: Bin locations and genotypes. Bin locations and genotypes of the BILs at each bin ordered by chromosome.

Supplemental Dataset 4: Annotated genes per BIN. Genes organized by bins and with functional annotations.

Supplemental Dataset 5: Marginal regression QTL mapping results. There is one table per trait. QTL are defined as bins significantly correlated to traits by marginal regression at a BH-adjusted p-value of less than 0.01.

Supplemental Dataset 6: SparseNet QTL mapping results. There is one table per trait, empty if there were no QTL found for a given trait. The SparseNet selected bin is shown for each QTL, along with its coordinates, the estimated regression coefficient, and the 0.95 and 0.9 bin correlation intervals. For the epistatic QTL this information there are two selected bins and two sets of correlated intervals.

Supplemental Dataset 7: Single locus QTL candidate genes. There is one table per trait. The SparseNet selected bin is shown, the ITAG gene number for the candidate genes, the gene coordinates, whether the gene has PROVEAN or *cis*-eQTL signal, and the closest Arabidopsis homolog of the candidate gene.

Supplemental Dataset 8: Epistatic QTL candidate genes. There is one table per trait. The SparseNet selected bins are shown, together with their 0.9 bin correlation intervals, and other data from the mapping results. For QTL with candidate genes, subsequent lines in the tables show: the bin that contains the candidate gene, its ITAG gene number, chromosome, gene coordinates, expression difference in the ILs with respect

to M82, the type of eQTL (*trans* vs *cis*), and the closest Arabidopsis homolog of the

candidate gene.

Supplemental dataset 9: IL-BIL fine-mapping tables. There is one table per trait

showing overlapping QTL between the IL and BIL results. For the IL QTL the bin and

sub-bin are shown, the bin coordinates, chromosome, slope (i.e., effect size), and

significance values. For the BIL QTL the SparseNet selected bin is shown for each QTL,

along with its coordinates, the estimated regression coefficient, and the 0.95 and 0.9 bin

correlation intervals. For the epistatic QTL this information there are two selected bins

and two sets of correlated intervals.

Acknowledgments

We thank Yaniv Brandvain and Paul Lott for discussions and assistance with early

versions of the HMM. We thank the Vincent J. Coates Genomics Sequencing

Laboratory at UC Berkeley (supported by NIH S10 Instrumentation Grants

S10RR029668 and S10RR027303), and computational resources/cyber infrastructure

provided by the iPlant Collaborative (www.iplantcollaborative.org), funded by the

National Science Foundation (Grant DBI-0735191).

Author's contributions

DHC, JNM and NRS conceived and designed the experiments. DF, AR, DHC, DW, YS

and LH performed the experiments. DF, AR, MFC, and DHC analyzed the data. DF

developed new statistical inference tools. DF, AR, IF, MFC, DHC, DZ, JNM contributed

reagent/materials/analysis tools. DF, AR, MFC, DHC, JNM, NRS wrote the paper. All

authors read and approved the final manuscript.

Figure and table legends:

Table 1. Per chromosome BIL introgression and bin counts.

Figure 1. Crossing scheme for the *S. pennellii* Backcrossed Introgression Lines.

Figure 2. Examples of hard to genotype lines. A. High noise. B. Low sequence

coverage. C. Heterozygous region with high noise. D. Narrow introgression

(chromosome 10). BILs 015 and 110 were resequenced to obtain cleaner data. The

boundaries obtained with the cleaner resequenced data did not differ from those shown

above, demonstrating the performance of the HMM. Plots of resequenced BILs 015 and

110 are at github.com/mfcovington/bil-paper.

Figure 3. Introgression and bin statistics. A. Distribution of introgressions across

bins. B. Introgression frequency per BIL. C. Introgression percentage per BIL. D.

Number of bins per chromosome. Panel A is plotted in genetic distance as

Supplemental Fig. S1.

Figure 4. Composite genotype map of the entire BIL population. See Supplemental

Fig. S2 for a version of this figure plotted by physical distance, and Supplemental Fig.

S3 for a version plotted with BILs clustered by each chromosome after removing

samples without an introgression.

Figure 5. QTL mapping results for additive and epistatic models. A. Primary

complexity. B. Total complexity. C. Aspect Ratio. D. Circularity. The bin selected by

SparseNet as a QTL is colored opaquely, and the nearby bins correlated to it at or

greater than 0.9 are taken as an approximate QTL interval and colored translucently.

Figure 6. *Punctate* fine-mapping. A. Adaxial leaf images showing presence and

absence of the "purple spot" phenotype. B. Comparison of the chromosome 10

introgressions of IL10-3 and BIL-228. C. Fine-mapping the *Punctate* locus using BILs;

purple-labeled BILs possess the "purple spot" phenotype, whereas black-labeled BILs

do not. D. Gene list of the fine-mapped region with candidate MYB tandem expansion

highlighted in purple.

Figure 7. Fine-mapping and confirmation of IL QTL using BILs. A. Flowering time,

showing the candidate gene *SP5G* in inset E. B. Total complexity. C. Circularity. The

BIL bin selected by SparseNet as a QTL is colored opaquely, and the nearby bins

correlated to it at or greater than 0.9 are taken as an approximate QTL interval and

colored translucently.

References

Andolfatto P, Davison D, Erezyilmaz D, Hu TT, Mast J, Sunayama-Morita T, Stern DL (2011)
Multiplexed shotgun genotyping for rapid and efficient genetic mapping. Genome Res **21**: 610–
617

Bates D, Mächler M, Bolker B, Walker S (2015) Fitting Linear Mixed-Effects Models Using lme4.
Journal of Statistical Software **67**: 1–48

Bharathan G, Goliber TE, Moore C, Kessler S, Pham T, Sinha NR (2002) Homologies in Leaf Form
Inferred from KNOXI Gene Expression During Development. Science **296**: 1858–1860

Carmel-Goren L, Liu YS, Lifschitz E, Zamir D (2003) The SELF-PRUNING gene family in tomato.
Plant Mol Biol **52**: 1215–1222

Carroll SB (2008) Evo-Devo and an Expanding Evolutionary Synthesis: A Genetic Theory of
Morphological Evolution. Cell **134**: 25–36

Chitwood DH, Headland LR, Filiault DL, Kumar R, Jiménez-Gómez JM, Schrager AV, Park DS,
Peng J, Sinha NR, Maloof JN (2012a) Native Environment Modulates Leaf Size and Response
to Simulated Foliar Shade across Wild Tomato Species. PLoS ONE **7**: e29570

Chitwood DH, Headland LR, Kumar R, Peng J, Maloof JN, Sinha NR (2012b) The Developmental
Trajectory of Leaflet Morphology in Wild Tomato Species. Plant Physiol **158**: 1230–1240

Chitwood DH, Kumar R, Headland LR, Ranjan A, Covington MF, Ichihashi Y, Fulop D, Jiménez-
Gómez JM, Peng J, Maloof JN, et al (2013) A Quantitative Genetic Basis for Leaf Morphology
in a Set of Precisely Defined Tomato Introgression Lines. Plant Cell **25**: 2465–2481

Chitwood DH, Ranjan A, Kumar R, Ichihashi Y, Zumstein K, Headland LR, Ostria-Gallardo E,
Aguilar-Martínez JA, Bush S, Carriedo L, et al (2014a) Resolving Distinct Genetic Regulators
of Tomato Leaf Shape within a Heteroblastic and Ontogenetic Context. Plant Cell **26**: 3616–3629

Chitwood DH, Ranjan A, Martinez CC, Headland LR, Thiem T, Kumar R, Covington MF, Hatcher
T, Naylor DT, Zimmerman S, et al (2014b) A Modern Ampelography: A Genetic Basis for
Leaf Shape and Venation Patterning in Grape. Plant Physiol **164**: 259–272

Choi Y, Sims GE, Murphy S, Miller JR, Chan AP (2012) Predicting the Functional Effect of Amino
Acid Substitutions and Indels. PLoS ONE **7**: e46688

Devisetty UK, Covington MF, Tat AV, Lekkala S, Maloof JN (2014) Polymorphism Identification and Improved Genome Annotation of Brassica rapa Through Deep RNA Sequencing. G3 **4**: 2065–2078

Doebley J, Wang R-L (1997) Genetics and the Evolution of Plant Form: An Example from Maize. Cold Spring Harb Symp Quant Biol **62**: 361–367

Efroni I, Eshed Y, Lifschitz E (2010) Morphogenesis of Simple and Compound Leaves: A Critical Review. Plant Cell **22**: 1019–1032

Eshed Y, Zamir D (1994) A genomic library of Lycopersicon pennellii in L. esculentum: A tool for fine mapping of genes. Euphytica **79**: 175–179

Fan P, Miller AM, Schilmiller AL, Liu X, Ofner I, Jones AD, Zamir D, Last RL (2016) In vitro reconstruction and analysis of evolutionary variation of the tomato acylsucrose metabolic network. PNAS **113**: E239–E248

Ferris KG, Rushton T, Greenlee AB, Toll K, Blackman BK, Willis JH (2015) Leaf shape evolution has a similar genetic architecture in three edaphic specialists within the Mimulus guttatus species complex. Ann Bot **116**: 213–223

Fournier-Level A, Cunff LL, Gomez C, Doligez A, Ageorges A, Roux C, Bertrand Y, Souquet J-M, Cheynier V, This P (2009) Quantitative Genetic Bases of Anthocyanin Variation in Grape (Vitis vinifera L. ssp. sativa) Berry: A Quantitative Trait Locus to Quantitative Trait Nucleotide Integrated Study. Genetics **183**: 1127–1139

Frary A, Nesbitt TC, Frary A, Grandillo S, Knaap E van der, Cong B, Liu J, Meller J, Elber R, Alpert KB, et al (2000) fw2.2: A Quantitative Trait Locus Key to the Evolution of Tomato Fruit Size. Science **289**: 85–88

Fridman E, Carrari F, Liu Y-S, Fernie AR, Zamir D (2004) Zooming In on a Quantitative Trait for Tomato Yield Using Interspecific Introgressions. Science **305**: 1786–1789

Fridman E, Pleban T, Zamir D (2000) A recombination hotspot delimits a wild-species quantitative trait locus for tomato sugar content to 484 bp within an invertase gene. PNAS **97**: 4718–4723

Goliber T, Kessler S, Chen JJ, Bharathan G, Sinha N (1999) Genetic, molecular, and morphological analysis of compound leaf development. Curr Top Dev Biol **43**: 259–290

Gonzalez A, Zhao M, Leavitt JM, Lloyd AM (2008) Regulation of the anthocyanin biosynthetic pathway by the TTG1/bHLH/Myb transcriptional complex in Arabidopsis seedlings. The Plant Journal **53**: 814–827

Hadfield J (2010) MCMC Methods for Multi-Response Generalized Linear Mixed Models: The MCMCglmm R Package. Journal of Statistical Software **33**: 1–22

Hoekstra HE, Coyne JA (2007) The Locus of Evolution: Evo Devo and the Genetics of Adaptation. Evolution **61**: 995–1016

Holtan HEE, Hake S (2003) Quantitative Trait Locus Analysis of Leaf Dissection in Tomato Using Lycopersicon pennellii Segmental Introgression Lines. Genetics **165**: 1541–1550

Ichihashi Y, Aguilar-Martínez JA, Farhi M, Chitwood DH, Kumar R, Millon LV, Peng J, Maloof JN, Sinha NR (2014) Evolutionary developmental transcriptomics reveals a gene network module regulating interspecific diversity in plant leaf shape. PNAS **111**: E2616–E2621

Iwata H, Ukai Y (2002) SHAPE: A Computer Program Package for Quantitative Evaluation of Biological Shapes Based on Elliptic Fourier Descriptors. J Hered **93**: 384–385

Jiang C, Wright RJ, Woo SS, DelMonte TA, Paterson AH (2000) QTL analysis of leaf morphology in tetraploid Gossypium (cotton). Theor Appl Genet **100**: 409–418

Jones CM, Rick CM, Adams D, Jernstedt J, Chetelat RT (2007) Genealogy and fine mapping of obscuravenosa, a gene affecting the distribution of chloroplasts in leaf veins, and evidence of selection during breeding of tomatoes (Lycopersicon esculentum; Solanaceae). Am J Bot **94**: 935–947

Kerstetter RA, Bollman K, Taylor RA, Bomblies K, Poethig RS (2001) KANADI regulates organ polarity in Arabidopsis. Nature **411**: 706–709

Keurentjes JJB, Bentsink L, Alonso-Blanco C, Hanhart CJ, Vries HB-D, Effgen S, Vreugdenhil D, Koornneef M (2007) Development of a Near-Isogenic Line Population of Arabidopsis thaliana and Comparison of Mapping Power With a Recombinant Inbred Line Population. Genetics **175**: 891–905

Kim M, McCormick S, Timmermans M, Sinha N (2003a) The expression domain of PHANTASTICA determines leaflet placement in compound leaves. Nature **424**: 438–443

Kim M, Pham T, Hamidi A, McCormick S, Kuzoff RK, Sinha N (2003b) Reduced leaf complexity in tomato wiry mutants suggests a role for PHAN and KNOX genes in generating compound leaves. Development **130**: 4405–4415

Kimura S, Koenig D, Kang J, Yoong FY, Sinha N (2008) Natural Variation in Leaf Morphology Results from Mutation of a Novel KNOX Gene. Current Biology **18**: 672–677

Koenig D, Bayer E, Kang J, Kuhlemeier C, Sinha N (2009) Auxin patterns Solanum lycopersicum leaf morphogenesis. Development **136**: 2997–3006

Koenig D, Jiménez-Gómez JM, Kimura S, Fulop D, Chitwood DH, Headland LR, Kumar R, Covington MF, Devisetty UK, Tat AV, et al (2013) Comparative transcriptomics reveals patterns of selection in domesticated and wild tomato. PNAS **110**: E2655–E2662

Koenig D, Sinha N (2010) Chapter Six - Evolution Of Leaf Shape: A Pattern Emerges. *In* MCP Timmermans, ed, Current Topics in Developmental Biology. Academic Press, pp 169–183

Koltai H, Bird DM (2000) Epistatic repression of PHANTASTICA and class 1 KNOTTED genes is uncoupled in tomato. The Plant Journal **22**: 455–459

Ku LX, Zhang J, Guo SL, Liu HY, Zhao RF, Chen YH (2012) Integrated multiple population analysis of leaf architecture traits in maize (Zea mays L.). J Exp Bot **63**: 261–274

Langlade NB, Feng X, Dransfield T, Copsey L, Hanna AI, Thébaud C, Bangham A, Hudson A, Coen E (2005) Evolution through genetically controlled allometry space. PNAS **102**: 10221–10226

Lawrence M, Huber W, Pagès H, Aboyoun P, Carlson M, Gentleman R, Morgan MT, Carey VJ (2013) Software for Computing and Annotating Genomic Ranges. PLoS Comput Biol **9**: e1003118

Liu YS, Zamir D Second generation L. pennellii introgression lines and the concept of bin mapping. Number 49-October 1999

Lott PC, Korf I (2014) StochHMM: a flexible hidden Markov model tool and C++ library. Bioinformatics **30**: 1625–1626

Mazumder R, Friedman JH, Hastie T (2011) SparseNet: Coordinate Descent With Nonconvex Penalties. Journal of the American Statistical Association **106**: 1125–1138

McConnell JR, Emery J, Eshed Y, Bao N, Bowman J, Barton MK (2001) Role of PHABULOSA and PHAVOLUTA in determining radial patterning in shoots. Nature **411**: 709–713

Monson-Miller J, Sanchez-Mendez DC, Fass J, Henry IM, Tai TH, Comai L (2012) Reference genome-independent assessment of mutation density using restriction enzyme-phased sequencing. BMC Genomics **13**: 72

Moyle LC (2008) Ecological and Evolutionary Genomics in the Wild Tomatoes (solanum Sect. Lycopersicon). Evolution **62**: 2995–3013

Muir CD, Pease JB, Moyle LC (2014) Quantitative Genetic Analysis Indicates Natural Selection on Leaf Phenotypes Across Wild Tomato Species (Solanum sect. Lycopersicon; Solanaceae). Genetics **198**: 1629–1643

Müller NA, Wijnen CL, Srinivasan A, Ryngajllo M, Ofner I, Lin T, Ranjan A, West D, Maloof JN, Sinha NR, et al (2016) Domestication selected for deceleration of the circadian clock in cultivated tomato. Nat Genet **48**: 89–93

Nath U, Crawford BCW, Carpenter R, Coen E (2003) Genetic Control of Surface Curvature. Science **299**: 1404–1407

Naz AA, Raman S, Martinez CC, Sinha NR, Schmitz G, Theres K (2013) Trifoliate encodes an MYB transcription factor that modulates leaf and shoot architecture in tomato. PNAS **110**: 2401–2406

Nicotra AB, Leigh A, Boyce CK, Jones CS, Niklas KJ, Royer DL, Tsukaya H (2011) The evolution and functional significance of leaf shape in the angiosperms. Funct Plant Biol **38**: 535–552

Ning J, Moghe GD, Leong B, Kim J, Ofner I, Wang Z, Adams C, Jones AD, Zamir D, Last RL (2015) A Feedback-Insensitive Isopropylmalate Synthase Affects Acylsugar Composition in Cultivated and Wild Tomato. Plant Physiol **169**: 1821–1835

Ori N, Cohen AR, Etzioni A, Brand A, Yanai O, Shleizer S, Menda N, Amsellem Z, Efroni I, Pekker I, et al (2007) Regulation of LANCEOLATE by miR319 is required for compound-leaf development in tomato. Nat Genet **39**: 787–791

Pérez-Pérez JM, Serrano-Cartagena J, Micol JL (2002) Genetic Analysis of Natural Variations in the Architecture of Arabidopsis thaliana Vegetative Leaves. Genetics **162**: 893–915

Ranjan A, Budke J, Rowland S, Chitwood DH, Kumar R, Ichihashi Y, Zumstein K, Maloof JN, Sinha NR eQTL in a Precisely Defined Tomato Introgression Population Reveal Genetic Regulation of Gene Expression Patterns Related to Physiological and Developmental Pathways. Plant Physiology In Revision:

Ranjan A, Ichihashi Y, Sinha NR (2012) The tomato genome: implications for plant breeding, genomics and evolution. Genome Biol **13**: 1–8

Rousseaux MC, Jones CM, Adams D, Chetelat R, Bennett A, Powell A (2005) QTL analysis of fruit antioxidants in tomato using Lycopersicon pennellii introgression lines. Theor Appl Genet **111**: 1396–1408

Schauer N, Semel Y, Balbo I, Steinfath M, Repsilber D, Selbig J, Pleban T, Zamir D, Fernie AR (2008) Mode of Inheritance of Primary Metabolic Traits in Tomato. Plant Cell **20**: 509–523

Schauer N, Semel Y, Roessner U, Gur A, Balbo I, Carrari F, Pleban T, Perez-Melis A, Bruedigam C, Kopka J, et al (2006) Comprehensive metabolic profiling and phenotyping of interspecific introgression lines for tomato improvement. Nat Biotech **24**: 447–454

Semel Y, Nissenbaum J, Menda N, Zinder M, Krieger U, Issman N, Pleban T, Lippman Z, Gur A, Zamir D (2006) Overdominant quantitative trait loci for yield and fitness in tomato. PNAS **103**: 12981–12986

Seymour DK, Filiault DL, Henry IM, Monson-Miller J, Ravi M, Pang A, Comai L, Chan SWL, Maloof JN (2012) Rapid creation of Arabidopsis doubled haploid lines for quantitative trait locus mapping. PNAS **109**: 4227–4232

Sharlach M, Dahlbeck D, Liu L, Chiu J, Jiménez-Gómez JM, Kimura S, Koenig D, Maloof JN, Sinha N, Minsavage GV, et al (2012) Fine genetic mapping of RXopJ4, a bacterial spot disease resistance locus from Solanum pennellii LA716. Theor Appl Genet **126**: 601–609

Steinhauser M-C, Steinhauser D, Gibon Y, Bolger M, Arrivault S, Usadel B, Zamir D, Fernie AR, Stitt M (2011) Identification of Enzyme Activity Quantitative Trait Loci in a Solanum lycopersicum × Solanum pennellii Introgression Line Population. Plant Physiol **157**: 998–1014

Stevens MA, Rick CM (1986) Genetics and breeding. *In* JG Atherton, J Rudich, eds, The Tomato Crop. Springer Netherlands, pp 35–109

Stevens R, Buret M, Duffé P, Garchery C, Baldet P, Rothan C, Causse M (2007) Candidate Genes and Quantitative Trait Loci Affecting Fruit Ascorbic Acid Content in Three Tomato Populations. Plant Physiol **143**: 1943–1953

Tian F, Bradbury PJ, Brown PJ, Hung H, Sun Q, Flint-Garcia S, Rocheford TR, McMullen MD, Holland JB, Buckler ES (2011) Genome-wide association study of leaf architecture in the maize nested association mapping population. Nat Genet **43**: 159–162

Timmermans MC, Hudson A, Becraft PW, Nelson T (1999) ROUGH SHEATH2: A Myb Protein That Represses knox Homeobox Genes in Maize Lateral Organ Primordia. Science **284**: 151–153

Townsley BT, Sinha NR (2012) A New Development: Evolving Concepts in Leaf Ontogeny. Annual
Review of Plant Biology **63**: 535–562

Tsiantis M, Hay A (2003) Comparative plant development: the time of the leaf? Nat Rev Genet **4**: 169–
180

Tsiantis M, Schneeberger R, Golz JF, Freeling M, Langdale JA (1999) The Maize rough sheath2
Gene and Leaf Development Programs in Monocot and Dicot Plants. Science **284**: 154–156

Tsukaya H (2013) Leaf Development. The Arabidopsis Book e0163

Waites R, Selvadurai HRN, Oliver IR, Hudson A (1998) The PHANTASTICA Gene Encodes a MYB
Transcription Factor Involved in Growth and Dorsoventrality of Lateral Organs in Antirrhinum.
Cell **93**: 779–789

Westerbergh A, Doebley J (2002) Morphological Traits Defining Species Differences in Wild Relatives
of Maize Are Controlled by Multiple Quantitative Trait Loci. Evolution **56**: 273–283

Xiao H, Jiang N, Schaffner E, Stockinger EJ, Knaap E van der (2008) A Retrotransposon-Mediated
Gene Duplication Underlies Morphological Variation of Tomato Fruit. Science **319**: 1527–1530

Chromosome	No. BILs with introgression	No. introgressions	No. bins
1	53	54	62
2	91	99	82
3	57	69	55
4	94	101	91
5	145	171	156
6	93	103	86
7	72	78	50
8	97	103	69
9	84	103	97
10	109	126	98
11	208	236	128
12	104	129	75

Table 1. Per chromosome BIL introgression and bin counts.

S. lycopersicum cv. M82 X *S. pennellii*

S. lycopersicum cv. M82 X F$_1$

BC$_1$

S. lycopersicum cv. M82 X   Backcrossing to M82 for
                                      2 or 3 generations

BC$_{2/3}$

                                      Selfing for 4 generations

BC$_{2/3}$S$_4$ lines

[BIL population, 545 individuals]

Figure 1. Crossing scheme for the *S. pennellii* Backcrossed Introgression Lines.

Figure 2. Examples of hard to genotype lines. A. High noise. B. Low sequence coverage. C. Heterozygous region with high noise. D. Narrow introgression (chromosome 10). BILs 015 and 110 were resequenced to obtain cleaner data. The boundaries obtained with the cleaner resequenced data did not differ from those shown above, demonstrating the performance of the HMM. Plots of resequenced BILs 015 and 110 are at github.com/mfcovington/bil-paper.

Figure 3. Introgression and bin statistics. A. Distribution of introgressions across bins. B. Introgression frequency per BIL. C. Introgression percentage per BIL. D. Number of bins per chromosome. Panel A is plotted in genetic distance as Supplemental Fig. S1.

Figure 4. Composite genotype map of the entire BIL population. See Supplemental Fig. S2 for a version of this figure plotted by physical distance, and Supplemental Fig. S3 for a version plotted with BILs clustered by each chromosome after removing samples without an introgression.

Figure 5. QTL mapping results for additive and epistatic models. A. Primary complexity. B. Total complexity. C. Aspect Ratio. D. Circularity. The bin selected by SparseNet as a QTL is colored opaquely, and the nearby bins correlated to it at or greater than 0.9 are taken as an approximate QTL interval and colored translucently.

Figure 6. *Punctate* fine-mapping. A. Adaxial leaf images showing presence and absence of the "purple spot" phenotype. B. Comparison of the chromosome 10 introgressions of IL10-3 and BIL-228. C. Fine-mapping the *Punctate* locus using BILs; purple-labeled BILs possess the "purple spot" phenotype, whereas black-labeled BILs do not. D. Gene list of the fine-mapped region with candidate MYB tandem expansion highlighted in purple.

Figure 7. Fine-mapping and confirmation of IL QTL using BILs. A. Flowering time, showing the candidate gene *SP5G* in inset E. B. Total complexity. C. Circularity. The BIL bin selected by SparseNet as a QTL is colored opaquely, and the nearby bins correlated to it at or greater than 0.9 are taken as an approximate QTL interval and colored translucently.