Focused Ultrasound Enhances Decision-Making in Monkeys

Matthew E. Downs[1], Amanda Buch[1], Maria Eleni Karakatsani[1], Tobias Teichert[3], Carlos Sierra[1], Shangshang Chen[3], Elisa E. Konofagou[1,2], Vincent P Ferrera[3,4],

1. Department of Biomedical Engineering, Columbia University
2. Department of Radiology, Columbia University
3. Department of Neuroscience, Columbia University
4. Department of Psychiatry, Columbia University

Abstract

Noninvasive brain stimulation using ultrasound has many potential applications as a research and clinical tool. Here, we investigated the effect of focused ultrasound (FUS) with systemically administered microbubbles on perceptual decision-making behavior in monkeys. We targeted FUS to the putamen in one hemisphere to open the blood-brain barrier, and then tested behavioral performance 3-4 hours later. On days when the monkeys were treated with FUS, their decisions were faster and more accurate than days without sonication. The performance improvement was greater for responses made with the hand contralateral to the treated hemisphere. FUS also enhanced the effect of a low dose of haloperidol. The results suggest that a two-minute application of FUS can have a sustained impact on cognitive performance, and can increase the efficacy of psychoactive medications.

Key words

Ultrasound, stimulation, putamen, accuracy, response time.

Introduction

Brain stimulation is an essential tool for establishing causal brain-behavior relationships, mapping brain circuits, and treating neurological disorders. Current methods are either invasive (electrical or chemical stimulation, and optogenetics) or have limited penetrability (TMS) or localizability (TDCS) (Borchers et al. 2012; Calvo et al. 2006; Dubuisson et al. 1977; Kobyashi et al. 2003; Miller 1965; Nitsche et al. 2003). Focused ultrasound (FUS) is emerging as a non-invasive technology capable of penetrating the skull and meninges to reach deep brain structures. FUS with systemically administered microbubbles has been shown to open the blood-brain barrier (BBB) in various animal models, and may also directly modulate neural activity (Marquet et al. 2011; Tung et al. 2011; Downs et al. 2015; Chu et al. 2015; McDannold et al. 2006). A few pioneering studies in monkeys (Deffieux et al.2013) and humans (Hameroff et al.2013; Legon et al. 2014) have provided evidence that FUS alone can be used to modify perception and behavior. Deffieux et al. found that FUS can increase the latency of antisaccades in monkeys. Tactile discrimination was enhanced during FUS stimulation of the somatosensory cortex in human subjects (Legon et al.2014), while overall mood improved when the frontal-temporal cortex was stimulated with FUS (Hameroff et al.2013). Further investigation using different species, brain targets, and behavioral tasks is warranted to establish the effectiveness and range of applications for this approach.

Here, we trained monkeys to perform a common perceptual decision-making task that

involves the detection of coherent visual motion (Lappin & Bell, 1976). The advantage of this

task is that it allows quantitative measures of response time and accuracy. Such measures have

been used to develop sophisticated computational models of decision-making in both humans

and monkeys (Luce 1986; Ratcliff 1978). Electrophysiological studies point to a critical role of

the striatum (caudate and putamen) in similar tasks (Ding & Gold, 2013). We used the coherent

motion task to investigate the effect of FUS on decision-making and motor performance.

Rhesus monkeys were treated with FUS and microbubbles to open the BBB 3-4 hours prior to

behavioral testing. FUS was targeted to the putamen, a part of the basal ganglia involved in

cognition, reward, and motor control. This study also investigated the interaction of FUS with a

low dose of haloperidol as this technique could be used to non-invasively facilitate the drug

effects, or to deliver drugs that cannot readily cross the intact BBB.

On days when the monkeys received the FUS treatment, their decisions were faster and

more accurate, particularly when responses were made with the contralateral hand. A threshold

dose of haloperidol also reduced response time, but impaired accuracy. FUS enhanced these

drug effects. These results indicate that FUS can be used alone or in combination with

psychoactive drugs to enhance or modify cognitive performance.

Results

<u>Effects of FUS on blood-brain barrier</u>

The BBB was successfully opened in the putamen region of the basal ganglia for all FUS

procedures. In Fig. 1, the red/yellow areas indicate where the contrast agent was able to pass

through the BBB, indicating successful BBB opening. The blue shaded regions indicate the

region targeted by the FUS transducer. All openings achieved within this study fell within the

targeted region and no untargeted BBB openings were observed. No damage from the FUS

procedure was detected; the T2-weighted MRI and susceptibility weighted imaging scans did

not display any hyper- or hypointense voxels in the targeted regions, which could indicate

edema (Supplementary Figures 1 & 2).

Figure 1. Contrast-enhanced (gadodiamide) MRI showing target of FUS in putamen of a monkey.
Top row shows sagittal, coronal and horizontal slices through the brain of monkey N. Blue oval
indicates the targeted region. Red and orange voxels indicate BBB opening. Bottom row shows the
same for monkey O.

<u>Effects of FUS on decision-making behavior</u>

The two monkeys performed 31 behavioral sessions (16 for monkey N, 15 for monkey O) of the

coherent motion detection task. N completed an average of 1385 trials per session (22,154 total

trials), while O averaged 931 trials (13,960 total). Behavior was quantified in terms of response

time and accuracy. Response time was measured as the interval between motion

stimulus/target onset and the first touch. Decision accuracy was measured as the percent

correct choices relative to total correct and incorrect responses. Trials in which the monkey

failed to respond were

disregarded. Results for the two

monkeys were qualitatively

similar, except that monkey N (the

younger of the pair) tended to

respond faster and more

accurately overall. The results

were therefore combined across

subjects after applying a

correction for overall response

time and accuracy (see Methods).

97   Figs. 2a and 3 summarize

Figure 2. Effects of motion coherence, reward, sonication, and haloperidol on response time (RT) for single target trials. A) Response time for single target trials as a function of motion coherence, offered reward, and sonication. Error bars omitted for clarity. B) RT averaged over coherence levels. Each pair of circles connected with a line compares average RT during haloperidol sessions (black border) with saline sessions (grey border). Error bars represent ±1 s.e.m. Legend applies to both subplots.

the behavioral results for all 31 sessions, including those with and without sonication and with

haloperidol or saline (effects of drug and sonication are considered separately below.) We first

consider performance on trials with only one target. The monkeys were 100% correct on these

trials because there was only one choice and trials without a response were not counted; hence,

only response times were analyzed. Statistical results (ANOVA and GLM) are given in

Supplementary Table 1 and main effects are plotted in Fig. 2a. Mean response time was faster

for large reward trials than small rewards. There was also an effect of motion strength even

though the motion stimulus was irrelevant; response times were faster on trials with high

coherence stimuli and increased with decreasing coherence. Haloperidol ("drug") tended to

shorten response time across all conditions (Supplementary Table 1 and Fig. 2b; symbols with

black borders are from haloperidol session, symbols with gray borders are from saline

sessions). The effect of haloperidol was not significant in sessions without sonication

(Supplementary Table 2.)

Figure 3. Effects of motion coherence, reward, and sonication on response time and accuracy for choice (2-target) trials. A) Accuracy (percent correct) for two-target trials. B) Response time for two-target trials. C) Accuracy vs. response time for two-target trials. Dashed horizontal line is 75% correct level. Legend in A applies to all subpanels. Error bars are omitted for clarity.

Sonication did not have a significant main effect on response time for all sessions

(Supplementary Table 1), but did have a significant effect for sessions without haloperidol

(Supplementary Table 3.) Furthermore, there was a significant interaction of sonication and drug

(Supplementary Table 1.) Post-hoc analysis showed that the effect of haloperidol was significant

only on days with sonication (Supplementary Table 4). Haloperidol-associated reduction of

response time on sonication days appeared to be greater for responses with the ipsilateral than

contralateral hand (Fig. 2b), possibly because responses with the contralateral hand were

already as fast as possible (floor effect).

For trials with two targets, both performance accuracy and response time were

analyzed. Accuracy improved with increasing motion coherence (Supplementary Table 5 and

Fig. 3a) and response times were reduced (Supplementary Table 1 and Fig. 3a), as shown in

previous studies (Roitman & Shadlen 2002). Offered reward size was associated with increased

response time, i.e. the monkeys were slower to respond when there was a larger reward at

stake (Supplementary Table 1). This was in contrast to their behavior on single-target trials

where responses tended to be faster on large reward trials. Larger offered reward size was

associated with slightly better performance accuracy, particularly on days without sonication

(Supplementary Table 5.)

Sonication was associated with slightly improved performance accuracy (Supplementary

Table 5 and Fig 3a). Sonication also had significant effects on response time (Supplementary

Table 1). These effects were significant when considering all sessions (Supplementary Table 1:

response time, Supplementary Table 5: accuracy), or only sessions without haloperidol

(Supplementary Table 3: response time, Supplementary Table 5: accuracy.) Sonication was

associated with faster response times for large rewards (Fig. 3b). For small rewards, sonication

sped up responses with the contralateral hand but slowed responses for the ipsilateral hand

(Fig. 3b).

Accuracy is plotted against response time in Fig. 3c to show the amount of time taken to

reach a given level of performance. This can be taken as a measure of decision-making

efficiency. For small rewards, monkeys achieved 75% correct performance in less time than for

large rewards; i.e. when a large

reward was at stake, their

responses were slowed even

though this gained them little in

accuracy. Sonication alone made

the monkeys more efficient for

large rewards with the contralateral

hand and less efficient for small

rewards with the ipsilateral hand.

Haloperidol also had

significant effects on accuracy and

Figure 4. Effects of sonication and haloperidol on performance for two-target trials. A) Effects on accuracy. B) Effects on response time. Error bars represent ±1 s.e.m. Black borders are sessions with haloperidol, grey borders are saline controls. All conventions same as **Fig. 3**. Data in each column are plotted separately for each coherence level.

response time compared to saline controls (Supplementary Tables 1 and 5). Haloperidol was

generally associated with higher accuracy on trials with small rewards, but had mixed or

equivocal effects on trials with large rewards (Fig. 4a). Sonication enhanced the negative

effects of haloperidol on accuracy (Supplementary Table 6.) Haloperidol was associated with

faster response times for both large and small rewards, with or without sonication (Fig. 4b).

However, sonication and drug had a significant interaction for response times (Supplementary

Table 1.) Sonication and haloperidol were both associated with reduced response times and

their interaction was strongest for the ipsilateral hand (Supplementary Table 4), though

response times were fastest overall for the contralateral hand (Fig. 4b).

Overall, sonication tends to improve accuracy and shorten response time, while

haloperidol speeds responses but reduces accuracy. When used in combination, sonication

enhances the effects of haloperidol.

164   Discussion

Targeting FUS to the putamen of monkeys resulted in significant improvements in decision-

making performance. Monkeys responded faster and more accurately when tested on days with

sonication than on days without. The physiological and psychological mechanisms underlying

this effect are not known. The effects depended on the hand used to respond and are therefore

unlikely to be due to general arousal. The effects also depended on reward size. When

monkeys had to make a choice and a large reward was available, their response times in the

absence of sonication were up to 200 ms slower than on small reward trials. Sonication reduced

this difference to 100 ms while slightly improving accuracy. Thus, sonication appears to have

improved the efficiency of decision-making. The performance enhancement was found even

though animals were tested 3-4 hours after sonication, suggesting that there may be a

persistent effect on the activity or responsiveness of putamen neurons, which, in turn, may be

due to a direct effect of ultrasound or an indirect effect of opening the blood-brain barrier.

Further experiments are needed to ascertain the temporal window within which performance

improvements are obtained. Such experiments should be done by sonicating subjects while

they are alert (Downs et al. 2015) to avoid any confounding effects of anesthesia.

Sonication also enhanced the effect of a low dose of haloperidol, a D2 dopamine

antagonist, given 5 minutes prior to behavioral testing. Previous studies of the effects of

haloperidol on response times have reported mixed results depending on species, task and

dosage (Brockel et al, 1995; Blokland & Honig 1999; Kern et al. 1998.) In the current study, low

dose haloperidol tended to shorten response time and reduce decision accuracy. Both effects

were enhanced by sonication. This result indicates that FUS can be used to significantly

enhance dopaminergic medications to modulate cognitive performance. The results also

demonstrate that the systemic dose of a drug necessary to achieve a desired pharmacological

effect can be reduced by increasing BBB permeability through the application of FUS to a

targeted brain region, even if the drug in question readily crosses the BBB. This would allow for

smaller systemic doses, and thus reduction of potential side effects of already available drugs

for therapies to treat neurological diseases and disorders.

There are few previous studies investigating the effect of FUS without BBB opening on

alert subjects completing behavioral tasks. Deffieux and colleagues applied FUS to monkeys

performing an antisaccade task by targeting the left frontal eye field (FEF) and the premotor

cortex (Deffieux et al, 2013). Ipsilateral antisaccade latencies were significantly slowed while

targeting the FEF but not the premotor cortex. Two other groups investigated the effects of FUS

on human subjects (Legon et al, 2014; Hameroff et al, 2013). Subjects tested by Legon et al.

exhibited enhanced sensitivity to the frequency of air puffs and discrimination at a two-point

touch test while FUS was applied to their somatosensory cortex. FUS was applied to the frontal-

temporal cortex in subjects of the Hameroff et al. study and unlike the other two studies with

simultaneous/immediate behavioral testing, results were determined 10 and 40 minutes after

application. Subjects reported a significant improvement on the Global Affect test, as well as

slightly reduced pain levels 40 minutes after the application of FUS. These studies demonstrate

that FUS is capable of affecting the function of the brain depending on the targeting area, while

the Hameroff et al. study shows the effects could be time sensitive. A key difference from the

current study is that in the aforementioned studies the BBB remained undisrupted in the

targeted region to the knowledge of the experimenters.

Recently, our group applied the FUS BBB opening procedure to awake, behaving

monkeys performing a reaching task with variable reward magnitude (Downs et al. 2015). That

study found a slight increase in response time as well as a significant improvement in the

accuracy of reaching to visual stimuli during a 2-minute application of FUS and throughout the

remaining 2 hours of behavioral testing. Another group, Chu et al, investigated the effects of

BBB opening via FUS opening on somatosensory evoked potentials (SSEPs) and blood-

oxygen-level dependent (BOLD) responses when targeting the left primary somatosensory

cortex in anesthetized rats (Chu et al.2015). Results showed both a decrease in SSEP and

BOLD signals within 10 minutes after finishing the FUS procedure. Their results highlighted the

impact of sonication parameters utilized, as lower acoustic pressures resulted in little to no

neurological effect, while higher acoustic pressures created sustained neurological effects. Our

study utilized an acoustic pressure found to be safe during prior studies conducted within our

lab, which was greater than the pressure used by Chu and colleagues. The exact mechanisms

behind the excitation or inhibition of neurons via FUS is currently unknown, but a current theory

is that mechanical forces emitted by the transducer during sonication affect mechanoreceptors

in the cell membrane (Ostrow et al.2011; Wahab et al.2012; Tyler et al.2008). However, this

explanation seems to be limited to cases in which the sonication is applied simultaneously

during the behavioral testing. Our results, along with the studies conducted by the Hameroff and

Chu studies, demonstrate that the effects of FUS sonication can persist after the time of

application. Further studies plan to determine the optimal time after FUS application to open the

BBB for maximal behavior modulation. Understanding the relationship between treatment time

and behavioral effects will help distinguish the mechanical effect of the sonication from the other

potential neurological effects of the BBB being opened at the target region.

In conclusion, opening the BBB via FUS with microbubbles can have a significant effect on the

behavioral responses of monkeys 3-4 hours after the end of the sonication. The BBB opening

also facilitated the delivery of a low dose of haloperidol demonstrating that therapeutic doses of

a drug can be reduced to mitigate the potential side-effects after opening the BBB at the target

region for therapy. Overall, our results demonstrate the potential for FUS BBB opening to

enhance behavioral performance in monkeys.

Methods

The procedures with monkeys were approved by the Institutional Animal Care and Use

Committees (IACUC) of Columbia University and the New York State Psychiatric Institute

(NYSPI). Two adult male Macaca mulatta (N, O) were used in all experiments (9 and 20 years

old, 5.5 and 9.5 kg). Monkeys were provided daily rations of vitamin enriched dry primate

biscuits, as well as enrichment toys and allowed access to play modules. Monkeys were trained

using operant conditioning to perform a decision-making task. On behavioral testing days,

monkeys performed the task for fluid reward until satiated. After behavioral testing, Monkeys

were given a fruit treat (banana, apple, or orange). On days when behavioral testing was not

conducted, monkeys were given a liter of water.

Focused Ultrasound and Drug Delivery

On selected days, monkeys received a FUS with microbubble treatment 3-4 hours prior to

behavioral testing. For the FUS procedures, subjects were sedated with ketamine (10 mg/kg)

and atropine (0.04 mg/kg) and placed into a stereotaxic positioning frame under general

anesthesia (isoflurane 1-2%) to ensure accurate targeting. Microbubbles (4-5 um, in-house

prepared, Feshitan et al. 2009) were administered intravenously at the onset of the FUS

application (single element transducer, 500 kHz, 400 kPa, 10 ms pulse length, 120 second

duration; H-107, Sonic Concepts, WA, USA). The putamen region of the basal ganglia was

targeted for all experiments. Throughout the procedure, the monkeys' vital signs were

continuously monitored (heart rate, SPO2, mean arterial pressure, respiratory rate and end tidal

CO_2). After the FUS procedure there was a 3 to 4 hour recovery period allowing the monkeys to

fully recover from anesthesia.

Haloperidol, a D2 antagonist (R&D Systems, Inc., Minneapolis, MN), was used to

augment neuromodulation. Haloperidol powder was dissolved in saline and titrated to the

concentration of 0.01mg/kg. On selected days, before the task began, monkeys were

administered either saline or haloperidol (0.01mg/kg) intramuscularly. The injection was given 5

minutes prior to the start of behavioral testing. The threshold dose of haloperidol was

determined as the maximum dose that had a minimal effect on behavioral results when the BBB

was intact. The timing of events during the FUS procedure, recovery, drug injection and

behavioral testing is shown in Fig. 5a.

MRI Analysis

One day after the FUS procedure, BBB opening and safety of the procedure was verified with

contrast enhanced T1-weighted as well as T2- weighted MRI and susceptibility weighted

imaging scans respectively. All MRI scans (3T, Philips Medical Systems, MA, USA) were

acquired 36 hours after the FUS procedure. T2-weighted (TR = 10ms, TE = 27ms, flip angle =

90°, spatial resolution = 400 x 400 µm2, slice thickness = 2mm with no interslice gap) and

susceptibility-weighted image (TR = 19ms, TE = 27ms, flip angle = 15°, spatial resolution = 400

x 400 µm2, slice thickness = 1 mm with no interslice gap) scans were used to verify the safety

of the procedure. Contrast enhanced T1-weighted (TR = 19ms, TE = 27ms, flip angle = 15°,

spatial resolution = 400 x 400 µm^2, slice thickness = 1 mm with no interslice gap) scans were

acquired 30 minutes after IV administration of 0.2ml/kg gadodiamide (Omniscan®, 573.66 DA,

GE, Healthcare, Princeton, NY, USA). Gadodiamide was selected as it does not cross the intact

BBB. All acquired scans were aligned with a previously acquired stereotactically aligned

structural T1-weighted MRI scan to verify opening in the targeted region (Marquet et al, 2015).

The contrast enhanced T1-weighted scans were then post processed to quantify the volume of

opening. This process has been thoroughly discussed elsewhere (Downs et al. 2015).

Behavioral Testing

Monkeys sat in a custom-made polycarbonate primate chair that allowed them to reach out to

visual stimuli presented on a 20-inch LCD touchscreen monitor (NEC 2010x with 3M SC4

resistive touchscreen) placed directly in front of the chair. The resolution of the LCD was 1280

horizontal x 1024 vertical pixels with a refresh rate of 60 Hz. The touchscreen had a resolution

of 1024 x 1024 pixels and a sampling rate of 60 Hz. The primate chair incorporated a

polycarbonate midline divider so that stimuli presented on the right side of the touchscreen

could only be reached by the right hand, and likewise for the left side. The viewing distance was

12 inches. Behavior was reinforced with drops of fluid delivered by a juice tube mounted on the

chair. The monkeys were free to move their head and eyes, though they tended to face directly

forward so that they could continuously lick the juice tube in anticipation of the reward.

The behavioral task was presented as discrete trials lasting roughly 5 seconds each.

Each trial began with a cue stimulus presented on the left or right side of the monitor (Fig. 5b,

"Cue"). The cue was a vertically or horizontally oriented yellow bar. The monkey touched the

cue with the corresponding hand to initiate the trial. After a short delay, the cue was replaced by

a random dot motion stimulus (Fig 5b, "Choice.") The motion stimulus consisted of 100 dots

moving within a circular aperture. Some of the dots moved in random directions while others

moved coherently in a single direction. The coherent direction, either leftward or rightward,

varied from trial to trial. The proportion of coherently moving dots defined the motion strength.

The strength of the motion stimulus (aka motion coherence) varied from 0 to 0.7 in steps of 0.1.

A particular coherence level was presented randomly on each trial. The motion stimulus was

flanked on either side by two target stimuli that appeared simultaneously with the motion

stimulus. The target stimuli were yellow bars that had the same orientation as the cue. The

direction of the coherent dots indicated which target would be rewarded. The monkey was

reinforced with drops of water for touching the appropriate target (Fig 5b, "Reward.") There was

no punishment for incorrect responses or failure to respond. There was no signal instructing the

monkey when to respond; rather, he was allowed to touch at any time after the motion stimulus

and targets appeared.

There were two reward conditions: small offered reward (1 drop of water) and large

offered reward (5 drops). Offered reward level on each trial was signaled by the orientation of

the cue and target stimuli. Horizontal indicated large reward, vertical indicated small reward.

One seventh of the trials were controls that were identical to the other trials except that

the target for the incorrect response was not presented. On these trials, no decision was

required; the monkey could ignore the motion stimulus and simply touch the correct target to

receive a reward.

Figure 5. Behavioral task and experimental timeline. A) Timeline of sonication and behavioral testing. B) Decision task sequence. The monkey initiates a trial by touching the cue. A random dot motion stimulus appeared moving to the left or right, flanked by two targets. The monkey touched the target toward which the dots were moving to receive a reward. Stimuli were displayed on the right or left of the screen. A physical barrier forced the monkey to respond with the corresponding hand. Only the yellow bars and dot stimulus were visible to the monkey, not the blue arrow, dotted line, or hand, which are used to indicate the motion of the dots, the physical barrier separating the two halves of the screen, and the manual response, respectively.

The complete task design thus had the following variables: display side (left or right),

cue/target orientation (vertical or horizontal, corresponding to small and large reward), motion

direction (left or right), motion coherence (0.0 to 0.7), and number of targets (1 or 2). This

resulted in a balanced design comprising 48 conditions per block of trials. All conditions were

randomly interleaved.

<u>Statistics</u>

Quantitative analyses were performed using Matlab 8.3 with the Statistics 9.0 toolbox

(Mathworks, Natick MA.) Response times were analyzed with multivariate ANOVA and

generalized linear model regression. Performance accuracy or outcome (correct, incorrect) was

analyzed with multivariate ANOVA and logistic regression. The explanatory variables used in all

analyses were: motion coherence (0 to 0.7, 8 levels), offered reward (1 or 5 drops), presence of

sonication, sonicated hemisphere (ispsilateral or contralateral to responding hand), and drug

treatment (saline or haloperidol).

Response times were normalized by subtracting from the response time for each trial the

difference between the average overall response time for each monkey and the average across

both monkeys (the size of this correction was +/-55 ms, i.e. all of N's response times were

decreased by 55 ms, while O's were increased by the same amount). Performance accuracy

(percent correct) was similarly normalized by calculating the correction needed to equalize the

accuracy at 0 coherence across monkeys (this correction was +/-2.2%), and adjusting the

accuracy at all coherences by that amount. The accuracy correction was not applied for logistic

regression analyses in which the dependent variable was trial-by-trial success or failure.

References

Blokland A, Honig W. (1999) Intra-striatal haloperidol and scopolamine injections: effects on

choice reaction time performance in rats. Eur Neuropsychopharmacol. 9(6):523-31.

Brockel BJ, Fowler SC. (1995) Effects of chronic haloperidol on reaction time and errors in a

sustained attention task: partial reversal by anticholinergics and by amphetamine. J Pharmacol

Exp Ther. 275(3):1090-8.

Borchers, S., Himmelbach, M., Logothetis, N. & Karnath, H.-O. Direct electrical stimulation of

human cortex - the gold standard for mapping brain functions? *Nat. Rev. Neurosci.* **13,** 63–70

(2012).

Calvo, F. & Coimbra, N. C. Interactions between opioid-peptides-containing pathways and

GABAA-receptors-mediated systems modulate panic-like-induced behaviors elicited by electric

and chemical stimulation of the inferior colliculus. *Brain Res.* **1104,** 92–102 (2006).

Deffieux T, Younan Y, Wattiez N, Tanter M, Pouget P, Aubry JF (2013) Low-intensity focused

ultrasound modulates monkey visuomotor behavior. Curr Biol. 23(23):2430-3.

Ding L, Gold JI (2013) The basal ganglia's contributions to perceptual decision making. Neuron

79(4):640-9.

Downs ME, Buch A, Karakatsani ME, Konofagou EE, Ferrera VP (2015) Blood-brain barrier

opening in behaving non-human primates via focused ultrasound with systemically administered

microbubbles. Nature Scientific Reports 5:15076.

Dubuisson, D. & Dennis, S. G. The formalin test: A quantitative study of the analgesic effects of

morphine, meperidine, and brain stem stimulation in rats and cats. *Pain* **4,** 161–174 (1977).

Feng S, Holmes P, Rorie A, Newsome WT. (2009) Can monkeys choose optimally when faced

with noisy stimuli and unequal rewards? PLoS Comput Biol. 5(2):e1000284.

Hameroff, S. et al. Transcranial ultrasound (TUS) effects on mental states: A pilot study. Brain

Stimul. 6, 409–415 (2013).

Hikosaka, O. Basal ganglia mechanisms of reward-oriented eye movement. Ann. N. Y. Acad.

Sci. 1104, 229–249 (2007).

Hikosaka, O., Nakamura, K. & Nakahara, H. Basal ganglia orient eyes to reward. J.

Neurophysiol. 95, 567–584 (2006).

Ibsen S, Tong A, Schutt C, Esener S, Chalasani SH (2015) Sonogenetics is a non-invasive

approach to activating neurons inn Caenorhabditis Elegans. Nature Communications. 6:8264.

Kern RS, Green MF, Marshall BD Jr, Wirshing WC, Wirshing D, McGurk S, Marder SR, Mintz J

(1998) Risperidone vs. haloperidol on reaction time, manual dexterity, and motor learning in

treatment-resistant schizophrenia patients. Biol Psychiatry. 1998 Oct 15;44(8):726-32.

King RI, Brown JR, Newsome WT, Butts Pauly K (2012) Effective parameters for ultrasound-

induced in vivo neurostimulation. Ultrasound in Med. & Biol.

Kobayashi, M. & Pascual-Leone, A. Transcranial magnetic stimulation in neurology. *Lancet*

Neurol. **2,** 145–156 (2003).

Lauwereyns, J., Watanabe, K., Coe, B. & Hikosaka, O. A neural correlate of response bias in

monkey caudate nucleus. Nature 418, 413–417 (2002).

Legon W, Sato TF, Opitz A, Mueller J, Barbour A, Williams A, Tyler WJ (2014) Transcranial

focused ultrasound modulates the activity of primary somatosensory cortex in humans. Nat

Neurosci. 17(2):322-9.

Lappin JS, Bell HH (1976) The detection of coherence in moving random dot pattens. Vision

Research. 16(2):161-168.

Luce, R. Response Times: Their Role in Inferring Elementary Mental Organization. Oxford Univ.

Press New York (1986).

Marquet F, Teichert T, Wu SY, Tung YS, Downs M, Wang S, Chen C, Ferrera V, Konofagou

EE.(2014) Real-time, transcranial monitoring of safe blood-brain barrier opening in non-human

primates. PLoS One. 9(2):e84310

McDannold N, Arvanitis CD, Vykhodtseva N, Livingstone MS. (2012) Temporary disruption of

the blood-brain barrier by use of ultrasound and microbubbles: safety and efficacy evaluation in

rhesus macaques. Cancer Res. 72(14):3652-63.

McDannold, N., Vykhodtseva, N., Raymond, S., Jolesz, F. A. & Hynynen, K. MRI-guided

targeted blood-brain barrier disruption with focused ultrasound: Histological findings in rabbits.

Ultrasound Med. Biol. **31,** 1527–1537 (2005).

Miller, N. E. Chemical Coding of Behavior in the Brain. *Science* **148,** 328–38 (1965).

Nitsche, M. a *et al.* Facilitation of implicit motor learning by weak transcranial direct current

stimulation of the primary motor cortex in the human. *J. Cogn. Neurosci.* **15,** 619–626 (2003).

Ratcliff, R. (1978). A theory of memory retrieval. Psychological Review, 85, 59-108.

Roitman JD, Shadlen MN (2002) Responses of neurons in the lateral intraparietal area during a

combined visual discrimination reaction time task. Journal of Neuroscience 22(21):9475-9489.

Schlaepfer TE, Bewernick BH. (2013) Deep brain stimulation for major depression. Handbook

Clin Neurol. 116:235-43.

Teichert T, Ferrera VP. (2010) Suboptimal integration of reward magnitude and prior reward

likelihood in categorical decisions by monkeys. Front Neurosci. 4:186.

Teichert T, Yu D, Ferrera VP. (2014) Performance monitoring in monkey frontal eye field.

J Neurosci. 34(5):1657-71.

Tung, Y.-S., Marquet, F., Teichert, T., Ferrera, V. & Konofagou, E. E. Feasibility of noninvasive

cavitation-guided blood-brain barrier opening using focused ultrasound and microbubbles in

nonhuman primates. Appl. Phys. Lett. 98, 163704 (2011).

Velling VA, Shklyaruk SP (1988) Modulation of the functional state of the brain with the aid of

focused ultrasound action. Neurosci Behav Physiol. 18(5), 369-375.

Feshitan, J. A., Chen, C. C., Kwan, J. J. & Borden, M. A. Microbubble size isolation by

differential centrifugation. *J. Colloid Interface Sci.* **329,** 316–324 (2009).

Acknowledgements: NIH R01MH059244 (VPF); Kavli Foundation (VPF); NIH R01AG038961

(EK); NIH R01EB009041 (EK); Wallace H. Coulter Foundation (EK); FUS Foundation (EK);

Alfonso Martin Escudero Foundation, Spain (CS)

Contributions: V.P.F. and E.K. conceived the study. V.P.F. and M.E.D. processed the

behavioral and MRI data. M.E.D. conducted all ultrasound and behavioral experiments. T.T.,

A.B., and M.K., assisted with ultrasound and behavioral experiments. S.C. assisted with

behavioral experiments. C.S. manufactured the microbubbles and assisted with ultrasound

experiments. All authors participated in manuscript editing.

Competing financial interests:

The authors declare no competing financial interests.

Figures

Figure 1. Contrast enhanced (gadodiamide) MRI of BBB opening in putamen. Top row shows

sagittal, coronal and horizontal slices through the brain of monkey N. Blue oval indicates the

targeted region. Red and orange voxels indicate BBB opening. Bottom row shows the same for

monkey O.

Figure 2. Effects of motion coherence, reward, sonication, and haloperidol on response time

(RT) for single target trials. A) Response time for single target trials as a function of motion

coherence, offered reward, and sonication. Error bars omitted for clarity. B) RT averaged over

coherence levels. Each pair of circles connected with a line compares average RT during

haloperidol sessions (black border) with saline sessions (grey border). Error bars represent ±1

s.e.m. Legend applies to both subplots.

Figure 3. Effects of motion coherence, reward, and sonication on response time and accuracy

for choice (2-target) trials. A) Accuracy (percent correct) for two-target trials. B) Response time

for two-target trials. C) Accuracy vs. response time for two-target trials. Dashed horizontal line

is 75% correct level. Legend in A applies to all subpanels. Error bars are omitted for clarity.

Figure 4. Effects of sonication and haloperidol on performance for two-target trials. A) Effects

on accuracy. B) Effects on response time. Error bars represent ±1 s.e.m. Black borders

sessions with haloperidol, grey borders are saline controls. All conventions same as **Fig. 3**.

Data in each column are plotted separately for each coherence level.

Figure 5. Behavioral task and experimental timeline. A) Timeline of sonication and behavioral

testing. B) Decision task sequence. The monkey initiates a trial by touching the cue. A random

dot motion stimulus appeared moving to the left or right, flanked by two targets. The monkey

touched the target toward which the dots were moving to receive a reward. Stimuli were

displayed on the right or left of the screen. A physical barrier forced the monkey to respond with

the corresponding hand. Only the yellow bars and dot stimulus were visible to the monkey, not

the blue arrow, dotted line, or hand, which are used to indicate the motion of the dots, the

physical barrier separating the two halves of the screen, and the manual response, respectively.