Trait Heritability in Major Transitions

Matthew D. Herron[1,*] and William C. Ratcliff[2]

[1] Division of Biological Sciences, University of Montana. Missoula, MT 59812.

[2] School of Biology, Georgia Institute of Technology. Atlanta, GA 30332.

*Correspondence: xprinceps@gmail.com

Abstract

A crucial component of major transitions theory is that after the transition, adaptation

occurs primarily at the level of the new, higher-level unit. For collective-level adaptations

to occur, though, collective-level traits must be heritable. Since collective-level traits are

functions of lower-level traits, collective-level heritability is related to particle-level

heritability. However, the nature of this relationship has rarely been explored in the

context of major transitions. We examine relationships between particle-level heritability

and collective-level heritability for several functions that express collective-level traits in

terms of particle-level traits. When this relationship is linear, the heritability of a

collective-level trait is never less than that of the corresponding particle-level trait and is

higher under most conditions. For more complicated functions, collective-level

heritability is higher under most conditions, but can be lower when the function relating

particle to cell-level traits is sensitive to small fluctuations in the state of the particles

within the collective. Rather than being an impediment to major transitions, we show that

collective-level heritability superior to that of the lower-level units can often arise 'for

free', simply as a byproduct of collective formation.

Introduction

Major transitions, or evolutionary transitions in individuality, are a framework for

understanding the origins of life's hierarchy and of biological complexity (Maynard

Smith & Szathmáry 1995; Michod & Roze 1997). During such a transition, a new unit of

evolution emerges from interactions among previously existing units. Thus the primary

level of selection shifts from the particle (lower-level unit) to the collective (higher-level

unit), for example from cells to multicellular organisms or from insects to eusocial

societies.

Evolution by natural selection requires heritable variation in phenotypes that

affect fitness at the level at which selection occurs (Lewontin 1970; Damuth & Heisler

1988). When collective-level traits are exposed to selection, it is collective-level

heritability that determines the magnitude of the response. Collective-level heritability of

traits is thus necessary for collective-level adaptations, and this has often been assumed to

be difficult. For example, Michod claims that collective-level heritability can only arise

when particles are specialized in different tasks (Michod & Roze 1997; Michod 2005,

2011). Simpson says that "From the view of some standard theory, these transitions are

impossible," in part because particle-level heritability greatly exceeds collective-level

heritability.

Major transitions can be conceptualized as a shift from MLS1 to MLS2, in the

sense of Damuth and Heisler (1988), as in Okasha (2005) (see also Godfrey-Smith

(2009), Shelton & Michod (2009)). In MLS1, properties of the particles are under

selection; in MLS2, it is the properties of the collectives. We follow Okasha (2005) in

referring to the lower-level units in a transition 'particles' and the higher-level units

'collectives.' Although our biological analogies are presented in terms of cells as

particles and multicellular organisms as collectives, in principle our results should hold

for any pair of adjacent levels.

According to Michod (2005 p. 971), "…the challenge of ETI [evolutionary

transitions in individuality] theory is to explain how fitness at the group level in the sense

of MLS2 emerges out of fitness at the group level in the sense of MLS1." But fitness, or

selection, is only half of the breeder's equation. Predicting the response to selection

requires an estimate of heritability.

Whether or not collective-level fitness in MLS2 is a function of particle-level

fitness is a matter of some disagreement (for example, Rainey & Kerr 2011 say no).

However, collective-level traits must be functions of particle-level traits, unless we

accept strong emergence, a philosophical position tantamount to mysticism (Bedau

1997). The function may be complex and involve cell-cell communication, feedbacks,

environmental influences, etc., but it is still a function that is, in principle, predictable

from particle-level traits. Nevertheless, the relationship between heritability of particle-

level traits and that of collective-level traits has rarely been considered in the context of

major transitions, leading Okasha (2006 p. 47) to wonder, "Does variance at the particle

level necessarily give rise to variance at the collective level? Does the heritability of a

collective character depend somehow on the heritability of particle characters? The

literature on multi-level selection has rarely tackled these questions explicitly, but they

are crucial."

We examine relationships between particle-level heritability and collective-level

heritability for several functions that express collective-level traits in terms of particle-

level traits. For the simplest (linear) function, we derive an analytical solution for the

relationship. For more complex functions, we employ a simulation model to explore the

relationship over a range of conditions.

Analytical model

There are several ways to estimate heritability. If the strength of selection is known,

heritability can be estimated by back-calculating from the breeder's equation: $R = h^2S$,

where R is the response to selection and S the selection differential. This can be

rearranged as $h^2 = S/R$. Another method is to compare parent and offspring traits: the

slope of the parent-offspring regression is an estimator of heritability (Falconer 1960).

We use the latter method in the simulations described in the next section.

Since heritability can be defined as the proportion of phenotypic variance

explained by genetic variance, one method of estimation is to partition total variance into

its components using an analysis of variance. We employ this approach in an analytical

model to derive the relationship between the heritability of a collective-level trait and that

of the particle-level trait from which it arises. For the sake of tractability, we begin with

the simplest case, assuming that the size (number of particles) of collectives is fixed and

that the collective-level trait is a linear function of the particle-level traits. We further

assume that reproduction is asexual, so the proper measure of heritability is broad-sense

heritability, H^2 (Lynch & Walsh 1998).

We imagine a population in which collectives are made up of particles and

genetically distinct clones are made up of collectives. As a concrete example, we can

think of a population of undifferentiated volvocine algae, such as *Gonium*, in which case

the particles are cells and the collectives are colonies. Because of asexual reproduction,

many genetically identical collectives may comprise a clone. Genetic variation among

clones may arise through mutation or because the population is facultatively sexual, in

which case these results will only hold for evolution within the asexual phase (in the

Gonium example, during the summer bloom that precedes autumn mating and winter

dormancy).

Broad-sense heritability is the ratio of genetic variance (V_G) to total phenotypic

variance (V_P), estimated as the ratio of among-clone variance to total phenotypic variance

(Lynch & Walsh 1998). Treating particles and collectives separately, the phenotype of

particle k in collective j within clone i can be expressed as

$$y_{ijk} = m + A_i + B_{j(i)} + C_{k(ij)} \qquad (1)$$

where m is the mean genotypic value of all clones, A_i is the deviation of clone i from m,

$B_{j(i)}$ is the deviation of collective j from the mean of clone i, and $C_{k(ij)}$ is the deviation of

particle k from the mean of colony j within clone i. The genetic variance is the sum of

squared deviations of A from m:

$$\mathbf{SSA} = bc \sum_{i=1}^{a}\left(\bar{y}_{i..} - \bar{y}_{...}\right)^2, \qquad (2)$$

where a, b, and c are the number of clones, collectives within a clone, and particles

within a collective, respectively. Phenotypic variance of collectives within clones is

$$\mathbf{SS(B/A)} = c \sum_{i=1}^{a}\sum_{j=1}^{b}\left(\bar{y}_{ij.} - \bar{y}_{i..}\right)^2, \qquad (3)$$

that among particles within collectives is

$$\mathbf{SS(C/B)} = \sum_{i=1}^{a}\sum_{j=1}^{b}\sum_{k=1}^{c}\left(y_{ijk} - \bar{y}_{ij.}\right)^2, \qquad (4)$$

and total phenotypic variance is

$$SST_y = SSA + SS(B/A) + SS(C/B). \tag{5}$$

Broad-sense heritability of a particle-level trait, H_y^2, is the ratio of genetic variance to

total phenotypic variance:

$$H_y^2 = \frac{V_{G_y}}{V_{P_y}} = \frac{SSA}{SSA+SS(B/A)+SS(C/B)}. \tag{6}$$

The phenotype of collective j within clone i can be expressed as

$$z_{ij} = \mu + \alpha_i + \beta_{j(i)}, \tag{7}$$

where μ is the mean genetic value of all clones, α_i is the deviation of clone i from μ, and

$\beta_{j(i)}$ is the deviation of collective j from the mean of clone i. The genetic variance is the

sum of squared deviations of α from μ:

$$SS\alpha = b \sum_{i=1}^{a} (\bar{z}_{i\cdot} - \bar{z}_{\cdot\cdot})^2. \tag{8}$$

Phenotypic variance among colonies within clones is

$$SS(\beta/\alpha) = \sum_{i=1}^{a} \sum_{j=1}^{b} (z_{ij} - \bar{z}_{i\cdot})^2. \tag{9}$$

and total phenotypic variance is

$$SST_z = SS\alpha + SS(\beta/\alpha). \tag{10}$$

Broad-sense heritability of a collective-level trait, H_z^2, is the ratio of genetic variance to

total phenotypic variance,

$$H_z^2 = \frac{V_{G_z}}{V_{P_z}} = \frac{SS\alpha}{SS\alpha+SS(\beta/\alpha)}. \tag{11}$$

If colony-level trait is the average of cell-level traits, $z_{ij} = y_{ij\cdot}$, $\bar{z}_{i\cdot} = \bar{y}_{i\cdot\cdot}$, and

$\bar{z}_{\cdot\cdot} = \bar{y}_{\cdot\cdot\cdot}$. Thus $SS\alpha = cSSA$, and $SS(\beta/\alpha) = cSS(B/A)$. Substituting into (11),

we get

$$H_z^2 = \frac{SSA}{(SSA+SS(B/A))}. \tag{12}$$

The ratio of collective-level heritability to particle-level heritability is thus

$$\frac{H_z^2}{H_y^2} = \frac{SSA + SS(B/A) + SS(C/B)}{SSA + SS(B/A)}.$$ **(13)**

Collective-level heritability is therefore never less than particle-level heritability, and is

greater unless $SS(C/B) = 0$, in other words unless particles within each collective have

identical phenotype. Although we have derived this relationship assuming that the

collective-level trait is the average of particle-level traits, the result holds for any linear

function.

Simulation model

The correspondence between particle-level and collective-level traits is likely to be more

complicated than a linear relationship for many interesting cases. Here we explore more

complicated trait mapping functions using a simulation model. As above, particles grow

in clonal collectives, which reproduce by forming two new collectives, each with as

many particles as its parent. The initial population is founded by ten genetically distinct

clones, each of which has a different genetically determined mean particle phenotype

(spaced evenly between 1 and 2). These are grown for at least 7 generations, resulting in

at least 127 collective-level reproductive events per genotype and $127n$ (where n is

particle number per collective) particle-level reproductive events per genotype.

In this model, we consider two sources of non-genetic effects on particle

phenotype (Figure 1), each of which should lower the heritability of both particle- and

collective-level traits. The first is intrinsic particle reproductive stochasticity, analogous

to developmental instability (Palmer 1994). In the model, we determine the phenotype of

daughter cells by sampling from a distribution centered on the parent's genetic mean,

with standard deviation σ. As shown in the analytical model above, by averaging out this

variation, collectives can gain a heritability advantage over cells.

Figure 1. Two non-genetic modifiers to cell size. There are two nongenetic influences
on cell size in our model: developmental instability, a stochastic effect that varies a cell's
phenotype from its genetic mean size (with standard deviation σ), and environmental
effects, which modify the size of all cells in a collective (with standard deviation σ').
168
Our simulation also considers the phenotypic effects of environmental

heterogeneity. Here, we model collectives as independently experiencing different

environmental conditions that affect the phenotypes of all cells within them in the same

manner. To extend the biological analogy offered above, *Gonium* colonies growing near

the surface of a pond (where light and CO_2 are abundant) may form colonies with larger

cells than clonemates near the bottom. We implemented this in our model by assigning a

size modifier, drawn from a normal distribution centered on 1 with standard deviation σ',

to each collective. We then multiplied the phenotype of each particle within the collective

by this modifier. This source of phenotypic heterogeneity should reduce the heritability

of collectives more than particles, simply because collectives experience a relatively

higher frequency of stochastic events than particles do (each collective gets assigned a

different size multiplier, but every particle within that collective experiences the same

size multiplier).

We examine the effect of each of the above sources of phenotypic variation

independently for the example of cells (particles) within nascent multicellular organisms

(collectives). For a linear relationship, collective size is simply the sum of the sizes of

cells within the collective. For both cells and collectives, heritability is assessed by

calculating the slope of a linear regression on parent and offspring phenotype (Falconer

1960). In this simple case, mean collective-level heritability is always greater than or

equal to cell-level heritability. Only when $\sigma = 0$ (*i.e.*, when all cells within a collective

have identical phenotype) are cell- and collective-level heritability equal, in agreement

with the analytical model. Greater developmental instability for cell size increases the

advantage of collective-level heritability over cell-level heritability (Figure 2a). Larger

collectives, which average out cellular stochasticity more effectively, experience a greater

benefit than smaller collectives (Figure 2a). Note that the simulations run in Figure 2a

reflect a very patchy environment in which environmental effects on cell size within

collectives are large ($\sigma' = 0.25$). Increasing the magnitude of these environmental effects

on cell size diminishes the difference in heritability between collectives and cells, but

mean collective-level heritability is still greater than cell-level heritability for all

parameter combinations (Figure 2b).

Figure 2. Collective-level heritability of size is greater than cell-level heritability for size. In **a**), we hold the effect of the environment fixed (standard deviation $\sigma' = 0.25$), and vary the degree of developmental instability σ: 10^{-4} (purple), 0.0625 (blue), 0.125 (green), 0.1875 (yellow), 0.25 (red). In the absence of developmental instability, collective and cell-level heritabilites are identical. Greater developmental instability increases relative collective-level heritability. **b**) Here we hold developmental instability fixed at $\sigma = 0.25$, and vary between-collective environmental effects on cell size from $\sigma' = 10^{-4}$ (purple) to 0.25 (red). When developmental instability is nonzero, larger collectives improve collective-level heritability. Ten replicates were run of each parameter combination. Populations were simulated for nine generations of growth.

The volume of the cellular collective (Figure 2, Figure 3a), which is simply the

sum of the cell volumes within it, represents the simplest function mapping cellular to

multicellular traits. We now consider more complicated nonlinear functions relating

cellular to multicellular traits, some of which have biological relevance to the evolution

of multicellularity. For each function, we calculated the relative heritability of collective-

to cell-level traits for 32-celled collectives across 1024 combinations of σ and σ' ranging

from 0 to 0.25. The first nonlinear collective-level trait we consider is its diameter. Large

size is thought to provide a key benefit to nascent multicellular collectives when they

become too big to be consumed by gape-limited predators (Stanley 1973; Bell 1985). For

a collective that is approximately spherical, the trait that actually determines the

likelihood of being eaten is diameter. For geometric simplicity we assume that the cells

within the collective are pressed tightly together into a sphere, allowing us to calculate

collective radius as $d = 2\left(\frac{3V}{4\pi}\right)^{\frac{1}{3}}$, where V is the sum of the cell volumes within the

collective. Collective volume (Figure 3a) and diameter (Figure 3b) exhibit similar

dynamics, with collective-level heritability always exceeding cell-level heritability, and

being maximized under conditions of strong cell size stochasticity (high σ) and no

environmental heterogeneity (low σ').

228
229

Figure 3. Relative heritability of various collective-level traits to cell-level heritability for size. Here we examine the heritability of four multicellular traits that depend on the size of their constituent cells, relative to cellular heritability for size. The relationship between the size of the cells within collectives and the multicellular trait are shown as insets. We consider three biologically-significant traits with different functions mapping the size of cells within the collective onto collective phenotype. The heritability of collective size (**a**) and diameter (**b**) is always higher than cell-level heritability for size, and is maximized when cellular developmental noise is greatest and among-collective environmental effects are smallest (lower right corner). We modeled cooperative metabolism (**c**) with a logistic function, such that there is a threshold over which collectives are large enough to perform some metabolic task. We also considered a multicellular trait that does not monotonically increase with greater collective size, but instead oscillates with varying collective size (**d**). Like a and b, collective-level heritability is highest relative to cell-level heritability when environmental heterogeneity is minimal. Pink contours denote relative heritability of 1. In these simulations we consider 32 cell collectives grown for 7 generations. The colormap denotes collective-level heritability divided by cell-level heritability for size across 1024 σ, σ' combinations.

Next, we consider a logistic function describing among-cell cooperation (*e.g.*,

production of costly extracellular metabolites), in which the extent of cooperative

metabolism depends nonlinearly on collective volume. Specifically, we calculated the

extent of cooperative metabolism as $c = 64/(1 + e^{(-.2(V-48))})$. Here, the center of the

collectives' size distribution (*i.e.*, the volume of a collective with all 32 cells having size

1.5) lies at the function's inflection point. As with the previous two functions, collective-

level heritability is greater than cell-level heritability for much of the trait space, and is

maximized under conditions of high cellular stochasticity and low environmental

heterogeneity.

Finally, we consider a collective-level trait that oscillates between -1 and 1 with

increased cell size: $O = \sin\left(\frac{V}{8}\right)$, where V is the sum of cell volumes. This trait has no

obvious biological interpretation, but is distinct from the linear and nonlinear (but

monotonically increasing) functions described previously. While the general relationship

observed for these other functions still holds (collectives have greater heritability when

cellular stochasticity is high and environmental heterogeneity low), we now find that

much of the trait space now favors cellular heritability over that of the collective (Figure

3d, upper left). This appears to be due to the sensitivity of the function relating collective

to particle-level traits. We explore this further in Figure S1, where we consider four

versions of the model presented in Figure 3d, varying the sensitivity of the collective's

response to its lower-level composition. The heritability of collective-level traits is

minimal when small differences in cellular phenotype generate radically different

collective-level phenotypes. This makes sense: there is little potential for collective-level

heritability when tiny differences in cellular phenotype within collectives generate

drastically different collective-level phenotypes.

Discussion

Using a quantitative genetics framework, we have derived an analytical solution for the

relationship between particle-level and collective-level heritability for a limited case.

When the organismal trait is a linear function of the cell level traits, the organismal

heritability turns out to be a simple function of the cell-level heritability. In contrast to

claims that particle-level heritability is always higher than collective-level heritability

(*e.g.*, Simpson 2011), we have shown that collective-level heritability is higher over a

wide range of conditions. Furthermore, we have shown that, contrary to Michod's results

for fitness heritability (Michod & Roze 1997; Michod 2005, 2011), specialization of

particles is not necessary for collective-level trait heritability to arise.

This analytical result is a step toward understanding the relationship between

heritabilities at two adjacent hierarchical levels, but the assumption that the collective-

level trait is a linear function of the particle-level trait is restrictive. The simulation model

shows that the results are dependent on the function relating the traits at the two levels.

Even under the conditions we model, which are favorable to collective-level heritability

(clonal reproduction, negligible within-collective mutation, and fixed cell number per

collective), collective-level heritability was not always higher than cell-level heritability

(Figure 3d, Figure S1). Specifically, the relative heritability of collectives to cells was

below 1 when the collective-level phenotype was extremely sensitive to small changes in

the phenotype of cells within the collective (Figure S1). It is important to note, however,

that we only saw this high sensitivity in a function with little biological relevance

(collective-level phenotype oscillated between -1 and 1 with increased cell volume).

Collective-level heritability was higher than cell-level heritability for most of the trait

space in the other three biologically significant functions we considered (Figure 3a-c).

In our simulation, we examined the effects of two independent sources of

phenotypic variation affecting the relative heritability of particle and collective-level

traits. Stochastic variation in cell size around the clone's genetic mean (σ) reduces the

absolute heritability of cells and collectives by introducing non-heritable phenotypic

variation. By averaging across multiple cells, however, collectives reduce the effects of

this phenotypic variation, providing them with a relative heritability advantage over cells.

We also considered the effect of environmental heterogeneity in which all of the cells

within a collective are affected in the same manner (σ'). Collectives are

disproportionately affected: each group is assessed a different size modifier, but all of the

cells within these groups are affected in the same manner. As a result, collectives

experience n-fold (where n is the number of cells per collective) more stochastic events,

which reduces their heritability relative to cells. The influence of these sources of

variation is evident in the contour plots of Figures 3 and S1: the relative heritability of

collectives to cells is maximized when cellular stochastic variation is high, and

environmental heterogeneity low (lower right corner of the plots).

Our results differ from previous considerations of heritability in important

respects. For example, Queller (1992) presents a useful reformulation of the Price

equation for selection at two levels:

$$\Delta \overline{G} = S_b h_b^2 + S_w h_w^2,$$

in which $\Delta \bar{G}$ is the change in average trait value, S_b and S_w are the selection differentials

between groups and within groups, respectively, and $h^2{}_b$ and $h^2{}_w$ are the heritabilities of

the group-level and individual-level traits, respectively. This formulation partitions the

response to selection on a particle-level trait into within- and among-collective change,

but the focus is still on particle-level traits. Our focus is on the evolution of collective-

level traits. In the terminology of Damuth and Heisler (1988), our focus is on MLS2,

while Queller's is on MLS1. In addition, Queller makes no attempt to relate collective-

level heritability to particle-level heritability.

Michod and Roze (1997) have previously modeled the relationship between

particle-level and collective-level heritability of fitness during a major transition.

However, as Okasha (2006) points out, heritability of fitness only ensures that mean

population fitness will increase over time. For selection to result in directional phenotypic

change, it is phenotypes that must be heritable. Futhermore, Michod and Roze focused on

within-organism genetic change. Our models assume that such change is negligible, as is

likely to be true early in a transition, when collectives (*e.g.*, nascent multicellular

organisms) presumably include a small number of clonally-replicating particles (*e.g.*,

cells).

Okasha (2003) considers heritability in MLS1 (which he refers to as group

selection 2) and MLS2 (his group selection 1) but does not attempt to derive a

relationship between heritabilities at two levels. We have focused on just this

relationship, because knowing the ratio of heritabilities is necessary to predict the

outcome of opposing selection at two levels. This has important implications for

collective-level traits that arise from cooperation among particles. The presumed higher

heritability of the particle-level traits has been seen as a problem for the evolution of

cooperation that benefits the collective (Buss 1986, 1987; Michod 1997; Michod & Roze

1997; Simpson 2011). Our results show that this problem does not always exist.

Several simplifying assumptions underlie our models, but these are not extreme

departures from some biological systems. For example, the volvocine algae, an important

model system for understanding major transitions, undergo clonal reproduction only

occasionally punctuated by sex, are small enough that within-collective mutation is

probably negligible, and have cell numbers that are under tight genetic control.

A great deal of work has gone into understanding the selective pressures that may

have driven major evolutionary transitions. However, heritability is just as important as

the strength of selection in predicting evolutionary outcomes. We have shown that, given

some simplifying assumptions, heritability of collective-level traits comes 'for free'; that

is, it emerges as an inevitable consequence of group formation. Understanding the

emergence of trait heritability at higher levels is necessary to model any process

involving multilevel selection, so our results are relevant to a variety of other problems.

Acknowledgments

This work was supported by grants from NASA (Cooperative Agreement Notice 7,

Exobiology award #NNX15AR33G), NSF (DEB-1457701), and the John Templeton

Foundation (43285). We would like to thank Sam Brown, Peter Conlin, Michael Doebeli,

and Rick Michod for helpful discussions.

Figure S1. **Multicellular traits that are very sensitive to underlying cell-level traits**
are less heritable. Here we vary the sensitivity of the multicellular trait to variation in
collective volume (x), ranging from very sensitive (sin(x/2), upper left panel) to relatively
insensitive (sin(x/16), lower right panel). Highly sensitive multicellular mapping
functions exhibit reduced multicellular heritability relative to cell-level heritability for
size. The colormap denotes collective heritability divided by cell-level heritability for
size across 1024 x,y combinations. Pink line denotes relative heritability of 1.
Populations were simulated for 7 generations.

374
375

References

Bedau MA (1997) Weak emergence. *Noûs*, **31**, 375–379.

Bell G (1985) *The origin and early evolution of germ cells as illustrated by the Volvocales* (HO Halvorson, A Monroy, Eds,). Alan R. Liss, New York.

Buss LW (1986) The uniqueness of the individual revisited. In: *Population Biology and Evolution of Clonal Organisms*, pp. 467–505.

Buss LW (1987) *The Evolution of Individuality*. Princeton University Press, Princeton.

Damuth J, Heisler IL (1988) Alternative formulations of multilevel selection. *Biology and Philosophy*, **3**, 407–430.

Falconer DS (1960) *Introduction to Quantitative Genetics*. Ronald Press Co., New York.

Godfrey-Smith P (2009) *Darwinian Populations and Natural Selection*. Oxford University Press, Oxford.

Lewontin RC (1970) The units of selection. *Annual Review of Ecology and Systematics*, **1**, 1–18.

Lynch M, Walsh B (1998) *Genetics and Analysis of Quantitative Traits*. Sinauer Associates, Inc., Sunderland, MA.

Maynard Smith J, Szathmáry E (1995) *The Major Transitions in Evolution*. Oxford University Press, Oxford.

Michod RE (1997) Evolution of the individual. *The American Naturalist*, **150 Suppl**, S5–S21.

Michod RE (2005) On the transfer of fitness from the cell to the multicellular organism. *Biology & Philosophy*, **20**, 967–987.

Michod RE (2011) Evolutionary transitions in individuality: multicellularity and sex. In: *The Major Transitions in Evolution Revisited* (eds Calcott B, Sterelny K), pp. 167–197. MIT Press, Cambridge, MA.

Michod RE, Roze D (1997) Transitions in individuality. *Proceedings: Biological Sciences*, **264**, 853–857.

Okasha S (2003) The concept of group heritability. *Biology and Philosophy*, **18**, 445–461.

Okasha S (2005) Multilevel selection and the major transitions in evolution. *Philosophy of Science*, **72**, 1013–1025.

Okasha S (2006) *Evolution and the Levels of Selection*. Oxford University Press, Oxford.

Palmer AR (1994) Fluctuating asymmetry analyses: A primer. In: *Developmental Instability: Its Origins and Evolutionary Implications* (ed Markow TA), pp. 335–364. Kluwer, Dordrecht, Netherlands.

Queller DC (1992) Quantitative genetics, inclusive fitness, and group selection. *The American Naturalist*, **139**, 540–558.

Rainey PB, Kerr B (2011) Conflicts among levels of selection as fuel for the evolution of individuality. In: *The Major Transitions in Evolution Revisited* (eds Calcott B, Sterelny K), pp. 141–162. The MIT Press, Cambridge.

Shelton DE, Michod RE (2009) Philosophical foundations for the hierarchy of life. *Biology & Philosophy*, **25**, 391–403.

Simpson C (2011) How many levels are there? How insights from evolutionary
transitions in individuality help measure the hierarchical complexity of life. In: *The*
Major Transitions in Evolution Revisited (eds Calcott B, Sterelny K), pp. 199–225.
The MIT Press, Cambridge.
Stanley SM (1973) An ecological theory for the sudden origin of multicellular life in the
Late Precambrian. *Proceedings of the National Academy of Sciences, USA*, **70**,
1486–1489.
425