1 **Trait Heritability in Major Transitions**
2
M. D. Herron[1,3], S. A. Zamani-Dahaj[2], and W. C. Ratcliff[1]

[1] School of Biology, Georgia Institute of Technology. North Avenue, Atlanta, GA 30332

[2] School of Physics, Georgia Institute of Technology. North Avenue, Atlanta, GA 30332

[3] Correspondence: xprinceps@gmail.com, phone 520-820-6698, fax 406-243-4184

Abstract: A crucial component of major transitions theory is that after the transition,

adaptation occurs primarily at the level of the new, higher-level unit. For collective-level

adaptations to occur, though, collective-level traits must be heritable. Since collective-

level trait values are functions of lower-level trait values, collective-level heritability is

related to particle-level heritability. However, the nature of this relationship has rarely

been explored in the context of major transitions. We examine relationships between

particle-level heritability and collective-level heritability for several functions that

express collective-level trait values in terms of particle-level trait values. When a

collective-level trait value is a linear function of particle-level trait values and collective

size is fixed, the heritability of a collective-level trait is never less than that of the

corresponding particle-level trait and is higher under most conditions. For more

complicated functions, collective-level heritability is higher under most conditions, but

can be lower when the environment experienced by collectives is heterogeneous. Within-

genotype variation in collective size reduces collective-level heritability, but it can still

exceed particle-level heritability when phenotypic variance among particles within

collectives is large. These results hold for a diverse sample of biologically relevant traits.

Rather than being an impediment to major transitions, we show that collective-level

heritability superior to that of the lower-level units can often arise 'for free', simply as a

byproduct of collective formation.

Keywords: Evolution; Heritability; Major Transitions; Multicellularity; Quantitative

genetics; Simulations

Introduction

Major transitions, or evolutionary transitions in individuality, are a framework for

understanding the origins of life's hierarchy and of biological complexity [1,2]. During

such a transition, a new unit of evolution emerges from interactions among previously

existing units. This new unit, or collective, has traits not present before the transition and

distinct from those of the units that comprise it (particles; see [3] for an in-depth

discussion of collective-level traits). These collective-level traits are potentially subject to

selection. Over the course of the transition, the primary level of selection shifts from the

particle (lower-level unit) to the collective (higher-level unit), for example from cells to

multicellular organisms or from individual insects to eusocial societies.

Evolution by natural selection requires heritable variation in phenotypes that

affect fitness at the level at which selection occurs [4,5]. The breeder's equation of

quantitative genetics shows that heritability and strength of selection contribute equally to

the adaptive response (see Analytical model below). When a collective-level trait is

exposed to selection, it is collective-level heritability (the heritability of the collective-

level trait) that determines the magnitude of the response. Collective-level heritability of

traits is thus necessary for collective-level adaptations, but the emergence of collective-

level heritability during a major transition has often been assumed to be difficult. For

example, Michod considers the emergence of collective-level heritability through conflict

mediation a crucial step in major transitions [2,6,7]. Simpson says that "From the view of

some standard theory, these transitions are impossible," in part because particle-level

heritability greatly exceeds collective-level heritability [8].

Major transitions can be conceptualized as a shift from MLS1 to MLS2, in the

sense of Damuth and Heisler [5], as in Okasha [9] (see also Godfrey-Smith [10], Shelton

& Michod [11]). In MLS1, properties of the particles are under selection; in MLS2, it is

the properties of the collectives. We follow Okasha [9] in referring to the lower-level

units in a transition as 'particles' and the higher-level units as 'collectives.' Although our

biological analogies are presented in terms of cells as particles and multicellular

organisms as collectives, in principle our model could be extended to any pair of adjacent

levels.

According to Michod [6], "…the challenge of ETI [evolutionary transitions in

individuality] theory is to explain how fitness at the group level in the sense of MLS2

emerges out of fitness at the group level in the sense of MLS1." But fitness, or selection,

is only half of the breeder's equation. Predicting the response to selection requires an

estimate of heritability.

Whether or not collective-level fitness in MLS2 is a function of particle-level

fitness is a matter of some disagreement (for example, Rainey and Kerr say no [11]).

However, collective-level phenotypes must be functions of particle-level trait

phenotypes, unless we accept strong emergence, a philosophical position tantamount to

mysticism [13]. The function may be complex and involve cell-cell communication,

feedbacks, environmental influences, etc., but it is still a function that is, in principle,

predictable from particle-level trait values.

Nevertheless, the relationship between the heritability of particle-level traits and

that of collective-level traits has rarely been considered in the context of major

transitions, leading Okasha [14] to wonder, "Does variance at the particle level

necessarily give rise to variance at the collective level? Does the heritability of a

collective character depend somehow on the heritability of particle characters? The

literature on multi-level selection has rarely tackled these questions explicitly, but they

are crucial." Similarly, Goodnight [15] says, "...we really do not have a good

understanding of what contributes to group heritability, how to measure it, or even how to

define it."

While the role of selection has often been considered in the context of major

transitions, the role of trait heritability has been relatively neglected. We examine

relationships between particle-level heritability and collective-level heritability for

several functions that express collective-level trait values in terms of particle-level trait

values. For the simplest (linear) function, we derive an analytical solution for the

relationship. For more complex functions, we employ a simulation model to explore the

relationship over a range of conditions.

Analytical model

There are several ways to estimate heritability, the proportion of phenotypic variation

explained by genetic variation. If the strength of selection is known, heritability can be

estimated by back-calculating from the breeder's equation: $R = h^2 S$, where R is the

response to selection, S the selection differential, and h^2 the narrow-sense heritability (i.e.

the proportion of phenotypic variation explained by additive genetic variation). This can

be rearranged as $h^2 = S/R$. Another method is to compare parent and offspring trait

values: the slope of the parent-offspring regression is an estimator of heritability [16]. We

use the latter method in the simulations described in the next section.

Since heritability can be defined as the proportion of phenotypic variance

explained by genetic variance, one method of estimation is to partition total variance into

its components using an analysis of variance. We employ this approach in an analytical

model to derive the relationship between the heritability of a collective-level trait and that

of the particle-level trait from which it arises. For the sake of tractability, we begin with

the simplest case, assuming that the size (number of particles) of collectives is fixed and

that the collective-level trait value is a linear function of the particle-level trait values.

We further assume that reproduction is asexual, so the proper measure of heritability is

broad-sense heritability, H^2 [17]. Broad-sense heritability describes the proportion of

phenotypic variation explained by all genetic variation, including both additive and non-

additive components.

We imagine a population in which collectives are made up of particles and

genetically distinct clones are made up of collectives. As a concrete example, we can

think of a population of undifferentiated volvocine algae, such as *Gonium*, in which case

the particles are cells and the collectives are colonies. Because of asexual reproduction,

many genetically-identical collectives may comprise a clone. Genetic variation among

clones may arise through mutation or because the population is facultatively sexual, in

which case these results will only hold for evolution within the asexual phase (in the

Gonium example, during the summer bloom that precedes autumn mating and winter

dormancy).

Broad-sense heritability is the ratio of genetic variance (V_G) to total phenotypic

variance (V_P), estimated as the ratio of among-clone variance to total phenotypic variance

[17]. Inherent in this concept is that genetically identical individuals are not always

phenotypically identical; V_P includes both genetic and non-genetic variation. Non-genetic

variation can arise from maternal effects, environmental (including microenvironmental)

effects, and random developmental noise. Phenotypic variation among genetically

identical individuals has been extensively documented, including in bacteria [18,19],

unicellular eukaryotes [20], plants [21], animals [17], and volvocine algae [22].

In this section, we use an ANOVA framework to estimate heritability as a ratio of

sums of squares. Strictly speaking, heritability is a ratio of variances, not of sums of

squares. However, the ratios of the relevant sums of squares converges to that of the

variances as the number of categories increases (see Supplemental Information), and for

all but tiny or genetically uniform biological populations, the difference between the two

ratios is negligible.

Treating particles and collectives separately, the phenotype of particle k in

collective j within clone i can be expressed as

$$y_{ijk} = m + A_i + B_{j(i)} + C_{k(ij)} \tag{1}$$

where m is the mean genotypic value of all clones, A_i is the deviation of clone i from m,

$B_{j(i)}$ is the deviation of collective j from the mean of clone i, and $C_{k(ij)}$ is the deviation of

particle k from the mean of collective j within clone i. The model in (1) describes a nested

ANOVA framework, in which the sums of squared deviations from the population mean

is partitioned into among-clone, among collectives within clone, and within-collective

components. The among-clone component, the sum of squared deviations of A from m, is

$$\text{SSA} = bc \sum_{i=1}^{a}\left(\bar{y}_{i\cdot\cdot} - \bar{y}_{\cdots}\right)^2 \tag{2}$$

where *a, b,* and *c* are the number of clones, collectives within a clone, and particles

within a collective, respectively. The sum of squared deviations of collectives within

clones is

$$\text{SS(B/A)} = c \sum_{i=1}^{a}\sum_{j=1}^{b}\left(\bar{y}_{ij\cdot} - \bar{y}_{i\cdot\cdot}\right)^2, \tag{3}$$

that among particles within collectives is

$$\text{SS(C/B)} = \sum_{i=1}^{a}\sum_{j=1}^{b}\sum_{k=1}^{c}\left(y_{ijk} - \bar{y}_{ij\cdot}\right)^2, \tag{4}$$

and total sum of squares is

$$\text{SST}_y = \text{SSA} + \text{SS(B/A)} + \text{SS(C/B)}. \tag{5}$$

Broad-sense heritability of a particle-level trait, H_y^2, is the ratio of genetic variance to

total phenotypic variance:

$$H_y^2 = \frac{V_{G_y}}{V_{P_y}} \approx \frac{\text{SSA}}{\text{SSA}+\text{SS(B/A)}+\text{SS(C/B)}}. \tag{6}$$

We now turn our attention to collective-level traits. The phenotype of collective *j*

within clone *i* can be expressed as

$$z_{ij} = \mu + \alpha_i + \beta_{j(i)}, \tag{7}$$

where μ is the mean genetic value of all clones, α_i is the deviation of clone *i* from μ, and

$\beta_{j(i)}$ is the deviation of collective *j* from the mean of clone *i*. The sum of squared

deviations of α from μ is

$$\text{SS}\alpha = b \sum_{i=1}^{a}(\bar{z}_{i\cdot} - \bar{z}_{\cdot\cdot})^2. \tag{8}$$

The sum of squares among colonies within clones is

$$\text{SS}(\beta/\alpha) = \sum_{i=1}^{a}\sum_{j=1}^{b}\left(z_{ij} - \bar{z}_{i\cdot}\right)^2, \tag{9}$$

and the total sum of squares is

$$\mathbf{SST_z = SS\alpha + SS(\beta/\alpha).} \tag{10}$$

Broad-sense heritability of a collective-level trait, H_z^2, is the ratio of genetic variance to

total phenotypic variance,

$$\mathbf{H_z^2 = \frac{V_{G_z}}{V_{P_z}} \approx \frac{SS\alpha}{SS\alpha + SS(\beta/\alpha)}.} \tag{11}$$

If collective-level trait value is the average of cell-level trait values, $z_{ij} = y_{ij\cdot}$,

$\bar{z}_{i\cdot} = \bar{y}_{i\cdot\cdot}$, and $\bar{z}_{\cdot\cdot} = \bar{y}_{\cdot\cdot\cdot}$. Thus $SS\alpha = cSSA$, and $SS(\beta/\alpha) = cSS(B/A)$. Substituting into

(11),

we get

$$\mathbf{H_z^2 \approx \frac{SSA}{(SSA + SS(B/A))}.} \tag{12}$$

The ratio of collective-level heritability to particle-level heritability is thus

$$\mathbf{\frac{H_z^2}{H_y^2} \approx \frac{SSA + SS(B/A) + SS(C/B)}{SSA + SS(B/A)}.} \tag{13}$$

Collective-level heritability is therefore never less than particle-level heritability (i.e., the

ratio of heritabilities is never less than 1), and is greater unless $SS(C/B) = 0$, in other

words unless particles within each collective have identical phenotype.

Although we have derived this relationship assuming that the collective-level trait

value is the average of particle-level trait values, the result holds for any linear function.

The substitution that gets us from (11) to (12) introduces the constant c, which scales

both numerator and denominator and therefore cancels out. Different linear functions

would change the magnitude of the constant relating $SS\alpha$ to $cSSA$ and $SS(\beta/\alpha)$ to

$cSS(B/A)$ but not the fact that numerator and denominator are scaled by the same

constant.

The approximations in (6) and (11), which express ratios of variances as ratios of

sums of squares, hold when the number of clones (a) and the number of genetically

identical collectives within a clone (b) are large (Electronic Supplement 1). For example,

at $a = b = 10$, the approximation differs from the true value by less than 1%. Thus the

results of the analytical model hold for all but tiny and/or extremely genetically

depauperate populations. The number of particles within a collective (c) does not play a

role, so our results are relevant even early in a major transition, when the collectives are

likely to be small. For most real biological populations, the difference between the true

heritability and the sums of squares approximation will be negligible (see Electronic

Supplement 1 for a simple numerical example).

Simulation model

The correspondence between particle-level and collective-level trait values is likely to be

more complicated than a linear relationship for many interesting and biologically relevant

cases. Here we explore more complicated trait mapping functions using a simulation

model. As above, particles grow in clonal collectives, which reproduce by forming two

new collectives, each with as many particles as its parent. The initial population is

founded by ten genetically distinct clones, each of which has a different genetically

determined mean particle phenotype (spaced evenly between 1 and 2). These are grown

for at least 7 generations, resulting in at least 127 collective-level reproductive events per

genotype and $127n$ (where n is particle number per collective) particle-level reproductive

events per genotype. Simulation models are provided as Electronic Supplements 2-8.

In this model, we consider two sources of non-genetic effects on particle

phenotype (Figure 1), each of which should lower the heritability of both particle- and

collective-level traits. The first is intrinsic reproductive stochasticity in particle

phenotype, analogous to developmental instability [23]. In the model, we determine the

phenotype of daughter cells by sampling from a distribution centered on the parent's

genetic mean, with standard deviation σ. As shown in the analytical model above, by

averaging out this variation, collectives can gain a heritability advantage over cells.

Figure 1. Two non-genetic modifiers to cell phenotype. There are two non-genetic
influences on particle phenotype (cell size in this example) in our model: developmental
instability, a stochastic effect that varies a particle's phenotype from its genetic mean
(with standard deviation σ), and environmental effects, which modify the phenotype of all
particles in a collective by the same amount (with standard deviation $\sigma\square$).
222
Our simulation also considers the phenotypic effects of environmental

heterogeneity. Here, we model collectives as independently experiencing different

environmental conditions that affect the phenotypes of all cells within them in the same

manner. To extend the biological analogy offered above, *Gonium* colonies growing near

the surface of a pond (where light and CO_2 are abundant) may form colonies with larger

cells than clonemates near the bottom. We implemented this in our model by assigning a

size modifier, drawn from a normal distribution centered on 1 with standard deviation

σ, to each collective. We then multiplied the phenotype of each particle within the

collective by this modifier. This source of phenotypic heterogeneity should reduce the

heritability of collectives more than particles, simply because collectives experience a

relatively higher frequency of stochastic events than particles do (each collective gets

assigned a different size multiplier, but every particle within that collective experiences

the same size multiplier).

236   We examine the effect of each of the above sources of phenotypic variation

independently for the example of cells (particles) within nascent multicellular organisms

(collectives). For a linear relationship, collective size is simply the sum of the sizes of

cells within the collective. For both cells and collectives, heritability is assessed by

calculating the slope of a linear regression on parent and offspring phenotype [16]. In this

simple case, mean collective-level heritability is always greater than or equal to cell-level

heritability. Only when $\sigma = 0$ (*i.e.*, when all cells within a collective have identical

phenotype) are cell- and collective-level heritability equal, in agreement with the

analytical model. Greater developmental instability for cell size increases the advantage

of collective-level heritability over cell-level heritability (Figure 2a). Larger collectives,

which average out cellular stochasticity more effectively, experience a greater increase in

heritability than smaller collectives (Figure 2a). Note that the simulations run in Figure 2a

reflect a very patchy environment in which environmental effects on cell size within

collectives are large ($\sigma\square = 0.25$). While our model is not explicitly spatial, when $\sigma\square$ is

high, different collectives experience different environmental effects on their mean cell

size, simulating the effects of a patchy environment. Increasing the magnitude of these

environmental effects on cell size diminishes the difference in heritability between

collectives and cells, but mean collective-level heritability is still greater than cell-level

heritability for all parameter combinations (Figure 2b).

Figure 2. Collective-level heritability of size is greater than particle-level heritability
for size. In **a**), we hold the effect of the environment fixed (standard deviation
$\sigma\square = 0.25$), and vary the degree of developmental instability for particle size σ: 10^{-4}
(purple), 0.0625 (blue), 0.125 (green), 0.1875 (yellow), 0.25 (red). In the absence of
developmental instability for size, collective and cell-level heritabilities are identical.
Greater developmental instability increases relative collective-level heritability. **b**) Here
we hold developmental instability fixed at $\sigma = 0.25$, and vary between-collective
environmental effects on cell size from $\sigma\square = 10^{-4}$ (purple) to 0.25 (red). When
developmental instability is nonzero, larger collectives improve collective-level
heritability. We ran ten replicates of each parameter combination and simulated
populations for nine generations of growth.
267
The volume of the cellular collective (Figure 2, Figure 3a), which is simply the

sum of the cell volumes within it, represents the simplest function mapping cellular to

multicellular trait values. We now consider more complicated nonlinear functions

relating cellular to multicellular trait values, some of which have biological relevance to

the evolution of multicellularity. For each function, we calculated the relative heritability

of collective- to cell-level traits for 32-celled collectives across 1024 combinations of σ

and $\sigma\square$ ranging from 0 to 0.25.

The first nonlinear collective-level trait we consider is its diameter. Large size is

thought to provide a key benefit to nascent multicellular collectives when they become

too big to be consumed by gape-limited predators [24,25]. For a collective that is

approximately spherical, the trait that actually determines the likelihood of being eaten is

diameter, which is therefore an important component of fitness. For geometric simplicity

we assume that the cells within the collective are pressed tightly together into a sphere,

allowing us to calculate collective radius as $d = 2\left(\frac{3V}{4\pi}\right)^{\frac{1}{3}}$, where V is the sum of the cell

volumes within the collective. Collective volume (Figure 3a) and diameter (Figure 3b)

exhibit similar dynamics, with collective-level heritability always exceeding cell-level

heritability, and being maximized under conditions of strong cell size stochasticity (high

σ) and no environmental heterogeneity (low   ).

286
287

288
Figure 3. Relative heritability of various collective-level traits to cell-level
heritability for size. Here we examine the heritability of four multicellular traits that
depend on the size of their constituent cells, relative to cellular heritability for size. The
relationship between the size of the cells within collectives and the multicellular trait are
shown as insets. We consider three biologically-significant traits with different functions
mapping the size of cells within the collective onto collective phenotype. The heritability
of collective size (**a**) and diameter (**b**) is always higher than cell-level heritability for size,
and is maximized when cellular developmental noise is greatest and among-collective
environmental effects are smallest (lower right corner). We modeled swimming speed (**c**)
based on the model of Solari *et al.* (2006) for volvocine green algae. We also considered
survival rate under predation as a logistic function of radius (**d**). Like a and b, collective-
level heritability is highest relative to cell-level heritability when environmental
heterogeneity is minimal. Pink contours denote relative heritability of 1. In these

simulations we consider 32 cell collectives grown for 7 generations. The colormap
denotes collective-level heritability divided by cell-level heritability for size across 1024
$\sigma, \sigma\square$ combinations.
305

Next, we consider swimming speed as a function of cell radius. We based this

simulation on the hydrodynamics model of volvocine green algae derived by Solari *et al.*

[26]. For simplicity, we modeled 32-celled, undifferentiated collectives (GS colonies in

[26]), which would be similar to extant algae in the genus *Eudorina*. Given these

assumptions, the function relating cell radius to upward swimming speed (Equation 4

from [26]) can be simplified to

$$V_{up} = \left(\frac{f N^{0.5}}{3\pi\eta_w}\right)r^{-1} - \left(\frac{g\Delta\rho_c(4/3)N^2}{3\eta_w}\right)r^2 \qquad\qquad (14)$$

where *f* is average effective upward swimming force per cell, N is the number of cells per

collective, η_w is water viscosity, r is the average radius of cells in the collective, and $\Delta\rho_c$

is the density difference between cells and water. Electronic Supplement 9 provides a

more detailed description of the derivation of Equation 14.

Using the numerical values in Solari *et al* [26], $\eta_w = 0.01$ g/cm·s, $\Delta\rho_c =$

0.047g/cm^3, and $f = 2.4 \times 10^{-7}$ g·cm/s^2, so

$$V_{up} = \frac{0.02}{\pi}r^{-1} - \frac{400}{3}r^2 \qquad\qquad (15)$$

In this model, the swimming force of cells is independent of cell size, so, as cells get

larger the collective will become heavier (more negatively buoyant) without a

corresponding increase in total swimming force, and therefore its upward swimming

speed will decrease. Thus upward swimming speed is a monotonically declining function

of cell radius (Fig. 3c inset), unlike the functions for volume and diameter (Fig. 3a, 3b

insets), both of which are monotonically increasing. Nevertheless, the general behavior of

heritability is very similar to the previous ones and for a wide range of parameter values,

the collective-level trait has a higher heritability than the cell-level trait (Fig. 3c).

Next, we consider a function describing a collective's survival rate in the presence

of a predator that can only consume collectives below a certain size. We calculated the

survival rate (*c*) as a logistic function of the collective's radius, effectively assuming that

predation efficiency drops off quickly when collectives reach a threshold size (Fig. 3d

inset):

$c = \dfrac{1}{1+e^{-0.5(0.5rN^{0.5}-25)}}$ (**16**)

As with the previous functions (Fig. 3a-c), collective-level heritability is greater

than cell-level heritability for much of the trait space and is maximized under conditions

of high cellular stochasticity (σ) and low environmental heterogeneity ($\sigma\square$; Fig. 3d).

Finally, we consider the case in which the simplifying assumption of constant cell

number does not hold. Instead, the number of cells per collective fluctuates around the

genetic mean $\overline{N}$. In this case, each collective reproduces two new collectives, but the

number of cells per new collective is a random variable drawn from a normal distribution

with mean $\overline{N}$ and coefficient of variation CV_N (the coefficient of variation for a normal

distribution is the ratio of standard deviation to the mean). We chose to represent

variation in the number of cells per collective as CV_N instead of standard deviation so that

the range of variation would not change with the size of the collective.

Variation in cell number, unlike the developmental and the environmental

variation, does not affect the heritability of cells, only that of collectives. Therefore, we

expected that increasing CV_N would decrease the ratio of collective-level to cell-level

heritability. To test this effect, we calculated the relative heritability of size (volume) for

collectives and cells across 1024 combinations of and CV_N ranging from 0 to 0.25

(   ). The simulation shows that the CV_N has a strong effect on collective-level

heritability (Fig. 4). As CV_N increases, the ratio of collective- to cell-level heritabilities

decreases, falling below one when the magnitude of is similar to or smaller than that of

CV_N (Figure 4).

355
Figure 4. Relative heritability of collective size to cell size when the number of cells
per collective varies. When the coefficient of variation for cell number per collective
(CV_N) is low, collective-level heritability is always higher than cell-level heritability, but
this advantage is undercut by increased variation in cell number. The ratio of collective-
to cell-level heritability is maximized when developmental variation in cell size (σ) is
large and variation in the number of cells per collective is low. The pink contour denotes
a ratio of collective-level to cell-level heritability of 1. In these simulations, we consider
collectives with a genetic mean of 32 cells grown for 7 generations. The colormap
denotes collective-level heritability divided by cell-level heritability for size across 1024
σ, CV_N combinations.

Discussion

Using a quantitative genetics framework, we have derived an analytical solution for the

relationship between particle-level and collective-level heritability for a limited case.

When particle number is constant and the collective-level trait value is a linear function

of the particle-level trait values, the organismal heritability turns out to be a simple

function of the cell-level heritability. In contrast to claims that particle-level heritability is

always higher than collective-level heritability (e.g. [8]), we have shown that collective-

level heritability is higher over a wide range of conditions. Because this result depends on

the number of clones and the number of colonies within a clone, it may not hold for very

small populations or those with little genetic variation. This is not a major limitation,

though, since tiny, genetically homogeneous populations are unlikely to be the ones

experiencing selectively driven evolutionary transitions in individuality.

This analytical result is a step toward understanding the relationship between

heritabilities at two adjacent hierarchical levels, but the assumptions of constant particle

number and linear function are restrictive. The simulation model shows that the results

are somewhat dependent on the function relating the trait values at the two levels.

However, these functions were chosen to be diverse, and the behavior of the relative

heritabilities is nevertheless qualitatively similar, increasing with cellular developmental

variation (σ), decreasing with environmental heterogeneity ($\sigma_\square$), and exceeding 1 for

most of the parameter space.

Of course, we have not (and cannot) comprehensively explored the universe of

possible functions relating collective-level traits to particle-level traits. What we have

done is explore a small sample of this space, with functions ranging from extremely

390 simple (volume) to somewhat more complex (swimming speed, survival under

391 predation). We do not claim that the high heritabilities estimated for these collective-level

392 traits would apply to all such traits, and a full accounting of possible functions is beyond

393 the scope of this (or any) study. Rather, we have shown that for at least some such

394 functions, the resulting collective-level traits can have high heritability, and thus be

395 altered by selection, early in an evolutionary transition in individuality.

396   All four of the collective-level traits in the simulation models are potentially

397 biologically relevant. Volume and diameter are both aspects of size, which can be an

398 important component of fitness both in evolutionary transitions in individuality [27] and

399 in life history evolution [28]. Swimming speed is a measure of motility, which has

400 selective consequences for a wide range of organisms, including many animals and

401 microbes. For planktonic organisms, a positive upward swimming speed provides active

402 control of depth, allowing some control over light intensity (for autotrophs) and prey

403 abundance (for heterotrophs). Survival under predation obviously has important fitness

404 implications for many organisms, and both theoretical and experimental evidence

405 implicate predation as a possible selective pressure driving the evolution of

406 multicellularity. Kirk, for example, suggests that a "predation threshold" above which

407 algae are safe from many filter feeders may have driven the evolution of multicellularity

408 in the volvocine algae [29]. Microbial evolution experiments in the algae *Chlorella* and

409 *Chlamydomonas* have shown that predation can drive the evolution of undifferentiated

410 multicellular clusters [30–32].

411   In our simulations, we examined the effects of three independent sources of

412 phenotypic variation affecting the relative heritability of particle and collective-level

413   traits. Stochastic variation in cell size around the clone's genetic mean (σ) reduces the

414   absolute heritability of cells and collectives by introducing non-heritable phenotypic

415   variation. By averaging across multiple cells, however, collectives reduce the effects of

416   this phenotypic variation, providing them with a relative heritability advantage over cells.

417   We also considered the effect of environmental heterogeneity in which all of the

cells within a collective are affected in the same manner (σ'). Collectives are

disproportionately affected: each collective is assessed a different size modifier, but all of

the cells within these collectives are affected in the same manner. As a result, collectives

experience n-fold more stochastic events (where n is the number of cells per collective),

which reduces their heritability relative to cells. The influence of these sources of

variation is evident in the contour plots of Figure 3: the relative heritability of collectives

to cells is maximized when cellular stochastic variation is high and environmental

heterogeneity low (lower right corner of the plots). The effect of environmental

heterogeneity in our simulations is consistent with the empirical finding of Goodnight

[33] that group selection of *Arabidopsis* was more effective when among-deme

environmental variance was low.

429   Finally, we considered variation in the number of particles per collective. Such

variation substantially reduces the heritability of a collective-level trait. Even with

reasonably large variation in collective size, though, the collective-level trait retains most

of the heritability of the particle-level trait on which it is based (for example, ~55% at a

CV_N in particle number of 0.25).

Our results differ from previous considerations of heritability in important

respects. For example, Queller [34] presents a useful reformulation of the Price equation

for selection at two levels:

$$\Delta \overline{G} = S_b h_b^2 + S_w h_w^2,$$

in which $\Delta \overline{G}$ is the change in average trait value, S_b and S_w are the selection differentials

between collectives and within collectives, respectively, and h_b^2 and h_w^2 are the

heritabilities of the collective-level and individual-level traits, respectively. This

formulation partitions the response to selection on a particle-level trait into within- and

among-collective change, but the focus is still on particle-level traits. Our focus is on the

evolution of collective-level traits. In the terminology of Damuth and Heisler [5], our

focus is on MLS2, while Queller's is on MLS1. In addition, Queller makes no attempt to

derive the relationship between collective-level heritability and particle-level heritability.

Michod and Roze [2] have previously modeled the relationship between particle-

level and collective-level heritability of fitness during a major transition. However, as

Okasha [14] points out, heritability of fitness only ensures that mean population fitness

will increase over time. For selection to result in directional phenotypic change, it is

phenotypes that must be heritable. Futhermore, Michod and Roze focused on within-

organism genetic change. Our models assume that such change is negligible, as is likely

to be true early in a transition, when collectives (*e.g.*, nascent multicellular organisms)

presumably include a small number of clonally-replicating particles (*e.g.*, cells).

Okasha [35] considers heritability in MLS1 (which he refers to as group selection

2) and MLS2 (his group selection 1) but does not attempt to derive a relationship between

heritabilities at two levels. We have focused on just this relationship, because knowing

the ratio of heritabilities is necessary to predict the outcome of opposing selection at two

levels. This has important implications for collective-level traits that arise from

cooperation among particles. The presumed higher heritability of the particle-level traits

has been seen as a problem for the evolution of cooperation that benefits the collective

[2,8,36–38]. Our results show that this problem does not always exist.

Several previous papers have shown that group-level heritability (collective-level

heritability in our terminology) exists and can be substantial. Slatkin [39], for example,

showed that one measure of group-level heritability, fraction of total variance between

lines, is substantial both in an additive model and in the *Tribolium* experiments of Wade

and McCauley [40]. Under some conditions, the between-line variance of a linear trait

such as the one we consider in our analytical model exceeds the within-line variance.

Bijma, Wade and colleagues [41–43] showed that variance in the total breeding

value of a population can be increased, even to the point of exceeding phenotypic

variance, by interactions among individuals. Our model does not consider (or require)

interactions among individuals. Further, their model and empirical example are

exclusively concerned with individual-level traits (particle-level traits in our

terminology), for example survival days in chickens. They do not estimate group

heritability as such, and judge that "it is unclear how this parameter should be defined or

estimated."

Goodnight [15] considers the ratio of group-level heritability to individual-level

heritability (in the narrow sense) using contextual analysis. Although this paper does not

provide a formula to calculate this ratio, its inequality sets a minimum bound (with the

assumption that selection at the two levels is in opposition). As in our analyses,

Goodnight shows that group-level heritability can exceed individual-level heritability in

some circumstances.

Several simplifying assumptions underlie our models, most importantly the

genetic identity of particles within collectives. This condition only applies to a subset of

the major transitions. Queller recognized two subcategories within Maynard Smith and

Szathmáry's [1] list of transtions, which he called "egalitarian" and "fraternal" transitions

[44]. Briefly, egalitarian transitions involve particles that may be genetically distinct, or

even from different species, such as the alliance of a bacterium with an Archaean that

gave rise to the eukaryotic cell. Fraternal transitions are those in which the particles are

genetically similar or identical, such as the origins of eusociality and of most

multicellular lineages.

Because of our assumption of genetic identity among particles, we cannot

generalize our results to all types of major transitions. Egalitarian transitions will not

normally meet this criterion. A possible exception is aggregative multicellularity, as seen

in cellular slime molds and myxobacteria, when assortment is so high that fruiting bodies

are genetically uniform. This is probably uncommon [45], but it does happen [46,47].

Transitions in which reproduction of particles is obligately sexual, such as the origins of

eusociality, also violate this assumption.

A better fit for our models is clonal multicellularity, which is probably the most

common type of major transition. An incomplete list of independent origins of clonal

multicellularity includes animals; streptophytes; chytrid, ascomycete, and basidiomycete

fungi; florideophyte and bangiophyte red algae; brown algae; peritrich ciliates; ulvophyte

green algae; several clades of chlorophyte green algae; and filamentous cyanobacteria

[48–51]. In most cases the early stages in these transitions probably violated the

assumption of uniform particle number per collective, but our simulations show that our

main results are robust to reasonable violations of this assumption.

506   One example that does approximate all of our assumptions is that of the volvocine

green algae, an important model system for understanding the evolution of

multicellularity. Volvocine algae undergo clonal reproduction only occasionally

punctuated by sex, are small enough that within-collective mutation probably has

negligible phenotypic effects, and have cell numbers that are under tight genetic control.

Conclusion

A great deal of work has gone into understanding the selective pressures that may have

driven major evolutionary transitions. However, heritability is just as important as the

strength of selection in predicting evolutionary outcomes. We have shown that, given

some simplifying assumptions, heritability of collective-level traits comes 'for free'; that

is, it emerges as an inevitable consequence of group formation. Qualitatively, this result

holds across a wide range of parameters and for a diverse sample of biologically relevant

traits. Collective-level heritability is maximized (relative to particle-level heritability)

when phenotypic variation among particles is high and when environmental

heterogeneity and variation in collective size are low. Understanding the emergence of

trait heritability at higher levels is necessary to model any process involving multilevel

selection, so our results are relevant to a variety of other problems.

Declarations

Ethics approval and consent to participate

Not applicable

Consent for publication

Not applicable

Availability of data and material

All data generated or analysed during this study are included in this published article [and

its supplementary information files].

Competing interests

The authors declare that they have no competing interests

Funding

This work was supported by grants from NASA (NNA17BB05A, NNX15AR33G), NSF

(DEB-1457701, DEB-1723293, DEB-1456652), and the John Templeton Foundation

(43285).

Authors' contributions

MDH conceived the project, developed the analytical model, contributed to the

simulation models, and contributed to writing the manuscript. SAZ-D and WCR

developed the simulation models and contributed to writing the manuscript. All authors

read and approved the final manuscript.

Acknowledgements

We would like to thank Sam Brown, Peter Conlin, Michael Doebeli, Rick Michod, and

Deborah Shelton for helpful discussions.

References

1. Maynard Smith J, Szathmáry E. The Major Transitions in Evolution [Internet]. Oxford:
Oxford University Press; 1995 [cited 2011 Jun 3]. Available from:
books.google.com/books?isbn=019850294X

2. Michod RE, Roze D. Transitions in individuality. Proc Biol Sci [Internet]. 1997 [cited
2011 Mar 2];264:853–7. Available from: http://www.jstor.org/stable/info/50748

3. Ratcliff WC, Herron M, Conlin PL, Libby E. Nascent life cycles and the emergence of
higher-level individuality. Philos Trans R Soc B Biol Sci. 2017;372:20160420.

4. Lewontin RC. The units of selection. Annu Rev Ecol Syst [Internet]. 1970 [cited 2011
Jun 21];1:1–18. Available from:
http://www.annualreviews.org/doi/abs/10.1146/annurev.es.01.110170.000245

5. Damuth J, Heisler IL. Alternative formulations of multilevel selection. Biol Philos
[Internet]. Springer Netherlands; 1988 [cited 2011 Jul 4];3:407–30. Available
from: http://www.springerlink.com/content/h2l02q8729931l36/

6. Michod RE. On the transfer of fitness from the cell to the multicellular organism. Biol
Philos [Internet]. Springer Netherlands; 2005 [cited 2011 Mar 2];20:967–87.
Available from: http://www.springerlink.com/content/111hx10587083p24/

7. Michod RE. Evolutionary transitions in individuality: multicellularity and sex. In:
Calcott B, Sterelny K, editors. Major Transitions Evol Revisit. MIT Press; 2011.
p. 167–97.

8. Simpson C. How many levels are there? How insights from evolutionary transitions in
individuality help measure the hierarchical complexity of life. In: Calcott B,
Sterelny K, editors. Major Transitions Evol Revisit. Cambridge: The MIT Press;
2011. p. 199–225.

9. Okasha S. Multilevel selection and the major transitions in evolution. Philos Sci.
2005;72:1013–25.

10. Godfrey-Smith P. Darwinian Populations and Natural Selection. Oxford: Oxford
University Press; 2009.

11. Shelton DE, Michod RE. Philosophical foundations for the hierarchy of life. Biol
Philos [Internet]. 2010 [cited 2011 Jan 5];25:391–403. Available from:
http://www.springerlink.com/content/d56j414582nr2172/

12. Rainey PB, Kerr B. Conflicts among levels of selection as fuel for the evolution of
individuality. In: Calcott B, Sterelny K, editors. Major Transitions Evol Revisit.
Cambridge: The MIT Press; 2011. p. 141–62.

13. Bedau MA. Weak emergence. Noûs [Internet]. 1997;31:375–9. Available from:
http://onlinelibrary.wiley.com/doi/10.1111/0029-4624.31.s11.17/abstract

14. Okasha S. Evolution and the Levels of Selection. Oxford: Oxford University Press;
2006.

15. Goodnight CJ. Multilevel selection: the evolution of cooperation in non-kin groups.
Popul Ecol [Internet]. Springer Japan; 2005 [cited 2011 May 3];47:3–12.
Available from: http://www.springerlink.com/content/j1362gv1440l410x/

16. Falconer DS. Introduction to Quantitative Genetics. New York: Ronald Press Co.;
1960.

17. Lynch M, Walsh B. Genetics and Analysis of Quantitative Traits. Sunderland, MA:
Sinauer Associates, Inc.; 1998.

18. Elowitz MB, Levine AJ, Siggia ED, Swain PS. Stochastic gene expression in a single
cell. Science (80-). 2002;297:1183–6.

19. McAdams HH, Arkin A. Stochastic mechanisms in gene expression. Proc Natl Acad
Sci USA. 1997;94:814–9.

20. Blake WJ, Kærn M, Cantor CR, Collins JJ. Noise in eukaryotic gene expression.
Nature   [Internet].   2003;422:633–7.   Available   from:
http://www.ncbi.nlm.nih.gov/pubmed/22839658

21. Burton GW, DeVane EH. Estimating heritability in tall fescue (*Festuca
Arundinaceae*) from replicated clonal material. Agron J. 1953;45:478–81.

22. Herron MD, Ghimire S, Vinikoor CR, Michod RE. Fitness trade-offs and
developmental constraints in the evolution of soma: An experimental study in a
volvocine alga. Evol Ecol Res [Internet]. 2014;16:203–21. Available from:
http://www.evolutionary-ecology.com/abstracts/v16/2917.html

23. Palmer AR. Fluctuating asymmetry analyses: A primer. In: Markow TA, editor. Dev
Instab Its Orig Evol Implic [Internet]. Dordrecht, Netherlands: Kluwer; 1994. p.
335–64.   Available   from:
http://www.biology.ualberta.ca/palmer.hp/pubs/94Primer/Primer.pdf

24. Stanley SM. An ecological theory for the sudden origin of multicellular life in the
Late Precambrian. Proc Natl Acad Sci USA [Internet]. 1973;70:1486–9. Available
from: http://www.pnas.org/cgi/content/abstract/70/5/1486

25. Bell G. The origin and early evolution of germ cells as illustrated by the Volvocales
[Internet]. Halvorson HO, Monroy A, editors. Orig. Evol. Sex. New York: Alan
R. Liss; 1985. Available from: books.google.com/books?id=OcsdAQAAIAAJ

26. Solari CA, Kessler JO, Michod RE. A hydrodynamics approach to the evolution of
multicellularity: flagellar motility and germ-soma differentiation in volvocalean
green algae. Am Nat [Internet]. The University of Chicago Press; 2006 [cited
2011   Jun   7];167:537–54.   Available   from:
http://www.jstor.org/stable/info/10.1086/501031

27. Bonner JT. Perspective: the size-complexity rule. Evolution (N Y). 2004;58:1883–90.

28. Peters RH. The Ecological Implications of Body Size. Cambridge: Cambridge
University Press; 1983.

29. Kirk DL. Volvox: Molecular-Genetic Origins of Multicellularity [Internet].
Cambridge: Cambridge   University   Press;   1998.   Available   from:
books.google.com/books?isbn=0521019141

30. Boraas ME, Seale DB, Boxhorn JE. Phagotrophy by a flagellate selects for colonial
prey: a possible origin of multicellularity. Evol Ecol. 1998;12:153–64.

31. Herron MD, Borin JM, Boswell JC, Walker J, Chen I-CK, Knox CA, et al. De novo
origins   of   multicellularity   in   response   to   predation.   bioRxiv.
2018;10.1101/247361.

32. Becks L, Ellner SP, Jones LE, Hairston NG. Reduction of adaptive genetic diversity
radically alters eco-evolutionary community dynamics. Ecol Lett [Internet]. 2010

[cited 2013 Aug 6];13:989–97. Available from: http://www.ncbi.nlm.nih.gov/pubmed/20528898

33. Goodnight CJ. The influence of environmental variation on group and individual selection in a cress. Evolution (N Y) [Internet]. 1985;39:545–58. Available from: http://links.isiglobalnet2.com/gateway/Gateway.cgi?GWVersion=2&SrcAuth=me kentosj&SrcApp=Papers&DestLinkType=FullRecord&DestApp=WOS&KeyUT =A1985AJY2600006%5Cnpapers2://publication/uuid/FDE6CE8D-EA49-47C3-9CC5-9956F9BDD9B8

34. Queller DC. Quantitative genetics, inclusive fitness, and group selection. Am Nat [Internet]. 1992 [cited 2011 Apr 14];139:540–58. Available from: http://www.jstor.org/stable/info/2462497

35. Okasha S. The concept of group heritability. Biol Philos [Internet]. 2003 [cited 2011 Mar 28];18:445–61. Available from: http://www.springerlink.com/content/rtx271pm8877276p/

36. Michod RE. Evolution of the individual. Am Nat [Internet]. 1997 [cited 2011 Mar 2];150 Suppl:S5–21. Available from: http://www.jstor.org/stable/info/10.1086/286047

37. Buss LW. The uniqueness of the individual revisited. Popul Biol Evol Clonal Org. 1986. p. 467–505.

38. Buss LW. The Evolution of Individuality [Internet]. Princeton: Princeton University Press; 1987. Available from: books.google.com/books?isbn=0691084696

39. Slatkin M. Populational heritability. Evolution (N Y) [Internet]. 1981 [cited 2011 May 3];35:859–71. Available from: http://www.jstor.org/stable/info/2407856

40. Wade MJ, McCauley DE. Group selection: the phenotypic and genotypic differentiation of small populations. Evolution (N Y) [Internet]. 1980 [cited 2011 May 3];34:799–812. Available from: http://www.jstor.org/stable/info/2408034

41. Bijma P, Muir WM, Ellen ED, Wolf JB, Van Arendonk JAM. Multilevel selection 2: Estimating the genetic parameters determining inheritance and response to selection. Genetics. 2007;175:289–99.

42. Bijma P, Muir WM, Van Arendonk JAM. Multilevel selection 1: Quantitative genetics of inheritance and response to selection. Genetics. 2007;175:277–88.

43. Wade MJ, Bijma P, Ellen ED, Muir W. Group selection and social evolution in domesticated animals. Evol Appl. 2010;3:453–65.

44. Queller DC. Cooperators since life began. Q Rev Biol [Internet]. 1997;72:184–8. Available from: http://www.jstor.org/stable/3036338

45. Sathe S, Kaushik S, Lalremruata A, Aggarwal RK, Cavender JC, Nanjundiah V. Genetic heterogeneity in wild isolates of cellular slime mold social groups. Microb Ecol. 2010;60:137–48.

46. Gilbert OM, Foster KR, Mehdiabadi NJ, Strassmann JE, Queller DC. High relatedness maintains multicellular cooperation in a social amoeba by controlling cheater mutants. Proc Natl Acad Sci USA [Internet]. 2007;104:8913–7. Available from: http://www.pubmedcentral.nih.gov/articlerender.fcgi?artid=1885602&tool=pmce

ntrez&rendertype=abstract

47. Gilbert OM, Queller DC, Strassmann JE. Discovery of a large clonal patch of a social
amoeba: Implications for social evolution. Mol Ecol. 2009;18:1273–81.

48. Parfrey LW, Lahr DJG. Multicellularity arose several times in the evolution of
eukaryotes. BioEssays. 2013;35:339–47.

49. Niklas KJ, Newman SA. The origins of multicellular organisms. Evol Dev [Internet].
2013 [cited 2013 May 24];15:41–52. Available from:
http://www.ncbi.nlm.nih.gov/pubmed/23331916

50. Herron MD, Rashidi A, Shelton DE, Driscoll WW. Cellular differentiation and
individuality in the "minor" multicellular taxa. Biol Rev. 2013;88:844–61.

51. Melkonian M. Phylum Chlorophyta. In: Margulis L, Corliss JO, Melkonian M,
Chapman DJ, editors. Handb Protoctista. Boston: Jones and Bartlett Publishers;
1990. p. 597–660.