Compound biological signatures facilitate phenotypic screening and target elucidation

Alvaro Cortes Cabrera[1], Daniel Lucena-Agell[2], Mariano Redondo-Horcajo[2], Isabel Barasoain[2], Fernando Diaz[2], Bernhard Fasching[3], Paula Petrone*[1]

[1]Pharma Research & Early Development Informatics (pREDi). Roche Innovation Center Basel, Basel, Switzerland.

[2]Laboratory of microtubule stabilizing agents. Department of Physico-chemical Biology. Center for Biological Research. CSIC. Madrid. Spain.

[3]Medicinal Chemistry. Pharma Research & Early Development (pRED). Roche Innovation Center Basel, Basel, Switzerland.

We developed a statistical method, BIOSEA, able to identify molecules that are capable of reproducing a desired cellular phenotype, by scanning a large compound collection based on biological similarity. Our method leverages highly incomplete and noisy compound bioactivity signatures from historical high-throughput screening campaigns. We applied our approach in a phenotypic screening workflow and found novel nanomolar inhibitors of cell division that reproduce the mode of action of reference natural products. In a drug discovery setting, our biological hit expansion protocol revealed new inhibitors of the NKCC1 co-transporter for autism spectrum disorders. Furthermore, we demonstrate BIOSEA's capabilities to predict novel targets for old compounds. We report new activities for the drugs nimedipine, fluspirilene and pimozide applicable for compound repurposing and rationalizing drug side effects. Our results highlight the opportunities of reusing public bioactivity data for prospective drug discovery applications where the target or mode of action is not known.

The last two decades have witnessed a dramatic change in our understanding of the interactions between small molecules and biological systems[1]. We are currently experiencing an important shift from reductionism[2], to a more pragmatic vision of living organisms as complex entities whose study cannot be effectively tackled by treating each part in isolation[3]. In this new framework, phenotypic screening has reemerged as a practical solution to finding novel chemical entities, which provide insights into the underlying cellular processes leading to new opportunities for drug discovery[4]. However, phenotypic screening brings about its own challenges such as the identification of the mechanism of action (MoA) and the effective target or targets responsible for these phenotypes[5]. Target and MoA identification play a fundamental role, paving the way to lead optimization via structural approaches, structure-activity relationships (SAR) rationalization and also by improving awareness about ADME (absorption, distribution, metabolism and excretion), safety liabilities and drug-drug interactions. The chemical biology and genomics fields have stood up to the challenge producing elegant solutions for these problems[6], such as CRISPR/Cas9 gene editing[7] and chemical proteomics-based techniques[8] that yet, bear inherent limitations such as chemical modification of the compounds, challenging specificity or high costs[8]. In the light of these advances, novel computational efforts are now in order to support phenotypic screening. In this work, we demonstrate how to leverage historical bioassay data as compound biological signatures to facilitate phenotypic deconvolution tasks, such as target identification and phenotypic library design. We report here a novel target and chemical structure-independent screening technique to find new chemical compounds to modulate a desired phenotype in a drug discovery setting.

Large amounts of pharmacological and bioactivity data accumulate both in public repositories and in proprietary databases in the pharmaceutical industry. A few approaches have attempted to use historical compound annotation data for target identification. For example, the TarFishDock[9], TargetHunter[10], Similarity Ensemble Approach[11,12], ChemMapper[13] and HitPick[14] methods, rely on either 2D or 3D chemical information of ligands or targets and an established molecular structure

similarity metric. These methods have been exploited with success in tasks such as the prediction of targets responsible for drug side effects[15]. However, an important limitation is the fact that these methods depend on the availability of crystal structures for the target, or the structural similarity between the compounds and previously known target ligands which confines their performance to well-explored areas in chemical space.

In recent years, it has been demonstrated that the cellular phenotype of compounds can be described, and even predicted, by a compound's biological signature: an array of numbers representing the perturbation caused by a chemical entity of interest measured throughout several cellular and biochemical assays[14]. In this context, compounds sharing a similar biological profile tend to modulate similar targets, without necessarily having similar chemical structure[16]. In the past, we and others[17-22] have proposed the use of historical high-throughput screening (HTS) activity data to represent molecules in the biological space. The so-called HTS fingerprints (HTSFP) have proven successful for the identification of new targets for known compounds[22]. Since they are structure-independent, HTSFP enable scaffold-independent linking of drug-like molecules to less frequently explored regions in chemical space such as those of complex natural products[16,22].

By integrating external pharmacological sources of information and HTS data, we have implemented BIOSEA – Biological Similarity Ensemble Approach – a novel statistical classifier that is able to establish new relationships between targets and small molecules on the basis of the mutual information between compound biological signatures. Algorithmically, BIOSEA was inspired by the method developed by Pearson[23] for sequence similarity searches (BLAST) and the Similarity Ensemble Approach (SEA) developed by Keiser et al.[11] to compare chemical structures.

In this work, first we report an unprecedented phenotypic screening workflow in which BIOSEA was applied to find compounds that modulate a desired cellular phenotype across large molecular libraries. We validated this procedure in two projects: (1) the discovery of drug-like inhibitors of cell division

and (2) the identification of novel chemical entities to alter intracellular ion concentration by

inhibiting the $Na^+K^+2Cl^-$ co-transporter (NKCC1) for the indication of autism. In a following section,

we demonstrate the application of BIOSEA for target identification, validated with both retrospective

data and prospective experiments that shed light on the polypharmacology of known drugs.

Consistently throughout these applications, BIOSEA revealed new relationships between molecules

and targets, which are difficult to detect even by a trained chemist's eye.

RESULTS

BIOSEA identifies molecules that reproduce a desired cellular phenotype

BIOSEA enables phenotypic screening without the need of information about the chemical structure

of the target or associated ligands (Fig. 1a). This capability was used in two drug discovery workflows

to find novel chemotypes that: (i) inhibit cell division and (ii) block NKCC1 mediated modulation of

intracellular ion concentration. In the first scenario, the cellular phenotype was characterized, and

both target and MoA of the proposed compounds were identified using cellular and biochemical

assays. In the second scenario, we applied BIOSEA for biological hit expansion with a NKCC1

cellular assay and compared its performance to standard methods that rely on chemical structure.

Phenotypic screening with BIOSEA: inhibition of cell division

As a reference set, we have selected six well-known natural products that perturb cell division in two

different ways: paclitaxel acts as a microtubule-stabilizing agent while the remaining compounds have

an inhibitory effect on tubulin polymerization. Virtual screening using these reference compounds and

a library of circa 365,000 compounds was performed to predict 20 biologically similar compounds

that would reproduce the phenotype of cell cycle arrest.

In a first validation step, using a cell cycle arrest screen in A549 cells, we could observe that 5 out 20

compounds were able to produce visible effects (rounding of cells) at a concentration of 5 μM which

was indicative of a cytostatic behavior. The cytotoxic activity of the positive compounds (hereby

named CT1 to CT5) was verified in the same cell line, with most IC_{50} values in the nanomolar range

(Table 1). Additionally, it was confirmed that all 5 compounds completely depolymerized the

interphasic cellular microtubule network (Fig. 2a). To establish tubulin as the target that modulated

microtubule dynamics, the time-course of the assembly of tubulin was recorded in excess

concentration of the hit compounds (Fig. 2b). All compounds were found to inhibit tubulin

polymerization *in vitro* and in cells. Binding of active compounds in known tubulin depolymerizing

binding sites (i.e. colchicine, vinca, maytansine, and eribulin) was investigated using various

competition assays (Fig. 2c, Supplementary Fig. 1 and Supplementary Fig. 2). CT5 was found to bind

to the eribulin biding site. The low solubility and weak activity of CT2 prevented any competition

assay. CT1, CT3 and CT4, were confirmed to bind to the colchicine site and a pharmacophore model

was proposed on the basis of colchicine-tubulin crystallographic information[24] to rationalize the

binding mode of these structurally diverse chemotypes (Fig. 2d, Supplementary Information).

Although the compounds do not share the same scaffold, the model showed a clear superimposition of

the di- and tri-methoxyphenyl moieties and good shape and chemical complementarity with the

reference compound, which indicates a similar binding pattern.

To deconvolute the information encoded in the HTSFP that leads to successful target-specific

compound predictions, we evaluated the individual contribution of every HTSFP assay to the global

biological similarity to the reference compounds. Thus, for each of the 5 hits, the top 10 HTSFP

components contributing most to the similarity were identified by a leave-one-out procedure

(Supplementary Fig. 3, Methods) and classified into biochemical or cell-based assays (Supporting

Table 1). Even though mutual information is a metric that is calculated based on the full range of the

HTSFP fingerprint, this approach provided the following observations on these highly informative

assays (HIA): **a) there is no unique pattern for HIA that is common for all active compounds**;

instead, each compound attributes its biological similarity to a different part of the fingerprint. For

example, two compounds that were found to bind in the colchicine pocket, CT1 and CT4, do not have

any HIA in common (Supplementary Fig. 3), and their biologically closest reference compounds are

also different (vincristine and vinblastine, respectively); **b) no HIA exists for the phenotype of**

interest (i.e. cell cycle arrest); only 2 assays measure unspecific cytotoxicity endpoints for *Giardia*

lamblia and *Escherichia coli* (Supplementary Table 1). The remaining HIA cover a diverse range of

biochemical and cell-based assays for targets such as the sentrin-specific protease 7, the sialic acid

acetylesterase or the vainilloid receptor 1, which are not associated to microtubule formation. It is

important to note that there is no tubulin-related assay in HTSFP; **c) cell-based and biochemical**

assays seem to contribute alike although most compounds resulted inactive in these HIA. While

compound activity in cell-based assays (Z-Score>2) may provide information on general cellular

perturbations (e.g. cytotoxicity, signaling pathways, etc.), inactivity throughout biochemical or

cellular assays (|Z-Score|<2) may bring information relevant for ruling out competing mechanisms

leading to a given phenotype. As an example, among its HIA the molecule CT2 has resulted inactive

for procaspases 3 and 7 -two biochemical assays- and inactive in a bacterial cell death screen in *E-*

coli.

Hit expansion on NKCC1

The primary goal of hit expansion efforts following an HTS campaign is to leverage informative lead

molecules to rescue other active compounds that were neglected in the hit triage or missing in the

primary screening library. In a phenotypic screen, this approach is specially challenging due to the

diversity in chemical leads which may modulate several unknown targets – a context in which the

chemical similarity principle for hit expansion may not apply.

The Roche NKCC1 inhibitor program aimed at correcting imbalances in the GABA developmental

switch[25], which controls the intracellular chloride concentration and shifts the functional role of

GABA between neonatal and adult stages from excitatory to inhibitory. Mediated by the change in

expression of two major ion co-transporters NKCC1 and KCC2, alterations in this mechanism

possibly result in CNS disorders including epilepsy, autism spectrum disorder, schizophrenia and

effects of traumatic brain injury[26]. The goal of the project was to identify NKCC1 inhibitors with high

selectivity over peripherally expressed NKCCs and largely improved brain exposure compared to

available compounds from the loop diuretic class such as bumetanide.

To this end, a primary screen of more than 1.2 million compounds within the Roche library was

conducted. The screen had a low hit rate of 0.06% confirmed active compounds covering several

compound classes. A standard approach that relies on the generation of 3D shapes and surface charge

models (ROCS[27]) was applied and 430 compounds pharmacophorically similar to selected active

compounds were submitted for profiling. This procedure resulted in a single novel hit with an affinity

of 9 μM. For biological hit expansion using BIOSEA, 12 confirmed hits with IC_{50} in the low

micromolar to submicromolar range for NKCC1 were used as the reference set for the virtual

screening of the Roche 1.2-million-compound library. Compounds were sorted by biological

similarity (i.e. e-value) to the reference set and the top 44 compounds were tested for their NKCC1

activity in the profiling cellular assay. This resulted in the identification of three molecules with an

IC_{50} below 30 μM (Fig. 3). Not only was there a significant improvement in the hit rate by using

BIOSEA compared to traditional approaches based on 2D or 3D chemical information[28], but, in

addition, two of these molecules provided novel chemotypes (Fig. 3) useful for further optimization

efforts. The remaining hit had a very similar chemotype to the reference compounds, which served as

a control result given that BIOSEA operates independently from chemical structure.

BIOSEA can identify new targets for old compounds

Using BIOSEA, we investigated the polypharmacology of known drugs and evaluated the method's

potential to identify novel targets following the procedure described in Fig. 1b.

Our efforts were focused on 711 FDA-approved, withdrawn or experimental drugs. BIOSEA was

used to assess their possible associations with public domain targets from the ChEMBL database[29].

Each molecule was assigned a maximum of 10 predicted targets above a minimum threshold of e-

value, building a pool of 1,124 predictions, out of which 407 could be either experimentally validated

or found in the literature. None of the reported target-compound associations were part of the training

set and none of the attributed targets were part of the HTS fingerprint. Noteworthy, we predicted and

experimentally confirmed 5 novel and unreported compound-target associations for drugs: pimozide,

fluspirilene, reserpine, and nimodipine (Table 2). These examples show how this technology can

provide: (i) deeper insights into the compound's pharmacological profile responsible for its MoA, (ii)

a rationale for previously identified side effects and (iii) opportunities for drug repurposing. **(i)**

Polypharmacology of pimozide and fluspirilene: Pimozide and fluspirilene (Table 2) are

antipsychotic drugs belonging to the diphenylbutylpiperidines (DPBPs) class, used conventionally for

the treatment of schizophrenia and anxiety disorders. The established MoA of DPBPs is attributed to

their potent blockade of the dopamine D_2 receptors. Using BIOSEA, we correctly predicted activities

against 7 other targets reported in the literature of which the serotonin transporter (SERT) and

dopamine D_3 receptor are common to both compounds (Supplementary Data Set 1). Previously

unreported norepinephrine transporter (NET) activity for both pimozide and fluspirilene ($IC_{50} = 110$

nM, 480 nM respectively) was predicted and *in vitro* confirmed. NET is a well-known target for

multiple mood and behavioral disorders such as depression, anxiety or attention deficit hyperactivity.

Given a pimozide plasma concentration of 500 nM (recorded *in vivo* studies[30]), our reported

submicromolar DPBPs activity in the NET is clinically relevant and contributes to the unique

pharmacological profile of DPBPs in their treatment of intricate imbalances of the central nervous

system. In particular, NET modulation may confer DPBPs with stimulant properties that are

advantageous in the treatment of the negative symptoms of schizophrenia (e.g. emotional withdrawal).

Additionally, BIOSEA was able to broadly capture the pharmacological profile of other

antipsychotics (i.e. prochlorperazine and perphenazine), for which it could detect four or more annotations (Supplementary Data Set 1). The efficacy of those drugs in the treatment of schizophrenia and other mental illnesses has been previously linked to their ability to modulate multiple targets[31] predominantly receptors subtypes and transporters of neurotransmitters (adrenaline, dopamine and serotonin). **(ii) Target identification provides a rationale for side effects:** In addition to NET, fluspirilene activity on the adrenergic α_{2B} receptor (ADRA2B, IC_{50} = 95 nM), a target related to vasoconstriction and blood pressure regulation[32] was predicted and experimentally confirmed. Reducing blood pressure in patients with hypertension is documented[33] as a side effect of fluspirilene, to which this ADRA2B activity could contribute. **(iii) Compound repurposing opportunity:** we predicted and experimentally confirmed the glucocorticoid receptor (GR) activity (IC_{50} = 640 nM) for the neuroprotective agent nimodipine. Established as a selective L-type voltage-dependent calcium channel antagonist, nimodipine is often used in the treatment of subarachnoid hemorrhage[34]. Animal studies have shown that nimodipine is able to improve cognitive performance[35], block memory destabilization and prevent memory reconsolidation[36]. On the other hand, it is well established that glucocorticoids influence cognitive performance and have impairing effects on memory function and consolidation[37]. Post-traumatic stress disorder (PTSD) is reported to be associated with increased glucocorticoid receptor sensitivity[38]. Our finding, therefore, suggests opportunities in using nimodipine in connection with psychological disorders associated with GR such as PTSD, although this possibility should be studied in more detail.

Taken together, 31% of validated target-compound relationships could be either positively confirmed, or putatively confirmed when a positive result was found in the literature for an orthologue target or convincing evidence of the relationship has been reported but is not quantified (Fig. 4a). Among compounds with validated predictions, 70 out of 193 were successfully assigned to at least one confirmed or putative target using BIOSEA. The same target identification benchmarking protocol was repeated using an in-house version of SEA[12], which is based on chemical fingerprints

(Supplementary Information). Only 4% of BIOSEA's total predictions were shared with SEA, which

hints at a high degree of complementarity between both methodologies. However, while

approximately 80% of the positive results from SEA can be explained by the high structural similarity

between the query compounds and the training set, only 32% of BIOSEA's confirmed predictions

could have been inferred in this way (Fig. 4c). The remaining predictions are not structurally obvious

and suggest that HTSFP encode information that goes beyond a pure pharmacophore description and

captures more general biological phenomena (see examples Fig. 4d).

Positive predictions from BIOSEA span both functional assays (63%) and binding assays (37%) (Fig.

4b). These percentages are in agreement with the biases in our database regarding the ratio of binding

vs. functional annotations, indicating that the method performs equally well independently to the type

of readout by which these predictions are confirmed. This result is consistent with previous

observations making use of HTSFP which contain both cell-based and biochemical HTS assays[16,22].

An interesting example among the literature-confirmed predictions is the inhibitory activity of

amoxicillin (IC_{50} = 3.47 μM) on carbonic anhydrase XII[39]. This prediction illustrates the ability of

BIOSEA to capture bioisosterism, since it was mostly based in the biologically similarity to

compounds in the training set such as 4-sulfamoylbenzoic acid, sulpiride or indapamide, all of them

sulfonamides. Though structurally unrelated to amoxicillin, all these compounds have in common a

chemical group that is able to interact with the zinc atom in the enzyme's active site (carboxylate or

sulfonamide moieties respectively).

DISCUSSION

Our results demonstrate that BIOSEA is able to classify compounds according to their biological

signature from HTS data. Applied to phenotypic screening, this approach can identify compounds that

are likely to produce a desired phenotype as captured in the biological signature of several reference

compounds. We show that this procedure is possible even in the absence of information on the

particular target(s) responsible for the compounds' mode of action. The results for tubulin and

NKCC1 show that the applicability domain of the method goes beyond the target, target families and

phenotypes that have been historically screened and represented in the HTSFP. HTSFP are unique in

the fact that they capture information on compound inactivity as well as compound activity, both

equally relevant to the mutual information between fingerprints. While the desired phenotype, cell

cycle arrest in this case, can be triggered by multiple mechanisms of action, interestingly, our hits

bind to the same target and follow the mechanism of action of their reference compounds. This

suggests that HTSFP are able to encode target-specific information (i.e. tubulin binding). The

opportunity of using this approach for finding compounds that can reverse a particular disease

phenotype remains an exciting avenue for further exploration.

In addition, the NKCC1 results suggest that the success of the screening effort can be evaluated not

only in terms of hit enrichment but also in the novelty and variety of active chemical structures. For

this reason, BIOSEA becomes an effective tool for building screening libraries for drug discovery

programs, even if the target is novel and challenging, as is the case for NKCC1.

In target identification, our method can be used to discriminate among different drug targets by the

similarity in the biological signatures of their corresponding ligands. Using our protocol we have

explored the pharmacological profile of known drugs. Despite the fact that most of them have been

optimized for selectivity, BIOSEA was able to accurately predict drug polypharmacology in many G-

protein coupled receptors and a wide variety of enzyme classes, consistent with the annotation biases

in public[40] and industrial databases[41] towards specific target families. In particular, we demonstrate

the applicability of our method for drug repurposing and rationalization of drug side effects.

Moreover, the majority of the confirmed predictions cannot be rooted in obvious chemical similarity

to reference compounds. This shows that BIOSEA provides complementary information to traditional

computational methodologies.

Despite its advantages, the approach has some inherent limitations, such as the availability of the HTS

signature for a particular compound and the existence of suitable reference compounds for a given

target or phenotype of interest. Additionally, for target identification, BIOSEA restricts its findings to

the target domain that is known and has several annotated ligands. Nonetheless, with its performance

validated in these in-house drug discovery and target identification scenarios, BIOSEA, offers a

complementary and alternative source for hypothesis generation by capitalizing on already existing

information resources.

Trained solely on public data, our innovative approach is within the reach of both academic and

industrial research groups. We invite the scientific community at large to contribute and explore this

technology by providing the computer code and the HTSFP public data set for download

(https://github.com/accsc/BIOSEA), in the hope that it will foster present and future drug discovery

efforts.

Author contributions: A.C.C. and P.P. designed the BIOSEA method and planned the target

identification experiments. D.L, M. R., I.B. and F.D. designed and performed the cell cycle arrest

phenotypic screening, the tubulin polymerization assay and the tubulin binding competition

experiments. B.F. was the NKCC1 project leader. A.C., P.P, B.F. designed the NKCC1 hit expansion

campaign.

METHODS

Biological fingerprints from HTS: HTS fingerprints (HTSFP) were built using screening data from

PubChem, consisting of 95 primary screening assays over a library of 365,231 molecules in a similar

way to what was described in earlier works[21] (Supplementary Information). HTS results were

normalized using the average response and the standard deviation to transform all the readouts to Z-

Score values. HTSFP-Roche were built in a similar fashion using Roche proprietary assay data.

Biosimilarity metric. A derived version of the mutual information (I) was introduced to estimate the

similarity between two biological fingerprints Z_{NMI}:

3
$$Z_{NMI}(X;Y) = \frac{I(X;Y)\sigma(n)}{\sqrt{H(X)H(Y)}}$$

where X and Y are the HTSFP reduced to the assays in common for compounds x and y; $I(X;Y)$ is the

estimated mutual information using the Kraskov entropy estimator from k-nearest neighbor

distances[42]; $H(X)$ and $H(Y)$ are the Shannon entropies[43]; and $\sigma(n)$ was introduced to correct for the

dependency in fingerprint-size n, parametrized based on a background distribution of random

fingerprint comparisons (a more detailed procedure can be found in Supplementary Information). An

optimal value of $k = 10$ was selected for the calculation of $I(X;Y)$ based on previous parameter

analysis (data not shown) and as trade-off between minimal fingerprint length required and a reliable

Z_{NMI} calculation.

BIOSEA. A statistical method to classify compounds into target classes using biological signatures

from HTS was developed, inspired by the BLAST[23] and SEA[11] algorithms. BIOSEA provides an e-

value that represents the likelihood of a compound to be related by chance to a given compounds set,

by aggregating the biological similarities (Z_{NMI}) between the compound and the members of the set,

and comparing the aggregated value to a random distribution of sums. To build the background

distribution, random lists of HTSFP were generated with sizes in the range $s = 1$ to $s = 100$. The

aggregated biological similarity between every two pairs of lists of compounds with sizes 1 and m

respectively was calculated. The average and standard deviation of the random sums of Z_{NMI} per

number of comparisons ($l \times m$) were calculated and fitted to two power-law equations (Supplementary

Table 2). Aggregated biological similarities were transformed to Z-Scores (Z_{AG}) and plotted in a

histogram that conformed to an Extreme Value Distribution (EVD), whose location, scale and shape

parameters were estimated (Supplementary Table 3). This distribution was used as a background for

probability calculation and its conversion to an expectation value (e-value) that evaluates the significance of the relationships between given sets of fingerprints. The choice of Z_{NMI} threshold between fingerprints selects which compounds are relevant for the aggregated biological similarity and has an effect on the Z_{AG} distribution. Z_{NMI} values of 0.5 and 4.5 in increments of 0.1 were tested and minimal ($Z_{NMI} = 2.6$) and optimal values ($Z_{NMI} = 4.0$) were obtained. More detailed information can be found on Supplementary Information.

Individual assay contributions to the aggregated similarity. For each compound with tubulin depolymerizing activity, we carried out a leave-one-out procedure in which the aggregated biological similarity of an individual compound to the reference compounds (Z_{AG}) was calculated after removing one assay at a time. The difference in Z_{AG} between using the complete fingerprint and the version with a removed component was computed and used to rank the assays. The first 10 assays with the highest similarity difference were selected for analysis.

Target identification database. For the training of BIOSEA, a database of 265 targets and their known ligands was built using dose-response annotations (IC_{50}, EC_{50}, AC_{50}, K_i, K_d, Potency) from ChEMBLdb release 20[29]. Molecules with at least one measurement below 5 μM and none higher than 15 μM were considered active. Only targets with at least 10 annotated ligands were included. Frequent binders (actives in more than 5 assays) were removed from the datasets. A total of 21,157 annotations for 19,808 molecules were collected as the training set for target identification (Supplementary Data Set 2).

Predictions for target identification. 711 known drugs were extracted from DrugBank[44] and ChEMBLdb release 20. Predictions were carried out by BIOSEA trained on the target identification database. The compound biosimilarity threshold was defined as $Z_{NMI}>2.6$. The threshold for compound-target predictions was e-value $> 2.72\text{x}10^{-5}$ (equivalent to $Z_{AG} > 8.0$) resulting in a total pool of 1,124 predictions. The 164 predictions for which the target is represented in the HTSFP were

eliminated from the statistics. A total of 407 remaining predictions were literature or experimentally

verified. ChEMBLdb release 20, DrugBank and Thomsons Reuters Integrity databases were scanned

for activity confirmation. PubMed and EMBASE databases were inspected with the text mining

Linguamatics I2E tool for additional evidence. Predictions were confirmed as active by a retrieved

dose-response value below a threshold of 20 μM, or they were supported by literature annotations

without a numeric readout. A subset of 169 unreported testable hypotheses (accessible compounds

and available assays, see Supplementary Data Set 1) was extracted for experimental evaluation.

Analysis of the 2D compound similarity was performed using ECFP4 fingerprints generated by

Pipeline Pilot 9.2 with a Tanimoto coefficient threshold of 0.8. 3D compound similarity was

calculated using a ROCS-clone[27,45] based on Gaussian shape similarity and a Tanimoto combo

threshold of 1.5.

NKCC1 assay. HEK293 cells conditionally expressing NKCC1 were grown under doxycycline

selection. Cell cultures were maintained at 37 °C in the presence of 5% CO_2 and 85% of humidity. A

volume of 50 μl containing 25,000 cells was plated on poly-D-lysine dishes and induced for 24h.

Cells were washed and incubated at 37°C in the presence of 5% CO_2 for 50 min with a hypotonic

buffer. The supernatant from the cells was removed to a final 21 μl volume and 25 μl of compound

solution diluted in rubidium containing buffer was added. After 10 min incubation, cells were washed

with a hypotonic buffer and 21 μl of 2% tergitol NP-40 was added per well. Plates were sealed and

stored overnight at room temperature. Finally, 30 μl of supernatant for each well were diluted in 170

μl of water and the rubidium content was measured using an atomic absorption spectrophotometer at

780 nm (Aurora Biomed, ICR 12000). Bumetanide was used as the reference compound for

calculation of percentage of inhibition for each compound.

Morphology and cytotoxicity cellular assay. Cytotoxicity evaluation was performed on A549

human lung carcinoma cells (ATCC), employing a modified MTT (3-(4,5-dimethylthiazol-2-yl)-2,5-

diphenyltetrazolium bromide) assay[46]. Indirect immunofluorescence was performed in A549 cells, as

previously described[47]. An absorbance spectrum in methanol was made for each compound in a

spectrophotometer Evolution 201 (Thermo Scientific) in order to identify the absorbance maxima.

Tubulin polymerization assay. The polymerization of 25 M tubulin in glycerol-assembling buffer,

GAB (3.4 M glycerol, 10 mM sodium phosphate, 1 mM EGTA, 1 mM GTP, pH 6.5), in the presence

of 27.5 M of the ligand studied (colchicine and compounds CT1, CT2, CT3, CT4 and CT5) or 2.75 l

of DMSO (vehicle) was monitored along time by turbidity using a Varioskan Flash multimode

microplate reader (Thermo Scientific) at a wavelength of 350 nm and 37 ºC.

Tubulin binding assays. The effect of compounds CT1 and CT4 in the binding of 2-methoxy-5-

(2,3,4-trimethoxyphenyl)- 2,4,6-cycloheptatrien-1-one MTC[48] was studied as previously

described[49,50]. The effect of compound CT3 in the binding of (R)-(+)-ethyl 5-amino 2-methyl-1,2-

dihydro-3-phenylpyrido[3,4-b]pyrazin-7-yl carbamate (R-PT), a well characterized reversible

colchicine-binding were performed as described[51]. The binding of CT5 at the eribulin site was studied

by monitoring the influence of eribulin in the kinetics of binding of the compound to tubulin. The

compound undergoes a large increase in fluorescence at 465 nm (excitation 357 nm) upon incubation

with tubulin. Their kinetics of binding of the compound to 10 M tubulin in NaPi buffer (10 mM

Phosphate Buffer pH 7.0, 0.1 mM GTP) was measured in a Fluoromax-2 fluorimeter (Horiba) in the

presence or absence of 50 M of vincristine, podophyllotoxin, maytansine or eribulin.

Molecular candidates for phenotypic screening. 6 different tubulin binders (i.e. colchicine,

paclitaxel, vinblastine, vincristine, vinpocetine and dolastatin 12) were used as reference compounds

to search in the HTSFP PubChem collection with a Z_{NMI} threshold of 4.0. The resulting 1,559

molecules were structurally clustered using the ECFP4 fingerprints generated by Pipeline Pilot 9.2

and sorted according to Z_{AG}. Known tubulin inhibitors and their chemically similar cluster members

were removed and first 20 compounds were selected for confirmation.

Target identification assays. A detailed list of the assay conditions, detection methods and reference

compounds for the different biochemical and cell-based assay employed in this work can be found in

the Supplementary Table 4.

Code availability. Computer code for the BIOSEA implementation and HTSFP public data set are

available for download from the authors' repository at https://github.com/accsc/BIOSEA.

ACKNOWLEDGEMENTS

A.C. gratefully acknowledges the Roche Postdoctoral Fellowship Program for financial support. The

authors thank specially Dr. Martin Erkens, Dr. Rene Wyler, Dr. Thilo Enderle, Dr. Lisa Sach-

Peltason, Dr Samuel Croset and Dr. Oliv Eidam for their help and thoughtful comments and to Dr.

Klaus Muller for his support and invaluable insight. Eurofins' (CEREP) is gratefully acknowledged

for providing us with single concentration activities from the BioPrint database in advantageous

conditions.

FIGURES:

Figure 1. (a) For virtual screening, BIOSEA calculates the aggregated biological similarity (Z_{AG}) of
each compound in a library against a set of biological signatures from HTS (HTSFP) corresponding to
compounds that are active in a cellular assay. The query library is sorted by Z_{AG} and the top ranking
compounds are selected for a confirmatory screen. (b) For target identification, BIOSEA calculates
the Z_{AG} of a query compound against a list of HTSFP sets $T_1…T_N$. Each fingerprint set represents the
compounds that are active in a given target. Target sets are ranked according to Z_{AG} and the top 10
targets are defined as target predictions for the query compound.

Figure 2. (a) Upon treatment with the compounds, the A549 cells showed type IV aberrant mitotic spindles[52] consisting of a ball of condensed chromosomes and no microtubules. Upper-left: Control; upper-right: colchicine effects at 5 μM; lower-left: CT3 effects at 5 μM; and lower-right: CT5 effects at 25 μM (b)Time-course tubulin polymerization. (c) Displacement curves for the MTC probe in the colchicine binding pocket for compounds CT1, CT4 and podophyllotoxin. (d) Structural model of colchicine-site binders and summary of tubulin binding pockets for the active compounds (Colchicine in green, CT3 in yellow and CT4 in blue).

Figure 3. (Top) Phenotypic screening workflow for NKCC1. The primary screen of a 1.2 million compounds provided starting points for hit expansion within a 2-million-compound virtual library. Hit expansion was carried out by three complementary approaches: chemical, pharmacophore and biological similarity which produced in total ~500 compounds to be tested in a confirmatory dose-response assay, resulting in 4 leads with IC50<30 μM. (Bottom) Chemical similarity (Tanimoto distance of ECFP4 fingerprints) vs. biological similarity (represented by). The regions relevant for chemical and biological expansion are highlighted in red and green respectively. Green and orange dots represent NKCC1 tested biological and pharmacophore similarity predictions.

Figure 4. (a) Performance of BIOSEA in terms of verified predictions. (b) Categorization of positive results by types of assay. Assays were labeled as either functional (e.g. assay description provides a functional end-point for the compound's effect), or binding (e.g. evidence for ligand binding); and biochemical or cell-based if the assay included only the target in an *in vitro* setup, or the whole cell in a more physiologic environment, respectively. (c) Biological similarity goes beyond 2D and 3D similarity. (d) Examples of similarity between predicted and reference molecules. From left to right: 2D similarity, 3D similarity and chemically unrelated (e) Previously unreported results of the target identification protocol for pimozide, fluspirilene, reserpine, and nimodipine.

1   Graphical abstract.

1 **TABLES**

2 Table 1: Phenotypic screening reference compounds (colchicine, vinblastine and paclitaxel) and cell
3 arrest hits (CT1-CT5).

Compound (PubChem CID)	IC$_{50}$ nM	Binding site	Biological similarity reference/s
Colchicine	55.9	Colchicine	-
Vinblastine	4.2	Vinca alkaloid	-
Paclitaxel	4.3	Taxanes	-
CT1 (988603)	577.2	Colchicine	Vincristine
CT2 (6515017)	2,487.7	Unknown	Vinblastine Vincristine
CT3 (5342152)	203.3	Colchicine	Colchicine Vinblastine
CT4 (3244178)	621.1	Colchicine	Vinblastine
CT5 (1211576)	273.0	Eribulin	Colchicine Vinblastine

Table 2: Novel targets for old drugs. Experimental conformation of the target identification
predictions for pimozide, fluspirilene, reserpine, and nimodipine. Chemical structures, novel targets
and experimental activities are included.

Molecule	Target	IC$_{50}$ (nM)	e-value
Fluspirilene	α2B adrenergic receptor	95	2.36x10-6
	Norepinephrine transporter	480	2.85x10-12
Nimodipine	Glucocorticoid receptor	110	1.80x10-5
Reserpine	α1D adrenergic receptor	20,000	8.40x10-7
Pimozide	Norepinephrine transporter	640	3.88x10-6

REFERENCES

Hopkins, A. L. Network pharmacology. *Nature biotechnology* **25**, 1110-1111 (2007).

Medina-Franco, J. L., Giulianotti, M. A., Welmaker, G. S. & Houghten, R. A. Shifting from the single to the multitarget paradigm in drug discovery. *Drug discovery today* **18**, 495-501 (2013).

Hopkins, A. L. Network pharmacology: the next paradigm in drug discovery. *Nature chemical biology* **4**, 682-690 (2008).

Swinney, D. Phenotypic vs. target-based drug discovery for first-in-class medicines. *Clinical Pharmacology & Therapeutics* **93**, 299-301 (2013).

Schenone, M., Dančík, V., Wagner, B. K. & Clemons, P. A. Target identification and mechanism of action in chemical biology and drug discovery. *Nature chemical biology* **9**, 232-240 (2013).

Molina, D. M. *et al.* Monitoring drug target engagement in cells and tissues using the cellular thermal shift assay. *Science* **341**, 84-87 (2013).

Smurnyy, Y. *et al.* DNA sequencing and CRISPR-Cas9 gene editing for target validation in mammalian cells. *Nature chemical biology* **10**, 623-625 (2014).

Rix, U. & Superti-Furga, G. Target profiling of small molecules by chemical proteomics. *Nature chemical biology* **5**, 616-624 (2009).

Li, H. *et al.* TarFisDock: a web server for identifying drug targets with docking approach. *Nucleic acids research* **34**, W219-W224 (2006).

Wang, L. *et al.* TargetHunter: an in silico target identification tool for predicting therapeutic potential of small organic molecules based on chemogenomic database. *The AAPS journal* **15**, 395-406 (2013).

Keiser, M. J. *et al.* Relating protein pharmacology by ligand chemistry. *Nature biotechnology* **25**, 197-206 (2007).

Keiser, M. J. *et al.* Predicting new molecular targets for known drugs. *Nature* **462**, 175-181 (2009).

Gong, J. *et al.* ChemMapper: a versatile web server for exploring pharmacology and chemical structure association based on molecular 3D similarity method. *Bioinformatics* **29**, 1827-1829 (2013).

Liu, X., Vogt, I., Haque, T. & Campillos, M. HitPick: a web server for hit identification and target prediction of chemical screenings. *Bioinformatics*, btt303 (2013).

Lounkine, E. *et al.* Large-scale prediction and testing of drug activity on side-effect targets. *Nature* **486**, 361-367 (2012).

Petrone, P. M. *et al.* Rethinking molecular similarity: comparing compounds on the basis of biological activity. *ACS chemical biology* **7**, 1399-1409 (2012).

Dančík, V. *et al.* Connecting small molecules with similar assay performance profiles leads to new biological hypotheses. *Journal of biomolecular screening* **19**, 771-781 (2014).

Bender, A. *et al.* Chemogenomic data analysis: Prediction of small-molecule targets and the advent of biological fingerprints. *Combinatorial chemistry & high throughput screening* **10**, 719-731 (2007).

Wassermann, A. M., Lounkine, E. & Glick, M. Bioturbo similarity searching: combining chemical and biological similarity to discover structurally diverse bioactive molecules. *Journal of chemical information and modeling* **53**, 692-703 (2013).

Helal, K. Y., Maciejewski, M., Gregori-Puigjané, E., Glick, M. & Wassermann, A. M. Public Domain HTS Fingerprints: Design and Evaluation of Compound Bioactivity Profiles from PubChem's Bioassay Repository. *Journal of chemical information and modeling* (2016).

Riniker, S., Wang, Y., Jenkins, J. L. & Landrum, G. A. Using Information from Historical High-Throughput Screens to Predict Active Compounds. *Journal of chemical information and modeling* **54**, 1880-1891 (2014).

Wassermann, A. M. *et al.* A Screening Pattern Recognition Method Finds New and Divergent Targets for Drugs and Natural Products. *ACS chemical biology* (2014).

Pearson, W. R. Empirical statistical estimates for sequence similarity searches. *Journal of molecular biology* **276**, 71-84 (1998).

Ravelli, R. B. *et al.* Insight into tubulin regulation from a complex with colchicine and a stathmin-like domain. *Nature* **428**, 198-202 (2004).

Ben-Ari, Y. Excitatory actions of GABA during development: the nature of the nurture. *Nature Reviews Neuroscience* **3**, 728-739 (2002).

Khazipov, R., Tyzio, R. & Ben-Ari, Y. Effects of oxytocin on GABA signalling in the foetal brain during delivery. *Progress in brain research* **170**, 243-257 (2008).

Rush, T. S., Grant, J. A., Mosyak, L. & Nicholls, A. A shape-based 3-D scaffold hopping method and its application to a bacterial protein-protein interaction. *Journal of medicinal chemistry* **48**, 1489-1495 (2005).

Cortés-Cabrera, A., Morris, G. M., Finn, P. W., Morreale, A. & Gago, F. Comparison of ultra-fast 2D and 3D ligand and target descriptors for side effect prediction and network analysis in polypharmacology. *British journal of pharmacology* **170**, 557-567 (2013).

Bento, A. P. *et al.* The ChEMBL bioactivity database: an update. *Nucleic acids research* **42**, D1083-D1090 (2014).

Finder, R. *et al.* Pimozide: a review of its pharmacological properties and therapeutic uses in psychiatry. *Drugs* **12**, 1-40 (1976).

Peters, J.-U. Polypharmacology–foe or friend? *Journal of medicinal chemistry* **56**, 8955-8971 (2013).

Muszkat, M. *et al.* Variation in the α2B-adrenergic receptor gene (ADRA2B) and its relationship to vascular response in vivo. *Pharmacogenetics and genomics* **15**, 407-414 (2005).

Cottrell, W. & Magnus, R. V. Fluspirilene in schizophrenia. *Journal of International Medical Research* **1**, 630-633 (1973).

Allen, G. S. *et al.* Cerebral arterial spasm–a controlled trial of nimodipine in patients with subarachnoid hemorrhage. *New England Journal of Medicine* **308**, 619-624 (1983).

Levy, A. *et al.* Nimodipine improves spatial working memory and elevates hippocampal acetylcholine in young rats. *Pharmacology Biochemistry and Behavior* **39**, 781-786 (1991).

Haubrich, J. *et al.* Reconsolidation allows fear memory to be updated to a less aversive level through the incorporation of appetitive information. *Neuropsychopharmacology* **40**, 315-326 (2015).

Roozendaal, B. Stress and memory: opposing effects of glucocorticoids on memory consolidation and memory retrieval. *Neurobiology of learning and memory* **78**, 578-595 (2002).

Labonte, B., Azoulay, N., Yerko, V., Turecki, G. & Brunet, A. Epigenetic modulation of glucocorticoid receptors in posttraumatic stress disorder. *Translational psychiatry* **4**, e368 (2014).

Ozensoy, O., Arslan, O. & Kockar, F. Differential in vitro inhibition effects of some antibiotics on tumor associated carbonic anhydrase isozymes of hCA-IX and hCA-XII. *Journal of enzyme inhibition and medicinal chemistry* **23**, 579-585 (2008).

Overington, J. P., Al-Lazikani, B. & Hopkins, A. L. How many drug targets are there? *Nature Reviews Drug Discovery* **5**, 993-996 (2006).

Peters, J. U., Schnider, P., Mattei, P. & Kansy, M. Pharmacological promiscuity: dependence on compound properties and target specificity in a set of recent Roche compounds. *ChemMedChem* **4**, 680-686 (2009).

Kraskov, A., Stögbauer, H. & Grassberger, P. Estimating mutual information. *Physical review E* **69**, 066138 (2004).

Shannon, C. E. A mathematical theory of communication. *The Bell System Technical Journal* **27**, 379 - 423 (1948).

Law, V. *et al.* DrugBank 4.0: shedding new light on drug metabolism. *Nucleic acids research* **42**, D1091-D1097 (2014).

Cabrera, A. l. C. s. *et al.* CRDOCK: An Ultrafast Multipurpose Protein–Ligand Docking Tool. *Journal of chemical information and modeling* **52**, 2300-2309 (2012).

Yang, C. *et al.* Overcoming Tumor Drug Resistence Mediated by P-glycoprotein Overexpression with high affinity taxanes: A SAR study of C-2 Modified 7-Acyl-10-Deacetil Cephalomannines. *ChemMedChem* **2**, 691-701 (2007).

de Ines, C. *et al.* Inhibition of microtubules and cell cycle arrest by a new 1-deaza-7, 8-dihydropteridine antitumor drug, CI 980, and by its chiral isomer, NSC 613863. *Cancer research* **54**, 75-84 (1994).

Fltzgerald, T. J. Molecular features of colchicine associated with antimitotic activity and inhibition of tubulin polymerization. *Biochemical pharmacology* **25**, 1383-1387 (1976).

Andreu, J. M., Perez-Ramirez, B., Gorbunoff, M. J., Ayala, D. & Timasheff, S. N. Role of the colchicine ring A and its methoxy groups in the binding to tubulin and microtubule inhibition. *Biochemistry* **37**, 8356-8368 (1998).

La Regina, G. *et al.* Arylthioindole inhibitors of tubulin polymerization. 3. Biological evaluation, structure-activity relationships and molecular modeling studies. *Journal of medicinal chemistry* **50**, 2865-2874 (2007).

Canela, M.-D. *et al.* Novel colchicine-site binders with a cyclohexanedione scaffold identified through a ligand-based virtual screening approach. *Journal of medicinal chemistry* **57**, 3924-3938 (2014).

Jordan, M. A., Thrower, D. & Wilson, L. Effects of vinblastine, podophyllotoxin and nocodazole on mitotic spindles. Implications for the role of microtubule dynamics in mitosis. *J cell Sci* **102**, 401-416 (1992).