Oomycetes during 120,000 years of temperate rainforest ecosystem development

Ian A. Dickie[1*], Angela M. Wakelin[2], Laura Martinez-Garcia[3], Sarah J. Richardson[4], Andreas

Makiola[2], Jason M. Tylianakis[1,5]

*Corresponding author: Bio-Protection Research Centre, School of Biological Sciences,

Canterbury University, Christchurch, New Zealand.

[1] Bio-Protection Research Centre, School of Biological Sciences, University of Canterbury,

Christchurch, New Zealand.

[2] Bio-Protection Research Centre, Lincoln University, Lincoln, New Zealand.

[3] Department of Soil Quality, Wageningen University, Wageningen, The Netherlands.

[4] Landcare Research, Lincoln, New Zealand.

[5] Department of Life Sciences, Silwood Park Campus, Imperial College London, Ascot, UK.

author emails: Ian.Dickie@canterbury.ac.nz; Angela.Wakelin@lincoln.ac.nz;

LauraBeatrizMartinez@gmail.com; RichardsonS@landcareresearch.co.nz;

Andreas.Makiola@lincolnuni.ac.nz; Jason.Tylianakis@canterbury.ac.nz

Running title: **Oomycetes and ecosystem development**

Suggested subject category: **Microbial ecology and functional diversity of natural**

habitats

Keywords: *Forest diversity, molecular ecology, Oomycetes, pathogen ecology, pedogenesis,*

plant–soil (below-ground) interactions, retrogression, succession

Abstract

The occurrence of plant-associated oomycetes in natural ecosystems and particularly during

long-term ecosystem development is largely unknown, despite the importance of many

oomycetes as plant pathogens. Using DNA sequencing from roots, we investigated the

frequency and host relationships of plant-associated oomycete communities along a 120 000

year glacial chronosequence, comprising site ages with rapid compositional change ("early

succession"; 5, 15, and 70 years old soil); relatively stable higher-diversity sites ("mature",

280, 500, 1000, 5000, 12000 years); and ancient, nutrient-limited soils with declining plant

diversity and stature ("retrogression", 60 000, 120 000 years). Plant-associated oomycetes

were frequent in all three early successional sites, occurring in 38 – 65% of plant roots, but

rare (average 3%) in all older ecosystems. Oomycetes were highly host specific, and more

frequent on those plant species that declined most strongly in abundance between ecosystem

ages. The results support the particular importance of plant-associated oomycetes in early

succession (up to 70 years). High host specificity and correlations of abundance of oomycete

inside roots with declining plant species are consistent with oomycete-driven successional

change.

Introduction

As ecosystems develop, biotic communities undergo succession both in composition and in the nature of mutualistic and antagonistic interactions. Many of these changes have been well studied, including changes in plant communities and traits, competition intensity, and mutualisms (Lambers et al, 2008). Nonetheless, long-term successional changes in composition and community interactions of many other organisms, such as oomycetes, remain poorly understood. Indeed, oomycetes in natural ecosystems are often out-of-sight out-of-mind, until large-scale forest diebacks occur (Scott & Williams, 2014). This lack of information is particularly apparent in forest disease outbreaks, where it is often unclear whether the oomycete is native, from where it may have originated, or what background levels of oomycetes are typical in healthy ecosystems (Podger & Newhook, 1971). Oomycetes occur as part of healthy ecosystems and play important roles in ecosystem processes (Castello et al, 1995; Gómez-Aparicio et al, 2012; Bever et al, 2015; Geisen et al, 2015). For example, *Pythium* and other oomycetes (Oomycota = Peronosporomycota) can drive negative density dependence in plants (Packer & Clay, 2000) and hence are believed to contribute to the maintenance of forest plant diversity (Mangan et al, 2010a). Oomycetes are also an important component of soil biodiversity, as pathogens of plant roots and soil invertebrates, and as saprotrophs (Arcate et al, 2006; Geisen et al, 2015). One way to better understand the role of oomycetes in natural ecosystems is to look at gradients of ecosystem development, which have a long history of providing insight into plant communities.

A useful way to consider ecosystem development, at least from a plant-community perspective, is based on compositional change through three phases: early succession, mature, and retrogression. Early succession (both primary and secondary) can be defined as the

period in which communities undergo major compositional change. In the context of plant

communities, this means that early successional plants are unlikely to regenerate under

conspecifics and are instead adapted for dispersal into new sites. Mature ecosystems can be

defined by compositional stability over timespans greater than one generation, such that

species are regenerating in the presence of established conspecifics (Veblen, 1992). Finally,

over very long time periods ecosystem retrogression eventually develops, where the

progressive loss of nutrients leads to a decline in plant diversity and a loss of biomass and

canopy stature (Peltzer et al. 2010). This shift to retrogression entails slow compositional

change, favouring species with life history traits that are adapted to stressful, low-nutrient

environments.

Although oomycetes can be saprotrophic, they have been proposed to play particularly

important roles at all stages of ecosystem development as potential plant pathogens. One

view of soil pathogens in relation to ecosystem development is that they play a critical role in

driving compositional change in early succession (Van der Putten et al, 1993; Van Der

Putten, 2003). This view is based, in part, on the assumption that newly establishing plant

species will encounter only low levels of pathogens in the absence of conspecific host plants

during succession. Over time, these established plants accumulate pathogens, which prevent

establishment of conspecific seedlings but have less negative effects on seedlings of other

plant species. Part of the role of pathogens in succession is driven by plant age, as pathogens

that kill seedlings are often tolerated by established plants (Martin & Loper, 1999; Van Der

Putten, 2003). This implies that early-successional (r-selected) plants may have limited

selection for pathogen resistance. Instead, these species may accumulate and tolerate

pathogens as mature plants, and rely on seed dispersal into new habitats to escape pathogens

during the vulnerable seedling stage (Van Der Putten, 2003). Support for the role of

antagonists in early succession comes from studies of invertebrate antagonists in grasslands,

where dominant early-successional plants are more suppressed than subordinate or late-

successional species (De Deyn et al, 2003) and from studies showing negative plant-soil-

feedback early in succession (Kardol et al, 2006). Oomycete pathogens have also been shown

to be important in the mortality of the early-successional tree *Prunus serotina* (Packer &

Clay, 2000; Packer & Clay, 2003). These examples span from short-term grassland to decadal

scale forest patterns, suggesting a widely applicable ecological pattern.

While much evidence supports the role of pathogens in early succession, there are also

suggestions that pathogens, in general, increase in mature ecosystems (Reynolds et al, 2003;

Bardgett et al, 2005; Peltzer et al, 2010). Soil biotic communities as a whole increase in

biomass and species diversity over periods of tens to hundreds of years (Bardgett et al, 2005),

and pathogen communities may follow similar patterns of increasing biomass and diversity.

Further, plant competition is high in mature ecosystems, and Gilbert (2002) suggests that

competition may stress plants and increase susceptibility to pathogens. Pathogen host-

specificity in these systems leads to density-dependent mortality (Augspurger, 1984).

Evidence for negative density dependence has come from both tropical and temperate forests

(Packer & Clay, 2003; Mangan et al, 2010b), although other studies have found positive

density dependence or no effect (Reinhart et al, 2012).

Finally, a recently proposed third view of the role of pathogens in ecosystem development is

that soil pathogens may be particularly critical in retrogressive ecosystems (Laliberté et al,

2015), where declining soil nutrient availability results in declining vegetation stature and

diversity (Richardson et al, 2004; Porder et al, 2007; Peltzer et al, 2010). Laliberté et al.

(2015) suggest that the highly weathered, P-limited soils of retrogressive ecosystems favour

species with ephemeral roots to maximize nutrient uptake, but that this imposes a trade-off

with increased susceptibility to pathogens. Ecosystem retrogression has been linked to

increased specific root length (length per unit mass), thinner roots, and increased root

branching (Holdaway et al, 2011), all of which may increase susceptibility to root pathogens

(Laliberté et al, 2015).

Based on the three views from the literature, we tested the non-exclusive hypotheses that

oomycetes would be associated with (1) early-successional, (2) mature, and/or (3)

retrogressive ecosystems. We used an extensively studied soil and ecosystem development

chronosequence created by the Franz Josef glacier in New Zealand, where glacial advances

and retreats have created a series of soils of varying age in close proximity (Walker & Syers,

1976; Richardson et al, 2004), allowing space-for-time substitution as an approach to infer

change over long time periods (Pickett, 1989). We focussed on oomycetes in living plant

roots, in order to link oomycete communities to particular hosts. The presence of an

oomycete in roots of specific plant species does not necessarily indicate that it causes disease

in that plant. Indeed, mature plants may often tolerate organisms that would kill conspecific

seedlings. It is also conceivable that oomycetes are living within the roots of plants on which

they would never cause disease, regardless of plant age. Nonetheless, while the presence of

oomycetes does not necessarily indicate disease, the absence of oomycetes from a plant can

be taken as an indication of absence of oomycete-driven disease.

Materials and Methods

Study site and sampling

The Franz Josef chronosequence includes early-successional (5, 15, 70 years of

development), mature (280, 500, 1000, 5000, 12 000 years), and retrogressive (60 000, 120

000 years) sites. As soils age, there are dramatic changes in nutrient availability (declining P,

increasing and then decreasing N), pH, and physical properties (Walker & Syers, 1976).

These changes are linked to changes in plant communities, with plant biomass increasing

through succession to mature stages, and then entering retrogression where declining soil

nutrients result in a concomitant decline in plant biomass, stature, and diversity (Richardson

et al, 2004) and shifts in root traits (Holdaway et al, 2011). Vegetation shifts from sparse

herbaceous plants and sub-shrubs (5 years) to shrub land (15 years) to angiosperm forest (70

years), followed by an increasing dominance of large gymnosperm trees (Podocarpaceae)

through mature stages, with an eventual decline in plant biomass, canopy height, and canopy

closure in retrogression. Rainfall is high along the entire chronosequence, ranging from 3.5 to

6.5 m, and all sites are below 365 m elevation (Richardson et al, 2004).

In a previous study, we collected 510 individual plant roots from ten sites along the

chronosequence and characterised both plant identity and arbuscular mycorrhizal fungal

communities in those roots (Martínez-García et al, 2015). Here we use these same samples

but used taxon-specific primers to amplify DNA from oomycetes. The collection of samples

is described in full in Martinez-García *et al.* (2015). In brief, we collected 51 or 52 (at two

sites) individual fine root (< 2 mm diameter) fragments, taking a single root fragment

(approx. 15 mg dry weight) every 2 m along two parallel transects with equal sampling from

three depths (up to 20 cm depth max). This depth is likely representative of the established

roots a seedling might encounter during the first year of growth and could be sampled in a

standardised way at even our youngest sites, but may omit deeper roots in older sites. For the

5-year-old site, which comprised sparse vegetation in rocks, most soil was root free. We

therefore collected the nearest plant to the sample point and sampled one root from that plant.

Roots from all sites were thoroughly rinsed in water and dried before DNA extraction. DNA

was extracted using a MoBio soil DNA kit, and plant species identified by PCR amplification

and DNA sequencing of the tRNL gene region, except for the 5-year-old site, where the plant

was already known due to collection method. The one difference in the sampling between

this study and the prior study (Martínez-García et al, 2015) was that in the earlier paper a

spare ("B") sample was used if the first ("A") sample failed to yield both plant and

arbuscular-mycorrhizal fungal PCR products. In the present study we did not use the "B"

samples, as it would have made quantification difficult.

Oomycete PCR and identification

We primarily based our oomycete detection, operational taxonomic unit (OTU) clustering,

and identification on a nested PCR of large-subunit (28S) DNA (supplementary methods).

Initial results using T-RFLP suggested that most samples contained either no oomycetes or

only a single OTU, hence direct sequencing was used for identification. A negative (omitting

DNA template) and a positive control (including genomic DNA from a *Pythium* species)

were included in every PCR. PCR products were purified using DNA Clean and Concentrator

Kit (Zymo Research Corporation) prior to performing Sanger sequencing (Canterbury

Sequencing and Genotyping, University of Canterbury, New Zealand).

Some ecosystem ages had a high proportion of samples that failed to amplify a product of the

expected size in the first-round PCR. In order to ascertain whether these were true absences

of oomycetes or due to PCR inhibition, nine samples from each of these sites with the highest

failure rate (500, 100, 1200, 60 000 and 120 000 yrs) were tested for PCR inhibitors by

repeating PCR reactions in duplicate for each sample, with one of the duplicate reactions

spiked with 20 ng positive control DNA. Only one out of the 45 samples spiked with positive

control DNA failed to amplify, suggesting PCR inhibitors were not likely to be causing low

detection rates.

To confirm identities of sequenced oomycetes, samples that produced positive large subunit

PCR products were sequenced for the internal transcribed spacer (ITS) 1 region

(supplementary material). As DNA sequences were obtained from environmental samples

(root fragments) potentially containing multiple oomycetes, the ITS sequence may or may not

represent the same species as the 28S sequence. ITS sequences were therefore used to help

inform identification, but frequency analyses were based on the 28S results.

Vegetation cover

The percent cover of all vascular plants within sites was measured following the standard

relevé (or 'recce') method for describing New Zealand vegetation (Hurst & Allen, 2007) in a

20 x 20 m area located within 50 m of the root sample transects. For testing the correlation of

plant cover change with oomycete frequency we used the mid-point of vegetation cover

classes summed across height tiers as a measure of the abundance of each plant species.

Sequence bioinformatics

All DNA sequences were matched against GenBank using BLAST to find the closest

matching sequence, and, where the closest match had no reliable taxonomic identity, the

closest matching sequence associated with a taxonomic identity. Sequences that matched

non-oomycete specimens were discarded. Three sequences had a closest match to a

Spongospora (Cercozoa) sequence, but only at 78% identity. We therefore considered these

three sequences to likely be root-associated protists (Bulman & Braselton, 2014) and retained

them in the analysis. Out of 510 root samples, 122 had positive PCR products with oomycete

primers. After filtering for sequence quality and matching to an oomycete sequence, the final

dataset contained 91 DNA sequences of oomycetes with a mean sequence length of 310 bp.

Plant IDs were obtained for 458 samples overall and 86 of the 91 samples with oomycete

products through DNA sequencing of the trnL gene (Martínez-García et al, 2015). We

clustered sequences into OTUs using BLASTn to merge any sequences with > 97% similarity

over at least 95% of the shorter sequence length based on single-linkage clustering. This

allowed us to form species concepts for testing host specificity, resolved some matches to

"uncultured oomycete" and suggested where the gene region used lacked sufficient resolution

to distinguish species.

Statistics

Most of our analyses are based on the frequency of roots with oomycetes, testing the

prevalence of oomycetes within different plant species and sites, rather than quantifying

abundance within any particular root fragment. Changes in the frequency of oomycetes as a

function of successional stage were tested by treating age as a three-level factor. Sites were

allocated to successional stage levels on the basis of soil nutrient concentrations, vegetation

height, species richness and biomass, and plant traits (Walker & Syers, 1976; Richardson et

al, 2004). Early-successional sites (5, 15, 70 years) had abundant N-fixing trees; vascular

plant traits of high foliar nutrient concentrations, low leaf mass per unit area; very high soil P,

rapidly accumulating biomass and species richness of vascular plants, and increasing nitrate-

N. Mature ecosystems (280 to 12 000 years) had slow rates of change in species richness,

height and biomass among sites. These sites also had the highest biomass and plant diversity.

Retrogressive sites (60 000, 120 000 years) had a decline in vegetation height, richness and

biomass and exceptionally proficient phosphorus resorption (Richardson et al, 2005; Peltzer

et al, 2010; Vitousek et al, 2010).

We tested changes in the frequency of oomycetes as a function of successional stage

as a factor and log-transformed ecosystem age as a linear variate using a binomial glm, fitted

with quasi-likelihood to account for over-dispersion. The difference between the successional

stage and ecosystem age models was tested using analysis of deviance, using an F test as

recommended for quasibinomial fit. For these analyses $n = 10$ based on 512 individual

samples.

We wanted to test whether declining plant species had higher than random oomycete

levels, as would be expected if oomycetes are an important driver of community changes

with ecosystem development. To do so, we calculated the change in cover of each plant

species from one time point (t_0) to the next (t_1) as a log ratio ($\log((\text{cover } t_0 + 1) / (\text{cover } t_1$

$+ 1)))$ and then used a binomial mixed effects model to test whether the presence / absence of

oomycetes in root fragments at t_0 could be predicted by change in cover from t_0 to t_1 across

the nine intervals between site ages. This test was carried out using lmer in the R package

lme4, with site age and plant species (to account for species occurring across multiple sites)

as random effects and the canonical logit link function. Host-specificity was tested within

ecosystem age using chi-square tests.

Results

Oomycete frequency as a function of stage of ecosystem development and ecosystem age

In early-successional ecosystems (5, 15 and 70 years), 38 to 65% of root samples had

oomycete DNA detected (Figure 1). In contrast, in mature and retrogressive ecosystems (280

to 120 000 years) an average of only 2% and never more than 8% of roots had oomycete

DNA found (Figure 1). Treating the sites as representing three stages, the early successional

stages had a significantly higher frequency of oomycetes than mature stages (quasibinomial

family glm; $t = -6.6$, $P = 0.00050$) or retrogressive stages ($t = -4.6$, $P = 0.0025$), and the

difference between mature and retrogressive ecosystems was not significant ($t = -1.1$, $P =$

0.32).

Oomycete frequency was also significantly related to log-transformed ecosystem age

(quasibinomial family glm; $t = -2.75$, $P = 0.025$). While statistically significant, this

relationship was a poor fit to the observed pattern of an increase for the first 3 site ages

followed by uniformly low levels of oomycetes at all subsequent site ages (Figure 1).

Analysis of deviance strongly supported the ecosystem-stage categorical model over the

linear ecosystem-age model ($F = 35.11$, $P = 0.00058$).

Those plant species that showed the largest decline in cover between ecosystem ages also had

the highest frequency of oomycetes in their roots (Figure 2, $z = -2.4$, $P = 0.018$).

Oomycete diversity, host specificity and ecosystem age

A total of 37 different OTUs were found, with 10 OTUs found more than once (Table 1). The

most frequent OTU occurred 17 times, and had DNA-sequence affinities to *Lagena*

radicicola and an uncultured oomycete from New York, USA agricultural soils (Table 1).

Within ecosystem ages, there was a significantly non-random distribution of oomycete OTU

identity across plant species at 5 (5 oomycete OTUs with 20 occurrences across 4 plant

species, χ^2 = 42.96, df = 12, p-value = 0.000022) and 15 years (15 oomycete OTUs, 28

occurrences, 14 plant species, χ^2 = 225.87, df = 182, p-value = 0.015), but only marginally

significant at 70 years (16 oomycete OTUs, 30 occurrences, 7 plant species, χ^2 = 109.85, df =

90, p-value = 0.076; Figure 3). There were too few observations at the ecosystem stages after

70 years to meaningfully test for plant by OTU host-specificity at later ecosystem stages.

Discussion

Our results strongly support the importance of ecosystem age as a determinant of oomycete

communities in plant roots, with high frequencies of oomycetes observed in the three earliest

ecosystem stages (5, 15, and 70 years) but only very low levels thereafter. Although the

importance of oomycetes as pathogens is often discussed in the ecological literature (Bagchi

et al, 2014), relatively few studies have actually quantified oomycete frequency in natural

ecosystems (Gómez-Aparicio et al, 2012). Further, we believe our results are the first study of

oomycetes throughout a complete ecosystem development sequence. The community of

oomycetes was diverse, including multiple *Pythium* and *Phytophthora* species and,

surprisingly, a very frequent sequence matching *Lagena* on two small sub-shrubs: *Raolia*

hookeri (Asteraceae) and *Epilobium glabellum* (Onograceae). *Lagena* has previously only

been reported as a pathogen on grass (Blackwell, 2011).

In terrestrial ecosystems, oomycetes occur as pathogens and commensals of plants

and other eukaryotes and, less commonly, as saprotrophs (Marano et al, 2016). By sampling

plant roots, our results focussed primarily on endophytes. Our observation of high host-

specificity is also consistent with a biotrophic interaction and the correlation of oomycete

frequency with subsequent plant decline is consistent with pathogenicity. However, while we

believe that the oomycetes we observed were primarily biotrophic, the gene regions we

sequenced lacked sufficient resolution to rule out the possibility that some of the

Phytophthora we observed could be in clade 9, which contains some known saprotrophs

(Marano et al, 2016). The other clades of oomycetes where saprotrophy is known to occur

(Marano et al, 2016) were not observed.

Ecological role of oomycetes in plant roots in early successional ecosystems

It has been suggested that pathogens, including oomycetes, drive early-successional

change in plant communities (Kardol et al, 2006). The high frequency of oomycetes in early

successional ecosystems is consistent with this hypothesis. In addition, those plant species

that had the highest frequency of oomycetes at a given ecosystem age declined in percent

cover most strongly before the next age. The higher frequency of oomycetes on plant roots in

early-successional ecosystems compared to mature and retrogressive ecosystems was not due

to PCR inhibition in older sites, as samples spiked with positive control oomycete DNA

showed no evidence of inhibition.

Oomycetes, particularly *Pythium*, are often tolerated as endophytes by established

plants, but prevent seedling establishment of the same plant species (Martin & Loper, 1999;

Van Der Putten, 2003). On that basis, we do not argue that the observed oomycetes were

necessarily having any direct negative effect on established plants. Instead, we suggest that

the correlation of oomycete frequency with decline in cover between ecosystem ages is

congruent with the suggestion that these pathogens are primarily preventing re-establishment

of plants and hence contributing to vegetation change across cohorts (Van Der Putten, 2003).

Our definition of "early succession" comprises the major transition from rock field (5

years) to shrub land (15 years) to forest (70 years). Therefore, our finding of high pathogen

levels is potentially consistent with Packer & Clay (2000), who found strong negative

feedback in *Prunus serotina* in a forest described as "at least 70 years old" (Packer & Clay,

2000), despite the differences between our primary succession and what was likely a

secondary succession in the Packer & Clay study.

We believe most of the change in oomycete frequency across ecosystem stages

reflects changes in plant species composition. While oomycete frequency was significantly

correlated with site age as a linear predictor, treating sites as three distinct stages resulted in a

much stronger model fit. Nonetheless, changes in oomycete frequency may also, in part,

reflect a substantial site-age effect and direct effects of changing soil environments on

oomycete populations (akin to the "habitat" hypothesis for mycorrhizal fungi of Zobel &

Opik 2014). Meaningful testing of soil variables as predictors of oomycetes was not possible,

as soil properties were strongly correlated with soil age.

Absence of oomycetes in plant roots in older ecosystems

It has been suggested that oomycetes are important in mature (Bagchi et al, 2010;

Bagchi et al, 2014) and retrogressive ecosystems. While Packer and Clay (2000) clearly

demonstrated oomycetes determined seedling dynamics in a relatively young temperate

forest, a more recent study by Bagchi and colleagues (2014) in a tropical seasonal forest

found that eliminating fungi alone had a stronger effect on negative density-dependence than

eliminating both fungi and oomycetes. Laliberté et al. (2015) suggested that root traits

adapted to highly P-limited retrogressive ecosystems would increase susceptibility to root

pathogens, and that oomycetes could drive plant diversity in these ecosystems (Albornoz et

al, 2017).

Our results do not support an important role of root associated oomycetes in this

retrogressive sequence, which contrasts with the recent findings from a western Australian

system where oomycetes play key roles in older ecosystems (Albornoz et al, 2017). In part

this may reflect the importance of non-mycorrhizal Proteaceae in Australia. The very fine

roots of the Proteaceae may make this plant family particularly susceptible to pathogens

(Laliberté et al, 2015). Further comparative studies across multiple retrogressive ecosystems

are needed to resolve the circumstance under which each pattern is more common.

Novel organisms or native?

Having baseline data on oomycete ecology may be important for understanding and

managing future pathogen outbreaks. For example, *Phytophthora agathidicida* (=

Phytophthora taxon *Agathis*) has been implicated as a cause of forest dieback of the iconic

New Zealand-endemic *Agathis australis*, but, like many other oomycete disease outbreaks,

the origin of *Phytophthora agathidicida* remains uncertain (Weir et al, 2015). The single-

direction sequencing we performed directly from environmental samples was not designed

for accurate phylogenetic classification, even with two gene regions sequenced per sample,

but it does suggest considerable diversity of *Phytophthora*. We found sequences with

affinities to a number of species already known to be present in New Zealand, including *P.

cinnamomi*, *P. cactorum*, *P. infestans* and *P. kernoviae* (Scott & Williams, 2014). Given that

our samples came from relatively pristine ecosystems, we believe these OTUs are likely to be

from native *Phytophthora*.

More surprising was that the most common OTU in our data had affinities to *Lagena

radicicola* in both 28S and many of the ITS sequence results. *Lagena* is presently considered

to be a monotypic genus widespread as a pathogen of grasses in North America (Blackwell,

2011), but Barr and Désaulniers (1990) suggest it may be under-reported due to being

morphologically very similar to *Pythium*, and not being easily culturable. Spores described as

"resembling *Lagenocystis* [syn. *Lagena*] spp." were noted by Skipp and Christensen (1989) in

New Zealand *Lolium perenne* pastures, but our finding is the first report of a putative *Lagena*

species outside of grasses. We believe this most likely reflects a lack of prior knowledge, as

very few prior studies have used molecular methods to detect oomycetes from healthy forest

ecosystems, and none of those have taken place in temperate southern hemisphere rainforest.

Conclusions

Much of the Earth's surface is covered in young, early successional ecosystems

comparable to our early successional sites of 5, 15, and 70 years (Vitousek et al, 1997;

Haddad et al, 2015). The strong plant-host specificity and correlation of plant decline and

oomycete frequency is consistent with the suggestion that these oomycetes as pathogens

contribute to vegetation succession. While we do not find frequent oomycetes in older

ecosystems, oomycete DNA was detected in all but one site. These low levels of oomycetes

in roots, along with oospores in soil, may still be sufficient to prevent establishment of

susceptible plant species or to drive pathogen outbreaks following external climatic stressors,

but seem incompatible with density dependent mortality being driven by oomycetes on

established plant roots. More broadly, our results support the concept that oomycetes are a

diverse part of native ecosystems. Understanding this native diversity in is critical to better

understand, prevent, and manage future oomycete disease outbreaks (Dickie et al, 2017).

Acknowledgements. We thank D. Peltzer for help with locating plots, characterising

vegetation, and editorial input, R. Holdaway for helping with data and other advice, H.

Ridgway for providing positive control DNA, N. Merrick for sequencing, R. Buxton and C.

Morse for collecting additional plant trait data, and S. Wakelin, L. Condron, J. Bufford, S.

Bulman and the Mycorum discussion group at Lincoln University who provided valued

advice and input. Funding was provided by Tertiary Education Commission funding for the

Bio-Protection Research Centre (I.A.D., AMW, JMT), Lincoln University Research Fund

(IAD, AMW), a Rutherford Fellowship (JMT), and Core Funding to Crown Research

Institutes (LMG, SR).

Data Accessibility

- DNA reference sequences: GenBank (accession numbers in Table 1)

- Vegetation data are archived in the New Zealand National Vegetation Survey Databank

(https://nvs.landcareresearch.co.nz; last accessed 8 Feb 2016).

Table 1. Identity of non-singleton OTUs as clustered based on 28S, with accession numbers and best matching 28S and best matching ITS1

sequences, the inferred identity, and the number of sequences in the OTU.

28S accession	Best match based on 28S (% identity / % length; sequence length)	ITS1 accession	Best match based on ITS (% identity / % length; sequence length)	Inferred identity (clustering based on 28S)	n
KU863596	Uncultured oomycete 98%/98% 407 bp	KU863605	*Lagena radicicola* 93%/99% 205 bp	*Lagena* sp.	17
KU863595	*Pythium dissotocum* 97%/100% 379 bp	KU863604	*Pythium intermedium* 90%/47% 302 bp	*Pythium* aff. *aquatile/dissotocum*	13
KU863588	*Phytophthora cinnamomi* 99%/100% 385 bp	KU863598	*Phytophthora cinnamomi* 99%/100% 254 bp	*Phytophthora* aff. *cinnamomi*	8
KU863589	Uncultured oomycete 98%/100% 380 bp	KU863599	*Pythium* sp. 92%/100% 203 bp	*Pythium* sp.	6
KU863594	*Pythium tracheiphilum* 98%/99% 370 bp	KU863603	*Pythium tracheiphilum* 89%/98% 309 bp	*Pythium* aff. *tracheiphilum*	6
KU863593	*Pythium junctum*	KU863602	*Pythium irregulare*	*Pythium* aff. *junctum*	4

	98%/99% 378 bp		100%/28% 307 bp	
KU863597	*Phytophthora megasperma*	KU863606	*Phytophthora infestans*	*Phytophthora* sp. 1	4
	91%/100% 383 bp		95%/53% 447 bp	
KU863590	*Spongospora* sp.	N/A	No PCR product obtained	Chromista aff. *Spongospora*	3
	75%/79% 306 bp			
KU863591	*Phytophthora polonica*	KU863600	*Phytophthora kernoviae*	*Phytophthora* sp. 2	3
	99%/99% 384 bp		96%/99% 214 bp	
KU863592	*Pythium violae*	KU863601	*Pythium* sp.	*Pythium violae*	3
	98%/100% 368 bp		99%/99% 286 bp	

Figure 1. Frequency of oomycetes in plant roots (%) as a function of ecosystem age. The 5,

15, and 70-year-old sites represent a successional sere from rock field through shrubland to

angiosperm forest. Sites 280 to 5000 years of age are mature forest with high biomass and

plant diversity, while after 12 000 to 60 000 years plant biomass, stature, and diversity

decline due to retrogressive P-limitation.

Figure 2. Oomycete frequency as a function of vegetation decline with circles representing

each plant species between site transitions. Points scaled by number of samples for each plant

species within site. Lines indicate random effects to account for site in a binary mixed effects

model with random intercepts for site and plant species. Colour coding in reflects the site age

at t_0 for each transition, hence no points for 120 000 years are shown.

Figure 3. Plant by site age visualisation of oomycete frequency within sampled roots. Each

square represents a single root sample, with coloured squares indicating the presence of

oomycete pathogens with colours indicating OTU and black indicating an OTU found only

once. See Table 1 for OTU information. Grey dots indicate species not found in root samples

for that site.

Supplemental files S1: Detailed molecular methods and discussion.

References

Albornoz FE, Burgess TI, Lambers H, Etchells H, Laliberté E. (2017). Native soilborne

pathogens equalize differences in competitive ability between plants of contrasting

nutrient-acquisition strategies. *Journal of Ecology* 105: 549–557.

Arcate JM, Karp MA, Nelson EB. (2006). Diversity of peronosporomycete (oomycete)

communities associated with the rhizosphere of different plant species. *Microbial*

Ecology 51: 36–50.

Augspurger CK. (1984). Seedling survival of tropical tree species: interactions of dispersal

distance, light-gaps, and pathogens. *Ecology* 65: 1705–1712.

Bagchi R, Gallery RE, Gripenberg S, Gurr SJ, Narayan L, Addis CE *et al.* (2014). Pathogens

and insect herbivores drive rainforest plant diversity and composition. *Nature* 506: 85–

88.

Bagchi R, Swinfield T, Gallery RE, Lewis OT, Gripenberg S, Narayan L *et al.* (2010).

Testing the Janzen-Connell mechanism: pathogens cause overcompensating density

dependence in a tropical tree. *Ecology Letters* 13: 1262–1269.

Bardgett RD, Bowman WD, Kaufmann R, Schmidt SK. (2005). A temporal approach to

linking aboveground and belowground ecology. *TRENDS IN ECOLOGY &*

EVOLUTION 20: 634–641.

Barr DJS, Désaulniers NL. (1990). The life cycle of Lagena radicicola, an oomycetous

parasite of wheat roots. *Canadian Journal of Botany* 68: 813–824.

Bever JD, Mangan SA, Alexander HM. (2015). Maintenance of plant species diversity by

pathogens. *Annual review of ecology, evolution, and systematics* 46: 305–325.

Blackwell WH. (2011). The genus Lagena (Stramenopila: Oomycota), taxonomic history and

nomenclature. *Phytologia* 93: 157–167.

Bulman S, Braselton JP. (2014). 4 Rhizaria: Phytomyxea. eds. *Systematics and evolution*.

Springer, 99–112.

Castello JD, Leopold DJ, Smallidge PJ. (1995). Pathogens, patterns, and processes in forest

ecosystems. *Bioscience* 45: 16–24.

De Deyn GB, Raaijmakers CE, Zoomer HR, Berg MP, de Ruiter PC, Verhoef HA *et al.*

(2003). Soil invertebrate fauna enhances grassland succession and diversity. *NATURE*

422: 711–713.

Dickie IA, Bufford JL, Cobb RC, Desprez-Loustau M, Grelet G, Hulme PE *et al.* (2017). The

emerging science of linked plant–fungal invasions. *New Phytologist* 215: 1314–1332.

Geisen S, Tveit AT, Clark IM, Richter A, Svenning MM, Bonkowski M *et al.* (2015).

Metatranscriptomic census of active protists in soils. *The ISME journal* 9: 2178.

Gilbert GS. (2002). Evolutionary ecology of plant diseases in natural ecosystems. *Annual*

Review of Phytopathology 40: 13–43.

Gómez-Aparicio L, Ibánez B, Serrano MS, De Vita P, Avila JM, Pérez-Ramos IM *et al.*

(2012). Spatial patterns of soil pathogens in declining Mediterranean forests:

implications for tree species regeneration. *New Phytologist* 194: 1014–1024.

Haddad NM, Brudvig LA, Clobert J, Davies KF, Gonzalez A, Holt RD *et al.* (2015). Habitat

fragmentation and its lasting impact on Earth's ecosystems. *Science Advances* 1:

e1500052.

Holdaway RJ, Richardson SJ, Dickie IA, Peltzer DA, Coomes DA. (2011). Species- and

community-level patterns in fine root traits along a 120 000-year soil chronosequence in

temperate rain forest. *Journal of Ecology* 99: 954–963.

Hurst JM, Allen RB. (2007). The recce method for describing New Zealad vegetation: field

protocols.

Kardol P, Martijn Bezemer T, Van Der Putten WH. (2006). Temporal variation in plant–soil

feedback controls succession. *Ecology Letters* 9: 1080–1088.

Laliberté E, Lambers H, Burgess TI, Joseph, Wright S. (2015). Phosphorus limitation, soil-

borne pathogens and the coexistence of plant species in hyperdiverse forests and

shrublands. *New Phytologist* 206: 507–521.

Lambers H, Raven JA, Shaver GR, Smith SE. (2008). Plant nutrient-acquisition strategies

change with soil age. *Trends Ecol Evol* 23: 95–103.

Mangan SA, Herre EA, Bever JD. (2010a). Specificity between Neotropical tree seedlings

and their fungal mutualists leads to plant-soil feedback. *Ecology* 91: 2594–2603.

Mangan SA, Schnitzer SA, Herre EA, Mack KML, Valencia MC, Sanchez EI *et al.* (2010b).

Negative plant-soil feedback predicts tree-species relative abundance in a tropical

forest. *Nature* 466: 752–755.

Marano AV, Jesus AL, de Souza JI, Jerônimo GH, Gonçalves DR, Boro MC *et al.* (2016).

Ecological roles of saprotrophic Peronosporales (Oomycetes, Straminipila) in natural

environments. *Fungal Ecology* 19: 77–88.

Martin FN, Loper JE. (1999). Soilborne plant diseases caused by Pythium spp.: ecology,

epidemiology, and prospects for biological control. *Critical reviews in plant sciences*

18: 111–181.

Martínez-García LB, Richardson SJ, Tylianakis JM, Peltzer DA, Dickie IA. (2015). Host

identity is a dominant driver of mycorrhizal fungal community composition during

ecosystem development. *New Phytologist* 205: 1565–1576.

Packer A, Clay K. (2000). Soil pathogens and spatial patterns of seedling mortality in a

temperate tree. *Nature* 404: 278–281.

Packer A, Clay K. (2003). Soil pathogens and *Prunus serotina* seedling and sapling growth

near conspecific trees. *Ecology* 84: 108–119.

Peltzer DA, Wardle DA, Allison VJ, Baisden WT, Bardgett RD, Chadwick OA *et al.* (2010).

Understanding ecosystem retrogression. *Ecological Monographs* 80: 509–529.

Pickett STA. (1989). Space-for-time substitution as an alternative to long-term studies. *Long-*

term studies in ecology: approaches and alternatives 110–135.

Podger FD, Newhook FJ. (1971). Phytophthora cinnamomi in indigenous plant communities

in New Zealand. *New Zealand Journal of Botany* 9: 625–638.

Porder S, Vitousek PM, Chadwick OA, Chamberlain CP, Hilley GE. (2007). Uplift, Erosion,

and Phosphorus Limitation in Terrestrial Ecosystems. *Ecosystems* 10: 159–171.

Reinhart KO, Johnson D, Clay K. (2012). Conspecific plant-soil feedbacks of temperate tree

species in the southern Appalachians, USA. *PloS one* 7: e40680.

Reynolds HL, Packer A, Bever JD, Clay K. (2003). Grassroots ecology: Plant-microbe-soil

interactions as drivers of plant community structure and dynamics. *ECOLOGY* 84:

2281–2291.

Richardson SJ, Peltzer DA, Allen RB, McGlone MS. (2005). Resorption proficiency along a

chronosequence: responses among communities and within species. *Ecology* 86: 20–25.

Richardson SJ, Peltzer DA, Allen RB, McGlone MS, Parfitt RL. (2004). Rapid development

of phosphorus limitation in temperate rainforest along the Franz Josef soil

chronosequence. *Oecologia* 139: 267–276.

Scott P, Williams N. (2014). Phytophthora diseases in New Zealand forests. *NZ Journal of*

Forestry 59: 15.

Skipp RA, Christensen MJ. (1989). Fungi invading roots of perennial ryegrass (Lolium

perenne L.) in pasture. *New Zealand Journal of Agricultural Research* 32: 423–431.

Van der Putten WH, Dijk CV, Peters BAM. (1993). Plant-specific soil-borne diseases

contribute to succession in foredune vegetation. *Nature* 362: 53–56.

Van Der Putten WH. (2003). Plant defense belowground and spatiotemporal processes in

natural vegetation. *ECOLOGY* 84: 2269–2280.

Veblen TT. (1992). Regeneration dynamics. *Plant succession: Theory and prediction* 11:

152–187.

Vitousek PM, Porder S, Houlton BZ, Chadwick OA. (2010). Terrestrial phosphorus

limitation: mechanisms, implications, and nitrogen-phosphorus interactions. *Ecological*

Applications 20: 5–15.

Vitousek PM, Mooney HA, Lubchenco J, Melillo JM. (1997). Human domination of Earth's

ecosystems. *Science* 277: 494–499.

Walker TW, Syers JK. (1976). The fate of phosphorus during pedogenesis. *Geoderma* 15: 1–

19.

Weir BS, Paderes EP, Anand N, Uchida JY, Pennycook SR, Bellgard SE *et al.* (2015). A

taxonomic revision of Phytophthora Clade 5 including two new species, Phytophthora

agathidicida and P. cocois. *Phytotaxa* 205: 21–38.

564