Functional metagenomics using *Pseudomonas putida* expands the known diversity of

polyhydroxyalkanoate synthases and enables the production of novel polyhydroxyalkanoate

copolymers

Jiujun Cheng and Trevor C. Charles*

Department of Biology and Centre for Bioengineering and Biotechnology, University of Waterloo,

Waterloo, Ontario, Canada N2L 3G1

*Corresponding author:

Dr. Trevor C. Charles

Department of Biology, University of Waterloo, Waterloo, Ontario, Canada N2L 3G1.

E-mail: tcharles@uwaterloo.ca

Abstract

Bacterially produced biodegradable polyhydroxyalkanoates with versatile properties can be achieved

using different PHA synthase enzymes. This work aims to expand the diversity of known PHA

synthases via functional metagenomics, and demonstrates the use of these novel enzymes in PHA

production. Complementation of a PHA synthesis deficient *Pseudomonas putida* strain with a soil

metagenomic cosmid library retrieved 27 clones expressing either Class I, Class II or unclassified

PHA synthases, and many did not have close sequence matches to known PHA synthases. The

composition of PHA produced by these clones was dependent on both the supplied growth substrates

and the nature of the PHA synthase, with various combinations of SCL- and MCL-PHA. These data

demonstrate the ability to isolate diverse genes for PHA synthesis by functional metagenomics, and

their use for the production of a variety of PHA polymer and copolymer mixtures.

Keywords:

Functional metagenomics; soil metagenomic library; polyhydroxyalkanoate (PHA); bioplastics;

polyhydroxyalkanoate synthase; *Pseudomonas putida*; *Sinorhizobium meliloti*

Introduction

Polyhydroxyalkanoates (PHAs) are natural polyesters biosynthesized by a variety of bacteria

under unbalanced growth conditions. They serve as reserves of carbon and reducing power and aid

in survival during starvation or stress conditions (Verlinden et al. 2007). These biodegradable and

environmentally friendly polymers can be used as alternative materials to conventional

petrochemical-based plastics. PHAs are classified into short-chain-length (SCL, C3-C5) and

medium-chain-length (MCL, $\geq$C6), and copolymers (SCL + MCL) based on the number of carbon

atoms per monomer. MCL-PHA is generally more useful than SCL-PHA due to it being less brittle

and more flexible. To produce PHA with versatile properties cost-effectively, strategies have

involved mining new PHA synthase enzymes (PhaC), engineering PhaC proteins and modifying the

metabolic pathways of the production strains (Keshavarz and Roy 2010; Schallmey et al. 2011;

Cheema et al. 2012; Park et al. 2012; Tripathi et al. 2013; Meng et al. 2014).

PHA molecules consist of over 150 possible constituent monomers of hydroxyl fatty acids

(Meng et al. 2014) (Steinbüchel and Lütke-Eversloh 2003). PHA synthase enzymes catalyze the

joining of the hydroxyl group of one monomer with the carboxyl group of another by an ester bond

to form PHA polymers. The monomeric composition of PHA is determined primarily by PhaC,

although the available carbon source and metabolic pathways also influence the properties of PHA

(Verlinden et al. 2007). PhaC proteins are grouped into four classes based on amino acid sequence,

substrate specificity and subunit composition (Rehm 2003). Class I and II PhaC consist of one

subunit (PhaC), but those in Class III and IV are composed of two subunits (PhaC+PhaE and

PhaC+PhaR, respectively). Class I and IV PhaCs synthesize SCL-PHA whereas Class II polymerize

MCL-PHA. Class III PhaCs can synthesize both SCL and MCL monomers.

The key Class I PHA synthesis pathway genes include *phaC* (or *phbC*), *phaA* (or *phbA*) and

phaB (or *phbB*). The three genes are often, but not always, clustered in a single operon. The

acetoacetyl-CoA reductase (EC 2.3.1.9) encoded by the *phaA* and β-ketothiolase (EC 1.1.1.36)

encoded by the *phaB* convert acetyl-CoA to (R)-3-hydroxybutyryl-CoA (3HB-CoA) through

acetoacetyl-CoA. The Class-I PhaC (EC 2.3.1.-) then polymerizes 3HB-CoA into

polyhydroxybutyrate (PHB) (Rehm 2003). The Class II *pha* cluster is well conserved in

Pseudomonas and consists of two PHA synthase genes (*phaC1* and *phaC2*) flanking a PHA

depolymerase gene (*phaZ*), and *phaD* encoding a transcriptional activator of *pha* genes. Phasin-

encoding *phaF* and *phaI* gene are transcribed divergently to other *pha* genes. MCL monomers ((R)-

3-hydroxylacyl-CoA) are derived from β-oxidation of fatty acids or fatty acid *de novo* synthesis from

unrelated carbon sources (Tortajada et al. 2013).

Traditional culture-based strategies for obtaining new biocatalysts are limited by the inability

to cultivate the majority of environmental microbes. While sequence-based metagenomics can

identify genes homologous to those present in available sequence databases, it is difficult to reliably

predict the function of truly new genes through homology-based analysis. In contrast, functional

metagenomics involves the construction of gene libraries from microbial community genomic DNA

and functional screening for novel enzymes of interest or potential industrial applications (Simon

and Daniel 2011). Functional metagenomics has the potential to identify truly novel sequences for a

given function.

In previous work, functional metagenomics was used to isolate new Class I PhaC from soil

metagenomic clones by Nile red staining and phenotypic screening in *α-Proteobacteria*

Sinorhizobium meliloti (Schallmey et al. 2011). In another study, *phaC* genes encoding both Class I

and II PhaC proteins were PCR amplified from oil-contaminated soil library clones (Cheema et al.

2012). One of the isolated genes produced PHA copolymer when expressed in *Pseudomonas putida*.

Moreover, partial *phaC* genes were also obtained from metagenomic DNA via direct PCR

amplification (Foong et al. 2014; Pärnänen et al. 2015; Tai et al. 2015). Building on these previous

studies, we constructed a PHA$^-$ strain of *P. putida* KT2440 to use as a surrogate host and

functionally identified *phaC* genes after screening millions of agricultural wheat soil metagenomic

clones. A total of 27 PHA$^+$ clones were obtained. Accumulation and monomer composition of PHA

directed by seven of these clones were further examined.

Methods

Bacterial strains, plasmids, cosmids and growth conditions

Bacterial strains, plasmids and cosmids are listed in Table 1. *E. coli* strains were grown at

37°C in LB (Lennox) medium (1% tryptone (w/v), 0.5% yeast extract (w/v), and 0.5% NaCl (w/v),

pH 7). *Pseudomonas* strains were grown at 30°C in LB or 0.1N M63 minimal medium (Escapa et al.

2011) supplemented with 0.5% sodium octanoate (w/v), 0.5% nonanoic acid (v/v) or 1% gluconic

acid (w/v). *S. meliloti* was grown in LB or YM medium (Schallmey et al. 2011). Antibiotics were

used at the following concentrations: streptomycin, 200 µg/ml for *S. meliloti* and 100 µg/ml for *E.*

coli; kanamycin, 100 µg/ml for *Pseudomonas* and 50 µg/ml for *E. coli*; neomycin, 200 µg/ml;

rifampicin, 100 µg/ml; gentamicin, 10 µg/ml for *E. coli* and 100 for *P. putida*; and tetracycline 20

µg/ml for *E. coli* or 40 µg/ml for *P. putida*.

Construction of PHA$^-$ strain PpUW2

DNA oligonucleotides are listed in Table 2. A Rif^R spontaneous mutant PpUW1 of *P. putida*

KT2440 was generated by plating a culture of strain KT2440 on a LB Rif plate, followed by single

colony purification. To construct the PHA deficient strain PpUW2 (*phaC1ZC2*) of *P. putida* PpUW1,

a 943-bp DNA fragment containing 5′-*phaC1* region (766 bp) was PCR amplified using *P. putida*

KT2440 genomic DNA as a template and primer pair JC161-JC162, digested with HindIII-BamHI

and then cloned into the same sites in pJQ200-SK (Quandt and Hynes 1993), yielding plasmid

pJC63. Another 876-bp DNA fragment containing the 3′-*phaC2* gene (120 bp) was PCR amplified

using primers JC163-JC164, digested with BamHI-SalI and then cloned into the same sites in pJC63

to obtain plasmid pJC69. An omega-Km cassette was obtained from pHP45Ω-Km (Fellay et al. 1987)

by BamHI digestion and then inserted into the same site in pJC69 to obtain pJC71. Plasmid pJC71

was then conjugated into *P. putida* PpUW1 in a triparental mating using helper plasmid pRK600.

Single cross-over recombination of pJC71 into the *P. putida* chromosome was selected with Rif and

Gm. A double cross-over event was achieved by growing a single Rif^R Gm^R colony overnight in LB,

making serial dilutions and then spreading on LB Km with 5% sucrose. The resulting PHA$^-$ strain

PpUW2 was verified by examining Gm sensitivity (lost plasmid backbone) and absence of PHA

production in LB supplemented with 0.5% octanoate (w/v), and was further confirmed by PCR

amplification analysis.

Phenotypic screening for *phaC* genes from metagenomic library clones

117   Construction of the metagenomic DNA library (11AW) of agricultural wheat soil was

described previously (Cheng et al. 2014). The 11AW library contains 9×10^6 clones hosted in *E. coli*

HB101. The pooled library clones (0.5 ml of 300 ml stock) were conjugated *en masse* into *P. putida*

PpUW2 (PHA$^-$) with the helper plasmid pRK600. Mating mixture was diluted serially with 0.85%

NaCl and ~20,000 transconjugants recovered on each LB Tc plate (15 mm × 150 mm) supplemented

with 0.5% Na octanoate (w/v). The plates were incubated at 30$^°$C for 24 h and then kept at 22$^°$C for

2-6 days. Potential PHA$^+$ clones of opaque white colour were streak purified on the LB octanoate

plates, and verified on 0.1N M63 minimal medium plates (Escapa et al. 2011) supplemented with

0.5% octanoate (w/v) and Nile red (0.5 μg/ml). PHA$^+$ cosmids were transferred from *P. putida* to *E.*

coli DH5α(Rif^R), mobilized by pRK600. The cosmid DNA was isolated from *E. coli* strains using a

GeneJET Plasmid Miniprep Kit (Thermo Scientific), digested by EcoRI-HindIII-BamHI (Thermo

Scientific), and then resolved on 1% TAE agarose gels. Cosmids with distinct electrophoretic

patterns were conjugated back to *P. putida* PpUW2 (PHA$^-$) to confirm their PHA$^+$ phenotype.

Complementation of *S. meliloti* (*phbC*)

11AW cosmid DNA encoding Class I and II *phaC* genes was introduced into *S. meliloti*

Rm11476 (Schallmey et al. 2011) via triparental conjugation. Transconjugants were selected on LB

SmNmTc plates, and then streaked on YM plates containing Nile red (Schallmey et al. 2011)for

visualizing PHB production.

DNA sequencing and bioinformatics

KOD Xtreme DNA polymerase (Novagen) was used for all PCR. Primers are listed in Table

2. PCR reactions consisted of one cycle of 94°C for 5 min, 30 cycles of 94°C for 30 s, 53°C for 30 s

and 68°C for 30 s, and final extension at 68°C for 10 min. The internal regions of *phaC* genes were

PCR amplified from PHA$^+$ cosmids using primer phaCF1 and phaCR4 (Sheu et al. 2000). When no

PCR product of correct size was obtained, a semi-nested PCR was performed with the primers

phaCF2 and phaCR4 as described previously (Sheu et al. 2000). PCR products were resolved on 2%

agarose gels, isolated from the gels using EZ-10 Spin Column DNA Gel Extraction Kit (Bio Basic),

and then cloned into pJET1.2 vector (Thermo Scientific). Sequences of the cloned partial *phaC*

genes were obtained using pJET1.2 sequencing primers (Table 2).

End sequences of PHA[+] cosmid DNA were obtained by Sanger sequencing using universal

M13F and M13R primers. For high throughput sequencing, 68 cosmids of 11AW PHA[+] clones and

28 cosmids from other research projects were grouped into 24 pools. Tagmentation of pooled cosmid

DNA, PCR amplification and clean-up, and library normalization were performed with the Nextera

XT DNA library and index kits (Illumina), according to the supplier's recommendation. The library

was sequenced using MiSeq 500-cycle version 2 reagent (Illumina). DNA sequences were

assembled using SPAdes Genome Assembler 3.5.0 (BaseSpace, Illumina). Cosmid clones were

identified based on the available Sanger end sequences (Lam et al. 2014) and ORFs were annotated

with MetaGeneMark (Zhu et al. 2010). Predicted protein sequences were analyzed by BLAST

against the non-redundant protein databases (NCBI). Multiple-sequence alignment was performed

using MUSCLE (Edgar 2004). Phylogenetic analysis was performed using MEGA6 (Tamura et al.

2013). Origin of cloned metagenomic DNA was predicted by PhyloPythia (Patil et al. 2012).

Cloning of *phaC1$_{Pp}$* gene of *P. putida* KT2440

To construct a *phaC* expression cosmid for a positive control, the *phaC1$_{Pp}$* gene (*Pp_5003*)

of *P. putida* KT2440 was PCR amplified using primers JC159-JC160 and cloned into the HindIII

and BamHI sites in the broad-host-range vector pRK7813 (Cheng et al. 2014), to obtain construct

pJC67 (P*lac*::*phaC1$_{Pp}$*). A stop codon (TGA) for terminating the translation of *lacZα* upstream of the

phaC1$_{Pp}$, and a ribosome-binding site (AGGAG) were incorporated into the primer JC159 (Table 2).

Subcloning *phaC* genes of 11AW metagenomic clone 16

The Class II *phaC1$_{16}$* gene of 11AW clone 16 was PCR amplified with oligos JC301 and

JC302, and cloned into the HindIII-EcoRI sites in pRK7813 to obtain pJC123. The *phaC2$_{16}$* was

amplified using primers JC337 and JC334, and inserted into the EcoRI site in pJC123, yielding

pJC157. The orientation of cloned *phaC2$_{16}$* was verified by restriction enzyme mapping. The

phaC2$_{16}$ gene was also obtained by PCR using oligos JC333 and JC334, and cloned into the HindIII-

EcoRI sites in pRK7813 to yield pJC159. A stop codon (TGA) for terminating the translation of

lacZα gene and ribosome-binding sites were added in primers JC333 and JC337. The cloned *phaC*

genes were verified by DNA sequencing.

Estimation of PHA production by fluorescent spectrometry

A modification of the previously described method (Wu et al. 2003) was used. *S. meliloti*

cells (2.5%, v/v) were added to YM medium. All cultures were incubated at 30^{o}C and 200 rpm for

48-72 h. The OD$_{600}$ values of 200 μl cultures were measured in 96-well microtiter plates with the

Multiskan Spectrum spectrophotometer (Thermo Labsystems). Samples (180 μl) were stained with

20 μl of Nile red (2.5 μg/ml) in the dark for 1 h. Fluorescent intensity was measured at excitation

(485 nm) and emission (595 nm) with the FilterMax F5 Multi-mode microplate reader (Molecular

Devices). PHA content was calculated based on the equation (fluorescent intensity/OD$_{600}$×0.9).

PHA production, extraction and characterization

P. putida PpUW1 (PHA^{+}) or PpUW2 (PHA^{-}) carrying PHA^{+} clones were grown overnight in

LB with or without Tc, washed once with 0.85% NaCl, and then subcultured (2%, v/v) in the 0.1N

M63 medium with or without Tc, supplemented with 0.5% Na octanoate, 0.5% nonanoic acid or 1%

gluconic acid. The cultures were grown at 30$^{°}$C and at 200 rpm for 48 hrs. Cells were collected by

centrifugation at 20°C and at $9000 \times g$ for 15 min, washed once with deionized water. Cell dry

weight (CDW) was obtained after drying the cells at 95°C for 24 hr. For PHA methanolysis, cell

pellet (~15 mg) was suspended in 2 ml of chloroform and 2 ml of methanol with 15% H_2SO_4 (v/v),

and incubated at 100°C for 5 h. After the reaction mixture was cooled down to 20°C, 1 ml of

deionized water was added, and vortexed vigorously for 1 min. The chloroform phase was passed

through a cotton plug in a Pasteur glass pipette to remove any cell debris. Methanolyzed sample (1

µl) was analyzed with GC-MS (an Agilent 7975B GC equipped with Agilent 5975B inert XL EI/CI

MSD and an HP-5MS capillary column). Oven temperature was run at initial 50°C for 5 min with a

ramp of 20°C/min to 280°C, and then held for 10 min. The flow rate of Helium carrier gas was 1.2

ml/min. Methylated PHA monomers were identified using the Agilent enhanced MSD chemstation

(E.02.01.117). PHA standards were kindly provided by Dr. Bruce A. Ramsay (Polyferm Canada).

Results

Isolation of metagenomic clones for PHA production

To employ *P. putida* as a surrogate host to isolate 11AW (agricultural wheat soil)

metagenomic library clones encoding functional PhaC, it was first necessary to construct a PHA

synthesis mutant. Expression of the contiguous three genes encoding PHA synthase PhaC1$_{Pp}$

(Pp_5003), PHA depolymerase PhaZ$_{Pp}$ (Pp_5004) and PHA synthase PhaC2$_{Pp}$ (Pp_5005) in *P.*

putida KT2440 Rif^{R} derivative PpUW1 was disrupted by deletion of 3437-bp comprised of the 3´

region (911 bp) of *phaC1$_{Pp}$* (1677 bp), *phaZ$_{Pp}$* and the 5´ region (1560 bp) of *phaC2$_{Pp}$* (1680 bp) and

replacement with an omega-Km kanamycin resistance insert, resulting in the PHA^{-} strain PpUW2.

Transfer of the negative control cosmid vector pJC8 to *P. putida* PpUW2 did not result in detectable

PHA production. However, introduction of pJC67 (P*lac*::*phaC1$_{Pp}$*) *in trans* restored the PHA^{+}

phenotype (Fig. 1). These data suggested that the strain PpUW2 could be used for screening of

PhaC-encoding metagenomic clones. The 11AW library clones (Cheng et al. 2014) were transferred

to *P. putida* PpUW2 via en masse triparental conjugation. Following selection of *P. putida* PpUW2

transconjugants (~4 million) on LB Km Tc plates supplemented with 0.5% Na octanoate, we

obtained 72 clones that exhibited greater opacity than the PpUW2 PHA$^-$ recipient strain. The PHA-

producing phenotype of those clones was verified by visualizing the fluorescence of Nile red-stained

PHA in 0.1N M63 minimal medium (Escapa et al. 2011) supplemented with Na octanoate as the sole

carbon source (Fig. 1). Restriction digest of DNA from the 72 clones demonstrated 68 distinct

restriction patterns, suggesting the presence of a broad diversity of DNA origin in those PHA$^+$ clones.

Identification of *phaC* genes encoding PHA synthases

Internal regions (~500 bp) of the *phaC* genes of 18 distinct cosmid clones were initially

obtained by PCR amplification using degenerate primers PhaCF1 and PhaCR4 (Sheu et al. 2000).

For additional 4 clones, *phaC* fragments (PhaC$_3$, PhaC$_7$, PhaC$_{10}$, and PhaC$_{15}$) of ~400 bp were

generated by nested-PCR using primer pair PhaCF2-PhaCR4 as described previously (Sheu et al.

2000). BLASTP analysis of the cloned gene products indicated that 13 PhaC proteins could be

grouped into Class I while the other 9 were categorized as Class II PHA synthases.

To identify all *pha* genes on the isolated clones, DNA sequences were obtained by high

throughput sequencing, and additional *pha* gene loci were identified in partially and fully assembled

clones (Table 3; Supplementary 1). The identified PHA synthases could be classified into 3 groups

based on amino acid sequences (Fig. 2). Class I *phaC* genes were annotated in 17 clones. The *phaC*

and *phaAB* genes were adjacently located in 12 of these clones, whereas the *phaB* gene was located

distantly downstream of *phaCA* genes in clone 25 and clone P1N3 (Supplementary 1). A *phaR* gene

was located immediately downstream of *phaCB* genes in partially assembled clone P11N2. The

phaC and *phaB* genes flanked a *phaZ* gene in the partially assembled clone P2N8. The metagenomic

DNA in these Class I clones was predicted to originate from *Gemmatimonas*, *α-Proteobacteria*

(*Sphingomonadaceae*)*, β-Proteobacteria* (*Leptothrix, Rubrivivax, Janthinobacterium*) and γ-

Proteobacteria (*Xanthomonadaceae*) (Table 3).

Class II PHA genes are commonly clustered with *phaZ* flanked by two PHA synthase

encoding genes, *phaC1* and *phaC2*. Nine clones carried Class II *phaC* genes (Table 3). The *pha* gene

locus (*phaC1$_1$-phaZ-phaC2$_1$-phaD-phaF-phaI*) in clone 1 was similar to the canonical locus in

Pseudomonas including *P. putida* KT2440 (de Eugenio et al. 2010), except the presence of *orf7*

encoding a PHA granule associated protein (Pfam09650) (Fig. 3A). Clone 2 and clone 15 were

identical to the clone 1, except that they contained an insertion of the 17,013-bp transposon Tn4652

at 13,740 nt and 16,362 nt respectively (Supplementary 2). The Tn4652 duplicated target sequences

were AACTC in clone 2 and TAGGA in clone 15. The transposon insertions in these clones are in

opposite orientation. The same Tn4652 is present in *P. putida* KT2440 genome (3,366,550 -

3,383,562 nt, GenBank: AE015451), located distant from the *pha* gene operon. These insertions

likely occurred following introduction of the cosmid clones into *P. putida*.

Class II clones 5, 16, 20 and 50 had no genes encoding *phaD* or *phaFI* homologs. A *phaJ*

gene encoding R-specific enoyl-CoA hydratase was identified in those clones (Fig. 3A;

Supplementary 1). The *phaJ* gene was located immediately upstream of the *phaC* genes in clones 16,

20 and 50, but downstream of *phaC* in clone 5. In addition, a gene encoding PHA depolymerase

(PhaZ) was located downstream of the *phaC* genes in clones 16, 20 and 50. ORFs downstream of the

phaZ genes in clones 16 and 50 encoded proteins (PhaC2$_{16}$ and PhaC2$_{50}$) homologous to unclassified

PHA synthases (Fig. 2; Supplementary 1). The proteins shared conserved regions in PhaC enzymes,

but were ~30% shorter in N-terminal regions than in those of Class I/II PhaC (Supplementary 2).

Two ORFs encoding putative PhaC and PhaE proteins were identified in partially assembled

clone P2N10 (Table 3; Supplementary 1). BLASTP showed the P2N10 PhaC protein best matched

to the Class III PhaC [GenBank: WP_002708071] (43% identity) from γ-*Proteobacteria Thiothrix*

nivea (Table 3). The available C-terminal sequence of P2N10 PhaE (234 amino acids) only exhibited

26% identity to the Class III PhaE [GenBank: WP_002708072] of *T. nivea*. In addition, the P2N10

PhaC protein was only 29% identical to the Class IV PhaC [GenBank: AAD05620] of *Bacillus*

megaterium. Phylogenetic analysis suggested that P2N10 PHA synthase was distant from Class

III/IV PhaC and might represent a new subclass of PHA synthases (Fig. 2).

Multiple alignment of amino acid sequences of each of the identified 11AW PhaC proteins

showed the conserved catalytic triad (C287, D445 and H473 in $PhaC1_{16}$), a tryptophan essential for

dimerization (W391 in $PhaC1_{16}$), and the lipase box GXCXGG (Jia et al. 2000) except that Ala

replaced the first Gly residue in $PhaC_{20}$ and second Gly in clone P8N7 (Fig. 3B; Supplementary 3).

The serine residue (S229 in $PhaC1_{16}$) required for PhaC activity (Hoppensack et al. 1999) was also

conserved in all 11AW PhaC proteins. We chose the PhaC proteins in the Class I clones 14, 18 and

25, and the Class II clones 1, 16 and 20, for further study. We examined the nature of PHA produced

using different carbon sources that result in substrate production either through fatty acid synthesis

or β-oxidation.

Class II clone 1 synthesizes MCL PHA

The cloned metagenomic DNA in clone 1 contained 42,747 bp (Table 3) [GenBank:

KT944254] and most likely originated from γ-*Proteobacteria Pseudomonas*. The $PhaC1_1$, PhaZ,

PhaC2$_1$ and PhaD proteins [GenBank: ALV86243, ALV86281, ALV86274 and ALV86275] were

82%, 92%, 73% and 78% identical to the corresponding orthologs (PhaC1, PhaZ, PhaC2 and PhaD)

of *P. putida* KT2440. Clone 1 PhaC2$_1$ was phylogenetically related to the PHA synthase 2

[GenBank: BAA36202] of *Pseudomonas* sp. 61-3 (Fig. 2). In contrast to other Class II PhaC, the

conserved amino acid at position 129 of PhaC2$_1$ was Asp rather than Glu (Fig. 3B). It has been

previously demonstrated that substitution of the conserved Glu with Asp in Class II PhaC improves

PHA yield with an increase of 3-hydroxybutyrate monomer (Matsumoto et al. 2005).

PHA synthesis in clone 1 was likely contributed solely by PhaC2$_1$ because the 5'-region of

phaC1$_1$ was absent in the cloned DNA. Expression of the functional *phaZ* and *phaC2$_1$* might be

driven by the promoters upstream of the individual genes, as occurs in *P. putida* KT2440 (de

Eugenio et al. 2010). *P. putida* PpUW2 (PHA$^-$) carrying clone 1 synthesized 3-hydroxyhexanote

(3HHx, C6) and 3-hydroxyoctanoate (3HO, C8) copolymer of ~95% C8 monomer when grown with

octanoate, similar to the PHA produced by wild-type *P. putida* PpUW1 (Table 4). When PpUW2

(clone 1) was grown with nonanoic acid, 3-hydroxynonanoate (3HN, C9) and 3-hydroxyheptanoate

(3HP, C7) were incorporated into the PHA with greater 3HP than that synthesized by wild type

PpUW1 (Table 4). These data suggest that those monomers were derived from β-oxidation of

octanoate and nonanoic acid.

When gluconic acid was used as the sole carbon source, 3HO and 3-hydroxyldecanoate

(3HD, C10) copolymer was produced in PpUW2 (PHA$^-$) carrying clone 1, very similar to the

production of the parental strain PpUW1 (PHA$^+$) (Table 4). The monomer composition of PHA was

similar in both strains, but total PHA accumulation was ~4 fold higher in gluconic acid-grown

PpUW2 with clone 1 (Table 4). These results suggest that Class II PhaC1$_2$ was able to synthesize

MCL (C6-C10) PHA.

Class II clone 16 synthesizes SCL-MCL copolymer

Clone 16 contains 33,900-bp metagenomic DNA [GenBank: KT944263] probably originated

from *β-Proteobacteria Methylibium* (Table 3). A *phaJ* gene (*phaJ$_{16}$*) is predicted to encode a R-

specific enoyl-CoA hydratase [GenBank: ALV86416] (Fig. 3A), which converts the β-oxidation

intermediate 2-enoyl-CoA to (R)-3-hydroxyacyl-CoA used as PhaC substrate (Fukui et al. 1998).

The PhaJ$_{16}$ was phylogenetically related to the PhaJ4a$_{Re}$ and PhaJ4b$_{Re}$ of *Ralstonia eutropha*

(Kawashima et al. 2012), and PhaJ4$_{Pp}$ of *P. putida* (Sato et al. 2011) (Supplementary 4A). The

amino acid residues Asp39 and His44 conserved at the active site of the dehydratases were identified

in PhaJ$_{16}$ (Supplementary 3B), except that the Ser residue was replaced by Pro71 in PhaJ$_{16}$, same as

in the PhaJ orthologs of *R. eutropha* and *P. putida* PhaJ4.

The three ORFs downstream of the *phaJ$_{16}$* gene encoded PhaC1$_{16,}$ PhaZ and PhaC2$_{16}$ (Fig.

3A). PhaC1$_{16}$ of Class II PHA synthase most closely matched the *Pseudomonas* sp. F15 PhaC1

[GenBank: WP_021487788], exhibiting 48% amino acid identity. PhaC1$_{16}$ was also 42.5% and

43.4% identical to the PhaC1 and PhaC2 proteins of *P. putida* KT2440. Thr318 and Val478 in the

PhaC1$_{16}$ might favour the 3HB incorporation into PHA as occurred in the engineered PhaC proteins

(Takase et al. 2003; Chen et al. 2014). Another putative PHA synthase encoded by the *phaC2$_{16}$* gene

downstream of the *phaC1$_{16}$ phaZ* genes was homologous to unclassified PHA synthases (Fig. 2).

The quantity of PHA accumulated in *P. putida* UW2 carrying clone 16 was comparable to that

of parental PpUW1 when grown with octanoate, nonanoic acid or gluconic acid (Table 4). In

contrast to the MCL PHA (C6-C8) present in octanoate-grown PpUW2 (clone 1) or wild-type

PpUW1, about 20% 3HB was detected in the PHA (C4-C6-C8) accumulated in PpUW2 carrying

clone 16 under the same conditions (Table 4). When PpUW2 (clone 16) was grown with nonanoic

acid (C9), 3-hydroxyheptanoate (3HP, C7), 3-hydroxyvalerate (3HV, C5) and 3HB (C4) were

polymerized, which was in contrast to the 3HP-3HN PHA in nonanoic acid-grown PpUW with

PhaC1 or wild type PpUW1. Additionally, equal amounts of C6 and C8 monomers were detected in

the PHA from gluconic acid-grown PpUW2 with clone 16. However, PHA with C8 and C10

monomers was accumulated in PpUW2 (clone 1) and PpUW1 under the same growth conditions.

These data implied that the $PhaC1_{16}$ and/or $PhaC2_{16}$ prefer substrates with shorter carbon chains than

the $PhaC2_1$ in clone 1 and both PhaC proteins of *P. putida* KT2440.

335   To further elucidate the activity of $PhaC1_{16}$ and $PhaC2_{16}$, the genes were cloned

downstream of the constitutively active P*lac* promoter. When *P. putida* PpUW2 carrying pJC123

(P*lac*::*phaC1$_{16}$*) was grown with octanoate, SCL-MCL copolymer was of similar composition of C4-

C6-C8 monomers to that synthesized by clone 16 (Table 4). However, PHA was not detected in

strain PpUW2 carrying pJC159 (P*lac*::*phaC2$_{16}$*) under the same growth conditions. Coexpression of

phaC1$_{16}$ and *phaC2$_{16}$* in pJC157 (*Plac*::*phaC1$_{16}$-phaC2$_{16}$*) in octanoate-grown PpUW2 resulted in

synthesis of PHA with the similar monomer composition as those in the polymers produced in

PpUW2 carrying clone 16 or pJC123 (P*lac*::*phaC1$_{16}$*) (Table 4). These data indicated that only the

$PhaC1_{16}$ was involved in SCL-MCL PHA biosynthesis in clone 16.

344   The *pha* gene locus of clone 16 was very similar to that of clone 50 (Supplementary 1), but

the DNA in clone 50 probably originated from γ-*Proteobacteria Hydrocarboniphata* (Table 3). Both

the Class II PhaC1 and unclassified PhaC2 in clones 16 and 50 shared 84% identity respectively.

Expression of the *pha* genes was probably driven by a promoter upstream of *phaJ* genes.

Class II clone 20 synthesizes MCL or SCL-MCL PHA

PHA$^+$ clone 20 contained 30,964-bp metagenomic DNA [GenBank: KT944266], which

probably originated from *Salinibacter* within the *Bacteroidetes* (Table 3). The cloned DNA

contained only a single *phaC$_{20}$* gene (Fig. 3A). A putative fatty acid CoA ligase/ synthetase

[GenBank: ALV86493] (ORF9; Supplementary 1) might be involved in degradation of fatty acids to

acyl-CoA (the first step of β-oxidation cycle). The acyl dehydratase PhaJ$_{20}$ [GenBank: ALV86492]

was 56% identical to the PhaJ$_{16}$. Conserved amino acid residues Asp41 and His46 at the active sites

of the acyl dehydratases were present (Supplementary 4B) except that the Ser residue was replaced

by Pro73 in the PhaJ$_{20}$ protein. The *phaJ$_{20}$*, *phaC$_{20}$* and *phaZ* genes in clone 20 were similar to those

of the corresponding genes in clones 16 and 50 (Fig. 3A; Supplementary 1). In addition, the *pha*

gene cluster of clone 20 was similar to that in *Burkholderia* sp. 383 (1,576,365 -1,581,635 nt)

[GenBank: NC_007511], but a gene encoding a phasin protein is located between the *phaC* and

phaZ genes in the *Burkholderia* genome.

The single PhaC$_{20}$ [GenBank: ALV86493] best matched to its ortholog from *Burkholderia lata*

at 59% amino acid sequence identity [GenBank: WP_011354994]. PhaC$_{20}$ was 46-53% identical to

the PhaC1$_{16}$, PhaC2 and PhaC5 as well as PhaC1 and PhaC2 of *P. putida* KT2440. Presence of amino

acid residues Thr353 and Val516 in PhaC$_{20}$ (Fig. 3B) might favour substrates with fewer carbons

than those preferred by clone 1 and *P. putida* PhaC proteins, as previously demonstrated with

engineered Class II PHA synthases (Takase et al. 2003; Chen et al. 2014).

Clone 20 was able to synthesize 3HHx-3HO copolymer (C6-C8) with octanoate (Table 4), as

occurred in PpUW2 (clone 1) and *P. putida* wild type. However, the proportion of C6 monomer

increased 4-35 fold in the PHA isolated from PpUW2 (clone 20), compared to that in PpUW2 (clone

1) and wild-type PpUW1 (Table 4). When nonanoic acid was supplied, PpUW2 (clone 20) produced

PHA with 8% 3-hydroxyvalerate (3HV), which was absent in the PHA synthesized by clone 1 and

PpUW1 (Table 4). In contrast to the SCL-MCL PHA produced by clone 16, 3HB was not detected in

the nonanoic acid or octanoate-grown PpUW2 with clone 20.

In contrast to the absence of detectable C6 monomer in the PHA produced in PpUW2 with

clone 1 or *P. putida* wild type, 3HHx (~10%) was incorporated into the MCL PHA (C6-C8-C10) in

PpUW2 (clone 20) (Table 4) when gluconic acid was used as the sole carbon source. About half the

quantity of monomers was C10 in the PHA accumulated in PpUW2 (clone 20) grown with gluconic

acid. These results suggested that the carbon chain lengths of favourite monomers of $PhaC_{20}$ were

between those of $PhaC1_{16}$ and $PhaC2_{1}$.

Class I clone 14 synthesizes SCL-MCL PHA

The DNA in clone 14 probably originated from *β-Proteobacteria Methylibium* (Table 3). The

$PhaC_{14}$, PhaA and PhaB [GenBank: ALV86397, ALV86398 and ALV86399] were 65%, 76% and

74% identical to the PhbC1 (A16_1437), PhbA (A16_1438) and PhbB1 (A16_1439) of *C. necator*

H16, respectively. In addition, $PhaC_{14}$ was 48% identical to the PhbC protein (Smc002960) of *S.*

meliloti Rm1021. The best match of $PhaC_{14}$ was the PhaC protein of *Burkholderia* JOSHI_001 at

74% (Table 3). The $phaC_{14}AB$ locus of clone 14 is similar to that of clones 8, 21 and 23

(Supplementary 1). Expression of the $phaC_{14}$, *phaA* and *phaB* genes in clone 14 was most likely

driven by a promoter upstream of the *orf9* encoding a hypothetical protein.

When PpUW2 (clone 14) was grown with nonanoic acid, 7% 3HB was incorporated into

3HB-3HV copolymer (Table 4). 3HB was the primary monomer of PHA in octanoate- or gluconic

acid-grown PpUW2. Both C6 and C8 monomer were present in the PHA with gluconic acid as

carbon source. However, C8 monomer was absent in the PHA when octanoate was supplied (Table

4). *P. putida* PpUW2 carrying clone 14 produced the highest quantity of PHA among all the PHA[+]

clones (Table 4). These results indicated that $PhaC_{14}$ could synthesize SCL-MCL copolymer though

3HB was the dominant monomer.

Class I clone 18 synthesizes SCL and SCL-MCL PHA

Clone 18 contained 37,818-bp metagenomic DNA [GenBank: KT944264], which probably

originated from *β-Proteobacteria Rubrivirax* (Table 3). The annotated *phaC₁₈*, *phaA* and *phaB* were

organized in one operon (Fig. 3A), as also occurred in clones 27 and 51 (Supplementary 1). $PhaC_{18}$

[GenBank: ALV86462] shared 100% and 99% identity to $PhaC_{27}$ [GenBank: ALV86602] and

$PhaC_{51}$ [GenBank: ALV86651] respectively. The $PhaC_{18}$ protein best matched to its homolog in

Azohydromonas austrilica (Table 3), and was 71%, 49% and 61% identical to the $PhaC_{14}$, $PhbC_{Sm}$

and $PhbC1_{Re}$ respectively.

SCL-MCL PHA (C4-C6-C8) was produced when PpUW2 (clone 18) was grown with

octanoate (Table 4), as occurred in PpUW2 with clone 16. However, the fraction of C6-C8

monomers was only 3% in the PHA from clone 18, in contrast to the ~83% in the copolymer from

clone 16. The yield of PHA was ~7 fold less than that produced by clone 14, but C8 monomer was

absent (Table 4). PHA of 3HB-3HV monomers was accumulated in nonanoic acid-grown PpUW2

(clone 18). The ratio of 3HB to 3HV increased 5 fold compared to that in PpUW2 (clone 14) under

the same growth conditions (Table 4). When PpUW2 carrying clone 18 was grown on gluconic acid,

only PHB was produced. The yield of polymer increased 5 fold compared to that in octanoate-grown

cells, and also the second highest quantity of PHA produced among all strains grown on gluconic

acid. These results implied that $PhaC_{18}$ could synthesize SCL- and SCL-MCL PHA dependent of the

available carbon source.

Class I clone 19 synthesizes SCL and SCL-MCL PHA

Clone 19 contained 25,085-bp of DNA that might originate from *β-Protoebacteria Leptothrix*

(Table 3) [GenBank: KT944265]. Three ORFs encoded $PhaC_{19}$, PhaA and PhaB for PHA

biosynthesis (Fig. 3A). $PhaC_{19}$ shared 99% identity with $PhaC_6$ and $PhaC_3$ (Fig. 2). The best match

for $PhaC_{19}$ is the ortholog of *A. austrilica* at 73% (Table 3). The *$phaC_{19}AB$* genes in clone 19

probably consisted of one operon, and the promoter was located upstream of the *$phaC_{19}$*.

Clone 19 was able to synthesize SCL-MCL PHA (C4-C6-C8) with octanoate but only PHB

(C4) in gluconic acid-grown PpUW2 (Table 4), as occurred in PpUW2 carrying clone 18. However

~5 fold more PHA was synthesized by clone 19 than was produced by clone 18 in octanoate medium

(Table 4). In contrast, the quantity of PHA was lower in gluconic acid-grown PpUW2 with clone 19

than it was in the same host carrying clone 18. Moreover, PHA composed of 3HB-3HV monomers

was synthesized in noanoic acid-grown PpUW2 (clone 19), as occurred in PpUW2 (clone 18). These

data suggested that $PhaC_{19}$ was able to synthesize both SCL and SCL-MCL PHA.

Class I clone 25 synthesizes SCL and SCL-MCL PHA

Clone 25 contained 41,012-bp of DNA [GenBank: KT944269], which was predicted to

originate from *β-Proteobacteria Variovorax* (Table 3)*. The *$phaC_{25}$* and *phaA* genes were apparently

located in a single operon (Fig. 3A). A *phaB* gene was located 9 open reading frames downstream of

the *phaA*. The *$phaC_{25}A$* and *phaB* loci in clone 25 are similar to those in clone P1N3 (Supplementary

1), though the *phaB* gene in clone P1N3 was four open reading frames away from *$phaC_{P1N3}A$*. PhaC

proteins of clones P1N3 and 25 shared 77% identity. In addition, $PhaC_{25}$ was 35% and 61% identical

to $PhbC_{Sm}$ and $PhbC1_{Re}$. The conserved amino acid Asp130 affecting PhaC substrate specificity

(Matsumoto et al. 2005) was replaced by Ala152 (Fig. 3B).

SCL-MCL PHA (C4-C6-C8) was produced in PpUW2 harbouring clone 25 when grown on

octanoate, and only PHB (C4) was isolated from gluconic acid-grown cells (Table 4). Similar results

were obtained in PpUW2 with clone 18 or clone 19 under the same conditions. However, the

quantity of C6-C8 monomers was greater with clone 25 that those with clone 18 or 19. Nonanoic

acid-grown PpUW2 (clone 25) produced 3HB-3HV copolymer similar to that from PpUW2 with

clone 18 or clone 19 (Table 4). These data suggested that $PhaC_{25}$ had similar properties to those of

$PhaC_{18}$ and $PhaC_{29}$, but different from that of $PhaC_{14}$.

Complementation of _S. meliloti phaC_ mutant with 11AW PHA[+] clones

In an earlier study, _S. meliloti_ Rm11476 (_exoY_::Tn5 _phaC_::Tn5-233) had been employed for

functionally harvesting metagenomic library clones pCX92, pCX9M1, pCX9M3 and pCX9M5

expressing Class I PHA synthases (Schallmey et al. 2011). The PhaA/PhbB and PhaB/PhbB proteins

encoded by the cloned metagenomic DNA and/or _S. meliloti_ genome could supply (R)-3-

hydroxybutyrate for PHB biosynthesis catalyzed by the PhaC/PhbC enzymes. In order to examine

the possibility that the PHA[+] clones of 11AW metagenomic library were able to complement the

PHA[-] phenotype of _S. meliloti_ Rm11476, The Class I clones 14, 18, 19 and 25 (Table 3) were

introduced into _S. meliloti_ Rm11476. In each case, complementation was observed, indicating PhaC

function in the _S. meliloti_ background (Fig. 4), with greatest accumulation of PHA observed with

clone 14. However, when clones pCX92, pCX9M1, pCX9M3 and pCX9M5 were transferred to _P._

putida PpUW2 (PHA[-]), PHA was not detected in the recombinant strains grown with either

octanoate or gluconic acid (data not shown). These results strengthen the argument that multiple

hosts should be employed to screen for novel _phaC_ genes.

We then asked whether Class II *phaC* could function in the *S. meliloti* background. First, we

observed no detection of PHA when $phaC1_{Pp}$ (Plac::$phaC1_{Pp}$) was expressed in Rm11476 even

though the *lac* promoter is functional and 3-hydroxybutyrate is supplied by the $PhbAB_{Sm}$. Similarly,

PHA^+ clones 1 and 20 carrying Class II *phaC* genes also failed to complement the PHB^- phenotype

of *S. meliloti* Rm11476 (Fig. 4), implying that the $phaC2_1$ and $phaC_{20}$ genes were not expressed

and/or the substrate (R)-3-hydroxyalkanoic acids were absent in *S. meliloti*. However, when clone 16

was introduced into *S. meliloti* Rm111476, PHA was accumulated to a similar level as with

Rm11476 containing PHB^+ clones pCX92, pCX9M1, pCX9M3 or pCX9M5 (Fig. 4). These data

indicated that the $phaC1_{16}$ gene was expressed and its product was functional. These results suggest

that *P. putida* is a more permissive surrogate host than *S. meliloti* for screening of novel PHA

synthases.

Discussion

Biosynthesis of SCL-MCL PHA requires PHA synthases having a broad range of substrate

specificity and SCL and MCL precursors (R-3-hydroxyacyl-CoAs) must be available within the cell.

Building on earlier work that has sought to mine metagenomic DNA for modification of bacterial

PHA production, we have successfully isolated cosmid clones that are able to functionally

complement a *P. putida* PHA synthesis mutant. DNA sequence analysis revealed that the isolated

metagenomic DNA originated from a broad diversity of bacteria, and encoded either Class I or Class

II PHA synthases enzymes. Of note is that DNA from most of the isolated clones did not very

closely match known sequences. Also of interest is the influence that the clones had on the quality

and quantity of PHA produced in the *P. putida* surrogate host background.

A major appeal of PHA is that, as alternatives to conventional fossil fuel-derived plastics,

they can be produced from renewable resources. The potential to improve the efficiency and cost of

production, and expand the range of polymers and copolymers that are available for production,

should impact on competitiveness and adoption of these materials in the marketplace. In this context,

it is important to understand how the properties of PHA are determined by the PHA synthase

enzyme and the substrate that is available to those enzymes, which is in turn influenced by the

culture conditions and the metabolic pathways leading to substrate formation (Meng et al. 2014).

PHA consisting of both short-chain-length (SCL, $\leq$C5) and medium-chain-length (MCL, $\geq$C6)

monomers have great flexibility, decreased breakage and reduced melting point (Noda et al. 2005).

Most naturally occurring PhaC proteins polymerize either SCL- or MCL-monomers, but there are a

few know examples such as the Class I PhaC of *Aeromonas caviae* (Fukui and Doi 1997) and some

Pseudomonas Class II PhaC (Matsusaki et al. 1998) that can synthesize SCL-MCL copolymers.

In the present work, heterologous complementation of the *P. putida* PHA⁻ strain facilitated

efficient simultaneous screening of millions of 11AW metagenomic clones, resulting in recovery of

a greater number of novel Class I, Class II, and unclassified PHA synthases than from previous

reports (Schallmey et al. 2011; Cheema et al. 2012). Additionally, the clones carrying Class I *phaC*

isolated in *P. putida* were able to complement the PHA⁻ phenotype of *S. meliloti*, but the previously

isolated Class I clones isolated in *S. meliloti* (Schallmey et al. 2011) failed to synthesize PHA in *P.*

putida. These results support the development of multi-host systems to increase the chances of the

successful expression of *pha* genes of interest.

Biosynthesis of SCL-MCL PHA of various forms by Class I clones 14, 18, 19 and 25 and

Class II clones 16 and 20 suggested availability of a full range of R-3-hydroxyacyl-CoAs derived

from β-oxidation of fatty acids such as nonanoic acid and octanoate. The differential monomer

compositions in PHA synthesized by the Class II enzymes $PhaC_{16}$ and $PhaC_{20}$ are of particular

interest, especially since most Class II PHA synthases are unable to incorporate SCL monomers such

as 3HB and 3HV. Changes of conserved amino acid residues Ser325Thr, Leu484Val, and

Glu508Leu (conserved positions in *P. putida* PhaC1) have been identified to be involved in PhaC

substrate specificity and PHA yield (Takase et al. 2003; Shozui et al. 2010; Chen et al. 2014).

Presence of the amino acids Thr353 (corresponding to *P. putida* PhaC1 position 325), Val516

(corresponding to *P. putida* PhaC1 position 484) and Ala540 (corresponding to *P. putida* PhaC1

position 508) $PhaC_{20}$ might contribute to the enzyme's ability to incorporate SCL monomers

compared to those of 11AW $PhaC2_1$ and PHA synthases of *P putida* KT2440. Similarly, the

presence of three residues (Thr317, Val478 and Leu502) at these positions in $PhaC1_{16}$, most likely

resulted in increased 3HB content as demonstrated previously with engineered Class II PHA

synthases (Takase et al. 2003; Shozui et al. 2010; Chen et al. 2014).

Conclusions

We obtained 27 clones encoding Class I, II and unclassified PHA synthases by functional

screening of soil metagenomic cosmid clones in a *P. putida* PHA⁻ strain. Seven clones that were

characterized in more detail were able to produce a broad range of polymers and copolymers,

including SCL-MCL mixtures, depending on carbon source. Through this work we have

demonstrated the potential for using metagenome-derived clones for production of a variety of PHAs

of possible industrial utility. The collection of PHA metabolism genes from uncultivated organisms

provides not only a resource for production strain development, but also a series of sequence

templates that could prove useful in enzyme engineering efforts directed towards generation of PHA

products with desired properties.

Nucleotide sequence accession numbers

Complete sequences of 11AW metagenomic PHA$^+$ DNA have been deposited in GenBank

nucleotide sequence database [GenBank: KT944254-KT944278, KU728995 and KU728996].

Abbreviations

P. putida, *Pseudomonas putida*; PHA, polyhydroxyalkanoate; PhaA/PhbA, β-ketothiolase;

PhaB/PhbB, acetoacetyl-CoA reductase; PhaC/PhbC, PHA synthase; PhaZ, PHA depolymerase;

PhaJ, R-specific enoyl-CoA hydratase; 3HA, 3-hydroxyalkanoate; 3HB, 3-hydroxybutyrate; 3HV.

3-hydroxyvalerate; 3HHx, 3-hydroxyhexanoate; 3HP, 3-hydroxyheptanoate; 3HO, 3-

hydroxyoctanoate; 3HN, 3-hydroxynonanoate; 3HD, 3-hydroxydecanoate; SCL, short chain length;

MCL, medium chain length; *S. meliloti*, *Sinorhizobium meliloti*; GC-MS, gas chromatography-mass

spectrometry; CDW, cell dry weight.

Authors's contributions

JC and TCC conceived and designed this study, carried out all the experiments, and prepared

the manuscript. Both authors read and approved the final manuscript.

Acknowledgements

We are very grateful to Ricardo Nordeste for assisting with GC-MS analysis of PHA. We

wish to thank Dr. Bruce Ramsay (Polyferm Canada) for providing PHA standards, and Dr. Josh

Neufeld for use of the FilterMax F5 Multi-mode microplate reader.

Funding

This work was financially supported by a New Directions grant from Ontario Ministry of Agriculture, Food, and Rural Affairs (award number 381646-09), by a Strategic Projects grant from the Natural Sciences and Engineering Research Council of Canada (award number ND2012-1679), and by Genome Canada, through the Applied Genomics Research in Bioproducts or Crops (ABC) program for the grant titled "Microbial Genomics for Biofuels and Co-Products from Biorefining Processes" awarded to David Levin and Richard Sparling.

Conflict of interest

Trevor Charles received partial financial support for presenting some of this material at the Society for Industrial Microbiology and Biotechnology Annual Meeting in 2014 and 2015. Jiujun Cheng declares no conflict of interest.

Ethical statement

The authors certify that this manuscript has not been published previously, and not under consideration for publication elsewhere, in whole or in part. No data have been fabricated or manipulated (including images), and no data, text, or theories by others are presented as if they were the authors' own. Consent to submit has been received explicitly from all the authors listed. And authors whose names appear on the submission have contributed sufficiently to the scientific work and therefore share collective responsibility and accountability for the results. This article does not contain any studies with human participants or animals performed by any of the authors.

580 **References**
581

Cheema S, Bassas-Galia M, Sarma PM, Lal B, Arias S (2012) Exploiting metagenomic diversity for
novel polyhydroxyalkanoate synthases: Production of a terpolymer poly(3-hydroxybutyrate-co-
3-hydroxyhexanoate-co-3-hydroxyoctanoate) with a recombinant *Pseudomonas putida* strain.
Bioresource Technology 103:322–328. doi: 10.1016/j.biortech.2011.09.098

Chen Y-J, Tsai P-C, Hsu C-H, Lee C-Y (2014) Critical residues of class II PHA synthase for
expanding the substrate specificity and enhancing the biosynthesis of polyhydroxyalkanoate.
Enzyme and Microbial Technology 56:60–66. doi: 10.1016/j.enzmictec.2014.01.005

Cheng J, Pinnell L, Engel K, Neufeld JD, Charles TC (2014) Versatile broad-host-range cosmids for
construction of high quality metagenomic libraries. Journal of Microbiological Methods 99:27–
34. doi: 10.1016/j.mimet.2014.01.015

de Eugenio LI, Escapa IF, Morales V, Dinjaski N, Galàn B, García JL, Prieto MA (2010) The
turnover of medium-chain-length polyhydroxyalkanoates in *Pseudomonas putida* KT2442 and
the fundamental role of PhaZ depolymerase for the metabolic balance. Environ Microbiol
12:207–221. doi: 10.1111/j.1462-2920.2009.02061.x

Edgar RC (2004) MUSCLE: a multiple sequence alignment method with reduced time and space
complexity. BMC Bioinformatics 5:113–19. doi: 10.1186/1471-2105-5-113

Escapa IF, Morales V, Martino VP, Pollet E, Avérous L, García JL, Prieto MA (2011) Disruption of
β-oxidation pathway in *Pseudomonas putida* KT2442 to produce new functionalized PHAs with
thioester groups. Appl Microbiol Biotechnol 89:1583–1598. doi: 10.1007/s00253-011-3099-4

Fellay R, Frey J, Krisch H (1987) Interposon mutagenesis of soil and water bacteria: a family of
DNA fragments designed for in vitro insertional mutagenesis of gram-negative bacteria. Gene
52:147–154.

Foong CP, Lau N-S, Deguchi S, Toyofuku T, Taylor TD, Sudesh K, Matsui M (2014) Whole
genome amplification approach reveals novel polyhydroxyalkanoate synthases (PhaCs) from
Japan trench and nankai trough seawater. BMC Microbiol 14:e11836. doi: 10.1186/s12866-014-
0318-z

Fukui T, Doi Y (1997) Cloning and analysis of the poly(3-hydroxybutyrate-co-3-hydroxyhexanoate)
biosynthesis genes of *Aeromonas caviae*. J Bacteriol 179:4821–4830. doi: 10.1186/s12866-014-
0318-z

Fukui T, Shiomi N, Doi Y (1998) Expression and characterization of (R)-specific enoyl coenzyme A
hydratase involved in polyhydroxyalkanoate biosynthesis by Aeromonas caviae. J Bacteriol
180:667–673.

Hoppensack A, Rehm BH, Steinbüchel A (1999) Analysis of 4-phosphopantetheinylation of
polyhydroxybutyrate synthase from *Ralstonia eutropha*: generation of beta-alanine auxotrophic
Tn5 mutants and cloning of the *panD* gene region. J Bacteriol 181:1429–1435.

Jia Y, Kappock TJ, Frick T, Sinskey AJ, Stubbe J (2000) Lipases provide a new mechanistic model
for polyhydroxybutyrate (PHB) synthases: characterization of the functional residues in
Chromatium vinosum PHB synthase. Biochemistry 39:3927–3936. doi: 10.1021/bi9928086

Kawashima Y, Cheng W, Mifune J, Orita I, Nakamura S, Fukui T (2012) Characterization and
functional analyses of R-specific enoyl coenzyme A hydratases in polyhydroxyalkanoate-
producing *Ralstonia eutropha*. Appl Environ Microbiol 78:493–502. doi: 10.1128/AEM.06937-
11

Keshavarz T, Roy I (2010) Polyhydroxyalkanoates: bioplastics with a green agenda. Current
Opinion in Microbiology 13:321–326. doi: 10.1016/j.mib.2010.02.006

Lam KN, Hall MW, Engel K, Vey G, Cheng J, Neufeld JD, Charles TC (2014) Evaluation of a
pooled strategy for high-throughput sequencing of cosmid clones from metagenomic libraries.
PLoS ONE 9:e98968. doi: 10.1371/journal.pone.0098968

Matsumoto K, Takase K, Aoki E, Doi Y, Taguchi S (2005) Synergistic effects of Glu130Asp
substitution in the type II polyhydroxyalkanoate (PHA) synthase: enhancement of PHA
production and alteration of polymer molecular weight. Biomacromolecules 6:99–104. doi:
10.1021/bm049650b

Matsusaki H, Manji S, Taguchi K, Kato M, Fukui T, Doi Y (1998) Cloning and molecular analysis
of the Poly(3-hydroxybutyrate) and Poly(3-hydroxybutyrate-co-3-hydroxyalkanoate)
biosynthesis genes in *Pseudomonas* sp. strain 61-3. J Bacteriol 180:6459–6467.

Meng D-C, Shen R, Yao H, Chen J-C, Wu Q, Chen G-Q (2014) Engineering the diversity of
polyesters. Curr Opin Biotechnol 29:24–33. doi: 10.1016/j.copbio.2014.02.013

Noda I, Green PR, Satkowski MM, Schechtman LA (2005) Preparation and properties of a novel
class of polyhydroxyalkanoate copolymers. Biomacromolecules 6:580–586. doi:
10.1021/bm049472m

Park SJ, Kim TW, Kim MK, Lee SY, Lim S-C (2012) Advanced bacterial polyhydroxyalkanoates:
towards a versatile and sustainable platform for unnatural tailor-made polyesters. Biotechnology
Advances 30:1196–1206. doi: 10.1016/j.biotechadv.2011.11.007

Patil KR, Roune L, McHardy AC (2012) The PhyloPythiaS web server for taxonomic assignment of
metagenome sequences. PLoS ONE 7:e38581. doi: 10.1371/journal.pone.0038581

Pärnänen K, Karkman A, Virta M, Eronen-Rasimus E, Kaartokallio H (2015) Discovery of bacterial
polyhydroxyalkanoate synthase (PhaC)-encoding genes from seasonal Baltic Sea ice and cold
estuarine waters. Extremophiles 19:197–206. doi: 10.1007/s00792-014-0699-9

Quandt J, Hynes MF (1993) Versatile suicide vectors which allow direct selection for gene
replacement in Gram-negative bacteria. Gene 127:15–21. doi: 10.1016/0378-1119(93)90611-6

Rehm BHA (2003) Polyester synthases: natural catalysts for plastics. Biochem J 376:15–33. doi:
10.1042/BJ20031254

Sato S, Kanazawa H, Tsuge T (2011) Expression and characterization of (R)-specific enoyl
coenzyme A hydratases making a channeling route to polyhydroxyalkanoate biosynthesis in
Pseudomonas putida. Appl Microbiol Biotechnol 90:951–959. doi: 10.1007/s00253-011-3150-5

Schallmey M, Ly A, Wang C, Meglei G, Voget S, Streit WR, Driscoll BT, Charles TC (2011)
Harvesting of novel polyhydroxyalkanaote (PHA) synthase encoding genes from a soil
metagenome library using phenotypic screening. FEMS Microbiol Lett 321:150–156. doi:
10.1111/j.1574-6968.2011.02324.x

Sheu DS, Wang YT, Lee CY (2000) Rapid detection of polyhydroxyalkanoate-accumulating
bacteria isolated from the environment by colony PCR. Microbiology 146 (Pt 8):2019–2025.
doi: 10.1099/00221287-146-8-2019

Shozui F, Sun J, Song Y, Yamada M, Sakai K, Matsumoto K, Takase K, Taguchi S (2010) A new
beneficial mutation in *Pseudomonas* sp. 61-3 polyhydroxyalkanoate (PHA) synthase for
enhanced cellular content of 3-hydroxybutyrate-based PHA explored using its enzyme homolog
as a mutation template. Biosci Biotechnol Biochem 74:1710–1712. doi: 10.1271/bbb.100224

Simon C, Daniel R (2011) Metagenomic analyses: past and future trends. Appl Environ Microbiol
77:1153–1161. doi: 10.1128/AEM.02345-10

Steinbüchel A, Lütke-Eversloh T (2003) Metabolic engineering and pathway construction for
biotechnological production of relevant polyhydroxyalkanoates in microorganisms. Biochemical
Engineering Journal 16:81–96. doi: 10.1016/S1369-703X(03)00036-6

Tai YT, Foong CP, Najimudin N, Sudesh K (2015) Discovery of a new polyhydroxyalkanoate
synthase from limestone soil through metagenomic approach. Journal of Bioscience and
Bioengineering. doi: 10.1016/j.jbiosc.2015.08.008

Takase K, Taguchi S, Doi Y (2003) Enhanced synthesis of poly(3-hydroxybutyrate) in recombinant
Escherichia coli by means of error-prone PCR mutagenesis, saturation mutagenesis, and in vitro
recombination of the type II polyhydroxyalkanoate synthase gene. J Biochem 133:139–145.

Tamura K, Stecher G, Peterson D, Filipski A, Kumar S (2013) MEGA6: Molecular Evolutionary
Genetics Analysis version 6.0. Mol Biol Evol 30:2725–2729. doi: 10.1093/molbev/mst197

Tortajada M, da Silva LF, Prieto MA (2013) Second-generation functionalized medium-chain-length
polyhydroxyalkanoates: the gateway to high-value bioplastic applications. Int Microbiol 16:1–15.
doi: 10.2436/20.1501.01.175

Tripathi L, Wu L-P, Dechuan M, Chen J, Wu Q, Chen G-Q (2013) *Pseudomonas putida* KT2442 as
a platform for the biosynthesis of polyhydroxyalkanoates with adjustable monomer contents and
compositions. Bioresource Technology 142:225–231. doi: 10.1016/j.biortech.2013.05.027

Verlinden RAJ, Hill DJ, Kenward MA, Williams CD, Radecka I (2007) Bacterial synthesis of
biodegradable polyhydroxyalkanoates. J Appl Microbiol 102:1437–1449. doi: 10.1111/j.1365-
2672.2007.03335.x

Wu H-A, Sheu D-S, Lee C-Y (2003) Rapid differentiation between short-chain-length and medium-
chain-length polyhydroxyalkanoate-accumulating bacteria with spectrofluorometry. Journal of
Microbiological Methods 53:131–135. doi: 10.1016/S0167-7012(02)00232-4

Zhu W, Lomsadze A, Borodovsky M (2010) Ab initio gene identification in metagenomic sequences.
Nucleic Acids Research 38:e132–e132. doi: 10.1093/nar/gkq275

Figure legends

Fig. 1. Activity-based screening of 11AW metagenomic library clones carrying PHA biosynthesis

genes in *P. putida* PpUW2 (PHA$^-$). PpUW2 recombinant strains were grown in 0.1N M63 medium

containing 0.5% Na octanoate and Nile red (0.5 μg/ml). Cosmid pJC8 and pJC67 (P*lac*::*phaC1$_{Pp}$*)

were used as negative and positive controls respectively. 11AW clones 14 and 16 were streaked on

the same selection plate to verify the PHA$^+$ phenotype initially screened on LB with 0.5% (w/v)

octanoate.

Fig. 2. Phylogenetic analysis of the polyhydroalkanoate synthases (PhaC) of 11AW metagenomic

library clones. Protein sequence alignments were performed using MUSCLE (Edgar 2004).

Neighbour-jointing phylogenetic trees were generated with MEGA6 (Tamura et al. 2013). GenBank

accession numbers of 11AW PhaC proteins: PhaC2$_1$ (ALV86274), PhaC$_3$ (ALV86289), PhaC$_4$

(ALV86299), PhaC$_5$ (ALV86308), PhaC$_6$ (ALV86351), PhaC$_7$ (partial, ALV86358), PhaC$_8$

(ALV86364), PhaC$_{10}$ (partial, ALV86387), PhaC$_{14}$ (ALV86397), PhaC1$_{16}$ (ALV86417), PhaC2$_{16}$

(ALV86419), PhaC$_{18}$ (ALV86462), PhaC$_{19}$ (ALV86476), PhaC$_{20}$ (ALV86493), PhaC$_{21}$

(ALV86517), PhaC$_{23}$ (ALV86529), PhaC$_{25}$ (ALV86574), PhaC$_{27}$ (ALV86602), PhaC1$_{50}$

(ALV86626), PhaC2$_{50}$ (ALV86626), PhaC$_{51}$ (ALV86651), PhaC$_{P1N3}$ (ALV86715), PhaC$_{P2N8}$

(ALV86750), PhaC$_{P2N10}$ (partial, ALV86755), PhaC$_{P4N10}$ (ALV86768), PhaC$_{P8N7}$ (ALV86771).

Fig. 3. The *pha* genes and PhaC proteins in 11AW metagenomic DNA library clones. (A) The PHA$^+$

clones are classified into Class I and II based on the PhaC protein sequences. (B) Conserved amino

acids required for the activity of PHA synthases are indicated in closed triangles. Residues affecting

substrate specificity were marked by open triangles. The positions of amino acid residues were

numbered based on the sequence of PhaC1$_{16}$ [GenBank: ALV86417]. GenBank accession numbers

of PhaC proteins: PhaC$_{14}$, ALV86397; PhaC$_{18}$, ALV86462; PhaC$_{19}$, ALV86476; PhaC$_{25}$,

ALV86574; PhaC1$_2$, ALV86274; PhaC$_{20}$, ALV86493; PhaC1$_{16}$, ALV86417; *C. necator* PhbC1,

AAA21975; *P. putida* PhaC1, Q88D25; and *P.* sp. 61-3 PhaC1, BAA36200.

Fig. 4. Complementation of *S. meliloti* Rm11476 (*phbC*) with metagenomic clones. The cosmid

DNA of PHA$^+$ clones encoding Class I and II PHA synthases was transferred to *S. meliloti* Rm11476

by conjugation. Recombinant strains were grown in YM medium. PHA production was estimated by

measuring the fluorescence of Nile red stained cells. *S. meliloti* wild type Rm1021 and previously

isolated Class I clones (CX92, pCX92M1, pCX92M3, and pCX92M5; (Schallmey et al. 2011)) were

used as controls.

Figure 1.

Fig. 2

A

B

Fig. 3

Fig. 4

Table 1. Bacterial strains, plasmids and cosmids

Strain, plasmids and cosmids	Relevant characteristics*	Reference
E. coli		
DH5α	F⁻ ф80 *lacZΔM15Δ*(*lacZYA-argF*)*U169 recA1 endA1 hsdR17 phoA supE44 thi-1 gyrA96 relA1*	Life Technologies
DH5α (Rif^R)	A spontaneous Rif^r mutant of DH5α	(Cheng et al. 2014)
HB101	F⁻ *thi-1 hsdS20 supE44 recA13 ara-14 leuB6 proA2 lacY1 galK2 rpsL20 xyl-5 mtl-1*, Sm^R	Life Technologies
S. meliloti		
Rm1021	Wild type, Sm^R	Lab collection
Rm11476	Rm1021, *exoY*::Tn5 *phbC*::Tn5-233, Sm^R Nm^R	(Schallmey et al. 2011)
P. putida		
KT2440	Wild type	Lab collection
PpUW1	A spontaneous Rif^R mutant of KT2440, Rif^R	This study
PpUW2	PpUW1 (*phaC1-phaZ-phaC2*::ΩKm^R), Rif^R Km^R	This study
Plasmids/cosmids		
pRK600	pRK2013 *tra* Nm^R::Tn9, Cm^R	Lab collection
pRK7813	IncP *oriT cos lacZα*, Tc^R	Lab collection
pJC8	Broad-host-range Gateway® entry cosmid, Tc^R	(Cheng et al. 2014)
pJQ200-SK	Cloning vector, *sacB*, Gm^R	(Quandt and Hynes 1993)
pHP45-ΩKm^R	Ω-Km^R cassette, Km^R	(Fellay et al. 1987)
pJC63	pJQ200-SK carrying 5′-*phaC1* of *P. putida* KT2440, Gm^R	This study
pJC69	pJC63 carrying 3′-*phaC2* of *P. putida* KT2440, Gm^R	This study
pJC71	pJC69 carrying Ω-Km^R between the *phaC1′* and *phaC2′* fragments, Gm^R Km^R	This study
pJC67	pRK7813 carrying P*lac*::*phaC1$_{Pp}$*, Tc^R	This study
pJC123	pRK7813 carrying P*lac*::*phaC1$_{16}$*, Tc^R	This study
pJC157	pRK7813 carrying P*lac*::*phaC1$_{16}$*::*phaC2$_{16}$*, Tc^R	This study
pJC159	pRK7813 carying P*lac*::*phaC2$_{16}$*, Tc^R	This study
pCX92	pRK7813 carrying *phaC* and *phaP*, Tc^R	(Schallmey et al. 2011)
pCX9M1	pRK7813 carrying *phaC*, *phaA* and *phaB*, Tc^R	(Schallmey et al. 2011)
pCX9M3	pRK7813 carrying *phaC*, Tc^R	(Schallmey et al. 2011)
pCX9M5	pRK7813 carrying *phaC* and *phaB*, Tc^R	(Schallmey et al. 2011)

*Antibiotic resistances: Cm^r, chloramphenicol; Gm^R, gentamycin; Km^R, kanamycin; Nm^R, neomycin; Sm^R, streptomycin; Tc^R, tetracycline.

Table 2. DNA oligonucleotides

Oligo ID	Sequence
JC159	CGCC<u>AAGCTT</u>G**TG**A<u>AGGAG</u>CGTCGTAGATGAGTAAC
JC160	GCGC<u>GGATCC</u>TCAACGCTCGTGAACGTAGGT
JC161	GCGC<u>TCTAGA</u>CACGAAAGTCGGGTAACCAAT
JC162	GGTC<u>GGATCC</u>CGCCAGCTGATGATGAAGGTC
JC163	GCTG<u>GGATCC</u>GTTGTGGCTGGAGTGGATCAC
JC164	CGCC<u>CTCGAG</u>CACCGGTACCCTGATCTGATA
phaCF1	ATCAACAARTWCTACRTCYTSGACCT
phaCR2	GTSTTCRTSRTSWSCTGGCGCAACCC
phaCR4	AGGTAGTTGTYGACSMMRTAGKTCCA
JC202	CCGGATGCA**TGA**T<u>AGGA</u>CGCCGGATGACAGCAGAGAAGGCTGAG
KC228	CCCAGATCTGCAGTATTCTAGATTCAGGCGCGCACGCGCACGTA
JC204	CCGGTCTAG<u>AGGA</u>ACGTCCCATGAGCAATCC
JC205	GGGGATGCATGCCTCAGGCGAAGTACTGGCCGCCATTG
JC301	CGCG<u>AAGCTT</u>C**TGA**<u>AGGAG</u>AGAACACATGGCGAAGACCA
JC302	GGGG<u>GAATTC</u>TGAATATCAGCGTCGGAGCATG
JC333	CGCG<u>AAGCTT</u>G**TG**A<u>GGAG</u>AGATCTATGAGCTCCCTCCCCGTGGCCGAACGTTC
JC334	CGCT<u>GAATTC</u>TCTACGCCGCCTGACCCACGAT
JC337	ACCA<u>GAATTC</u>A<u>GGAG</u>ATCTATATGAGCTCCCTCCCCGTGGCCGAAC
pJET forward	CGACTCACTATAGGGAGAGCGGC
pJET reverse	AAGAACATCGATTTTCCATGGCAG

Restriction sites are underlined. Translation stop codon is in bold, and ribosomal binding site is

double underlined.

Table 3. 11AW metagenomic clones encoding PHA synthases (PhaC).

PHA⁺ Clones	DNA (bp)	GC content (%)	Possible origin of cloned DNA	Best match of PHA synthases	Identity (%)	Cosmid Genbank ID
Class I						
3	>16,077	67.5	*Leptothrix* (*â-Proteobacteria*)	Hypothetical protein (WP_046112351), *Aquincola tertiaricarbonis*	74	KT944255
4	40,585	65.7	*Leptothrix* (*â-Proteobacteria*)	Polyhydroxyalkanoate synthase (AAD10274), *Azohydromonas lata*	72	KT944256
6	>11,479	67.7	*Leptothrix* (*â-Proteobacteria*)	Polyhydroxyalkanoate synthase (AAD10274), *Azohydromonas lata*	73	KT944258
8	34,220	67.8	*Leptothrix* (*â-Proteobacteria*)	Poly-(R)-hydroxyalkanoic acid synthase (WP_009550222), Burkholderiales bacterium JOSHI_001	74	KT944260
14	29,565	69.1	*Methylibium* (*â-Proteobacteria*)	Poly-(R)-hydroxyalkanoic acid synthase (WP_009550222), Burkholderiales bacterium JOSHI_001	74	KT944262
18	37,718	69.4	*Rubrivirax* (*â-Proteobacteria*)	Hypothetical protein (WP_043458567), *Azohydromonas austrilica*	73	KT944264
19	25,085	68.1	*Leptothrix* (*â-Proteobacteria*)	Polyhydroxyalkanoate synthase (AAD10274), *Azohydromonas lata*	73	KT944265
21	33,064	69.3	*Gemmatimonas* (*Gemmatimonadetes*)	Poly-(R)-hydroxyalkanoic acid synthase (WP_009550222), Burkholderiales bacterium JOSHI_001	73	KT944267
23	42,865	69.6	*Leptothrix* (*â-Proteobacteria*)	Poly-(R)-hydroxyalkanoic acid synthase (WP_009550222), Burkholderiales bacterium JOSHI_001	74	KT944268
25	41,012	66.8	*Varivorax* (*â-Proteobacteria*)	Poly-(R)-hydroxyalkanoic acid synthase (WP_018902870), *Variovorax paradoxus*	93	KT944269
27	37,156	67.0	*Rubrivivax* (*â-Proteobacteria*)	Hypothetical protein (WP_043458567), *Azohydromonas austrilica*	73	KT944270
51	38,794	69.9	*Rubrivirax* (*â-Proteobacteria*)	Hypothetical protein (WP_043458567), *Azohydromonas australica*	73	KT944272
P2N8	22,625	66.3	*Xanthomondaceae* (*ã-Proteobacteria*)	Poly(3-hydroxyalkanonate) synthase (WP_028465014), *Nisaea denitrificans*	55	KT944272
P11N2	40,293	58.4	*Janthinobacterium* (*â-Proteobacteria*)	Poly(3-hydroxyalkanonate) synthase (WP_041099836), *Sulfuritalea hydrogenivorans*	73	KT944278
P1N3	32,797	68.6	*Variovorax* (*â-Proteobacteria*)	Poly(3-hydroxyalkanonate) synthase (WP_028253507), *Variovorax* sp. URB0020	86	KT944273
P4N10	>2,724	64.2	*Leptothrix* (*â-Proteobacteria*)	Polyhydroxyalkanonate synthase (AAD10274), *Azohydromonas lata*	72	KT944276
P8N7	>5,619	67.6	*Sphingomonadaceae* (*á-Proteobacteria*)	Poly-(R)-hydroalkanoic acid synthase (WP_033922275), *Sphingomonas* sp. 37zxx	67	KT944277
Class II						
16 (PhaC1₁₆)	33,900	66.1	*Methylibium* (*â-Proteobacteria*)	PhaC1 synthase (AHL43537), *Pseudomonas* sp.F15	48	KT944263
50 (PhaC1₅₀)	22,965	67.0	*Hydrocarboniphaga* (*ã-Proteobacteria*)	Polyhydroxyalkanoic acid synthase (WP_007929265), *Pseudomonas* sp. GM17	46	KT944271
1 (PhaC1₂)	42,747	60.0	*Pseudomonas* (*ã-Proteobacteria*)	Poly(R)-hydroxyalkanoic acid synthase (WP_033055349), *Pseudomonas mandelii*	98	KT944254
2	59,765	59.1	*Pseudomonas* (*ã-Proteobacteria*)	Poly(R)-hydroxyalkanoic acid synthase (WP_033055349), *Pseudomonas mandelii*	98	KU729001
15	59,765	59.1	*Pseudomonas* (*ã-Proteobacteria*)	Poly(R)-hydroxyalkanoic acid synthase (WP_033055349), *Pseudomonas mandelii*	98	KU728996
5	37473	66.3	*Burkholderia* (*â-Proteobacteria*)	Poly-beta-hydroxybutyrate polymerase (WP_051556988), *Geodermatophilaceae bacterium* URHB0062	53	KT944257
7	>407	65.6	*Pseudomonadaceae* (*ã-Proteobacteria*)	Poly(R)-hydroxyalkanoic acid synthase (WP_008439266), *Pseudomonas* sp. Ag1	100	KT944259
10	>404	63.9	*Pseudomonas* (*ã-Proteobacteria*)	PHA synthase (BAE07114), uncultured bacterium	98	KT944261
20	30,964	66.8	*Salinibacter* (*Bacteroidetes*)	Poly(R)-hydroxyalkanoic acid synthase (WP_011354994). *Burkholderia lata*	59	KT944266
Unclassified						
P2N10	>13,508	66.6	*Alcagenaceae* (*â-Proteobacteria*)	Poly-R-hydroxyalkonic acid synthease PhaE (EGV16635), *Thiocapsa marina* 5811	27	KT944275
				Poly-beta-hydroxybutyrate polymerase PhaC (WP_002708071), *Thiothrix nivea*	43	KT944275
16 (PhaC2₁₆)	33,900	66.1	*Methylibium* (*â-Proteobacteria*)	Easterase (WP_043292995), *Burkholderia vietnamiensis*	53	KT944263
50 (PhaC2₅₀)	22,965	67.0	*Hydrocarboniphaga* (*ã-Proteobacteria*)	Easterase (WP_043292995), *Burkholderia vietnamiensis*	53	KT944271

The PhaC proteins were classified based on their amino acid sequences. Possible origin of cloned metagenomic DNA was predicted by PhyloPythia.

Table 4. Biosynthesis of SCL- and/or -MCL PHAs by 11AW metagenomic clones in *Pseudomonas putida*.

P. putida	Carbon sources	PHA content (%)	PHA composition (mol%)						
			3HB	3HV	3HHx	3HP	3HO	3HN	3HD
Class II									
PpUW1	Octanoate	30.0			0.7		99.3		
	Nonanoic acid	45.0				19.1		80.9	
	Gluconic acid	1.7					21.1		76.8
PpUW2 (clone 1)	Octanoate	32.7			4.5		95.5		
	Nonanoic acid	23.4				42.0		58.1	
	Gluconic acid	8.2					20.4		79.6
PpUW2 (clone 16)	Octanoate	25.1	17.3		39.3		43.2		
	Nonanoic acid	46.5	3.4	40.9		55.4			
	Gluconic acid	4.5			47.6		52.4		
PpUW2 (pJC123)	Octanoate	17.5	13.9		30.3		51.3		
PpUW2 (pJC157)	Octanoate	32.5	14.5		33.9		49.3		
PpUW2 (clone 20)	Octanoate	17.6			17.3		82.7		
	Nonanoic acid	45.0		7.5		50.2		42.3	
	Gluconic acid	3.9			9.9		37.4		52.7
Class I									
PpUW2 (clone 14)	Octanoate	41.8	98.9		1.1				
	Nonanoic acid	59.4	6.8	93.2					
	Gluconic acid	81.2	97.9		0.8		1.3		
PpUW2 (clone 18)	Octanoate	6.4	97.9		0.5		2.4		
	Nonanoic acid	17.3	32.8	67.2					
	Gluconic acid	30.7	100.0						
PpUW2 (clone 19)	Octanoate	30.9	98.8		0.5		0.7		
	Nonanoic acid	38.1	20.9	79.2					
	Gluconic acid	12.3	100.0						
PpUW2 (PhaC25)	Octanoate	15.6	95.0		2.9		2.1		
	Nonanoic acid	43.1	25.2	74.9					
	Gluconic acid	4.1	100.0						

P. putida PpUW1 (PHA[+]) and PpUW2 (PHA[-]) carrying PHA[+] cosmids of 11AW metagenomic clones were grown at 30°C for 48 h in 0.1N M63 media supplemented with 0.5% octanoate, 0.5% nonanoic acid or 1% gluconic acid. PHA content was represented as relative to cell dry weight. Values were the averages of triplicate cultures. The standard deviations were <10% of the values. Abbreviations: 3HB, 3-hydroxybutyrate (C4); 3HHx, 3-hydroxyhexanoate (C6); 3HO, 3-hydroxyoctanoate (C8); 3HD, 3-hydroxydecanoate (C10).

Supplementary figures

Supplementary 1. The *pha* gene locus in 11AW metagenomic DNA clones encoding PHA synthases (PhaC) were classified based on the PhaC protein sequences.

Supplementary 2. Transposon Tn4652 in 11AW PHA+ clones 2 and 15. Insertion of the Tn4652 into the *orf17* of clone 1 resulted in clone 2. Clone 15 was derived from clone 1 by the transposon insertion into the *orf19*.

Supplementary 3. Multiple sequence alignment of PHA synthases of 11AW metagenomic DNA clones. Alignment was performed with MUSCLE. The conserved amino acid residues affecting substrate specificity in Class I and II PhaC are highlighted in purple and yellow respectively. The positions of amino acids essential for PhaC activity are highlighted in green. The position of amino acid residues are marked based on PhaC1$_{16}$ sequence. Names of Class I PhaC are in regular font, Class II PhaC in bold, and unclassified PhaC underlined. PhaC proteins with partial sequences are marked with a star.

Supplementary 4. R-specific enoyl-CoA hydratases PhaJ. (A) Phylogenetic tree was constructed with MEGA6. The bar represents substitution of amino acid residue. (B) Conserved amino acid Asp^{39}, His^{44} and Ser^{7} required for activity are marked.

Class I

Class II

Unclassifed

Supplementary 1.

Supplementary 2.

```
PhaC2₁₆     -----msslpvaers-----------------------------------
PhaC2₅₀     ----Mistkpvaers-----------------------------------
PhaC_P2N10* --------------------------------------------------
PhaC1₁₆     MAktkgsrrAriPsP----psrS-----------EmLrlAL---------
PhaC1₅₀     ----MaKtkApprwarlpspppsrs---------ElLrlALv--------
PhaC5       MAkreatknvAatev------------------vggGedVG---------
PhaC20      --mtatRrskArPde-prakkrv-----------rkVaaAVGapkpeagEArgfae----
PhaC10*     --------------------------------------------------
PhaC2₁      --mrektarefsPtPaafinaqS-----------aitGlrgr--------
PhaC7*      --------------------------------------------------
PhaC_P11N2  ----MqyaghAlfqs------fn-----------QlLaaqLqQ-------
PhaC25      ----MmKqqAtgada---fapfq-----------QALsegwnKAlesfqqSatqgasafn
PhaC_P1N3   ----MrQeqqAaada--tfapfq-----------QALtqgwe--------
PhaC14      ----MkskpAAsarPtkqvtDsA-----------QALGhAwG-------
PhaC8       ----MkskpAApsrP-dqvaEsA-----------QALGhAwG-------
PhaC23      ----MkskpAApPrP-dqvaEsA-----------QALGhAwG-------
PhaC21      ----MkskpAApPrP-dqvaEsA-----------QsLGhAwG-------
PhaC4       ----MpKtiASlPas-------A-----------QAyGaAVG-------
PhaC_P4N10  ----MpKtiASlPas-------A-----------QAyGaAVG-------
PhaC19      ----MsKnntklPhs-----EqA-----------dAfGhAVG------
PhaC6       ----MsKnntklPhs-----EqA-----------dAfGhAVG------
PhaC3       ----MsKnntklPhs-----EqA-----------dAfGhAVG------
PhaC51      ----MmtgdA------------------------dAfGkAMG------
PhaC27      -----mtgdA------------------------dAfGkAMG------
PhaC18      -----mtgdA------------------------dAfGkAMG------
PhaC_P8N7   MAdpdtKapeApP1P---slEem-----------QhwtwvMGRAqqmmlE----------
PhaC_P2N8   ----MqtenSApidPgavleElAqvaqqiqggqmrdALdggwqQpvvdg-----------

PhaC2₁₆     -----nLeKlL--------LAvP---------------------------ayW---
PhaC2₅₀     -----nLeKAL--------LAfP---------------------------pyW---
PhaC_P2N10* -----------------MiLP---------glAaAaL------QAE---------rer
PhaC1₁₆     -----vilgrLrGn-----pgLP-------------------------IRhiagfakef
PhaC1₅₀     -----ifgRlrrn------pALP-------------------------IRhiaafakel
PhaC5       -------vlsprSl-------Ieaa----------tsAMNpL--------tmaRESarLygew
PhaC20      -----rvvdAIpG-------pnPfvgfSvFdvlssaeQLvagalqQpglVlrnqaafaGd
PhaC10*     -----------------------------------------------------
PhaC2₁      -----dLFsTLrs-------------------------vaAhglr------npvhtarhalklgGq
PhaC7*      -----------------------------------------------------
PhaC_P11N2  -----ssveltnG------LSht-----gtaasasvmaL------QeEwqarhmQLWqGm
PhaC25      vtggtpLWqipqt------aSMP-elpkisidpekLqsi------QqQYVaEATELWrqg
PhaC_P1N3   -----kaleAfqslgKApqadMPwdmprfsfsasrLqEL------QrhYIeEATELWrqd
PhaC14      -----eMWKSavG------LSLP---------tdAMSEL------QtrYVKQATELWNqa
PhaC8       -----hMWKSVAG------MSLP---------atAMTEL------QSnYLKQATELWNqt
PhaC23      -----dMWKSVAG------MSLP---------atAMTEL------QSnYLKQATELWNqt
PhaC21      -----dMWKSVAG------MSLP---------atAMTEL------QSnYLKQATELWNqt
PhaC4       -----diWKSMSe------LNLP---------lPALSEL------QSnYVKQATEIWNGa
PhaC_P4N10  -----diWKSMSe------LNLP---------lPALSEL------QSnYVKQATEIWNGa
PhaC19      -----diWKSMgG------LSLP---------lPALSEL------QtEYLKDATaLWNGa
PhaC6       -----diWKSMgG------LSLP---------lPALSEL------QtEYLKDvTaLWNGa
PhaC3       -----diWKSMgG------LSLP---------lPALSEL------QtEYLKDATaLWNGa
PhaC51      -----rMlqAaAG------LTLP---------ptALTDL------QAEYLREATEMWNGa
PhaC27      -----rMlqAaAG------LTLP---------psALTDL------QAEYLREATEMWNGa
PhaC18      -----rMlqAaAG------LTLP---------psALTDL------QAEYLREATEMWNGa
PhaC_P8N7   -----hvaRAMgetaEAkapALP---wlTlFgdpakLAQq------QAElwtEglaIWqra
PhaC_P2N8   -----pvlnSMAG---gllqALavfnqSlFadparLlQa------QAKlwdgyaQLWqrg

PhaC2₁₆     ---------------------piEW-------------------------------ar
PhaC2₅₀     ---------------------piEW-------------------------------tk
PhaC_P2N10* LDaLlgasaea--------------vsWpdrsAAcLpv--------------------eg
PhaC1₁₆     LniLrgksdv----tpaagDkRFAdaaWSgNrlSgalvQvYiaasqeayaLvkeLEl-eg
PhaC1₅₀     LrvLrnksev----ApsagDkRFtdaaWSdrrwSSalvQvYLaasqeayaLvneLEl-Dg
PhaC5       LKiLvgrser----dvPvkDwRFAdptWhehPAykrTAQgYLafcdaidRviDgnp--Dw
PhaC20      LaRIlsgrstl---epeegDRRFqdptWkeNPfyraglQtYLawrkgvhRLvegagl-Da
PhaC10*     ---------------------------------------------------------
PhaC2₁      LgRvllgetlh---ptnpqDnRFAdpaWSlNPfyrrSlQaYLswqkevkhwiDdsnm-tp
PhaC7*      ---------------------------------------------------------
PhaC_P11N2  LQsnqaatPva--qslpgDRRFshpaWAesPiydylrQaYLINAdyLqRLADtap--eg
PhaC25      fa-----------AkPegDRRFASdaWgSNPlSAfSAavYLLNgRTLLnMADAIDa-De
PhaC_P1N3   ia-----------ArthgDRRFAAeaWgSNPvAAfSAavYLLNtRTLLgMAeAVEG-Da
PhaC14      LaRLqpDGakETAPAaPitDkRFAApEWSANPATAyTAQmYLLNARTLMQLAesMQG-Da
PhaC8       LsRLqpEGatETAaAtPitDkRFAApEWSANPASAyTAQmYLLNARTLMQLAesMQG-Da
PhaC23      LsRLqpEGatETAaAtPitDkRFAApEWSANPASAyTAQmYLLNARTLMQLAesMQG-Da
PhaC21      LsRLqpEGatETAPAtPitDkRFAApEWSAsPASAyTAQmYLLNARTLMQLAesMQG-Da
PhaC4       vERLqaDGets---kkPlpDRRFAAnEWAANPvSALTAQiYLLNARTLLQMADAIEG-De
PhaC_P4N10  vERLqaDGets---kkPlpDRRFAAnEWAANPvSALTAQiYLLNARTLLQMADAIEG-De
PhaC19      vERLktDesqt---AvaprDRRFAAdEWASNPAAAMaAQtYLLNARTLMKMADAVEG-Dt
PhaC6       vERLktDeskt---AvaprDRRFAAdEWASNPAAAMaAQtYLLNARTLMKMADAVEG-Dt
PhaC3       vERLktDeskt---AvaprDRRFAAdEWASNPAAAMaAQtYLLNARTLMKMADAVEG-Dt
PhaC51      vERfsgkaPeQ---AqkigDRRFAAnDWAANPATALaAQaYLLNsRTLMKMADAIQG-Da
PhaC27      vERfsgkaPeQ---ShkigDRRFAAnDWAANPATALaAQaYLLNsRTLMKMADAIQG-Da
PhaC18      vERfsgkaPeQ---ShkigDRRFAAnDWAANPATALaAQaYLLNsRTLMKMADAIQG-Da
```

```
PhaC_P8N7      Lg--vdsGPrtelvekadrDRRFAApQWreNPlfdMirQsYLLvserLLgsvDsIDGLDp
PhaC_P2N8      LgaLtgqvPpaagPptaptDRRFAqDWql-PAfdLirQwYLLtsqwLLEqvtgagdLDp

                         120
PhaC2₁₆        qiRkRgtklprdWlryVraagaItf-PD----------------------Rpd--wVt
PhaC2₅₀        qiRqRgmrlpldWVsyVreakaItf-PD----------------------KpG--wVt
PhaC_P2N10*    KaR-------------------------------------------------------
PhaC1₁₆        rdaSaaRFlstmlVeALaPSNnpltNPvvlKrvrETRGkSllrGfQHfLsDrldnaGmIS
PhaC1₅₀        rdaSsaRFlatmlVeAMaPSNnpltNPvvmqrvrETKGrSllrGLQHfIsDrldnaGmIS
PhaC5          rkReRaRFltgiltsAMaPtNtLigNPaAlKKAyETgGkSlvrGvQnLIDDVRhnkGmpS
PhaC20         KgteRaRFvaslmteALaPtNFflgNPaAiKrALETgGgSllrGLanLfDDVatngGmpS
PhaC10*        -----------------------------------------------------------
PhaC2₁         ddRARahFAfallnDAMaPSNsL-LNPlAiKeifnsgGnSlvrGisHLvDDllhndGlpr
PhaC7*         -----------------------------------------------------------
PhaC_P11N2     qiRnRMqFlthQaIDAMaPSNFaAtNPEfiKtALaTeGESIrlGiQnLLgDlEkG--rIS
PhaC25         KTkARLRFAVEQWmaAsSPSNsLAfNaEAQKKAIdTqGESIAkGiQnLLhDVKQG--HlS
PhaC_P1N3      KTRARLRFAVEQWVaAsSPSNYLAfNaEAQKKALEsgGESIAkGiQnLLkDVQQG--HlS
PhaC14         KTkARVRFAVQQWVDAMSPaNFMALNPEAQKKALdTKGESVSQGiQqLLNDlRkG--aVS
PhaC8          KTkARVRFAVQQWVDAMSPaNFMALNPEAQKKALdTKGESVAQGLQqLfNDlRkG--sVS
PhaC23         KTkARVRFAVQQWVDAMSPaNFMALNPEAQKKALdTKGESVAQGLQqLfNDlRkG--sVS
PhaC21         KTkARVRFAVQQWVDAMSPaNFMALNPEAQKKALdTKGESVAQGLQqLfNDlRkG--sVS
PhaC4          KTRqRIRFAVQQWVeAaSPSNYLALNPEAQrKALETKGESItQGLlHLwqDVQQG--HVS
PhaC_P4N10     KTRqRIRFAeQQWgeAarrSNYvALNPEAQrKALETKGESItQGLlHLwqDVQQG--HVS
PhaC19         KTRARIRFAVQQWVDAaSPSNYLALNPEAQrKALETRGESIAQGLRqLwDDIQQG--HVS
PhaC6          KTRARIRFAVQQWVDAaSPSNYLALNPEAQrKALETRGESIAQGLRqLwDDIQQG--HVS
PhaC3          KTRARIRFAVQQWVDAaSPSNYLALNPEAQrKALETRGESIAQGLRqLwDDIQQG--HVS
PhaC51         KTkARVRFAVQQWIDAaSPSNYLAfNPEAQrKALETKGESIAQGLaHLLgDIQQG--HVS
PhaC27         KTkARVRFAVQQWIDAaSPSNYLAfNPEAQrKALETKGESIAQGLaHLLgDIQQG--HVS
PhaC18         KTkARVRFAVQQWIDAaSPSNYLAfNPEAQrKALETKGESIAQGLaHLLgDIQQG--HVS
PhaC_P8N7      KqRekLRFmtrnfVDAMaPSNFaAtNPvvlerAMaTRGESllkGLEHmLDDVskG--qlt
PhaC_P2N8      aTRhkLaFytrQfaDALSPSNFvAtNPEvlratvEsgGqnllhGfRHvLEDmKrGgGrId

PhaC2₁₆        ------------------pntiayEsqtmlLrnfsdpvvsadtiPviidaPfaghPstIa
PhaC2₅₀        ------------------phtiaFEtqtmwLrdfsdadApadaiPvivdaPyaghPstIa
PhaC_P2N10*    ---------------------------LLcYpsLrepa-aRPLLlVyslvNr-ihvL
PhaC1₁₆        --mvDrtaFEVGRNVAaTpGdVVFrNEvleLLqYKPqsAeVyEiPLmvVPPqINK-YYvL
PhaC1₅₀        --mvDrtpFEVGRNVAcTpGdVVFrNEvcevLqYKPqTAeVysiPLvvVPPqINK-fYvL
PhaC5          --QvkrSdFkVGeNlAaTpGAVVlrtErleLLqYqPtTAKVyaiPtLiIPPpIgK-hYfi
PhaC20         --QvDkSaFrVGeNlAsTpGAVVFrNEvleviqYaPagAqVygRPLLlVPPqINK-YYvi
PhaC10*        -----------------------------------------------------------
PhaC2₁         --QvtkqaFEVGktVATTtGsVVFrNELleLiqYKPMseKqysKPLLvVPPqINK-YYIf
PhaC7*         -----------------------------------------------------------
PhaC_P11N2     --mTDEaaFEiGRNlAvTsGsVVFENELmQLiqYaPLTpKVaaRPLviVPPCINK-fYIL
PhaC25         --mTDESaFEVGRNVATTEGAVVFENELFQLLEYKPLTpKVyERPfLlVPPCINK-YYIL
PhaC_P1N3      --mTDESaFEVGRNVATTEGAVVFENELFQLLEYKPLTAKVyERPLLlVPPCINK-fYIL
PhaC14         --QTDEtVFEVGRNVATsEGAVVFENELFQLiEYKPLTgKIHtRPMLFVPPCINK-YYIL
PhaC8          --QTDESVFEVGRNVATsEGAVVFENELFQLiEYKPLTAKVHtRPMLFVPPCINK-fYIL
PhaC23         --QTDESVFEVGRNVATsEGAVVFENELFQLiEYKPLTAKVHtRPMLFVPPCINK-fYIL
PhaC21         --QTDESVFEVGRNVATsvGAVVFENELFQLiEYKPLTAKVHtRPMLFVPPCINK-fYIL
PhaC4          --QTDESVFEVGRNVATTEGAVVFENdLFQLiEYKPLTdKVHEtPfLFVPPCINK-YYIL
PhaC_P4N10     --QTDESVFEVGRNVATTEGAVVFENdLFQLiEYKPLTdKVHEtPfLFVPPCINK-YYIL
PhaC19         --QTDESVFEVGRNVATsEGAVVFENELFQLLEYKPLTAqVHERPMLFIPPCINK-YYIL
PhaC6          --QTDESVFEVGRNVATsEGAVVFENELFQLLEYKPLTAqVHERPMLFIPPCINK-YYIL
PhaC3          --QTDESVFEVGRNVATsEGAVVFENELFQLLEYKPLTAqVHERPMLFIPPCINK-YYIL
PhaC51         --QTDESVFEVGRNVATTEGAVVFENELFQLLEYKPLTAqVHERPMLFVPPCINK-YYIL
PhaC27         --QTDESVFEVGRNVATTEGAVVFENELFQLLEYKPLTAqVHERPMLFVPPCINK-YYIL
PhaC18         --QTDESVFEVGRNVATTEGAVVFENELFQLLEYKPLTAqVHERPMLFVPPCINK-YYIL
PhaC_P8N7      --hTDpeaFEVGRNiAvTpGkVVkrtpLYeLiqYsPtTetVygtPLvifPPwINr-fYIL
PhaC_P2N8      prmTDpnaFEVGaNiAaTpGkVVFqNgLmQLiqYtPvTdqVyrRPLLvVPPwINK-fYvm

                     229
PhaC2₁₆        DyadgqSLvqtlkagGCkRVyVtdWksatheMrnfgiDkYlsDld----QaIdAVgGRa-
PhaC2₅₀        DyadgqSLvqtlkagGCkRlyVtdWksathaMrnfgiDkYlaD--LR--daVdhVgGRa-
PhaC_P2N10*    DLarghSfIerlqglG-HpVyVldWgrPDaSdAglslaDYVcgylstAVRLVrrhgcaR-
PhaC1₁₆        DLsPgNSfvnYaVKQG-fqVFmVSWRNPgpeLAhlslDhYlr-GleeAhnaVmeITGaK-
PhaC1₅₀        DLsPgNSfvnYaVKQG-fqVFmVSWRNPgpeLAylslqhYlQ-aledAhRaVmeITGad-
PhaC5          DLsPgrSfveYaVakG-fqVFcsVSWRNPqveqgdwgiDDYI-gsmLeAVdaVckITGsK-
PhaC20         DLsPgkSLveYaVasG-iqfFaVSWRNPtvaerdwgletYll-alseAIdaVceITGsE-
PhaC10*        -------------------FmVSWRNPtkaqrewglstYIE-alkeAVdVVsAITGsK-
PhaC2₁         DLsPhNSfvqfalKnG-lqtFmISWRNPDvrhrewglssYVE-aveeAmnVcrAITGaR-
PhaC7*         -------------------VFVVSWRNPDvrhrewglstYVa-aleeAlnVtrAITGaR-
PhaC_P11N2     DLQPENSLvRYaVEQG-ntVFmVSWRNvqadLghlTWDDYIEqGaLKAIRVaQqIcrvp-
PhaC25         DLQPENSLIRYanEQG-HRVFVVSWRNPDeSLAnaTWDDYIEnaaIKAIhtVQdISGsK-
PhaC_P1N3      DLQPENSLIRYaVEQG-HRtFVVSWRNPDaSMsqyTWDDYIEDaaIKAIQVtreITGQdk
PhaC14         DLQPENSLIRYTVEQG-HRVFVmSWRNPDeSLAayTWDDYIEkGpIKAIeVVQqISGQK-
PhaC8          DLQPENSLIRYTVEQG-HRVFVmSWRNPDeSLAhfTWDDYIEkGpLKAIeVVQqIgGQK-
PhaC23         DLQPENSLIRYTVEQG-HRVFVmSWRNPDeSLAhfTWDDYIEkGpLKAIeVVQqIgGQK-
PhaC21         DLQPENSLIRYTVEQG-HRVFVmSWRNPDeSLAhfTWDDYIEkGpLKAIeMVQqIgGQK-
PhaC4          DLQPENSLIRwTVEQG-HRVFVmSWRNPDdSVAakTWnDYIEDGtIRAIeVVreITGaE-
PhaC_P4N10     DLQPENSLIRwTVEQG-HRlFemSWRNPDdSVAakTWnDYIEDGtIRAIeVVreITGaE-
PhaC19         DLQPDNSLIRYTVaQG-HRtFVVSWRNPDpSIAakTWDDYIDDGaIRAIReVQAISGQa-
```

```
PhaC6         DLQPDNSLIRYTVaQG-HRtFVVSWRNPDpSIAakTWDDYIDDGaIRAIReVQAISGQa-
PhaC3         DLQPDNSLIRYTVaQG-HRtFVVSWRNPDpSIAakTWDDYIDDGaIRAIReVQAISGQa-
PhaC51        DLQPDNSLIRwTVgQG-HRtFVVSWRNPDaSIAdkTWDDYIEDGaIRAIgeVQSISGQK-
PhaC27        DLQPDNSLIRwTVgQG-HRtFVVSWRNPDaSIAdkTWDDYIEDGaIRAIReVQSISGQK-
PhaC18        DLQPDNSLIRwTVgQG-HRtFVVSWRNPDaSIAdkTWDDYIEDGaIRAIReVQSISGQK-
PhaC_P8N7     DLnPKkSfIRwaVEQG-ltVFVVSWksaDeSLAgtTlDDYVlgGqVdAIdtIrellGvE-
PhaC_P2N8     DLQPKNSLvgwwlQQG-ytVFmISWiNPgpqLAeksfeDYmiEGpLaAldaIEqacGER-
```

                       287   318
                   Lipase box

```
PhaC2₁₆       -----hlvGlCqGGwLs----AMhAARfp-gkAvSLvLagapiDthAgh-Gpvtkmaqrt
PhaC2₅₀       -----hlvGlCqGGwLst----MLAARyp-dkvvSLvLagapiDthAgD-Gpvtklskrt
PhaC_P2N10*   ---tIdlLGiCqGGs-LAlcLAsLepRal----nrLvtLvIpvDFh--tpG--DVlaDlA
PhaC1₁₆       ---KVsfkGlCaGGltsAiALgryAdqGkLdtvntLTLnvTLLDiSgmEktnmgyFLtpe
PhaC1₅₀       ---KVNfkGlCaGGltLAiALgryAdqGkLdtvntLTLnvTLLDiAgmEktnmgyFLtpS
PhaC5         ---KVNisGFCaGGimttlmLsymAAvrD-krvnaaafgvmLLDFd--teapigalhsrk
PhaC20        ---RVNTLaaCaGGitLmllLsyLAAeGD-rrvnSvTLLvTLLDte--aeGvLgLFasEe
PhaC10*       ---dINmLGaCsGGitctalLghyAAlGE-kkvnaLTLLvsvLDtT--ldtqvaLFVDEq
PhaC2₁        ---EVNlMGaCaGGltIAalqghLqAkrQLrrvSSaTyLvsLLDsq--mdspatLFaDEq
PhaC7*        ---EVNlMGaCaGGltIAalqghLqAkrQLrriSSasyLvsLLDsq--idspatLFaDEq
PhaC_P11N2    ---QVNaLGFCVGGTILssALAVarlRGE-dPvASLTLLTTLLDFS--DsGeigyFVDEA
PhaC25        ---QVNvLGFCVGGTILsTALAVLAARGE-kPASSvTLLTTfLLDFS--DTGiLDIFVDEp
PhaC_P1N3     -kggQIdTLFCVGGTILsTALAVLAARGE-QPASSvTLLTTLLDFe--DTGiLDIFIDEg
PhaC14        ---tfNmLGFCVGGTLIATALAVLAARGQ-QPAhSLTLLTTLLDFk--hTGvLDVFVDEp
PhaC8         ---QfNmLGFCVGGTLIATALAVLAARGE-QPAhSLTLLTTLLDFk--hTGvLDVFVDEp
PhaC23        ---QfNmLGFCVGGTLIATALAVLAARGE-QPAhSLTLLTTLLDFk--hTGvLDVFVDEp
PhaC21        ---EfNmLGFCVGGTLIsTALAVLAARGE-QPAhSLTLLTTLLDFk--hTGvLDVFVDEp
PhaC4         ---RlNTLGFCVGGTILATALAVLAARGD-QPAASvTLLTaLLDFS--ETGvLDLFIDEp
PhaC_P4N10    ---RlNTLGFCVGGTILATALAVLAARGD-QPAASvTLLTaLLDFS--ETGvLDLFIDEp
PhaC19        ---tIdTLGFCVGGTILtTALAVLAARGE-QPAASLTLLTTLLDFA--DTGvLDIFIDEA
PhaC6         ---tIdTLGFCVGGTILtTALAVLAARGE-QPAASLTLLTTLLDFA--DTGvLDIFIDEA
PhaC3         ---tIdTLGFCVGGTILtTALAVLAARGE-QPAASLTLLTTLLDFg--DTGvLDIFIDEA
PhaC51        ---QIdvLGFCVGGTILATALAVLAARGE-QPAASLTLLTTLLDFS--NTGvLDIFIDEA
PhaC27        ---QIdvLGFCVGGTILATALAVLAARGE-QPAASLTLLTTLLDFS--NTGvLDIFIDEA
PhaC18        ---QIdvLGFCVGGTILATALAVLAARGE-QPAASLTLLTTLLDFS--NTGvLDIFIDEA
PhaC_P8N7     ---sVhaIGYCVaGTtLAatLALLeARGEadkvASaTffTaqvDFS--EaGdLnLFVade
PhaC_P2N8     ---EVNavGYCIGGTLLlstLAymtAvnD-ErikSaTtfasLLDFS--DpGdLgVFIDEA
```

```
PhaC2₁₆       pfnfyrslVa----mGgGrmRGrfmlqgwknmhPdehyWgkyidlfdnitnPqyveRARH
PhaC2₅₀       plsfyrsmVa----mGgGrmRGrfmlqgwknmhPdehyWgkyidlfdnitnPqyvsRARH
PhaC_P2N10*   rVldlpalLapd-----GnLpGavLAglFgmLRPlrgVgaarapwldgGaEPekleRlRr
PhaC1₁₆       glEtsmkrskKE-----GvLyGhEmAkmFawLRPNDLVWNYwVnNYLmGQKPaaFD----
PhaC1₅₀       glEksmkrskQE-----GvLyGhEmAkmFawLRPNDLVWNYwVnNYLmGKKPaaFD----
PhaC5         lIgVarsrsaKk-----GIhpasaLAqvFawmRPNDLVWNYwVnNYLtGQdPPsFD----
PhaC20        aVaLaklnswtk-----GvLpGEELgrvFawLRPNDLVWNYwVnNYLlGnaPPaFD----
PhaC10*       tlEaakrhsyQa-----GvLEGrDmAkvFawmRPNDLi--------------------
PhaC2₁        tlEaakrrsyQk-----GvLdGrDmAkvFawmRPNDLiWsYfVnNYLlGKEPPaFD----
PhaC7*        tlEaakrrsyQQ-----GvLdGrDmAkvFawmRPNDLi--------------------
PhaC_P11N2    sVaaREagIG----QG-GLLhGrELATvFSaLRaNDLiWqYVVGNYLKGgKPPaFD----
PhaC25        mVayREmqLG----kG-GLLpGmDLAsTFSFLRPNDLVWNYVVGNYLKGEtPPPFD----
PhaC_P1N3     fVrfREmqMn----aG-GLLpGsELAsTFSFLRPNDLVWNYVVGNYLKGEtPPPFD----
PhaC14        sVQLREvTIGEQAPnGPGLLKGkELATTFSFLRPNDLVWNYVVGNYLKGEaPPPFD----
PhaC8         aVQMREmTLGEQAPnGPaLLKGkELATTFSFLRPNDLVWNYVVGNYLKGEaPPPFD----
PhaC23        tVQMREmTLGEQAPnGPaLLKGkELATTFSFLRPNDLVWNYVVGNYLKGEaPPPFD----
PhaC21        sVQMREmTIGEQAPnGPaLLKGkELATTFSFLRPNDLVWNYVVGNYLKGEaPPPFD----
PhaC4         aVQMREaTIGpQsPnGcGLLKGQELATTFSFLRPNDLVWNYVVGNYLKGEaPPPFD----
PhaC_P4N10    aVQMREaTIGpQsPnGcGLLKGQELATTFSFLRPNDLVWNYVVGNYLKGEaPPPFD----
PhaC19        aVQLREmTLGEQAPQGPaLLKGkELATTFSFLRPNDLVWNYVVGNYLKGnKPPPFD----
PhaC6         aVQLREmTLGEQAPnGPaLLKGkELATTFSFLRPNDLVWNYVVGNYLKGnKPPPFD----
PhaC3         aVQLREmTLGEQAPQGPaLLKGQELATTFSFLRPNDLVWNYVVGNYLKGnKPPPFD----
PhaC51        qVQLREaTIGldAPQGPGLLKGQELATTFSFLRPNDLVWNYVVGNYLKGDKPPPFD----
PhaC27        qVQLREaTIGldAPQGPGLLKGQELATTFSFLRPNDLVWNYVVGNYLKGDKPPPFD----
PhaC18        qVQLREaTIGldAPQGPGLLKGQELATTFSFLRPNDLVWNYVVGNYLKGDKPPPFD----
PhaC_P8N7     tlQLvEqisaDk-----GyLdGrymAaTFnlLRgrDLiWsYVtnNYLmGEdyaPFD----
PhaC_P2N8     qIaalEgkMa-----GaGyLdsaDmATaFnlLRaNDLVWsffVnNYLlGKdPaPFD----
```

```
                       391                                  445
PhaC2₁₆       farWyehvvdLPGrMYleavkHlfklNrLar-GEfvalGERisLksIavPVYllAgadDd
PhaC2₅₀       fasWyehvvdLPGrLYleavkHlfklNrLar-GEfvalGERisLksItvPVYllAgadDd
PhaC_P2N10*   LvaWqgDypdqaGraWlefvtacYreNrLlk-GtLvldGhtVDLrRlrlPVlnvfaRaDH
PhaC1₁₆       vLYWNSDSTrLPaaLHhdfc-dmvakNELgngevyqlpdaqVDLsRvDvesfIvAgRtDH
PhaC1₅₀       vLfWNSDaTrLPaaLHrdfc-ellktNELgrgevyqVedaKVDLrKvDvdsfVvAgRtDH
PhaC5         iLaWsvDgTNLPGkLHgqfL-difeaNaLpkkGsinIlGkplDLesIkvetlVtggttDH
PhaC20        vLYWNnDtTrLParLHgefL-dlfLlNpfRnPdaLellGkpiDvanvscdaYvAgttDH
PhaC10*       ------------------------------------------------------------
PhaC2₁        iLYWNnDnTrLPaaLHgdlL-dffkhNpLshPGgLeVCGtpiDLqKvNvdsfsvAginDH
PhaC7*        ------------------------------------------------------------
PhaC_P11N2    LLYWNSDSTNLPGPfltWYLRnmYLeNnLRiPGKLamCGvKaDLGhvDmPsfVvAcREDH
PhaC25        LLYWNSDaTNLPGPfYtWYLRnTYheNkLakPnaLTVCGEKiDLGKIDiPaYIYgSREDH
PhaC_P1N3     LLYWNSDaTNLPGPMYCWYLRnTYheNkLavPGaLTVaGEqVDLGRIEAPVYlYgSREDH
PhaC14        LLYWNgDSTNLPGPMYCWYLRHTYLqNELRqPGKLTVCGEKiDLGaIDAPVYIYgSREDH
```

```
PhaC8      LLYWNgDSTNLPGPMYCWYLRHTYLtNELRqPGKLTVCGEKiDLGaIDAPVYIYgSREDH
PhaC23     LLYWNgDSTNLPGPMYCWYLRHTYLtNELRqPGKLTVCGEKiDLGaIDAPVYIYgSREDH
PhaC21     LLYWNgDSTNLPGPMYCWYLRHTYLtNELRqPGKLTVCGEKiDLGaIDAPVYIYgSREDH
PhaC4      LLYWNSDSTNLPGPflCWYLRnTYLeNkLKdPGavTVCGEKVDLGaIEAPVYlYsSREDH
PhaC_P4N10 LLYWNSDSTNLPGPflCWYLRnTYLeNkLKdPGavTVCGEKVDLGaIEAPVYlYsSREDH
PhaC19     LLYWNSDSTNLPGPMFCWYLRHTYLqNELRvPGKLvVCGEKVDLGKIkAPVfVYgSREDH
PhaC6      LLYWNSDSTNLPGPMFCWYLRHTYLqNELRvPGKLvVCGEKVDLGKIkAPVfVYgSREDH
PhaC3      LLYWNSDSTNLPGPMFCWYLRHTYLqNELRvPGKLvVCGEKVDLGKIkAPVfVYgSREDH
PhaC51     LLYWNgDSTNLPGPMFCWYLRHTYLsNELRqPGKLTVCGQKVDLGKIkAPVfVYASREDH
PhaC27     LLYWNgDSTNLPGPMFCWYLRHTYLnNELRqPGKLTVCGQKVDLGKIkAPVfVYASREDH
PhaC18     LLYWNgDSTNLPGPMFCWYLRHTYLnNELRqPGKLTVCGQKVDLGKIkAPVfVYASREDH
PhaC_P8N7  LLhWNSDtTNLPakwHraYLRdfYrdNkLvraGELvVdGtpiDihKvktPtYVqAgREDH
PhaC_P2N8  LLfWNaDaTrmPaaMHsfYLRnmYLhNrLRePGgiTlaGvpinLsKItvPcYfvAtvEDH
```

                                                    473   478
```
PhaC2₁₆     IttpdqvfnaenLiGTPKdqiRkeLapgGHIglfmgaktlKqq-----------------
PhaC2₅₀     IttpeqvfnaenLvGTPRdqiRkeLapgGHIglfmgaktlneq-----------------
PhaC_P2N10* lVPaaAsqalgRai---gsrryqeLaieGghvGV--------------------------
PhaC1₁₆     ItPWdAcYKSvnLL---gGKsqFVLvnSGHIqtlvcPPgKgKaSyqT--aDvLPeDphEW
PhaC1₅₀     ItPWdgcYKSvnLL---gGesqFVLvnSGHIqtlvcPPgKgKaSyqT--gEtLPeDphEW
PhaC5       ltPWkgcYrtTQLL---gGpstFVLsnaGHvAslvNPPgnpKatyWl--gpKpddDpEaW
PhaC20      ItPWktcYatTQLL---tGereFVLsSSGHIqsIvNPPgnpKarflT--NpsfPaDpDDW
PhaC10*     ------------------------------------------------------------
PhaC2₁      ItPWdAvYrSTlLL---gGerRFVLsnSGHvqsIlNPPsnpKanfve--stKLsgDpraW
PhaC7*      ------------------------------------------------------------
PhaC_P11N2  IVPWrAsYrgrRLL---gGKsRFVLGASGHIAGVINPPAKgKRnyWvnqgDagkasADEW
PhaC25      IVPiggAYaSTQLL---pGKkRFVmGASGHIAGVINPPAKKKRSHWiRddgKfPktqaEW
PhaC_P1N3   IVPiggAYaSTQLL---KGKkRFVmGASGHIAGVINPPAKKKRSHWiRadgKLPktqaEW
PhaC14      IVPWhAAYESTtiL---ntKrRFVLGASGHIAGVIPaaKnKRSHWv--NDKLPpiADEW
PhaC8       IVPWhAAYESTtiL---nsKrRFVLGASGHIAGVINPasKnKRSHWT--NDKLPasADDW
PhaC23      IVPWhAAYESTtiL---nsKrRFVLGASGHIAGVINPasKnKRSHWT--NDKLPasADDW
PhaC21      IVPWqAAYESTtiL---nsKrRFVLGASGHIAGVINPasKnKRSHWT--NDKLPpsADDW
PhaC4       IVPWtgAYaSTaLL---RGKkRFVLGASGHIAGVINPPAKKKRSHWT--NtttPkDpqrW
PhaC_P4N10  IVPWtgAYaSpaLL---RGKkRFVLGASGHIAGVINPPAKKKRSHWT--NtttPkDpqrW
PhaC19      IVPWqAAfESTKim---KGKvRFVLGASGHIAGVINPPAaKKRSHWv--dgKLgatAqQW
PhaC6       IVPWqAAfESTKim---KGKvRFVLGASGHIAGVINPPAaKKRSHWv--dgKLgatAqQW
PhaC3       IVPWqAAfESTKim---KGKvRFVLGASGHIAGVINPPAaKKRSHWv--dgKLgatAqQW
PhaC51      IVPWdAAfESTKiL---KGKvRFVLGASGHIAGVINPPAaKKRSHWa--ggKVdgDAqQW
PhaC27      IVPWdAAfESTKiL---KGKvRFVLGASGHIAGVINPPAaKKRSHWv--ggKVggDAqQW
PhaC18      IVPWdAAfESTKiL---KGKvRFVLGASGHIAGVINPPAaKKRSHWv--ggKVggDAqQW
PhaC_P8N7   IaPaasvwKiThyf---QGplRFVLagSGHIAGVvNPPeaQKyqyWT--NDgkaetlDQf
PhaC_P2N8   IaPWrsvYagaRLp---gGKtRFVLsgSGHIAGIINPPAahKyqHWT--cDsLPtspDEW
```

```
PhaC2₁₆     ----------WariSAWivgqAa-------------------------------------
PhaC2₅₀     ----------WariSAWivqqg--------------------------------------
PhaC_P2N10* -------------------fAGR------------hglaiVpPAisRfltAtsiKKpr
PhaC1₁₆     LehsKattGSWWemWikWaserhGpkkPA--PKApGs-aRfpSlgaAPGeYVrA-------
PhaC1₅₀     LeqsKptiGSWWdlWvkWasqrsGakrPA--PKApGs-aKfppIgaAPGdYVrt-------
PhaC5       LqeAtkhtGtWWevWvdWagkrsGRqklA--PKAlGN-atYKArEkAPGtYVmerA-----
PhaC20      aAgAtphtGSWWehWTdWLatrsGerkaA--PKSlGs-ERHqplaaAPGtYVhg-------
PhaC10*     ------------------------------------------------------------
PhaC2₁      yydAKQvdGSWWTqWlgWiQersGaLket--hmAlGN-QnYppmEaAPGtYVrvr------
PhaC7*      ------------------------------------------------------------
PhaC_P11N2  LeTAsEvsGSWWplWAeWLggfgGKkVtA--rrrlGs-adfsplEPAPGRYVKeKA-----
PhaC25      LAgAtEqPGSWWTDWAqWLKgHAGKqVPA--PKAYGdGKKYKAIEPAPGRYVKArA-----
PhaC_P1N3   LAgAtEhPGSWWTDWSnWLKtHAGKqIPA--PKSYGkGsaYKAIESAPGRYVKArA-----
PhaC14      fvSAtErPGSWWpDWSAWLKaqgGaMIaA--PKgYGN-KtfKAIEPAPGRYVKqKA-----
PhaC8       MASAiEkPGSWWSDWSAWLKtqgGpMVaA--PKgYGN-KKlKAIEPAPGRYVKqKA-----
PhaC23      MAgAiEkPGSWWSDWSAWLKtqgGpMVaA--PKgYGN-KmlKAIEPAPGRYVKqKA-----
PhaC21      MASAtEkPGSWWSDWSAWLKtqgGpMVaA--PKgYGN-KKlKAIEPAPGRYVKqKA-----
PhaC4       LAgAtEsPGSWWTDWAAWLaplsGKeIaA--PKAlGs-RKYKSIEPAPGRYVKqKA-----
PhaC_P4N10  LAgAtEsPGSWWTDWAAWLaplsGKeIaA--PKAlGs-RKYKSIEPAPGRYVKqKA-----
PhaC19      LdSAKEvPGSWWTDWSAWLKpHAGKqVPA--PKtYGN-KsHKvIEPAPGRYVKAKA-----
PhaC6       LdSAKEvPGSWWTDWSAWLKpHAGKqVPA--PKtYGN-KsHKvIEPAPGRYVKAKA-----
PhaC3       LdSAKEvPGSWWTDWSAWLKpHAGKqVPA--PKtYGN-KsHKvIEPAPGRYVKAKA-----
PhaC51      LeSAKDvPGSWWSDWAAWLKpHAGKMVPA--PKAYGd-RsHKAIEaAPGRYVKAKA-----
PhaC27      LeSAKDvPGSWWSDWAAWLKpHAGKMVPA--PKAYGd-RsHKAIEaAPGRYVKAKA-----
PhaC18      LeSAKDvPGSWWSDWAAWLKpHAGKMVPA--PKAYGd-RsHKAIEaAPGRYVKAKA-----
PhaC_P8N7   vAgAtEhkGSWWpDWieWigaqdGKrVPAkgarvpGk-gKlKAlEnAPGsYVKAr------
PhaC_P2N8   qAqAaahaGSWWpDWkfWLgrrAGgkVmA---rtpGs-ggltAIEdAPGsYVrmtshQE--
```

Supplementary 3

A.

0.1

B.

```
                                                                 39    44
R. eutropha PhaJ4a   -MrtiaSLeeLEglqGQEvavSDWIeVtQqqvnqFAdaTGDhQWIHIDVERAkKESPyGG
R. eutropha PhaJ4b   -MktyeNiadLQplVGevIGtsEWLaLDQARIntFAdaTGDhQWIHVDVERAkn-gPFGa
P. putida PhaJ4      --mphvpvteLsqyVGkELGhSEWLKIDQqRInlFAeaTGDfQfIHVDpEKAAK-tPFGG
PhaJ20               mteEaySvrtIpqfaGrELGvSDWLRIDQeRIdrFAACTdDrQWIHVDVERAeRESPFGt
PhaJ50               --mstlkMatLtekVGQELGttrWvaLDQSRIQaFAdCTGDqQWIHtDVERAkKESPFGG
PhaJ16               --msaltMqnIaakIGeELGvtgWttLDQARIQaFAdCTGDgQWIHtDVERArKESPFGG
PhaJ5                -MadaltLatIEsfVGrELGeSqWvRmDQqRIdaFAeCTGDrQWIHVDVERAAREgPFGG
A. caviae PhaJ       -------MsaqsleVGQkarlSk--RfgaAevaaFAAlseDfnplHlDpafAAt-taFer
P. putida PhaJ1      -MsQvtNtpyealeVGQkaeykk--sVEerdIQlFAAmsGDhnpvHlDaEfAAK-SmFre
P. aeruginosa PhaJ1  -MsQvqNipyaEleVGQkaeyts--sIaerdlQlFAAvsGDrnpvHlDaayAAt-tqFke

                            71
R. eutropha PhaJ4a   PIAHGfLTLSLLPkfmhNalhmPskI--GVNYGLnRVRFtAPVpVGSklRaRIKLLkver
R. eutropha PhaJ4b   PIAHGfLTLSLLPafthsayrirns-stGVNYGLDKVRFpAPVpVdSllRaqfKLMSyea
P. putida PhaJ4      tIAHGfLTLSLiPklieDIlvlPqgLkMvVNYGLDsVRFiqPVKVdSrVRlkVKLgevve
PhaJ20               tvAHGlLvLSLLPrfgfEVglvPpgVssalNYGfDgVRFvSPVKaGArVRdRVtLLeatd
PhaJ50               PvAHGMLTLSLLPmwlfDLpaaPddagailNYGfDKVRFlAPVKsGAkVRaRIKLLaatP
PhaJ16               PvAHGMLTLSLLPpwlfELpaaPldagailNYGfDKVRFlAPVKaGSrVRgRIKLLAaaP
PhaJ5                tIAHGfLTLSiigpaqlDVwiaPagIgtaVhYGLDKVRFlAPVpaGrnVRtRIKLaAvea
A. caviae PhaJ       PIvHGML-LSLfsgllgqqlpgkgsIyLG-----qslsFklPVfVGdeVtaeVevtAlre
P. putida PhaJ1      rIAHGMfsgaLisaavactlpgPgtIyLG-----qQmsFqkPVKIGdtltvRleiLeklP
P. aeruginosa PhaJ1  rIAHGMLsgaLisaaiatVlpgPgtIyLG-----qtlRFtrPVKLGddlkvelevLeklP

R. eutropha PhaJ4a   ldplpkspeLvgaqsTwevTVErEGsdrPvcVAEsItrryg--------
R. eutropha PhaJ4b   len-------GgaqFkvemmVERQGgSKPvcIAEsILrrfP---------
P. putida PhaJ4      Kkp-------GqwLlkaiaTlEIEGEeKPAyIAEsLsLcfv---------
PhaJ20               KgE-------GrlLvkarhTIEIEGESKPALVAEmLaMlItg-------
PhaJ50               KeK-------GrvLlTqeyTVEIEnEtKPALIAElLVMlIPkaa-----
PhaJ16               KdK-------GrlLlTqeyTVEIEnEtKPALIAElLVMlvPka------
PhaJ5                Kgs-------GrtLvTtenvVEIEGhdKPALIAtaLaMiMP--------
A. caviae PhaJ       dkp---iatLttrIFTqggalaVtGE----------VVkLP--------
P. putida PhaJ1      KfK----vRiatnVYnqndelvVaGEAeilaprkqqtVeLvspPnfvas
P. aeruginosa PhaJ1  KnR----vRmatrVFnqagkqvVDGEAeimapeEkLsVeLaelPpisig
```

Supplementary 4