Title:
Relic DNA is abundant in soil and obscures estimates of soil microbial diversity

Authors:
Paul Carini[1*], Patrick J. Marsden[2], Jonathan W. Leff[1,3], Emily E. Morgan[1], Michael S. Strickland[4], Noah Fierer[1,3*]

Affiliations:
[1]Cooperative Institute for Research in Environmental Sciences, University of Colorado, Boulder, CO 80309
[2]Department of Chemistry, Metropolitan State University of Denver, Denver, CO 80217
[3]Department of Ecology and Evolutionary Biology, University of Colorado, Boulder, CO 80309
[4]Department of Biological Sciences, Virginia Tech University, Blacksburg, VA 24061

*Corresponding Authors: Paul Carini (Paul.Carini@gmail.com) and Noah Fierer (Noah.Fierer@colorado.edu)

Abstract:
It is implicitly assumed that the microbial DNA recovered from soil originates from living cells. However, because relic DNA (DNA from dead cells) can persist in soil for weeks to years, it could impact DNA-based analyses of microbial diversity. We examined a wide range of soils and found that, on average, 40% of prokaryotic and fungal DNA was derived from the relic DNA pool. Relic DNA inflated the observed prokaryotic and fungal diversity by as much as 55%, and caused misestimation of taxon abundances, including taxa integral to key ecosystem processes. These findings imply that relic DNA can obscure treatment effects, spatiotemporal patterns, and relationships between taxa and environmental conditions. Moreover, relic DNA may represent a historical record of microbes formerly living in soil.

Main Text:
Microbes play critical roles in terrestrial biogeochemistry and the maintenance of soil fertility. Microbiologists, mycologists, biogeochemists, and ecologists now routinely use DNA-based approaches to determine the composition and diversity of soil microbial communities using molecular methods that include amplicon (marker gene) sequencing, quantitative polymerase chain reaction (qPCR), and shotgun metagenomics. These methods have advanced our understanding of terrestrial microbiology in a myriad of ways, by: *i)* revealing that thousands of unique microbial taxa can inhabit a single gram of soil (*1-3*); *ii)* uncovering novel soil microbial diversity (*4, 5*); and *iii)* identifying putative functions of uncultivated taxa (*6*). As DNA sequencing costs continue to plummet, the use of molecular methods to describe soil microbial diversity aimed at answering both basic and applied research questions will become even more commonplace.

Linking the activities of microbes to soil processes first necessitates distinguishing living cells (both metabolically active and dormant) from those that are dead. Most molecular investigations of soil microbial diversity make the implicit assumption that the total pool of DNA extracted from soil is derived exclusively from living cells contributing to, or potentially contributing to, biogeochemical transformations. However, total microbial DNA can originate from both living and dead cells. Previous work has shown that extracellular DNA and DNA from dead or partially degraded organisms can persist in soils for weeks to years (reviewed in (*7-9*)). The longevity and size of this extracellular DNA pool is controlled by a myriad of soil factors. For example, complex physical factors such as soil mineralogy, pH and ionic strength control the sorption of DNA to the soil matrix, as well as the molecular integrity of the DNA itself (*10, 11*). This sorbed DNA can be protected from removal by microbes that use it as a source of transformable genetic material or for nutrition (reviewed in (*12*)).

Given the potential for extracellular DNA to persist in soil, we hypothesized that DNA from dead microbes (which we term 'relic DNA') may obscure DNA-based estimates of the diversity and structure of soil microbial communities (*13, 14*). Such effects should be most apparent if relic DNA is abundant and if the taxa represented in the relic DNA pool are not reflective of the taxa present as living cells. It is important to distinguish living microbes from dead microbes because ecological definitions of community-level diversity and structure are meant to encompass organisms actually

alive at a site, not both the living and extinct organisms. Here, we used a propidium monoazide (PMA)-based approach (*15, 16*) to remove relic DNA from a broad range of soil types. We report the amount of microbial DNA derived from relic DNA pools and show how relic DNA affects the observed richness and composition of microbial communities. We also identify which soil characteristics are linked to greater relic DNA effects and discuss the implications of relic DNA on molecular analyses of microbial communities in soil and other environments.

We tested the effect of relic DNA on estimation of microbial diversity by analyzing 31 soils collected from a broad range of ecosystem types across the United States, selected to encompass a wide variety of edaphic characteristics (Supplementary Table S1). Subsamples of each soil were either treated with PMA (n=5) or left untreated (n=5). PMA is a DNA intercalating molecule that is generally excluded by cells with intact membranes, but binds extracellular DNA and DNA of cells with compromised cytoplasmic membranes (*15*). After exposure to light, intercalated PMA covalently binds and permanently modifies DNA, rendering it un-amplifiable by the polymerase chain reaction (PCR) (*15*). We used quantitative PCR (qPCR) to calculate the amount of relic DNA by subtracting the number of amplifiable prokaryotic 16S rRNA gene copies or fungal internal transcribed spacer 1 (ITS) amplicon copies in untreated samples (total DNA = DNA from living cells + relic DNA) from the number of gene copies in PMA-treated samples (DNA from living cells only). Microbial communities were characterized by high-throughput sequencing of amplified rRNA gene regions (16S rRNA for prokaryotes or the ITS region for fungi) from both PMA treated and untreated soils. We compared estimates of microbial richness, overall community composition, and taxon abundances after standardizing all libraries to equivalent sequencing depths (see materials and methods in Supplementary Materials).

Relic DNA represented a large fraction of microbial DNA in many soils. Across all 31 soils, $40.7 \pm 3.75\%$ (mean $\pm$ SE; n=155) of amplifiable prokaryotic 16S rRNA genes were derived from the relic DNA pool (Fig. 1A). Similar patterns were observed for fungi, where $40.5 \pm 4.12\%$ (mean $\pm$ SE; n=155) of fungal ITS amplicons originated from the relic DNA pool (Fig. 1B). There are a number of lines of evidence that suggest these estimates of the amounts of relic DNA found in soil are conservative. First, when we experimentally added naked DNA to soil, our approach completely removed the naked DNA from most soil samples, but only reduced (did not completely remove) naked DNA added to soils which were found to contain high levels of relic DNA (Fig. S1A). Second, because not all dead cells have compromised membranes (*17*), PMA may not infiltrate all dead cells. Conversely, it is unlikely our approach removes DNA from intact cells because: *i)* numerous studies have shown that the effects of PMA on intact microbial cells are minimal and PMA overwhelmingly targets extracellular DNA or DNA from dead cells (*15, 16, 18*); *ii)* our own tests confirmed PMA treatment did not lead to significant reductions in the amounts of DNA coming from live bacterial or fungal cells (Fig. S1B,C); and *iii)* if PMA inflated estimates of the amount of relic DNA by entering live cells, we would expect to detect relic DNA in all soils studied, which was not the case (Fig. 1A,B).

Removal of relic DNA significantly reduced estimates of microbial diversity.
Across all samples, 13.9 ± 1.20% (mean ± SE; n=155) of the total prokaryotic richness
(number of taxa) was found in the relic DNA pool (Fig.1C). In other words, nearly 14% of
the taxa were no longer living in soil and were exclusively recovered in the relic DNA
pool. In 24 of the soils tested, the prokaryotic richness was significantly lower, by as
much as 55%, after relic DNA was removed (two-tailed t test q value ≤ 0.05) (Fig. 1C).
The percent of prokaryotic taxa exclusively found in the relic DNA pool was positively
correlated with the proportional abundance of 16S rRNA genes present in the relic DNA
pool (Fig. S2A). That is, soils with more relic DNA tended to have lower richness once
relic DNA was removed. Similar results were observed when we analyzed fungal DNA.
The relic DNA pool contributed 12.4 ± 1.97% (mean ± SE; n=152) of the total fungal
richness across all soils (Fig. 1D). The relic DNA contribution to estimates of fungal
richness was significant in 14 soils (two-tailed t test q value ≤ 0.05), and removal of relic
DNA reduced estimates of fungal diversity by up to 52% (Fig. 1D).
126
The removal of relic DNA can substantially reduce estimates of soil microbial
diversity, indicating that the most commonly used molecular methods for assessing soil
microbial diversity lead to inflated richness estimates due to the detection of DNA from
dead cells. Although an average of ~40% of the total prokaryotic and fungal DNA was
derived from relic DNA pools (Fig. 1A,B), the effect of relic DNA removal on fungal
richness was variable across the soils examined (Fig. 1D) and there was no significant
correlation between the percent of fungal relic DNA and the percent of fungal taxa found
exclusively within the relic DNA pool (Fig. S2B). These between-sample differences in
the magnitude of the effects of relic DNA on estimates of fungal diversity may be a
product of differences in the temporal turnover of fungal communities at individual sites.
If turnover in the fungal community composition is minimal, removal of relic DNA should
have little effect on estimated fungal taxonomic richness as the relic DNA pool would
reflect the diversity found in the pool of DNA extracted from intact fungal cells. This
suggests that targeted analyses of the relic DNA pool could be used to identify taxa that
once lived in soil, but are no longer alive due to changes in soil conditions, or to
discriminate between endemic microbes and those microbes that have been
transported into soils that can not support their growth.
144
We found that estimates of microbial community composition were also
significantly influenced by the presence of relic DNA. Soil source was the strongest
predictor of community differences (PERMANOVA R^2=0.727, P ≤ 0.001 for prokaryotes;
R^2=0.646, P ≤ 0.001 for fungi). That is, we could discriminate between the distinct
microbial communities found in the different soils, regardless of whether relic DNA was
removed or not (Fig. S3A-D). However, the effect of relic DNA removal on community
composition was significant for both prokaryotic and fungal communities (PERMANOVA
R^2=0.004, P ≤ 0.001 for prokaryotes; R^2=0.002, P ≤ 0.001 for fungi). On an individual
soil basis, the composition of prokaryotic communities was significantly affected by the
removal of relic DNA in all 31 of the soils tested (PERMANOVA R^2=0.10-0.23, q value ≤
0.05) (Fig. 2A). In 21 of the 31 soils, removal of relic DNA also had a significant effect
on the composition of fungal communities (PERMANOVA R^2=0.10-0.22, q value ≤ 0.05)
(Fig. 2B). The effects of relic DNA on the composition of both prokaryotic and fungal

communities were positively correlated (Fig. 2C), highlighting that the magnitude of
these relic DNA effects on microbial community composition were similar for both
prokaryotic and fungal communities.
161
The relative abundances of numerous key microbial lineages changed after the
removal of relic DNA, but the taxa that changed, and the direction of observed shifts,
varied across soils. For example, in a New Hampshire grassland (soil 25),
Actinobacteria and α-Proteobacteria significantly increased in relative abundance after
relic DNA was removed, but Verrucomicrobia decreased (Fig. 3). In many cases, the
changes in estimated relative abundances after relic DNA removal were large, often
approaching or exceeding 25% (Fig. 3). The relative abundances of α-Proteobacteria
were consistently and significantly greater after relic DNA removal (Mann-Whitney U
two-tailed $P \leq 0.05$) (Fig. S4A), suggesting that the abundances of viable or dormant α-
Proteobacteria are underestimated in many soil studies. In contrast, agaricomycete
fungi were significantly less abundant after relic DNA was removed (Mann-Whitney U
two-tailed $P \leq 0.05$) (Fig. S4B), suggesting Agaricomycetes are less abundant than
commonly thought. Together, these results show that the effects of relic DNA removal
vary depending on the taxon in question, are not predictable *a priori,* and will vary
depending on the particular soil studied.
177
Relating the abundances of microbial taxa or protein-coding genes to soil
biogeochemical process rates has been challenging, hindering attempts to link microbial
communities to the ecosystem-level processes they can control (*19*). Because the
abundances of living microbial populations may be much higher or lower than is
apparent from estimates of relative abundances derived from total DNA analyses, our
results show that relic DNA likely obscures correlations between the abundances of
individual microbial taxa (or their functional genes) and key biogeochemical processes.
We found evidence of this when we compared the relative abundances of prokaryotes
integral to soil nitrification before and after relic DNA was removed. For instance,
ammonia oxidizing archaea classified as '*Ca.* Nitrosphaera' and nitrite oxidizing bacteria
classified as *Nitrospira spp.* changed by >25% in several soils after relic DNA was
removed (Fig. S5A,B). Similarly, the relative abundance of *Glomeraceae*, a family of
arbuscular mycorrhizal fungi, increased by >25% in 2 soils and decreased by >25% in 7
soils (Fig. S5C). Thus, by removing relic DNA prior to investigating relationships
between specific soil processes and DNA-based quantification of microbial abundances,
researchers may uncover more robust associations between microbes and key soil
processes.
195
Because relic DNA can result in the overestimation of soil microbial diversity (Fig.
1), change assessments of overall community composition (Fig. 2), and alter the
observed abundances of individual taxa (Fig. 3), we investigated which edaphic
characteristics were predictive of the presence of relic DNA. Consistent with previous
studies (reviewed in (*11*)), we show that edaphic characteristics influencing electrostatic
interactions between DNA and soil particles were significant predictors of the presence
of microbial relic DNA (Table 1). For example, soils with few exchangeable bases,
especially K^+ and Ca^{2+}, were likely to contain relic DNA from both prokaryotic and fungal

sources (Table 1, Fig S6). While soils with low pH, electrical conductivity and cation
exchange capacity were more likely to harbor relic DNA, this pattern was stronger for
prokaryotes than for fungi (Table 1). Moreover, pH predicted the change in both
prokaryotic and fungal community composition after relic DNA removal (Fig. S7). These
results highlight that, although the effects of relic DNA are variable across different soil
types, it is especially important to account for relic DNA in acidic soils, or in soils with
few exchangeable base cations (below *ca.* 40 meq 100 g^{-1}).

Our finding that relic DNA can lead to significant overestimation of soil microbial
diversity and reduce the ability to accurately quantify prokaryotic and fungal community
structure has several important implications. First, it suggests that the actual microbial
diversity in soil is lower than often reported. Second, relic DNA may obscure subtle
spatiotemporal patterns or treatment effects in microbial communities. For example,
shifts in soil microbial communities across seasons, or with plant species growing on a
site, are often difficult to detect from DNA-based analyses (*20, 21*). Similarly, long-term
soil transplant experiments designed to study effects of climate change on soil microbial
communities have showed little change in microbial community composition (*22*). Our
ability to detect such shifts in soil microbial communities should increase if the 'noise'
generated from non-living microbes is reduced by first removing relic DNA. Finally, the
extreme diversity of soil microbial communities presents multiple computational
problems for metagenomic assembly and analysis (*23*). Our data shows that a
significant portion of this diversity is coming from relic DNA pools, suggesting that
removal of relic DNA from samples prior to shotgun metagenomic analyses will result in
improved metagenomic assemblies and improve our ability to infer the genomic
attributes of undescribed soil microbes (*24*).

Relic DNA dynamics may have important ramifications for understanding
community composition and processes in other ecosystems besides soil. For example,
deep sea sediments also harbor large amounts of extracellular DNA (*25*), suggesting
the removal of relic DNA would also affect diversity estimates in the deep biosphere.
Specific analysis of the relic DNA pool in deep biosphere samples may have particular
utility as a 'fossil record' to distinguish extinct microbial taxa from living organisms and
more accurately reconstruct the subsurface paleome (*26, 27*). More generally, relic DNA
likely influences studies where DNA from dead organisms may be abundant or where
DNA is particularly resistant to degradation, including studies of the microbial diversity
found on aquatic particles, on mineral surfaces, in the human body, and in the built
environment.

References:

1. L. F. W. Roesch *et al.*, Pyrosequencing enumerates and contrasts soil microbial
diversity. *The ISME Journal.* **1**, 283–290 (2007).

2. J. Gans, M. Wolinsky, J. Dunbar, Computational improvements reveal great
bacterial diversity and high metal toxicity in soil. *Science.* **309**, 1387–1390 (2005).

3. N. Fierer *et al.*, Metagenomic and Small-Subunit rRNA Analyses Reveal the

Genetic Diversity of Bacteria, Archaea, Fungi, and Viruses in Soil. *Applied and Environmental Microbiology*. **73**, 7059–7066 (2007).

4. C. T. Brown *et al.*, Unusual biology across a group comprising more than 15% of domain Bacteria. *Nature*. **523**, 208–211 (2015).

5. M. D. M. Jones *et al.*, Discovery of novel intermediate forms redefines the fungal tree of life. *Nature*. **474**, 200–203 (2011).

6. R. Mackelprang *et al.*, Metagenomic analysis of a permafrost microbial community reveals a rapid response to thaw. *Nature*. **480**, 368–371 (2011).

7. K. M. Nielsen, P. J. Johnsen, D. Bensasson, D. Daffonchio, Release and persistence of extracellular DNA in the environment. *Environ. Biosafety Res.* **6**, 37–53 (2007).

8. G. Pietramellara *et al.*, Extracellular DNA in soil and sediment: fate and ecological relevance. *Biol Fertil Soils*. **45**, 219–235 (2008).

9. D. J. Levy-Booth *et al.*, Cycling of extracellular DNA in the soil environment. *Soil Biology and Biochemistry*. **39**, 2977–2991 (2007).

10. A. Ogram, G. S. Sayler, D. Gustin, R. J. Lewis, DNA adsorption to soils and sediments. *Environ. Sci. Technol.* **22**, 982–984 (1988).

11. K. Saeki, T. Kunito, Adsorptions of DNA molecules by soils and variable-charged soil constituents. *Current research* (2010).

12. K. M. Nielsen, L. Calamai, G. Pietramellara, in *Nucleic Acids and Proteins in Soil* (Springer Berlin Heidelberg, 2006), vol. 8 of *Soil Biology*, pp. 141–157.

13. A. Agnelli *et al.*, Distribution of microbial communities in a forest soil profile investigated by microbial biomass, soil respiration and DGGE of total and extracellular DNA. *Soil Biology and Biochemistry*. **36**, 859–868 (2004).

14. G. Dlott, J. E. Maul, J. Buyer, S. Yarwood, Microbial rRNA:rDNA gene ratios may be unexpectedly low due to extracellular DNA preservation in soils. *Journal of Microbiological Methods*. **115**, 112–120 (2015).

15. A. Nocker, C.-Y. Cheung, A. K. Camper, Comparison of propidium monoazide with ethidium monoazide for differentiation of live vs. dead bacteria by selective removal of DNA from dead cells. *Journal of Microbiological Methods*. **67**, 310–320 (2006).

16. A. Nocker, P. Sossa-Fernandez, M. D. Burr, A. K. Camper, Use of propidium monoazide for live/dead distinction in microbial ecology. *Applied and Environmental Microbiology*. **73**, 5111–5117 (2007).

283   17.   G. A. Cangelosi, J. S. Meschke, Dead or Alive: Molecular Assessment of
284   Microbial Viability. *Applied and Environmental Microbiology*. **80**, 5884–5891
285   (2014).

286   18.   A. Nocker, T. Richter-Heitmann, R. Montijn, F. Schuren, R. Kort, Discrimination
287   between live and dead cells in bacterial communities from environmental water
288   samples analyzed by 454 pyrosequencing. *Int. Microbiol.* **13**, 59–65 (2010).

289   19.   J. D. Rocca *et al.*, Relationships between protein-encoding gene abundance and
290   corresponding process are commonly assumed yet rarely observed. *The ISME*
291   *Journal*. **9**, 1693–1699 (2015).

292   20.   X. Zhang, A. Barberán, X. Zhu, G. Zhang, X. Han, Water Content Differences
293   Have Stronger Effects than Plant Functional Groups on Soil Bacteria in a Steppe
294   Ecosystem. *PLoS ONE*. **9**, e115798–12 (2014).

295   21.   K. M. Docherty *et al.*, Key Edaphic Properties Largely Explain Temporal and
296   Geographic Variation in Soil Microbial Communities across Four Biomes. *PLoS*
297   *ONE*. **10**, e0135352–23 (2015).

298   22.   B. Bond-Lamberty *et al.*, Soil Respiration and Bacterial Structure and Function
299   after 17 Years of a Reciprocal Soil Transplant Experiment. *PLoS ONE*. **11**,
300   e0150599–16 (2016).

301   23.   A. C. Howe *et al.*, Tackling soil diversity with the assembly of large, complex
302   metagenomes. *Proceedings of the National Academy of Sciences*. **111**, 4904–
303   4909 (2014).

304   24.   K. S. Ramirez et al., Biogeographic patterns in below-ground diversity in New
305   York City's Central Park are similar to those observed globally. *Proc. R. Soc. B.*
306   **281**, 20141988 (2014).
307

308   25.   A. Dell'Anno, R. Danovaro, Extracellular DNA plays a key role in deep-sea
309   ecosystem functioning. *Science*. **309**, 2179–2179 (2005).

310   26.   M. J. L. Coolen *et al.*, Evolution of the plankton paleome in the Black Sea from the
311   Deglacial to Anthropocene. *Proceedings of the National Academy of Sciences*.
312   **110**, 8609–8614 (2013).

313   27.   F. Inagaki, H. Okada, A. I. Tsapin, K. H. Nealson, Microbial survival: the paleome:
314   a sedimentary genetic record of past microbial communities. *Astrobiology*. **5**, 141–
315   153 (2005).

316   28.   W. D. Kemper, R. C. Rosenau, in *Methods of soil analysis.*, A. Klute, Ed.
317   (Madison, WI, ed. 2, 1986).

29.   J. W. Leff *et al.*, Consistent responses of soil microbial communities to elevated

nutrient inputs in grasslands across the globe. *Proc. Natl. Acad. Sci. U.S.A.* **112**,
10967–10972 (2015).

30. R. C. Edgar, UPARSE: highly accurate OTU sequences from microbial amplicon
reads. *Nat Meth*. **10**, 996–998 (2013).

31. R. C. Edgar, Search and clustering orders of magnitude faster than BLAST.
Bioinformatics. **26**, 2460–2461 (2010).

32. D. McDonald *et al.*, An improved Greengenes taxonomy with explicit ranks for
ecological and evolutionary analyses of bacteria and archaea. *The ISME Journal*.
6, 610–618 (2011).

33. K. Abarenkov *et al.*, The UNITE database for molecular identification of fungi-
recent updates and future perspectives. *New Phytol.* **186**, 281–285 (2010).

34. Q. Wang, G. M. Garrity, J. M. Tiedje, J. R. Cole, Naive Bayesian classifier for
rapid assignment of rRNA sequences into the new bacterial taxonomy. *Applied*
and Environmental Microbiology. **73**, 5261–5267 (2007).

35. J. W. Leff, mctoolsr: Microbial Community Data Analysis Tools, (available at
github.com/leffj/mctoolsr).

36. J. D. Storey, A direct approach to false discovery rates. *Journal of the Royal*
Statistical Society: Series B (Statistical Methodology). **64**, 479–498 (2002).

37. F. E. Harrell Jr, C. Dupont, Hmisc: Harrell Miscellaneous, (available at
http://CRAN.R-project.org/package=Hmisc).

38. S. J. Blazewicz, R. L. Barnard, R. A. Daly, M. K. Firestone, Evaluating rRNA as an
indicator of microbial activity in environmental communities: limitations and uses.
7, 2061–2068 (2013).

Acknowledgements:
Raw sequence data are available in the Sequence Read Archive at the National Center
for Biotechnology Information (project accession no. SRP070563). We thank Stuart
Grandy and Jörn Schnecker at University of New Hampshire, Cliff Bueno de Mesquita
and Lara Vimercati at the University of Colorado Boulder, Eric Skokan at Black Cat
Farms, and Kendra McLachlan at Kansas State University for soil collection or access
to collection sites. We also thank Jessica Henley, Robin Hacker-Cary and Kristen
Vaccarello for assistance with DNA extractions. Soil tests were preformed at Colorado
State University's Soil, Water and Plant Testing Laboratory. Sequencing was performed
at the University of Colorado BioFrontiers Institute's Next-Gen Sequencing Core
Facility. Funding to support this work was provided by the National Science Foundation
(DEB 0953331, EAR 1331828, DUE 1259336, EAR 1461281), and a Visiting

357
358
359
360   Supplementary Materials for
361

Relic DNA is abundant in soil and obscures estimates of soil microbial diversity

365   Paul Carini[1*], Patrick J. Marsden[2], Jonathan W. Leff[1,3], Emily E. Morgan[1], Michael S.
366   Strickland[4], Noah Fierer[1,3*]

*Corresponding Authors: Paul Carini (Paul.Carini@gmail.com) and Noah Fierer
(Noah.Fierer@colorado.edu)

Supporting Table 1: List of soil sample locations, descriptions and edaphic
characteristics.

Supporting Dataset 1: Full dataset for Fig. 1 and mean % each taxa per soil, per
treatment

Materials and Methods

Soil collection and edaphic characteristics: Thirty one surface soils (0-5 cm; mineral
soils only) were collected from locations in Colorado, New Hampshire, Virginia and
Kansas, USA in August or September 2015 (Table S1), sieved to 2.0 mm, homogenized
and stored at 4°C until PMA treatment. Soil characteristics were measured at the
Colorado State University Soil Water and Plant Testing Laboratory using their standard
protocols and included pH, electrical conductivity (mmhos cm^{-1}), % organic matter, NO_3-
N (ppm), P (ppm), K (ppm), Zn (ppm), Fe (ppm), Mn (ppm), Cu (ppm), texture (% sand,
silt, and clay), exchangeable bases (meq 100 g^{-1}), and cation exchange capacity (meq
100 g^{-1}). Percent moisture was determined gravimetrically on soils sieved to 2 mm
before and after drying at 80°C for 72 hours. Mean weight diameter (MWD_W; a proxy for
the amount of water stable aggregates) was determined using the wet sieving method
described in (*28*). Briefly, 2.0 g (W_t) wet soils were sieved in a water bath through a 0.25
mm (d) sieve for 3 minutes. Aggregates remaining on the filter were dried at 80°C for 72
h and weighed (W_s). Mean Weight Diameter after wet sieving (MWD_W) = stable
aggregates (W_s)* sieve diameter (d) / Initial weight (W_t).

Propidium monoazide treatment and DNA extraction: After homogenizing each soil, 10
replicate sub-samples of each soil type were resuspended (1% w/v) in sterile PBS (pH
7.4) in transparent screw cap tubes. PMA was added to 5 of the sub-samples (the PMA-

treated samples) at a final concentration of 40 µM in a dark room. The five PMA-treated
and the five untreated sub-samples were vortexed (setting 6 on VWR-brand vortexer)
for 4 minutes in the dark at room temperature. After the incubation, both PMA-treated
and untreated samples were exposed to a 650-watt halogen lamp placed 20 cm from
the sample tubes for four consecutive 30 s:30 s light:dark cycles, while continually
vortexing to ensure even light exposure throughout resuspension. After light exposure,
samples were frozen at -20°C until DNA extraction. DNA was extracted from 960 µL of
homogenized soil/PBS slurry using a MoBio PowerSoil DNA extraction kit, following
manufacturer's instructions. Experiments were conducted to determine the efficacy of
this approach and specifically determine whether: i) there was excess capacity of 40 µM
PMA to remove naked DNA; and ii) if PMA penetrated living *Escherichia coli* or
Saccharomyces cerevisiae, harvested in logarithmic-phase, and resuspended in PBS.
Details and results of these control experiments are provided in Fig. S1.
413
Quantitative PCR: qPCR reactions were conducted in 25 µL total volumes on a
Eppendorf realplex^2 Mastercycler ep gradient S. The reaction mixture was as follows:
12.5 µl of Absolute qPCR SYBR Green Mix (Thermo Scientific) master mix; 1.25 µl of
each primer (prokaryotic 16S rRNA: 515F 5'-GTGCCAGCMGCCGCGGTAA-3' & 806R
5'-GGACTACHVGGGTWTCTAAT-3'; fungal ITS: 5′-CTTGGTCATTTAGAGGAAGTAA-
3′ & ITS2 5′-GCTGCGTTCTTCATCGATGC-3'); 5 µl water; 5 µl of template DNA. All
reactions were run in triplicate in 96 well plates with triplicate standard curves containing
purified *E. coli* genomic DNA dilutions for 16S rRNA gene quantification and purified
Aspergillus fumigatus (ATCC MYA-4609D-2) genomic DNA dilutions for ITS amplicon
quantification. Program: 95°C for 15 min, followed by 40 cycles of (94°C 45 s; 50°C 60
s; 72°C 90 s) final extension 72°C 10 min. A qPCR 'no template' negative control was
included with each qPCR run.
426
Amplicon sequencing and analytical methods: For sequence-based analyses of 16S
rRNA and ITS gene regions (for prokaryotes and fungi respectively), we used the
approaches described previously (*29*). Briefly, we amplified each of the 310 DNA
samples in triplicate in 25 µL PCR reactions containing: 12.5 µl of Promega GoTaq Hot
Start Colorless Master Mix; 0.5 µl of each barcoded primer (bacterial 16S: 515F 5'-
GTGCCAGCMGCCGCGGTAA-3' & 806R 5'-GGACTACHVGGGTWTCTAAT-3'; fungal
ITS: 5′-CTTGGTCATTTAGAGGAAGTAA-3′ & ITS2 5′-GCTGCGTTCTTCATCGATGC-
3'); 10.5 µl water; 1 µl of template DNA. Program: 94°C for 5 min, followed by 35 cycles
of (94°C 45 s; 50°C 60 s; 72°C 90 s) and a final extension 72°C 10 min. Several
negative controls, including 'no template' controls and 'DNA extraction kit' controls, were
included alongside the soil DNA samples and sequenced. Amplicons were cleaned and
normalized using the ThermoFisher Scientific SequalPrep Normalization Plate kit.
Amplicons pools were spiked with phiX (15%) and sequenced on an Illumina MiSeq
using v2 500-cycle Paired End kits at the University of Colorado BioFrontiers Institute's
Next-Gen Sequencing Core Facility. Reads were processed as described in (*29*).
Briefly, raw amplicon sequences were demultiplexed according to the raw barcodes and
processed with the UPARSE pipeline (*30*). A database of ≥97% similar sequence
clusters was constructed in USEARCH (Version 8) (*31*) by merging paired end reads,
using a "maxee" value of 0.5 when quality filtering sequences, dereplicating identical

sequences, removing singleton sequences, clustering sequences after singleton
removal, and filtering out cluster representative sequences that were not ≥75% similar
to any sequence in Greengenes (for prokaryotes; Version 13_8) (*32*) or UNITE (for
fungi) (*33*) databases. Demultiplexed sequences were mapped against the *de novo*
constructed databases to generate counts of sequences matching clusters (i.e. taxa) for
each sample. Taxonomy was assigned to each taxon using the RDP classifier with a
threshold of 0.5 (*34*) and trained on the Greengages or UNITE databases. To normalize
the sequencing depth across samples, samples were rarefied to 6,000 and 1,100
sequences per sample for the 16S rRNA and ITS analyses, respectively.
455
Statistical analyses: The percent changes in marker gene copy number and microbial
richness in the relic DNA pool was calculated by comparing the mean number (n=5) of
amplicons or taxa from total soil DNA extracts to the number of amplicons or taxa from
each individual PMA treated subsample, using the formula (mean value in untreated
samples - value in each PMA treated sample)/(mean value in untreated samples).
Microbial richness and Bray-Curtis dissimilarities were calculated in the mctoolsr R
package (*35*). Bray-Curtis dissimilarities were calculated on square root transformed
taxa abundances. Where appropriate, statistical tests were corrected for multiple
comparisons using the qvalue R package (*36*) using a significance threshold of *q*
value ≤ 0.05. Spearman correlation coefficients and associated P values were
calculated with the Hmisc R package (*37*). Logistic regression models were used to
identify which edaphic characteristics predicted the presence of relic DNA (presence is
defined as >20% relic DNA).
469
Rationale for using PMA vs alternative methods: Several techniques have been
developed to investigate microbial viability in a variety of ecosystems (reviewed in (*17*)).
The approach we used was selected because *i*) it can be used for high-throughput
analyses; *ii*) the utility of PMA for the purposes of distinguishing extracellular DNA from
DNA contained within cells is well documented and robust (*15, 16*); and *iii*) key
assumptions in other methods are untested. For example, we did not use RNA-based
assays because it is impractical and prohibitively expensive to extract sufficient RNA
from ~300 samples, as we do here. In addition, although RNA-based approaches are
also thought to discriminate between metabolically active and inactive cells, this is not
necessarily true (*38*). Moreover, the assumption that extracellular RNA cannot persist in
soil has not been explicitly tested. It is possible that, as with DNA, there could be
substantial amounts of RNA in soil coming from extracellular pools or cells that are no
longer living, but additional studies are required to determine if this is the case.
483

Figure S1: PMA effectively removed experimentally added naked DNA from some, but
not all soils (A) and PMA did not penetrate live bacterial (B) or fungal (C) cells. A) Dark
and light shaded boxes show the range of 16S rRNA gene copies detected from DNA
extracts of untreated or PMA treated soils, respectively. Open circles are individual data
points within range. Blue bars represent the number of 16S rRNA gene copies
quantified in fresh soil spiked with ~8.8 × 10^4 copies of purified *E. coli* genomic DNA.
Red bars represent the number of 16S rRNA gene copies quantified in soil spiked with
~8.8 × 10^4 copies of purified *E. coli* genomic DNA and treated with 40 µM PMA as
described for soils in 'materials and methods'. PMA did not remove all of the spiked
DNA in Soil 12, which also contained the largest percent of extracellular DNA (Fig. 1),
suggesting additional PMA was necessary to remove the spiked DNA. B,C) Sterile PBS
was spiked with equal volumes of exponentially growing *E. coli* and *S. cerevisiae* cell
cultures. Tubes were either treated with 40 µM PMA (n=5) or left untreated (n=5) as
described for soils in 'materials and methods'. The number of 16S rRNA gene copies or
ITS amplicon copies were indistinguishable to untreated controls (two-tailed *t* test
P=0.96 and *P*=0.22, respectively), illustrating PMA did not penetrate live bacterial or
fungal cells.
505

Figure S2: The percent of marker genes in the relic DNA pool is positively correlated with the proportional richness in the relic DNA pool, but this correlation is only significant for prokaryotes. Points are the mean percent relic DNA and mean percent richness in relic DNA, taken from Fig. 1. Linear regressions, formulas, Spearman's ρ and P value are shown.

Figure S3: Differences in community composition between individual soil types is greater than the effect of relic DNA removal for any given soil. Dendrogram of prokaryotic (A) and fungal (B) community composition for all 31 soils. Multidimensional scaling plot of prokaryotic (C) and fungal (D) community composition for all 31 soils. Hulls in C & D connect the outermost points for each soil.

539
Figure S4: Fold changes in the estimated relative abundances of individual prokaryotic
(A) and fungal (B) taxa after removal of relic DNA across all soils. Only major taxonomic
groups comprising ≥5% of the total prokaryotic community or ≥3% of the total fungal
community are shown. Points are the $\log_2$ fold change in estimated mean relative
abundances after relic DNA removal. Points are not shown for some fungal taxa
because they were not detected in one of the two conditions, preventing calculation of
$\log_2$ fold change. See Supplemental Dataset S1 for full dataset. *Significant (two-tailed
Mann-Whitney U *P*≤0.05).

Figure S5: The relative abundances of prokaryotic nitrifiers and arbuscular mycorrhizal fungi change after relic DNA removal. Relative abundance of ammonia oxidizing archaea (A) or of nitrite oxidizing bacteria (B) before and after relic DNA removal (mean ± SE). C) Relative abundance of *Glomeraceae* (mycorrhizal fungi) before and after relic DNA removal (mean ± SE). In all plots, dashed line represents no change in relative abundance (1:1 line). Dark grey shaded region represents ±10% change; light grey shaded region represents ±25% change.

572
Figure S6: Logistic regressions fitting the presence of relic DNA to total exchangeable
base cations (Na$^+$, K$^+$, Ca^{2+} and Mg^{2+}) and exchangeable K$^+$. Relic DNA was considered
present if it represented ≥20% of the total DNA pool. Logistic regressions are shown in
blue. Associated *P* values are provided in Table 1.

577
578 **Figure S7:** Soil pH is correlated with the difference in community composition after relic
579 DNA removal. Points show the mean dissimilarity in soil microbial communities for
580 prokaryotes (A) and fungi (B) after relic DNA removal, relative to total DNA extracts.
581 Linear regressions, formulas, Spearman's ρ and P values are shown.
582