The evolutionary fates of a large segmental duplication in mouse

Andrew P Morgan[1], J Matthew Holt[2], Rachel C McMullan[1], Timothy A Bell[1], Amelia M-F Clayshulte[1],

John P Didion[1], Liran Yadgary[1], David Thybert[3], Duncan T Odom[4,5], Paul Flicek[3,5], Leonard McMillan[2],

Fernando Pardo-Manuel de Villena[1]

[1] Department of Genetics, Carolina Center for Genome Sciences and Lineberger Comprehensive Cancer Center, University of North Carolina, Chapel Hill, NC

[2] Department of Computer Science, University of North Carolina, Chapel Hill, NC

[3] European Molecular Biology Laboratory, European Bioinformatics Institute, Wellcome Genome Campus, Hinxton, Cambridge, CB10 1SD, United Kingdom

[4] University of Cambridge, Cancer Research UK Cambridge Institute, Robinson Way, Cambridge, CB2 0RE, United Kindgom

[5] Wellcome Trust Sanger Institute, Wellcome Genome Campus, Hinxton, Cambridge, CB10 1SA, United Kingdom

Corresponding author:

Fernando Pardo-Manuel de Villena

5049 Genetic Medicine Building

120 Mason Farm Road CB#7264

Chapel Hill, NC 27599-7264

ABSTRACT

Gene duplication and loss are major sources of genetic polymorphism in populations, and are important forces shaping the evolution of genome content and organization. We have reconstructed the origin and history of a 127 kbp segmental duplication, *R2d*, in the house mouse (*Mus musculus*). *R2d* contains a single protein-coding gene, *Cwc22*. *De novo* assembly of both the ancestral (*R2d1*) and the derived (*R2d2*) copies reveals that they have been subject to non-allelic gene conversion events spanning tens of kilobases. *R2d2* is also a hotspot for structural variation: its diploid copy number ranges from zero in the mouse reference genome to more than 80 in wild mice sampled from around the globe. Hemizgyosity for high-copy-number alleles of *R2d2* is associated in *cis* with meiotic drive, suppression of meiotic crossovers, and copy-number instability, with a mutation rate in excess of 1 per 100 transmissions in laboratory populations. We identify an additional 57 loci covering 0.8% of the mouse genome with patterns of sequence variation similar to those at *R2d1* and *R2d2*. Our results provide a striking example of allelic diversity generated by duplication and demonstrate the value of *de novo* assembly in a phylogenetic context for understanding the mutational processes affecting duplicate genes.

INTRODUCTION

Duplication is an important force shaping the evolution of plant and animal genomes: it provides both a substrate for evolution and transient relief from selective pressure (Lynch and Conery 2000). Segmental duplications (SDs), defined as contiguous sequences which map to more than one physical position (Bailey and Eichler 2006), are a common feature of eukaryotic genomes and particularly those of vertebrates.

Like any sequence variant, a duplication first arises in a single individual in a population. The distinction between such copy-number variants (CNVs) and SDs is fluid and somewhat arbitrary: tracts of SDs are highly polymorphic in populations in species from *Drosophila* (Dopman and Hartl 2007) to mouse (She *et al.* 2008) to human (Bailey and Eichler 2006). Studies of parent-offspring transmissions have shown that SDs are prone to recurrent *de novo* mutations including some implicated in human disease (reviewed in Stankiewicz and Lupski 2002). Bursts of segmental duplication have preceded dramatic species radiations in primates. More broadly, blocks of conserved synteny in mammals frequently terminate at SDs (Bailey and Eichler 2006), suggesting that SDs could mediate the chromosomal rearrangements through which karyotypes diverge and reproductive barriers arise.

Notwithstanding their evolutionary importance, SDs are difficult to analyze. Repeated sequences with period longer than the insert size in a sequencing library and high pairwise similarity are likely to be collapsed into a single sequence during genome assembly. Efficient and sensitive alignment of high-throughput sequencing reads to duplicated sequence remains challenging (Treangen and Salzberg 2011). Genotyping of sites within SDs is difficult because variants between copies (paralogous variants) are easily confounded with variants within copies between individuals at a given copy (allelic variants). Latent paralogous variation may bias interpretations of sequence diversity and haplotype structure (Hurles 2002).

Paralogy also complicates phylogenetic inference. Ancestral duplication followed by differential losses along separate lineages may yield a local phylogeny that is discordant with the genome-wide phylogeny (Goodman *et al.* 1979). Within each duplicate copy, local phylogenies for adjacent intervals may also be discordant due to non-allelic gene conversion between copies (Dover 1982; Nagylaki and Petes 1982). As a result, over some fraction of the genome, sequences from individuals of the same species may be more closely related to sequences from individuals of an outgroup species than they are to each other.

In this manuscript we present a detailed analysis of a segmental duplication, *R2d*, in the house mouse (*Mus musculus*). *R2d* is a 127 kbp unit which contains the protein-coding gene *Cwc22* and flanking intergenic sequence. Although the C57BL/6J reference strain and other classical laboratory strains have a single haploid copy of the *R2d* sequence (in the *R2d1* locus), the wild-derived CAST/EiJ, ZALENDE/EiJ, and WSB/EiJ strains have an additional 1, 16 and 33 haploid copies respectively in the *R2d2* locus. *R2d2* is the responder locus in a recently-described meiotic drive system on mouse chromosome 2 but is absent from the mouse reference genome (Waterston *et al.* 2002; Didion *et al.* 2015, 2016). We draw on a collection of species from the genus *Mus* sampled from around the globe to reconstruct the sequence of events giving rise to the locus' present structure (**Figure 1**). Using novel computational tools built around indexes of raw high-throughput sequencing reads, we perform local *de novo* assembly of phased haplotypes and explore patterns of sequence diversity within and between copies of *R2d*.

Both phylogenetic analyses and estimation of mutation rate in laboratory mouse populations reveal that *R2d2* and its surrounding region on chromosome 2 are unstable in copy number. Cycles of duplication, deletion and non-allelic gene conversion have led to complex phylogenetic patterns discordant with

species-level relationships within *Mus* which cannot be explained by known patterns of introgression
between *Mus* species (Bonhomme *et al.* 2007; Yang *et al.* 2011).

Finally, we identify 57 other loci, covering 0.8% of the mouse genome, which share the key features of
R2d2: elevated local sequence divergence; low recombination rate; and enrichment for segmental
duplications. Previous studies of sequence variation in the mouse (Keane *et al.* 2011) have attributed this
pattern to sorting of alleles segregating in the common ancestor of *M. musculus* and its sister species. We
suggest instead that these loci have been subject to independent cycles of duplication and loss along *Mus*
lineages. Marked enrichment for odorant, pheromone, and antigen-recognition receptors supports a role
for balancing selection on the generation and maintenance of the extreme level of polymorphism
observed at these loci.

RESULTS

R2d* was duplicated in the common ancestor of *M. musculus* and *M. spretus

In order to determine when the *R2d* CNV arose, we used quantitative PCR and/or depth of coverage in
whole-genome sequencing to assay *R2d* copy number in a collection of samples spanning the phylogeny
of the genus *Mus.* Samples were classified as having diploid copy number 2 (two chromosomes each with
a single copy of *R2d*) or >2 (at least one chromosome with an *R2d* duplication).

We find evidence for >2 diploid copies in representatives of all mouse taxa tested from the Palearctic
clade (Suzuki *et al.* 2004) (**Figure 1** and **Supplementary Table 1**): 236 of 525 *Mus musculus*, 1 of 1 *M.*
macedonicus, 1 of 1 *M. spicilegus*, 1 of 1 *M. cypriacus* and 8 of 8 *M. spretus* samples. However, we find no
evidence of duplication in species from the southeast Asian clade, which is an outgroup to Palearctic
mice: 0 of 2 *M. famulus*, 0 of 2 *M. fragilicauda*, 0 of 1 *M. cervicolor*, 0 of 1 *M. cookii* and 0 of 1 *M. caroli*
samples. Outside the subgenus *Mus*, we found evidence for >2 diploid copies in none of the 9 samples
tested from subgenus *Pyromys*. We conclude that the *R2d* duplication most likely occurred between the
divergence of southeast Asian from Palearctic mice (~3.5 million years ago [Mya]) and the divergence of
M. musculus from *M. spretus* (~2 Mya) (Suzuki *et al.* 2004; Chevret *et al.* 2005), along the highlighted
branch of the phylogeny in **Figure 1A**. If the *R2d* duplication is ancestral to the divergence of *M. musculus*,
then extant lineages of house mice which have 2 diploid copies of *R2d* — including the reference strain
C57BL/6J (of predominantly *M. musculus domesticus* origin (Yang *et al.* 2007)) — represent subsequent
losses of an *R2d* copy.

Duplication of the ancestral *R2d* sequence resulted in two paralogs residing in loci which we denote *R2d1*
and *R2d2* (**Figure 1B**). Only one of these is present in the mouse reference genome, at chr2: 77.87 Mbp; the
other copy maps approximately 6 Mbp distal (Didion *et al.* 2015), as we describe in more detail below.
The more proximal copy, *R2d1*, lies in a region of conserved synteny with rat, rabbit, chimpanzee and
human (Muffato *et al.* 2010) (**Supplementary Figure 1**); we conclude that it is the ancestral copy.

The sequence of the *R2d2* paralog was assembled *de novo* from whole-genome sequence reads (Keane *et al.*
2011) from the strain WSB/EiJ (of pure *M. m. domesticus* origin (Yang *et al.* 2011)), which has diploid *R2d*
copy number ~68 (Didion *et al.* 2015). We exploited the difference in depth of coverage for *R2d1* (1 haploid
copy) and *R2d2* (33 haploid copies) to assign variants to *R2d1* or *R2d2*. Pairwise alignment of the *R2d2*
contig against *R2d1* is shown in **Supplementary Figure 2**. The paralogs differ by at least 8 transposable-
element (TE) insertions: 7 LINE elements specific to *R2d1* and 1 endogenous retroviral element (ERV)
specific to *R2d2* (**Supplementary Table 2**). (Due to the inherent limitations of assembling repetitive

elements from short reads, it is likely that we have underestimated the number of young TEs in *R2d2*.)
The *R2d1*-specific LINEs are all < 2% diverged from the consensus for their respective families in the
RepeatMasker database (http://www.repeatmasker.org/cgi-bin/WEBRepeatMasker), consistent with
insertion within the last 2 My. The oldest *R2d2*-specific ERV we could detect is 0.7% diverged from its
family consensus. TE insertions occurring since the ancestral *R2d* duplication are almost certainly
independent, so these data are consistent with duplication <2 Mya. The *R2d* unit, minus paralog-specific
TE insertions, is 127 kbp in size. *R2d* units in the *R2d2* locus are capped on both ends by (CTCC)n
microsatellite sequences, and no read pairs spanning the breakpoint between *R2d2* and flanking sequence
were identified.

In order to obtain a more precise estimate of the molecular age of the duplication event we assembled *de*
novo a total of 16.9 kbp of intergenic and intronic sequence in 8 regions across the *R2d* unit from diverse
samples and constructed phylogenetic trees. The trees cover 17 *R2d1* or *R2d2* haplotypes, 13 from inbred
strains and 4 from wild mice. The sequence of *Mus caroli* (diploid copy number 2) is used as an outgroup.
A concatenated tree is shown in **Figure 1C**. Using 5.0 ± 1.0 million years before present (Mya) as the
estimated divergence date for *M. caroli* and *M. musculus* (Suzuki *et al.* 2004; Chevret *et al.* 2005), Bayesian
phylogenetic analysis with BEAST v1.8 (Drummond *et al.* 2012) yields 1.6 Mya (95% HPD $0.7 - 5.1$ Mya)
as the estimated age of the duplication event that gave rise to *R2d1* and *R2d2*. Although the assumption
of a uniform molecular clock may not be strictly fulfilled for *R2d1* and *R2d2*, the totality of evidence –
from presence/absence data across the mouse phylogeny, paralog-specific TE insertions, and sequence
divergence between paralogs – strongly supports the conclusion that R2d was first duplicated within the
last 2 My in the common ancestor of *M. musculus* and *M. spretus*.

For clarity, **Figure 1D** illustrates diploid copy number states that will be referenced in the remainder of
the manuscript. Hereafter we refer to diploid copy numbers except when discussing inbred strains
(which are effectively haploid).

R2d contains the essential gene *Cwc22*

The *R2d* unit encompasses one protein-coding gene, *Cwc22*, which encodes an essential mRNA splicing
factor (Yeh *et al.* 2010). The gene is conserved across eukaryotes and is present in a single copy in most
non-rodent species represented in the TreeFam database (http://www.treefam.org/family/TF300510 (Li
2006)). Five groups of *Cwc22* paralogs are present in mouse genomes: the copies in *R2d1* (*Cwc22^R2d1^*) and
R2d2 (*Cwc22^R2d2^*) plus retrotransposed copies in one locus at chr2: 83.9 Mbp and at two loci on the X
chromosome (**Figure 2A**).

The three retrogenes are located in regions with no sequence similarity to each other, indicating that each
represents an independent retrotransposition event. The copy on chr2 was subsequently expanded by
further segmental duplication and now exists (in the reference genome) in 7 copies with >99.9% mutual
similarity. The two retrotransposed copies on chrX are substantially diverged from the parent gene (<
90% sequence similarity), lack intact open reading frames (ORFs), have minimal evidence of expression
among GenBank cDNAs, and are annotated as likely pseudogenes (Pei *et al.* 2012). We therefore restricted
our analyses to the remaining three groups of *Cwc22* sequences, all on chr2.

The canonical transcript of *Cwc22^R2d1^* (ENSMUST00000065889) is encoded by 21 exons on the negative
strand. The coding sequence begins in the third exon and ends in the terminal exon (**Figure 2B**). Six of the
seven protein-coding *Cwc22^R2d1^* transcripts in Ensembl v83 use this terminal exon, while one transcript
(ENSMUST0000011824) uses an alternative terminal exon. Alignment of the retrogene sequence
(ENSMUST00000178960) to the reference genome demonstrates that the retrogene captures the last 19

161 exons of the canonical transcript — that is, the 19 exons corresponding to the coding sequence of the
162 parent gene.

Copy number at *R2d2* is highly polymorphic in *M. musculus*

164 We previously demonstrated that haploid copy number of *R2d* ranges from 1 in the reference strain
165 C57BL/6J and classical inbred strains A/J, 129S1/SvImJ, NOD/ShiLtJ, NZO/HILtJ; to 2 in the wild-derived
166 strain CAST/EiJ; to 34 in the wild-derived strain WSB/EiJ. Using linkage mapping in two multiparental
167 genetic reference populations, the Collaborative Cross (Collaborative Cross Consortium 2012) and
168 Diversity Outbred (Svenson *et al.* 2012), we showed that, for the two strains with haploid copy number
169 >1, one of the copies maps to *R2d1* while all extra copies map to the *R2d2* locus at chr2: 83 Mbp (Didion *et*
170 *al.* 2015). *Cwc22* was recently reported to have diploid copy number as high as 83 in wild *M. m. domesticus*
171 (Pezer *et al.* 2015). In whole-genome sequence data from more than sixty mice from both laboratory stocks
172 and natural populations (**Supplementary Table 1**), we have observed zero instances in which the *R2d*
173 copy in *R2d1* is lost. We conclude that diploid copy number >2 indicates at least one copy of *R2d* is
174 present in *R2d* (**Figure 1D**).

175 In order to understand the evolutionary dynamics of copy-number variation at *R2d2*, we investigated the
176 relationship between copy number and the local phylogeny in the *R2d2* candidate region. In particular,
177 we sought evidence for or against a single common origin for each of the copy-number states at *R2d2*
178 which are derived with respect to the *M. spretus* – *M. musculus* common ancestor (**Figure 1D**). If a derived
179 copy-number state has a single recent origin, it should be associated with a single haplotype at *R2d2*. If a
180 derived copy-number state arises by recurrent mutation, the same copy number should be associated
181 with multiple haplotype backgrounds and possibly in multiple populations.

182 The extent of *R2d* copy-number variation in *M. musculus*, as estimated on a continuous scale by qPCR, is
183 shown in **Figure 3A**. (Note that the qPCR readout is proportional to copy number on the log scale.
184 Extrapolation to integer copy number is imprecise for copy numbers greater than ~6.) We confirmed that
185 *R2d2* maps to chr2: 83 Mbp by performing association mapping between SNP genotypes from the
186 MegaMUGA array (Morgan *et al.* 2016) and the qPCR readout (**Figure 3B**).

187 To test the hypothesis that losses of *R2d2* (diploid copy number < 4, at least one chromosome with zero
188 copies in *R2d2*, **Figure 1D**) have a single origin, we examined their distribution across the three well-
189 differentiated subspecies of *M. musculus*. Losses of *R2d2* occur in all subspecies of *M. musculus*, in
190 populations that span its geographic range (**Supplementary Table 3**). Based on this distribution and a
191 previous observation that no common haplotype is shared in samples with low copy number in *M. m.*
192 *domesticus* (Didion *et al.* 2016), we reject the hypothesis of single origin and conclude that *R2d2* has been
193 lost multiple times on independent lineages in each subspecies.

194 We performed a similar analysis to test the hypothesis that *R2d2* alleles with high copy number (diploid
195 copy number >4, **Figure 1D**; hereafter "*R2d2^HC^*") have a single origin. First we observed that *R2d2^HC^*
196 alleles are confined with few exceptions to *M. m. domesticus* (**Supplementary Table 3**). The best-
197 associated SNP on the MegaMUGA array (JAX00494952) only weakly tags copy number ($r^2 = 0.137$), but
198 ascertainment bias on the MUGA platform (Morgan *et al.* 2016) makes local LD patterns difficult to
199 interpret. To examine further, we constructed a neighbor-joining phylogenetic tree for the region
200 containing *R2d2* (chr2: 83 – 84 Mb) using genotypes from the 600K-SNP Mouse Diversity Array (Yang *et*
201 *al.* 2011). We restricted our attention to inbred strains or wild mice with homozygous, non-recombinant
202 haplotypes in the target region. Twelve samples with *R2d2^HC^* alleles, both wild mice and laboratory

stocks, cluster in a single clade (**Figure 3C**). (A single *M. spretus* strain, SPRET/EiJ, also carries an *R2d2^HC^*
allele, but see **Discussion**).

Next we expanded the analysis to include an additional 11 samples with *R2d2^HC^* alleles and evidence of
heterozygosity around *R2d2*. The total set of 24 samples includes 7 wild-derived laboratory strains
(DDO, RBA/DnJ, RBB/DnJ, RBF/DnJ, WSB/EiJ, ZALENDE/EiJ and SPRET/EiJ), 4 classical inbred strains
(ALS/LtJ, ALR/LtJ, CHMU/LeJ and NU/J), a line derived from the ICR:HsD outbred stock (HR8; Swallow
et al. 1998) and 12 wild-caught mice. All 24 samples with *R2d2^HC^* alleles share an identical haplotype
across a single 21 kbp interval, chr2: 83,896,447 – 83,917,565 (GRCm38/mm10 coordinates) (**Figure 3D**).
These analyses support a single origin for *R2d2^HC^* alleles within *M. m. domesticus*.

Cwc22 is intact in and expressed from all *R2d* paralogs, and fast-evolving in rodents

To identify the coding sequence of *Cwc22^R2d2^* we first aligned the annotated transcript sequences of
*Cwc22^R2d1^* from Ensembl to our *R2d2* contig. All 21 exons present in *R2d1* are present in *R2d2*. We created a
multiple sequence alignment and phylogenetic tree of *Cwc22* cDNAs and predicted amino acid sequences
from *Cwc22^R2d1^*, *Cwc22^R2d2^*, retro-*Cwc22*, and *Cwc22* orthologs in 19 other placental mammals, plus
opossum, platypus and finally chicken as an outgroup (**Supplementary Figure 3**). An open reading frame
(ORF) is maintained in all three *Cwc22* loci in mouse, including the retrogene. Information content of each
column along the alignment (**Supplementary Figure 4**) reveals that sequence is most conserved in two
predicted conserved domains, MIF4G and MA3, required for *Cwc22*'s function in mRNA processing (Yeh
et al. 2010).

Next we examined public RNA-seq data from adult brain and testis in inbred strains with one or more
copies of *R2d2* for evidence of transcription of each *Cwc22* family member. We identified several novel
transcript isoforms specific to *R2d2* arising from two intron-retention events and one novel 3' exon
(**Figure 4A**). The 18th intron is frequently retained in *Cwc22^R2d2^* transcripts, most likely due to an A>G
mutation at the 5' splice donor site of exon 17 in *Cwc22^R2d2^*. The 12th intron is also frequently retained.
While we could not identify any splice-region variants near this intron, it contains an ERV insertion that
may interfere with splicing (**Figure 4A**). Both intron-retention events would create an early stop codon.
Finally, we find evidence for a novel 3' exon that extends to the boundary of the *R2d* unit and is used
exclusively by *Cwc22^R2d2^* (**Figure 4A**).

We estimated the expression of the various isoforms of *Cwc22^R2d1^*, *Cwc22^R2d2^* and retro-*Cwc22* in adult
brain and testis. For brain we obtained reads from 8 replicates (representing both sexes) on 3 inbred
strains, and for testis a single replicate on 23 inbred strains and estimated transcript abundance using the
kallisto package (Bray *et al.* 2015). Briefly, kallisto uses an expectation-maximization (EM) algorithm to
accurately estimate the abundance of a set of transcripts by distributing the "weight" of each read across
all isoforms with whose sequence it is compatible. *Cwc22* is clearly expressed from all three paralogs in
both brain and testis (**Figure 4B**). However, both the total expression and the pattern of isoform usage
differ by tissue and copy number.

Maintenance of an ORF in all *Cwc22* paralogs for >2 My is evidence of negative selection against
disrupting mutations in the coding sequence, but long branches within the rodent clade in
Supplementary Figure 3 suggest that *Cwc22* may also be under relaxed purifying selection or positive
selection in rodents. The rate of evolution of *Cwc22* sequences in mouse is faster than in the rest of the tree
($\chi^2 = 4.33$, df = 1, $p = 0.037$ by likelihood ratio test).

Phylogenetic discordance in *R2d1* is due to non-allelic gene conversion

The topology of trees across *R2d* is generally consistent: a long branch separating the single *M. caroli* sequence from the *M. musculus* sequences, and two clades corresponding to *R2d1*- and *R2d2*-like sequences. However, we observed that the affinities of some *R2d* paralogs change along the sequence (**Figure 5A**), a signature of non-allelic (*i.e.* inter-locus) gene conversion. In this context, we use "gene conversion" to describe a non-reciprocal "copy-and-paste" transfer of sequence from one donor locus into a different, homologous receptor locus, without reference to a specific molecular mechanism (Chen *et al.* 2007).

To investigate further, we inspected patterns of sequence variation in whole-genome sequencing data from 15 wild-caught mice, 2 wild-derived inbred strains, and 22 classical inbred strains of mice with diploid *R2d* copy number 2. We first defined 1,411 pairwise single-nucleotide differences between *R2d2* and *R2d1* for which *R2d2* has the derived allele with respect to *M. caroli*. Then we tested for the presence of the derived allele, ancestral allele or both at each site in each sample. Finally we identified conversion tracts by manual inspection as clusters of derived variants shared with *R2d2* (**Supplementary Figure 5**).

This analysis revealed non-allelic gene conversion tracts on at least 9 chromosomes (**Figure 5B**). The conversion tracts range in size from approximately 1.2 kbp to 119 kbp. The boundaries of several tracts are shared within populations, suggesting that they are identical by descent. We excluded the possibility of complementary losses from *R2d1* and *R2d2* – which would leave similar patterns of sequence variation – by finding read pairs spanning the boundary between *R2d1* and flanking sequence, and between *R2d1*-like and *R2d2*-like tracts on the same chromosome (examples shown in **Figure 5C**).

The conversion tracts we detected are orders of magnitude longer than the 15 to 750 bp reported in recent studies of allelic gene conversion at recombination hotspots in mouse meiosis (Cole *et al.* 2010, 2014). We require the presence of *R2d2*-diagnostic alleles at two or more consecutive variants to declare a conversion event, and these variants occur at a rate of approximately 1 per 100 bp, so the smallest conversion tracts we could theoretically detect are on the order of 200 bp in size. Even if we require only a single variant to define a conversion tract, all samples without a long conversion tract share fewer than 55 and most fewer than 10 (of 1,411) derived alleles with *R2d2*, of which all are also shared by multiple other samples from different populations (**Supplementary Figure 5**). This pattern indicates that those sites in fact represent either artifacts (from mis-assignment of ancestral and derived alleles) or homoplasy rather than short gene conversions.

Four conversion tracts partially overlap the *Cwc22* gene to create a sequence that is a mosaic of *R2d1*- and *R2d2*-like exons (**Figure 5B**). Recovery of *Cwc22* mRNA in an inbred strain with a mosaic sequence (PWK/PhJ, see section "*Cwc22* is intact and expressed") indicates that its exons are intact and properly oriented in *cis*. The presence of both *R2d1*- and *R2d2*-like sequence in extant *M. musculus* lineages with a 2 diploid copies of *R2d* further reinforces our conclusion that the duplication is indeed ancestral to the divergence of *M. musculus*.

In addition to exchanges between *R2d1* and *R2d2*, we identified an instance of exchange between *R2d2* and the adjacent retrotransposed copy of *Cwc22* in a single *M. m. domesticus* individual from Iran (IR:AHZ_STND:015; **Supplementary Figure 6**). This individual carries a rearrangement that has inserted a 30 kbp fragment corresponding to the 3' half of *Cwc22^{R2d2}* into the retro-*Cwc22* locus, apparently mediated by homology between the exons of *Cwc22^{R2d2}* and retro-*Cwc22*.

High copy number at *R2d2* suppresses meiotic recombination

Based on a previous observation that the rate of meiotic recombination is reduced near clusters of segmental duplications (Liu *et al.* 2014), we tested whether the region around *R2d2* has lower recombination when an *R2d2^HC* allele is present. Understanding patterns of recombination at *R2d2* is important for interpreting levels of sequence and haplotype diversity in the surrounding region.

First we analyzed local recombination rate in the Diversity Outbred (DO) population. The DO is an outbred stock derived from eight inbred founder strains (including one, WSB/EiJ, with an *R2d2^HC* allele) and maintained by random mating with 175 breeding pairs; at each generation, one male and one female offspring are chosen from each mating and randomly paired with a non-sibling to produce the next generation (Svenson *et al.* 2012). **Figure 6A** shows the cumulative distribution of 2,917 recombination events on central chromosome 2, stratified according to *R2d2* copy number of the participating haplotypes. The recombination map has a pronounced plateau in the region between *R2d1* and approximately 1 Mb distal to *R2d2* (dashed lines) for *R2d2^HC* haplotypes, but not *R2d2^LC* haplotypes. As a result, *R2d2^HC* haplotype blocks overlapping *R2d2* are significantly shorter than *R2d2^LC* haplotype blocks ($p < 0.01$ by Wilcoxon rank-sum tests with Bonferroni correction) in 8 of the 10 generations sampled (**Figure 6B**). The difference arose early in the breeding of the DO and persists through the most recent generation for the randomized breeding scheme was maintained (FPMdV, unpublished).

Second we re-examined genotype data from 11 published crosses in which at least one parent was segregating for an *R2d2^HC* allele. Whereas in the DO we used haplotype block length as a proxy for recombination rate, in these F$_2$ and backcross designs we can directly estimate the recombination fraction across *R2d2* and compare it to its expected value in the absence of an *R2d2^HC* allele (**Supplementary Figure 7**). In 9 of 11 crosses examined, the observed recombination fraction is lower than the expected ($p < 0.032$, one-sided binomial test).

The genomic region containing *R2d2* is structurally unstable but has low sequence diversity

The extent of copy-number polymorphism involving *R2d2* suggests that it is intrinsically unstable. Consistent with these observations, we find that the rate of *de novo* copy-number changes at *R2d2* is extremely high in laboratory populations (**Figure 7**). In 183 mice sampled from the DO population we identified and confirmed through segregation analysis 8 new alleles, each with distinct copy number and each occurring in an unrelated haplotype (**Supplementary Table 4**). Without complete pedigrees and genetic material from breeders a direct estimate of the mutation rate in the DO is not straightforward to obtain. However, since the population size is known, we can make an analogy to microsatellite loci (Moran 1975) and estimate the mutation rate via the variance in allele sizes: 3.2 mutations per 100 transmissions (3.2%) (95% bootstrap CI $1.1\% - 6.0\%$).

Structural instability in this region of chromosome 2 extends outside the *R2d2* locus itself. Less than 200 kbp distal to *R2d2* is another segmental duplication (**Figure 8B,** grey shaded region) — containing a retrotransposed copy of *Cwc22* — that is present in 7 tandem copies in the reference genome. That region, plus a further 80 kbp immediately distal to it, is copy-number polymorphic in wild *M. m. domesticus* and wild *M. m. castaneus* (**Figure 8B**). Instability of the region over longer timescale is demonstrated by the disruption, just distal to the aforementioned segmental duplication, of a syntenic block conserved across all other mammals (**Supplementary Figure 1**).

Despite the high mutation rate for structural variants involving *R2d2* and nearby sequences, sequence diversity at the nucleotide level is modestly reduced relative to diversity in *R2d1* and relative to the genome-wide average in *M. m. domesticus*. In a 200 kbp region containing the *R2d2* insertion site at its

proximal end, π (an estimator of average heterozygosity) in *M. m. domesticus* reduced from approximately
0.3% (comparable to previous reports in this subspecies, (Salcedo *et al.* 2007)) to nearly zero (**Figure 8B**).
Divergence between *M. musculus* and *M. caroli* is similar to its genome-wide average of ~ 2.5% over the
same region.

Estimation of diversity *within* a duplicated sequence such as *R2d* is complicated by the difficulty of
distinguishing allelic from paralogous variation. To circumvent this problem we split our sample of 26
wild *M. m. domesticus* into two groups: those having *R2d1* sequences only, and those having both *R2d1*
and *R2d2* sequences. Within each group we counted the number of segregating sites among all *R2d2*
copies, using nearby fixed differences between *R2d1* and *R2d2* to phase sites to *R2d2* (see **Methods** for
details), and used Watterson's estimator to calculate nucleotide diversity per site. Among *R2d1*
sequences, $\theta = 0.09\% \pm 0.03\%$ versus $\theta = 0.04\% \pm 0.02\%$ among *R2d2* sequences (**Figure 8C**) and
$\theta = 0.13\% \pm 0.04\%$ among *R2d2* sequences in *M. m. castaneus*.

The "revolving door" affects at least 0.8% of the mouse genome

The superposition of duplication, non-allelic gene conversion, and loss — dubbed the "genomic
revolving door" (Demuth *et al.* 2006) — can obscure the true history of multicopy sequences. Over short
evolutionary times, duplication and loss may be further confounded with incomplete lineage sorting, or
the stochasitic distribution of ancestral polymorphisms along descendant lineages (Pamilo and Nei 1988).

Most previous studies of gene gain and loss have focused on between-species comparisons using finished
genome assemblies (Demuth *et al.* 2006; Bailey and Eichler 2006). We sought evidence for the "revolving
door" effect over the last ~1 My of evolution within *M. musculus*. To do so we identified regions where
the divergence between sequenced samples and the reference genome assembly substantially exceeds
what is predicted by those samples' ancestry. We developed a simple and unbiased method to estimate
divergence from short sequencing reads without alignment. Briefly, we estimated the proportion of short
subsequences (k-mers, with $k = 31$) from windows along the reference assembly for which no evidence
exists in the reads generated for a sample. This quantity can be rescaled to approximate the divergence
between the reference sequence and the template sequence from which reads were generated (see
Methods). Applied genome-wide to 31 kbp ($= 1000{\times}k$) windows, this method captures the distribution
of sequence divergence between the reference assembly (chiefly *M. m. domesticus* in origin) in
representative samples from the three subspecies of *M. musculus* and the outgroups *M. spretus* and *M.*
caroli (**Figure 9A**).

Divergent regions were identified by fitting a hidden Markov model (HMM) to the windowed
divergence profiles for 7 wild or wild-derived samples with available whole-genome sequence (**Figure 9B**
and **Supplementary Table 6**). This analysis revealed a striking pattern: over most of the genome, the
divergence estimates hover around the genome-wide expectation, but high-divergence windows are
clustered in regions 100 kbp to 5 Mbp in size. Our method does not capture signal from structural
variation (besides deletions, see **Methods**), and so probably underestimates the true level of sequence
divergence. In any case the union of these divergent regions across all seven *M. musculus* samples
analyzed covers 0.82% of the reference genome (mean 0.58% per sample). Not surprisingly, divergent
regions are enriched for segmental duplications: 39.0% of sequence in divergent regions is comprised of
segmental duplications versus a median of 3.4% (central 99% interval $0.2\% - 16.7\%$) in random regions
of equal size ($p < 0.001$). Divergent regions also have significant overlap with previously-identified (Liu
et al. 2014) regions of low recombination ($p < 0.001$). Yet divergent regions are more gene-dense than the
genomic background: they contain $3.6{\times}10^{-2}$ genes/kbp relative to a median $1.6{\times}10^{-2}$ genes/kbp (central
99% interval $1.1{\times}10^{-2} - 2.7{\times}10^{-2}$ genes/kbp) genome-wide ($p = < 0.001$). Divergent regions are strongly

enriched for genes related to odorant and pheromone sensing ($p = 1.3 \times 10^{-8}$) and adaptive immunity
($p = 3.1 \times 10^{-3}$).

As a representative example we focus on a divergent region at chr4: 110 - 115 Mbp (**Figure 9B**). This
region contains the 11 members of the *Skint* family of T-cell-borne antigen receptors. The first member to
be described, *Skint1*, functions in negative selection in the thymus of T-cells destined for the epidermis
(Boyden *et al.* 2008). Coding sequence from 23 inbred strains (including CAST/EiJ, WSB/EiJ, PWD/PhJ and
SPRET/EiJ) was reported in Boyden *et al*. (2008). A phylogenetic tree constructed from those sequences,
plus *M. caroli* and rat (not shown) as outgroups, reveals the expected pattern of "deep coalescence"
(**Figure 9C**): the *M. m. domesticus* sequences are paraphyletic, and some (group A) are more similar to
SPRET/EiJ (*M. spretus*) than to their subspecific congeners (group B). Although the level of sequence
divergence in the *Skint1* coding sequence was attributed to ancestral polymorphism maintained by
balancing selection in the original report, selection would not be expected to maintain diversity in both
coding and non-coding sequence equally as we observe in **Figure 9B**. The best explanation for the
observed pattern of diversity at *Skint1* is therefore that groups A and B represent "pseudo-orthologs"
(Koonin 2005) descended from an ancestral duplication followed by subsequent deletion of different
paralogs along different lineages (**Figure 9D**). This conclusion is supported by the structure of the *Skint*
region in the mouse reference genome assembly, which reflects the superposition of many duplications
and rearrangements (**Figure 9E**).

DISCUSSION

In this manuscript we have reconstructed in detail the evolution of a multi-megabase segmental
duplication (SD) in mouse, *R2d2*. Our findings illustrate the challenges involved in accurately
interpreting patterns of polymorphism and divergence within duplicated sequence.

SDs are among the most dynamic loci in mammalian genomes. They are foci for copy-number variation
in populations, but the sequences of individual duplicates beyond those present in the reference genome
are often poorly resolved. Obtaining the sequence of this "missing genome," as we have done for *R2d2*, is
an important prerequisite to understanding the evolution of duplicated loci. Since each paralog follows a
partially independent evolutionary trajectory, individuals in a population may vary both quantitatively
(in the number of copies) and qualitatively (in which copies are retained). Cycles of duplication and loss
may furthermore lead to the fixation of different paralogs along different lineages. This "genomic
revolving door" leaves a signature of polymorphism far in excess of the genome-wide background, due
to coalescence between alleles originating from distinct paralogs. We identify 57 additional regions
covering 0.82% of the mouse genome with this property (**Figure 9** and **Supplementary Table 6**). These
regions have gene density similar to unique sequence and are strongly enriched for genes involved in
odorant sensing, pheromone recognition and immunity that play important roles in social behavior and
speciation (Hurst *et al.* 2001). Excess polymorphism at these loci in mouse has previously been attributed
to some combination of incomplete lineage sorting and diversifying selection (White *et al.* 2009; Keane *et*
al. 2011). Our results suggest that inferences regarding the strength of selection on highly polymorphic
loci in regions of subject to recurrent duplication and loss should be treated with caution.

Accurate deconvolution of recent duplications remains a difficult task that requires painstaking manual
effort. Clone-based and/or single-molecule long-read sequencing remain the gold standard techniques.
But short reads at sufficient depth nonetheless contain a great deal of information. We exploited the
specific properties of *R2d2* in the WSB/EiJ mouse strain — many highly-similar copies of *R2d2* relative to
the single divergent *R2d1* copy — to obtain a nearly complete assembly of *R2d2* from short reads

(**Supplementary Figure 8**). With the sequence of both the *R2d1* and *R2d2* paralogs in hand, we were able
to recognize several remarkable features of *R2d2* that are discussed in detail below.

Long-tract gene conversion.

Previous studies of non-allelic gene conversion in mouse and human have focused either on relatively
small (<5 kbp) intervals within species, or have applied phylogenetic methods to multiple paralogs from a
single reference genome (Dumont and Eichler 2013). This study is the first, to our knowledge, with the
power to resolve large (>5 kbp) non-allelic gene conversion events on an autosome in a population
sample. We identify conversion tracts up to 119 kbp in length, orders of magnitude longer than tracts
arising from allelic conversion events during meiosis. Gene conversion at this scale can rapidly and
dramatically alter paralogous sequences, including — as shown in **Figure 5** — the sequences of essential
protein-coding genes. This process has been implicated as a source of disease alleles in humans (Chen *et*
al. 2007).

Importantly, we were able to identify non-allelic exchanges in *R2d1* as such only because we were aware
of the existence of *R2d2* in other lineages. In this case the transfer of paralogous *R2d2* sequence into *R2d1*
creates the appearance of deep coalescence among *R2d1* sequences. Ignoring the effect of gene conversion
would cause us to overestimate the degree of polymorphism at *R2d1* by an order of magnitude, and
would bias any related estimates of population-genetic parameters (for instance, of effective population
size).

Our data are not sufficient to estimate the rate of non-allelic gene conversion between *R2d2* and
homologous loci. At minimum we have observed two distinct events: one from *R2d2* into *R2d1*, and a
second from *R2d2* into retro-*Cwc22*. From a single conversion event replacing most of *R2d1* with *R2d2*-like
sequence, the remaining shorter conversion tracts could be generated by recombination with *R2d1*
sequences. Because we find converted haplotypes in both *M. m. musculus* and *M. m. domesticus*, the single
conversion event would have had to occur prior to the divergence of the three *M. musculus* subspecies
and subsequently remain polymorphic in the diverged populations. We note that all conversion tracts we
observed are polarized: *R2d2* is always the donor.

The other possibility is that non-allelic gene conversion between *R2d* sequences is recurrent. Recurrent
gene conversion homogenizes duplicate sequences, coupling their evolutionary trajectories ("concerted
evolution", Dover 1982). The absolute sequence divergence (~2%) between *R2d1* and *R2d2* (**Figure 1B**)
argues against a hypothesis of ongoing gene conversion between these loci. However, we cannot rule out
a role for gene conversion in maintaining sequence identity between multiple copies of *R2d* located in
R2d2. This would help explain the reduced diversity within *R2d2* versus *R2d1* (**Figure 8C**). For
sequences with copy number greater than two – such as the *R2d2* casssette in *R2d2HC* alleles – gene
conversion tends to slow the accumulation of new mutations in a copy-number dependent manner
(Nagylaki and Petes 1982). New mutations arising in any single copy are prone to loss not only by drift
but also by being "pasted-over" by gene conversion from the intact copies which outnumber the mutant
(Melamed and Kupiec 1992).

R2d2 is not unlike the male-specific region of the Y chromosome in mouse (Soh *et al.* 2014) and human
(Rozen *et al.* 2003). The large palindromic repeats on chrY are homogenized by frequent non-allelic gene
conversion (Hallast *et al.* 2013) such that they have retained >99% sequence identity to each other even
after millions of years of evolution. Frequent non-allelic gene conversion has also been documented in
arrays of U2 snRNA genes in human (Liao 1997), and in rRNA gene clusters (Eickbush and Eickbush
2007) and centromeric sequences (Schindelhauer 2002; Shi *et al.* 2010) in several species.

Pervasive copy-number variation.

Clusters of segmental duplications have long been known to be hotspots of copy-number variation in populations (Bailey and Eichler 2006; She *et al.* 2008) and *de novo* mutations in pedigrees (Egan *et al.* 2007). Recent large-scale sequencing efforts have revealed the existence of thousands of multiallelic CNVs segregating in human populations (Handsaker *et al.* 2015).

We have surveyed *R2d2* copy number in a large and diverse sample of laboratory and wild mice, and have shown that it varies from 0 to >80 in certain *M. m. domesticus* populations (**Figure 8A**). In a cohort of outbred mice expected to be hemizygous for an *R2d2^HC* allele from WSB/EiJ (33 diploid copies) we estimate that large deletions, >2 Mbp in size, occur at a rate of 3.2% (95% bootstrap CI 1.1% − 6.0%) per generation. This estimate of the mutation rate for CNVs at *R2d2* should be regarded as a lower bound. The power of our copy-number assay to discriminate between copy numbers above ~25 is low, so that the assay is much more sensitive to losses than to gains. Even our lower-bound mutation rate exceeds that of the most common recurrent deletions in human (~1 per 7000 live births) (Turner *et al.* 2007) and is an order of magnitude higher than the most active CNV hotspots described to date in the mouse (Egan *et al.* 2007).

However, the structural mutation rate appears to depend strongly on the diplotype configuration at *R2d2*. As **Figure 1D** shows, individuals heterozygous for an *R2d2^HC* haplotype and an *R2d2*-null haplotype are in fact hemizygous for several megabases of DNA in *R2d2*. This has important consequences. High mutation rates are observed only in the context of populations in which hemizygosity for *R2d2^HC* is common (**Figure 7**): highest in the DO, and to a lesser extent in wild *M. m. domesticus* populations harboring both *R2d2^HC* and *R2d2*-null alleles. Homozygosity for *R2d2^HC* is not associated with mutability: in 8 recombinant inbred lines from the Collaborative Cross which are homozygous for an *R2d2^HC* haplotype, we observed zero new mutations in at least 400 meioses, through both the male and female germline (8 lines × 2 meioses/generation × 25 or more generations of inbreeding). Sex also appears to have a role in determining the mutation rate at *R2d2*: in a pedigree in which all females were hemizygous for *R2d2^HC*, zero new mutations were observed in 1256 meioses (data not shown).

Taken together, these observations hint at a common structural or epigenetic mechanism affecting the resolution of double-strand breaks in large tracts of unpaired (*i.e.* hemizygous) DNA during male meiosis. At least one other study in mouse has hinted that hemizygous SDs on the sex chromosomes are unstable in inter-subspecific hybrids (Scavetta and Tautz 2010). Both the obligate-hemizygous sex chromosomes and large unpaired segments on autosomes are epigenetically marked for transcriptional silencing during male meiotic prophase (Laan 2004; Baarends *et al.* 2005), and are physically sequestered into a structure called the sex body. Repair of double-strand breaks within the sex body is delayed relative to the autosomes (Mahadevaiah *et al.* 2001) and involves a different suite of proteins (Turner *et al.* 2004). We hypothesize that these male-specific pathway(s) are generally error-prone in the presence of non-allelic homologous sequences.

Origin and distribution of an allele subject to meiotic drive.

Females heterozygous for a high- and low-copy allele at *R2d2* preferentially transmit the high-copy allele to progeny via meiotic drive (Didion *et al.* 2015). Meiotic drive can rapidly alter allele frequencies in laboratory and natural populations (Lindholm *et al.* 2016), and we recently showed that high-copy alleles of *R2d2* (*R2d2^HC*) sweep through laboratory and natural populations despite reducing the fitness of heterozygous females (Didion *et al.* 2016). These "selfish sweeps" account for the marked reduction in within-population diversity in the vicinity of *R2d2* (**Figure 8B**).

The present study sheds additional light on the age, origins and fate of *R2d2^HC* alleles. We find that *R2d2^HC*
alleles have a single origin in *M. m. domesticus*. They are present in several different "chromosomal races"
— populations fixed for specific Robertsonian translocations between which gene flow is limited (Hauffe
and Searle 1993) — indicating that they were likely present at intermediate frequency prior to the origin
of the chromosomal races within the past 6,000 to 10,000 years (Nachman *et al.* 1994) and were dispersed
through Europe as mice colonized the continent from the south and east (Boursot *et al.* 1993). The
presence of *R2d2^HC* in a non-*M. m. domesticus* sample (SPRET/EiJ, *M. spretus* from Cadiz, Spain) is best
explained by recent introgression following secondary contact with *M. m. domesticus* (Bonhomme *et al.*
2007; Yang *et al.* 2011).

A new member of the *Cwc22* family.

The duplication that gave rise to *R2d2* also created a new copy of *Cwc22*. Based on our assembly of the
R2d2 sequence, the open reading frame of *Cwc22^R2d2* is intact and encodes a nearly full-length predicted
protein that retains the two key functional domains characteristic of the *Cwc22* family. Inspection of
RNA-seq data from samples with high copy number at *R2d2* reveals several novel transcript isoforms
whose expression appears to be copy-number- and tissue-dependent. In testis, the most abundant
isoform retains an intron containing an ERV insertion (red arrow in **Figure 4**), consistent with the well-
known transcriptional promiscuity in this tissue. The most abundant isoforms in adult brain is unusual in
that its stop codon is in an internal exon which is followed by a 7 kbp 3' UTR in the terminal exon.
Transcripts with a stop codon in an internal exon are generally subject to nonsense-mediated decay
(NMD) triggered by the presence of exon-junction complexes downstream the stop codon. Curiously,
Cwc22 is itself a member of the exon-junction complex (Steckelberg *et al.* 2012).

That an essential gene involved in such a central biochemical pathway should both escape NMD and be
overexpressed more than tenfold is surprising. Preliminary data from the Diversity Outbred population
shows that the *R2d2^HC* allele is associated with elevated levels of both *Cwc22* transcripts and protein in
adult liver (Gary Churchill, personal communication). Further studies will be required to determine the
distribution of transcription and translation of *Cwc22* across isoforms, tissues and developmental stages.

Conclusions and future directions

Our detailed analysis of the evolutionary trajectory of *R2d2* provides insight into the fate of duplicated
sequences over short (within-species) timescales. The exceptionally high mutation rate and low
recombination associated specifically with hemizygous *R2d2^HC* alleles motivate hypotheses regarding the
biochemical mechanisms which contribute to observed patterns of polymorphism at this and similar loci.
Finally, the birth of a new member of the deeply conserved *Cwc22* gene family in *R2d2* provides an
opportunity to test predictions regarding the evolution of young duplicate gene pairs.

METHODS

Mice

Wild *M. musculus* mice used in this study were trapped at a large number of sites across Europe, the
United States, the Middle East, northern India and Taiwan (**Figure 8A**). Trapping was carried out in
accordance with local regulations and with the approval of all relevant regulatory bodies for each locality
and institution. Trapping locations are listed in **Supplementary Table 1**. Most samples have been
previously published (Didion *et al.* 2016).

Tissue samples from the progenitors of the wild-derived inbred strains ZALENDE/EiJ (*M. m. domesticus*),
TIRANO/EiJ (*M. m. domesticus*) and SPRET/EiJ (*M. spretus*) were provided by Muriel Davisson, as
described in Didion *et al.* (2016).

Tissue samples from the high running (HR) selection and intercross lines were obtained as described in
Didion *et al.* (2016).

Female Diversity Outbred mice used for estimating mutation rates at *R2d2* were obtained from the
Jackson Laboratory and housed with a single FVB/NJ male. Progeny were sacrificed at birth by cervical
dislocation in order to obtain tissue for genotyping.

All live laboratory mice were handled in accordance with the IACUC protocols of the University of North
Carolina at Chapel Hill.

DNA preparation

High molecular weight DNA. High molecular weight DNA was obtained for samples genotyped with the
Mouse Diversity Array or subject to whole-genome sequencing. Genomic DNA was extracted from tail,
liver or spleen using a standard phenol-chloroform procedure (Sambrook and Russell 2006). High
molecular weight DNA for most inbred strains was obtained from the Jackson Laboratory, and the
remainder as a generous gift from Francois Bonhomme and the University of Montpellier Wild Mouse
Genetic Repository.

Low molecular weight DNA. Low molecular weight DNA was obtained for samples to be genotyped on the
MegaMUGA array (see "Microarray genotyping" below). Genomic DNA was isolated from tail, liver,
muscle or spleen using Qiagen Gentra Puregene or DNeasy Blood & Tissue kits according to the
manufacturer's instructions.

Whole-genome sequencing and variant discovery

Inbred strains. Sequencing data for inbred strains of mice except ZALENDE/EiJ and LEWES/EiJ was
obtained from the Sanger Mouse Genomes Project website (ftp://ftp-mouse.sanger.ac.uk/current_bams) as
aligned BAM files. Details of the sequencing pipeline are given in Keane *et al.* (2011). Coverage ranged
from approximately 25X to 50X per sample.

The strains LEWES/EiJ and ZALENDE/EiJ were sequenced at the University of North Carolina High-
Throughput Sequencing Facility. Libraries were prepared from high molecular weight DNA using the
Illumina TruSeq kit and insert size approximately 250 bp, and 2x100bp paired-end reads were generated
on an Illumina HiSeq 2000 instrument. LEWES/EiJ was sequenced to approximately 12X coverage and
ZALENDE/EiJ to approximately 20X. Alignment was performed as in Keane *et al.* (2011).

Wild mice. Whole-genome sequencing data from 26 wild *M. m. domesticus* individuals described in Pezer
et al. (2015) was downloaded from ENA under accession #PRJEB9450. Coverage ranged from
approximately 12X to 20X per sample. An additional two wild *M. m. domesticus* individuals, IT175 and
ES446, were sequenced at the University of North Carolina to approximate coverage 8X each. Raw reads
from an additional 10 wild *M. m. castaneus* described in Halligan *et al.* (2013), sequenced to approximately
20X each, were downloaded from ENA under accession #PRJEB2176. Reads for a single *Mus caroli*
individual sequenced to approximately 40X were obtained from ENA under accession #PRJEB2188.
Reads for each sample were realigned to the mm10 reference using bwa-mem v0.7.12 with default
parameters (Li 2013). Optical duplicates were removed with samblaster (Faust and Hall 2014).

Variant discovery. Polymorphic sites on chromosome 2 in the vicinity of *R2d2* (**Figure 8B**) were called
using freebayes v0.9.21-19-gc003c1e (Garrison and Marth 2012) with parameters "—standard-filters"
using the Sanger Mouse Genomes Project VCF files as a list of known sites (parameter "—@"). Raw calls
were filtered to have quality score > 30, root mean square mapping quality > 20 (for both reference and
alternate allele calls) and at most 2 alternate alleles.

Copy-number estimation

R2d copy number was estimated using qPCR as described in Didion *et al.* (2016). Briefly, we used
commercial TaqMan assays against intron-exon boundaries in *Cwc22* (Life Technologies assay numbers
Mm00644079_cn and Mm00053048_cn) to determine copy number relative to reference genes *Tert* (cat. no.
4458368, for target Mm00644079_cn) or *Tfrc* (cat. no. 4458366, for target Mm00053048_cn). Cycle
thresholds for *Cwc22* relative to the reference gene were normalized across assay batches using linear
mixed models with batch and target-reference pair treated as random effects. Control samples with
known haploid *R2d* copy numbers of 1 (C57BL/6J), 2 (CAST/EiJ), 17 (WSB/EiJxC57BL/6J)F_1 and 34
(WSB/EiJ) were included in each batch.

Samples were classified as having 1, 2 or >2 haploid copies of *R2d* using linear discriminant analysis. The
classifier was trained on the normalized cycle thresholds of the control samples from each plate, whose
precise integer copy number is known, and applied to the remaining samples.

Microarray genotyping

Genome-wide genotyping was performed using MegaMUGA, the second version of the Mouse Universal
Genotyping Array platform (Neogen/GeneSeek, Lincoln, NE) (Morgan *et al.* 2016). Genotypes were called
using the GenCall algorithm implemented in the Illumina BeadStudio software (Illumina Inc, Carlsbad,
CA). For quality control we computed, for each marker i on the array: $S\_i = X_i + Y_i$, where X_i and Y_i are
the normalized hybridization intensities for the two alleles. The expected distribution of Si was computed
from a large set of reference samples. We excluded arrays for which the distribution of S_i was
substantially shifted from this reference; in practice, failed arrays can be trivially identified in this manner
(Morgan *et al.* 2016). Access to MegaMUGA genotypes was provided by partnership between the
McMillan and Pardo-Manuel de Villena labs and the UNC Systems Genetics Core Facility.

Additional genotypes for inbred strains and wild mice from the Mouse Diversity Array were obtained
from Yang *et al.* (2011).

De novo **assembly of *R2d2***

Raw whole-genome sequencing reads for WSB/EiJ from the Sanger Mouse Genomes Project were
converted to a multi-string Burrows-Wheeler transform and associated FM-index (msBWT) (Holt and
McMillan 2014) using the msbwt v0.1.4 Python package (https://pypi.python.org/pypi/msbwt). The
msBWT and FM-index implicitly represent a suffix array of sequencing to provide efficient queries over
arbitrarily large string sets. Given a seed k-mer present in that string set, this property can be exploited to
rapidly construct a de Bruijn graph which can in turn be used for local *de novo* assembly of a target
sequence (**Supplementary Figure 8A**). The edges in that graph can be assigned a weight (corresponding
to the number of reads containing the $k + 1$-mer implied by the edge) which can be used to evaluate
candidate paths when the graph branches (**Supplementary Figure 8B**).

R2d2 was seeded with the 30 bp sequence (TCTAGAGCATGAGCCTCATTTATCATGCCT) at the
proximal boundary of *R2d1* in the GRCm38/mm10 reference genome. A single linear contig was

assembled by "walking" through the local de Bruijn graph. Because WSB/EiJ has ~33 copies of *R2d2* and a
single copy of *R2d1*, any branch point in the graph which represents a paralogous variant should having
outgoing edges with weights differing by a factor of approximately 33. Furthermore, when two (or more)
branch points occur within less than the length of a read, it should be possible to "phase" the underlying
variants by following single reads through both branch points (**Supplementary Figure 8B**). We used
these heuristics to assemble the sequence of *R2d2* (corresponding to the higher-weight path through the
graph) specifically.

After assembling a chunk of approximately 500 bp the contig was checked for colinearity with the
reference sequence (*R2d1*) using BLAT and CLUSTAL-W2 (using the EMBL-EBI web server:
http://www.ebi.ac.uk/Tools/msa/clustalw2/).

Repetitive elements such as retroviruses are refractory to assembly with our method. Upon traversing
into a repetitive element, the total edge weight (total number of reads) and number of branch points
(representing possible linear assembled sequences) in the graph become large. It was sometimes possible
to assemble a fragment of a repetitive element at its junction with unique sequence but not to assemble
unambiguously across the repeat. Regions of unassembleable sequence were marked with blocks of Ns,
and assembly re-seeded using a nearby *k*-mer from the reference sequence. The final contig is provided
in FASTA format in **Supplementary File 1**.

The final contig was checked against its source msBWT by confirming that each 30-mer in the contig
which did not contain an N was supported by at least 60 reads. A total of 16 additional haplotypes in 8
regions of *R2d* totaling 16.9 kbp (**Supplementary Table 6**) were assembled in a similar fashion, using the
WSB *R2d2* contig and the *R2d1* reference sequence as guides. Multiple sequence alignments from these
regions are provided in **Supplementary File 1**.

Sequence analysis of *R2d2* contig

Pairwise alignment of R2d *paralogs.* The reference *R2d1* sequence and our *R2d2* contig were aligned using
LASTZ v1.03.54 (http://www.bx.psu.edu/~rsharris/lastz/) with parameters "—step=10 —seed=match12 —
notransition —exact=20 —notrim —identity=95".

Transposable element (TE) content. The *R2d2* contig was screened for TE insertions using the RepeatMasker
web server (http://www.repeatmasker.org/cgi-bin/WEBRepeatMasker) with species set to "mouse" and
default settings otherwise. As noted previously, we could not assemble full-length repeats, but the
fragments we could assemble at junctions with unique sequence allowed identification of some candidate
TEs to the family level. *R2d1*-specific TEs were defined as TEs annotated in the RepeatMasker track at the
UCSC Genome Browser with no evidence (no homologous sequence, and no Ns) at the corresponding
position in the *R2d2* contig. Candidate *R2d2*-specific TEs were defined as gaps >= 100 bp in size in the
alignment to *R2d1* for which the corresponding *R2d2* sequence was flagged by RepeatMasker.

Gene conversion tracts. To unambiguously define gene conversion events without confounding from
paralogous sequence, we examined 15 wild *M. m. domesticus* samples and 37 laboratory strains with
evidence of 2 diploid copies of *R2d*. We confirmed that these copies of *R2d* were located at *R2d1* by
finding read pairs spanning the junction between *R2d1* and neighboring sequence. Gene conversion tracts
were delineated as clusters of derived alleles shared with *R2d2*. Using a pairwise of alignment of *R2d2*
and *R2d1* we identified single-nucleotide variants between the two sequences, and queried those sites in
aligned reads for *Mus caroli*. If the *Mus caroli* and *R2d1* shared an allele, we recorded the site as a derived
allele informative for the presence of *R2d2*. We used the resulting list of 1,411 informative sites to query
aligned reads for the samples of interest and recorded, for each site and each sample, whether the derived

allele (*R2d2*), ancestral allele (*R2d1*) or both alleles were present. Conversion tracts were then identified by

manual inspection. Boundaries of conversion tracts were defined at approximately the midpoint between

the first *R2d1*- (or *R2d2*-) specific variant and the last *R2d2*- (or *R2d1*-) specific variant.

Sequence diversity in R2d1 *and* R2d2. Assembling individual copies of *R2d2* is infeasible in high-copy

samples. Instead we treated each *R2d* unit as an independent sequence and used the number of

segregating sites to estimate sequence diversity. Segregating sites were defined as positions in a collection

of alignments (BAM files) with evidence of an alternate allele. To identify segregating sites we used

freebayes v0.9.21-19-gc003c1e (Garrison and Marth 2012) with parameters "-ui -Kp 20 —use-best-n-alleles

2 -m 8". These parameters treat each sample as having ploidy up to 20, impose an uninformative prior on

genotype frequencies, and limit the algorithm to the discovery of atomic variants (SNVs or short indels,

not multinucleotide polymorphisms or other complex events) with at most 2 alleles at each segregating

site. Sites in low-complexity sequence (defined as Shannon entropy < 1.6 in the 30 bp window centered on

the site) or within 10 bp of another variant site were further masked, to minimize spurious calls due to

ambiguous alignment of indels. To avoid confounding with the retrocopies of *Cwc22* outside *R2d*, coding

exons of *Cwc22* were masked. Finally, sites corresponding to an unaligned or gap position in the pairwise

alignment between *R2d1* and *R2d2* were masked.

To compute diversity in *R2d1* we counted segregating sites in 12 wild *M. m. domesticus* samples with 2

diploid copies of *R2d* (total of 24 sequences), confirmed to be in *R2d1* by the presence of read pairs

spanning the junction between *R2d1* and neighboring sequence. To compute diversity in *R2d2*, we

counted segregating sites in 14 wild *M. m. domesticus* samples with >2 diploid copies of *R2d* (range 3 — 83

per sample; total of 406 sequences) but excluded sites corresponding to variants among *R2d1* sequences.

Remaining sites were phased to *R2d2* by checking for the presence of a 31-mer containing the site and the

nearest *R2d1*-vs-*R2d2* difference in the raw reads for each sample using the corresponding msBWT.

Sequence diversity was then computed using Watterson's estimator (Watterson 1975), dividing by the

number of alignable bases (128973) to yield a per-site estimate. Standard errors were estimated by 100

rounds of resampling over the columns in the *R2d1*-vs-*R2d2* alignment.

Analyses of *Cwc22* expression

RNA-seq read alignment. Expression of *Cwc22* was examined in adult whole brain using data from

Crowley *et al.* (2015), SRA accession #SRP056236. Paired-end reads (2x100bp) were obtained from 8

replicates each of 3 inbred strains: CAST/EiJ, PWK/PhJ and WSB/EiJ. Raw reads were aligned to the

mm10 reference using STAR v2.4.2a (Dobin *et al.* 2012) with default parameters for paired-end reads.

Alignments were merged into a single file per strain for further analysis. Expression in adult testis was

examined in 23 wild-derived inbred strains from Phifer-Rixey *et al.* (2014) SRA accession #PRJNA252743.

Single-end reads (76bp) were aligned to the mm10 genome with STAR using default parameters for

single-end, non-strand-specific reads.

Transcript assembly. Read alignments were manually inspected to assess support for *Cwc22* isoforms in

Ensembl v83 annotation. To identify novel isoforms in *R2d2*, we applied the Trinity v0.2.6 pipeline

(Grabherr *et al.* 2011) to the subset of reads from WSB/EiJ which could be aligned to *R2d1* plus their mates

(a set which represents a mixture of *Cwc22^R2d1* and *Cwc22^R2d2* reads). De novo transcripts were aligned

both to the mm10 reference and to the *R2d2* contig using BLAT, and were assigned to *R2d1* or *R2d2* based

on sequence similarity. Because expression from *R2d2* is high in WSB/EiJ, *R2d2*-derived transcripts

dominated the assembled set. Both manual inspection and the Trinity assembly indicated the presence of

retained introns and an extra 3' exon, as described in the **Results**. To obtain a full set of Cwc22 transcripts

including those of both *R2d1* and *R2d2* origin, we supplemented the *Cwc22* transcripts in Ensembl v83

with their paralogs from *R2d2* as determined by a strict BLAT search against the *R2d2* contig. We
manually created additional transcripts reflecting intron-retention and 3′ extension events described
above, and obtained their sequence from the *R2d2* contig.

Abundance estimation. Relative abundance of *Cwc22* paralogs was estimated using kallisto v0.42.3 (Bray *et*
al. 2015) with parameters "—bias" (to estimate and correct library-specific sequence-composition biases).
The transcript index used for pseudoalignment and quantification included only the *Cwc22* targets.

Phylogenetic analyses

Tree for R2d. Multiple sequence alignments for 8 the regions in **Supplementary Figure 5** were generated
using MUSCLE (Edgar 2004) with default parameters. The resulting alignments were manually trimmed
and consecutive gaps removed. Phylogenetic trees were inferred with RAxML v8.1.9 (Stamatakis 2014)
using the GTR+gamma model with 4 rate categories and *M. caroli* as an outgroup. Uncertainty of tree
topologies was evaluated using 100 bootstrap replicates.

Divergence time. The time of the split between *R2d1* and *R2d2* was estimated using the Bayesian method
implemented in BEAST v1.8.1r6542 (Drummond *et al.* 2012). We assumed a divergence time for *M. caroli*
of 5 Mya and a strict molecular clock, and analyzed the concatenated alignment for our *de novo* assembled
regions under the GTR+gamma model with 4 rate categories and allowance for a proportion of invariant
sites. The chain was run for 10 million iterations with trees sampled every 1000 iterations.

Local phylogeny around R2d2. Genotypes for 173 SNPs in the region surrounding *R2d2* (chr2: 83 — 84 Mb)
were obtained for 90 individuals representing both laboratory and wild mice genotyped with the Mouse
Diversity Array (Yang *et al.* 2011) (**Supplementary Table 1**). Individuals with evidence of heterozygosity
(>3 heterozygous calls) were excluded to avoid ambiguity in phylogenetic inference. A distance matrix
for the remaining 62 samples was created by computing the proportion of alleles shared identical by state
between each pair of samples. A neighbor-joining tree (**Figure 3C**) was inferred from the distance matrix
and rooted at the most recent common ancestor of the *M. musculus*- and non-*M. musculus* samples.

Cwc22 *coding sequences*. To create the tree of Cwc22 coding sequences, we first obtained the sequences of
all its paralogs in mouse. The coding sequence of *Cwc22^{R2d1}* (RefSeq transcript NM_030560.5) was
obtained from the UCSC Genome Browser and aligned to our *R2d2* contig with BLAT to extract the exons
of *Cwc22^{R2d2}*. The coding sequence of retro-*Cwc22* (genomic sequence corresponding to GenBank cDNA
AK145290) was obtained from the UCSC Genome Browser. Coding and protein sequences of *Cwc22*
homologs from non-*M. musculus* species were obtained from Ensembl (Cunningham *et al.* 2014). The
sequences were aligned with MUSCLE and manually trimmed, and a phylogenetic tree estimated as
described above.

We observed that the branches in the rodent clade of the *Cwc22* tree appeared to be longer than branches
for other taxa. We used PAML (Yang *et al.* 2007) to test the hypothesis that *Cwc22* is under relaxed
purifying selection in rodents using the branch-site model (null model "model = 2, NSsites = 2, fix_omega
= 1"; alternative model "model = 2, NSsites = 2, omega = 1, fix_omega = 1") as described in the PAML
manual. This is a test of difference in evolutionary rate on a "foreground" branch (ω_1) — in our case, the
rodent clade — relative to the tree-wide "background" rate (ω_0). The distribution of the test statistic is an
even mixture of a χ^2 distribution with 1 df and a point mass at zero; to obtain the *p*-value, we calculated
the quantile of the χ^2 distribution with 1 df and divided by 2.

Genome-wide sequence divergence

The msBWT of a collection of whole-genome sequencing reads can be used to estimate the divergence between the corresponding template sequence (*i.e.* genome) and a reference sequence as follows. Non-overlapping k-mers from the reference sequence are queried against the msBWT. (The value of k is chosen such that nearly all k-mers drawn from genomic sequence exclusive of repetitive elements.) Let x be the count of reads containing an exact match to the k-mer or its reverse complement. If the template and reference sequence are identical, standard theory for shotgun sequencing (Lander and Waterman 1988) holds that $P(x > 0|\lambda) = 1 - e^{-\lambda}$, where λ is the average sequencing coverage. We assume $P(x > 0|\lambda) \approx 1$, which is satisfied in practice for high-coverage sequencing.

However, if a haploid template sequence contains at least one variant (versus the reference) within the queried k-mer, it will be the case that $x = 0$. We use this fact and assume that mutations arise along a sequence via a Poisson process to estimate the rate parameter α from the proportion of k-mers that have read count zero. Let m be the number of mutations arising between a target and reference in a window of length L, and y the number of k-mers in that window with nonzero read count. Then $P(m = 0|\alpha) = e^{-\alpha L}$ and a simple estimator for α is $\hat{\alpha} = -\log\left(\frac{1-y}{L}\right)$.

Interpretation of α is straightforward in the haploid case: it is the per-base rate of sequence divergence between the template sequence and the reference sequence. In the diploid case it represents a lower bound on the sequence divergence of the two homologous chromosomes.

We applied this estimator with $k = 31$ and $L = 1000 \times k = 31$ kbp to msBWTs for 7 inbred strains (3 *M. m. domesticus*, 1 *M. m. musculus*, 1 *M. m. castaneus*, 1 *M. spretus*, 1 *M. caroli*) and 2 wild *M. m. domesticus* individuals (IT175, ES446) using the GRCm38/mm10 mouse reference sequence as the source of k-mer queries. As shown in **Figure 9A**, the mode of the distribution of divergence values matches what is expected based on the ancestry of the samples with respect to the reference. To identify divergent regions, we fit a discrete-time hidden Markov model (HMM) to the windows divergence values. The HMM had two hidden states: "normal" sequence, with emission distribution $N(0.005, 0.005)$ and initial probability 0.99; and "divergent" sequence, with emission distribution $N(0.02, 0.005)$ and initial probability 0.01. The transmission probability between states was 1×10^{-5}. Posterior decodings were obtained via the Viterbi algorithm, as implemented in the R package HiddenMarkov (https://cran.r-project.org/package=HiddenMarkov).

Significance tests for overlap with genomic features were performed using the resampling algorithm implemented in the Genomic Association Tester (GAT) package for Python (https://pypi.python.org/pypi/gat). Segmental duplications were obtained from the genomicSuperDups table of the UCSC Genome Browser and genes from Ensembl v83 annotation.

Analyses of recombination rate at *R2d2*

To test the effect of *R2d2* copy number on local recombination rate examined recombination events accumulated during the first 16 generations of breeding of the Diversity Outbred (DO) population, in which the high-copy *R2d2* allele from WSB/EiJ is segregating. Founder haplotype reconstructions were obtained for 4,640 DO individuals reported in (Didion *et al.* 2016), and recombination events were identified as junctions between founder haplotypes. We compared the frequency of junctions involving a WSB/EiJ haplotype to junctions not involving a WSB/EiJ haplotype over the region chr2: 75-90 Mb. Within each generation we also tested for differences in the lengths of haplotype blocks overlapping *R2d2* using one-sided Wilcoxon rank-sum tests (alternative hypothesis: WSB/EiJ haplotypes longer than

others). Resulting p-values were subject to Bonferroni correction: for nominal significance level $\alpha = 0.01$,

the corrected threshold is $p = \frac{0.01}{12} = 8.3 \times 10^{-4}$.

We also estimated the difference between observed and expected recombination fraction in 11

experimental crosses in which one of the parental lines was segregating for a high-copy allele at *R2d2*. We

obtained expected recombination fractions from the standard mouse genetic map (Cox *et al.* 2009), which

was constructed from crosses between strains lacking *R2d2^HC* alleles. Genotype data was obtained from

The Jackson Laboratory's Mouse Phenome Database QTL Archive

(http://phenome.jax.org/db/q?rtn=qtl/home). Recombination fractions were calculated using R/qtl

(http://rqtl.org/). Confidence intervals for difference between observed and expected recombination

fractions were calculated by 100 iterations of nonparametric bootstrapping over individuals in each

dataset.

Results of these analyses are presented in **Supplementary Figure 7**.

DATA AVAILABILITY

804 All *de novo* assemblies used in this study are included in **Supplementary File 1**. The data structures on

805 which the assemblies are based, and the interactive computational tools used for assembly, are publicly

806 available at http://www.csbio.unc.edu/CEGSseq/index.py?run=MsbwtTools. Genotype data used for

807 mapping the location of R2d2 and defining associated haplotypes (**Figure 3B-C**) are available on Dryad

808 (accession XXX).

ACKNOWLEDGMENTS

We thank all the scientists and personnel who collected and processed the wild mouse samples used in

this study. In particular we thank Francois Bonhomme for providing samples from wild-derived inbred

strains housed at the University of Montpellier Wild Mouse Genetic Repository, and Ted Garland for

providing tissue samples from the HR selection lines and related crosses. This work was supported by

National Institutes of Health grants P50GM076468 (FPMdV), U19AI100625 (FPMdV, APM),

F30MH103925 (APM), T32GM067553 (JPD, APM), and by Vaadia-BARD Postdoctoral Fellowship Award

FI-12 478-13 to LY. Additional support was provided by Cancer Research UK, the European Research

Council, EMBO Young Investigator Programme (DTO), European Molecular Biology Laboratory (DTO,

PF) and the Wellcome Trust (WT095908) (PF) and (WT098051) (PF, DTO); and finally by the European

Union's Seventh Framework Programme (FP7/2007-2013) under grant agreement HEALTH-F4-2010-

241504 (EURATRANS).

REFERENCES

Baarends W. M., Wassenaar E., Laan R. van der, Hoogerbrugge J., Sleddens-Linkels E., Hoeijmakers J. H.

J., Boer P. de, Grootegoed J. A., 2005 Silencing of unpaired chromatin and histone H2A ubiquitination in

mammalian meiosis. Mol Cell Biol **25**: 1041–1053.

Bailey J. A., Eichler E. E., 2006 Primate segmental duplications: Crucibles of evolution, diversity and

disease. Nat Rev Genet **7**: 552–564.

Bonhomme F., Rivals E., Orth A., Grant G. R., Jeffreys A. J., Bois P. R., 2007 Species-wide distribution of
highly polymorphic minisatellite markers suggests past and present genetic exchanges among house
mouse subspecies. Genome Biol **8**: R80.

Boursot P., Auffray J. C., Britton-Davidian J., Bonhomme F., 1993 The evolution of house mice. Annu Rev
Ecol Syst **24**: 119–152.

Boyden L. M., Lewis J. M., Barbee S. D., Bas A., Girardi M., Hayday A. C., Tigelaar R. E., Lifton R. P., 2008
Skint1, the prototype of a newly identified immunoglobulin superfamily gene cluster, positively selects
epidermal T cells. Nat Genet **40**: 656–662.

Bray N., Pimentel H., Melsted P., Patcher L., 2015 Near-optimal RNA-seq quantification. arXiv:
1505.02710 [q-bio.QM]

Chen J.-M., Cooper D. N., Chuzhanova N., Férec C., Patrinos G. P., 2007 Gene conversion: Mechanisms,
evolution and human disease. Nat Rev Genet **8**: 762–775.

Chevret P., Veyrunes F., Britton-Davidian J., 2005 Molecular phylogeny of the genus Mus (Rodentia:
Murinae) based on mitochondrial and nuclear data. Biol J Linn Soc **84**: 417–427.

Cole F., Keeney S., Jasin M., 2010 Comprehensive, fine-scale dissection of homologous recombination
outcomes at a hot spot in mouse meiosis. Mol Cell **39**: 700–710.

Cole F., Baudat F., Grey C., Keeney S., Massy B. de, Jasin M., 2014 Mouse tetrad analysis provides insights
into recombination mechanisms and hotspot evolutionary dynamics. Nat Genet **46**: 1072–1080.

Collaborative Cross Consortium, 2012 The genome architecture of the Collaborative Cross mouse menetic
reference population. Genetics **190**: 389–401.

Cox A., Ackert-Bicknell C. L., Dumont B. L., Ding Y., Bell J. T., Brockmann G. A., Wergedal J. E., Bult C.,
Paigen B., Flint J., Tsaih S. W., Churchill G. A., Broman K. W., 2009 A new standard genetic map for
the laboratory mouse. Genetics **182**: 1335–1344.

Crowley J. J., Zhabotynsky V., Sun W., Huang S., Pakatci I. K., Kim Y., Wang J. R., Morgan A. P., Calaway
J. D., Aylor D. L., Yun Z., Bell T. A., Buus R. J., Calaway M. E., Didion J. P., Gooch T. J., Hansen S. D.,
Robinson N. N., Shaw G. D., Spence J. S., Quackenbush C. R., Barrick C. J., Nonneman R. J., Kim K.,
Xenakis J., Xie Y., Valdar W., Lenarcic A. B., Wang W., Welsh C. E., Fu C.-P., Zhang Z., Holt J., Guo Z.,
Threadgill D. W., Tarantino L. M., Miller D. R., Zou F., McMillan L., Sullivan P. F., Pardo-Manuel de
Villena F., 2015 Analyses of allele-specific gene expression in highly divergent mouse crosses identifies
pervasive allelic imbalance. Nat Genet **47**: 353–360.

Cunningham F., Amode M. R., Barrell D., Beal K., Billis K., Brent S., Carvalho-Silva D., Clapham P.,
Coates G., Fitzgerald S., Gil L., Giron C. G., Gordon L., Hourlier T., Hunt S. E., Janacek S. H., Johnson N.,
Juettemann T., Kahari A. K., Keenan S., Martin F. J., Maurel T., McLaren W., Murphy D. N., Nag R.,
Overduin B., Parker A., Patricio M., Perry E., Pignatelli M., Riat H. S., Sheppard D., Taylor K., Thormann
A., Vullo A., Wilder S. P., Zadissa A., Aken B. L., Birney E., Harrow J., Kinsella R., Muffato M., Ruffier M.,
Searle S. M. J., Spudich G., Trevanion S. J., Yates A., Zerbino D. R., Flicek P., 2014 Ensembl 2015. Nucleic
Acids Res **43**: D662–D669.

Dallas J. F., 1992 Estimation of microsatellite mutation rates in recombinant inbred strains of mouse.
Mamm Genome **3**: 452–456.

Demuth J. P., De Bie T., Stajich J. E., Christianini N., Hahn M. W., 2006 The evolution of mammalian gene
families. PLoS ONE **1**: e85.

Didion J. P., Morgan A. P., Clayshulte A. M.-F., Mcmullan R. C., Yadgary L., Petkov P. M., Bell T. A., Gatti
D. M., Crowley J. J., Hua K., Aylor D. L., Bai L., Calaway M., Chesler E. J., French J. E., Geiger T. R.,
Gooch T. J., Garland T., Harrill A. H., Hunter K., McMillan L., Holt M., Miller D. R., O'Brien D. A., Paigen
K., Pan W., Rowe L. B., Shaw G. D., Simecek P., Sullivan P. F., Svenson K. L., Weinstock G. M., Threadgill
D. W., Pomp D., Churchill G. A., Pardo-Manuel de Villena F., 2015 A multi-megabase copy number gain
causes maternal transmission ratio distortion on mouse chromosome 2. PLoS Genet **11**: e1004850.

Didion J. P., Morgan A. P., Yadgary L., Bell T. A., McMullan R. C., Solorzano L. O. de, Britton-Davidian J.,
Bult C. J., Campbell K. J., Castiglia R., Ching Y.-H., Chunco A. J., Crowley J. J., Chesler E. J., Förster D. W.,
French J. E., Gabriel S. I., Gatti D. M., Garland T., Giagia-Athanasopoulou E. B., Giménez M. D., Grize S.
A., Gündüz I., Holmes A., Hauffe H. C., Herman J. S., Holt J. M., Hua K., Jolley W. J., Lindholm A. K.,
López-Fuster M. J., Mitsainas G., Luz Mathias M. da, McMillan L., Ramalhinho M. G., Rehermann B.,
Rosshart S. P., Searle J. B., Shiao M.-S., Solano E., Svenson K. L., Thomas-Laemont P., Threadgill D. W.,
Ventura J., Weinstock G. M., Pomp D., Churchill G. A., Pardo-Manuel de Villena F., 2016 R2d2 drives
selfish sweeps in the house mouse. Mol Biol Evol: msw036.

Dobin A., Davis C. A., Schlesinger F., Drenkow J., Zaleski C., Jha S., Batut P., Chaisson M., Gingeras T. R.,
2012 STAR: Ultrafast universal RNA-seq aligner. Bioinformatics **29**: 15–21.

Dopman E. B., Hartl D. L., 2007 A portrait of copy-number polymorphism in drosophila melanogaster.
Proceedings of the National Academy of Sciences **104**: 19920–19925.

Dover G., 1982 Molecular drive: A cohesive mode of species evolution. Nature **299**: 111–117.

Drummond A. J., Suchard M. A., Xie D., Rambaut A., 2012 Bayesian phylogenetics with BEAUti and the
BEAST 1.7. Mol Biol Evol **29**: 1969–1973.

Dumont B. L., Eichler E. E., 2013 Signals of historical interlocus gene conversion in human segmental
duplications. PLoS ONE **8**: e75949.

Edgar R. C., 2004 MUSCLE: Multiple sequence alignment with high accuracy and high throughput.
Nucleic Acids Res **32**: 1792–1797.

Egan C. M., Sridhar S., Wigler M., Hall I. M., 2007 Recurrent DNA copy number variation in the
laboratory mouse. Nat Genet **39**: 1384–1389.

Eickbush T. H., Eickbush D. G., 2007 Finely orchestrated movements: Evolution of the ribosomal RNA
genes. Genetics **175**: 477–485.

Faust G. G., Hall I. M., 2014 SAMBLASTER: Fast duplicate marking and structural variant read
extraction. Bioinformatics **30**: 2503–2505.

Garrison E., Marth G., 2012 Haplotype-based variant detection from short-read sequencing. arXiv.

Goodman M., Czelusniak J., Moore G. W., Romero-Herrera A. E., Matsuda G., 1979 Fitting the gene
lineage into its species lineage, a parsimony strategy illustrated by cladograms constructed from globin
sequences. Syst Biol **28**: 132–163.

Grabherr M. G., Haas B. J., Yassour M., Levin J. Z., Thompson D. A., Amit I., Adiconis X., Fan L.,
Raychowdhury R., Zeng Q., Chen Z., Mauceli E., Hacohen N., Gnirke A., Rhind N., Palma F. di, Birren B.
W., Nusbaum C., Lindblad-Toh K., Friedman N., Regev A., 2011 Full-length transcriptome assembly from
RNA-seq data without a reference genome. Nat Biotechnol **29**: 644–652.

Hallast P., Balaresque P., Bowden G. R., Ballereau S., Jobling M. A., 2013 Recombination dynamics of a
human Y-chromosomal palindrome: Rapid GC-biased gene conversion, multi-kilobase conversion tracts,
and rare inversions. PLoS Genet **9**: e1003666.

Halligan D. L., Kousathanas A., Ness R. W., Harr B., Eöry L., Keane T. M., Adams D. J., Keightley P. D.,
2013 Contributions of protein-coding and regulatory change to adaptive molecular evolution in murid
rodents. PLoS Genet **9**: e1003995.

Handsaker R. E., Doren V. V., Berman J. R., Genovese G., Kashin S., Boettger L. M., McCarroll S. A., 2015
Large multiallelic copy number variations in humans. Nat Genet **47**: 296–303.

Hauffe H. C., Searle J. B., 1993 Extreme karyotypic variation in a Mus musculus domesticus hybrid zone:
The tobacco mouse story revisited. Evolution **47**: 1374.

Holt J., McMillan L., 2014 Merging of multi-string BWTs with applications. Bioinformatics **30**: 3524–3531.

Hurles M., 2002 Are 100,000 "SNPs" useless? Science **298**: 1509.

Hurst J. L., Payne C. E., Nevison C. M., Marie A. D., Humphries R. E., Robertson D. H. L., Cavaggioni A.,
Beynon R. J., 2001 Individual recognition in mice mediated by major urinary proteins. Nature **414**: 631–
634.

Keane T. M., Goodstadt L., Danecek P., White M. A., Wong K., Yalcin B., Heger A., Agam A., Slater G.,
Goodson M., Furlotte N. A., Eskin E., Nellåker C., Whitley H., Cleak J., Janowitz D., Hernandez-Pliego P.,
Edwards A., Belgard T. G., Oliver P. L., McIntyre R. E., Bhomra A., Nicod J., Gan X., Yuan W., Weyden L.
van der, Steward C. A., Bala S., Stalker J., Mott R., Durbin R., Jackson I. J., Czechanski A., Guerra-
Assunção J. A., Donahue L. R., Reinholdt L. G., Payseur B. A., Ponting C. P., Birney E., Flint J., Adams D.
J., 2011 Mouse genomic variation and its effect on phenotypes and gene regulation. Nature **477**: 289–294.

Koonin, E. V., 2005 Orthologs, paralogs and evolutionary genomics. Annu Rev Genet **39**: 309–338.

Laan R. van der, 2004 Ubiquitin ligase Rad18Sc localizes to the XY body and to other chromosomal
regions that are unpaired and transcriptionally silenced during male meiotic prophase. J Cell Sci **117**:
5023–5033.

Lander E. S., Waterman M. S., 1988 Genomic mapping by fingerprinting random clones: A mathematical
analysis. Genomics **2**: 231–239.

Li H., 2006 TreeFam: A curated database of phylogenetic trees of animal gene families. Nucleic Acids Res
34: D572–D580.

Li H., 2013 Aligning sequence reads, clone sequences and assembly contigs with BWA-MEM. arXiv.

Liao D., 1997 Concerted evolution of the tandemly repeated genes encoding human U2 snRNA (the
RUN2 locus) involves rapid intrachromosomal homogenization and rare interchromosomal gene
conversion. EMBO J **16**: 588–598.

Lindholm A. K., Dyer K. A., Firman R. C., Fishman L., Forstmeier W., Holman L., Johannesson H., Knief
U., Kokko H., Larracuente A. M., Manser A., Montchamp-Moreau C., Petrosyan V. G., Pomiankowski A.,
Presgraves D. C., Safronova L. D., Sutter A., Unckless R. L., Verspoor R. L., Wedell N., Wilkinson G. S.,
Price T. A., 2016 The ecology and evolutionary dynamics of meiotic drive. Trends Ecol Evol.

Liu E. Y., Morgan A. P., Chesler E. J., Wang W., Churchill G. A., Villena F. P.-M. de, 2014 High-resolution
sex-specific linkage maps of the mouse reveal polarized distribution of crossovers in male germline.
Genetics **197**: 91–106.

Lynch M., 2000 The evolutionary fate and consequences of duplicate genes. Science **290**: 1151–1155.

Mahadevaiah S. K., Turner J. M., Baudat F., Rogakou E. P., Boer P. de, Blanco-Rodríguez J., Jasin M.,
Keeney S., Bonner W. M., Burgoyne P. S., 2001 Nat Genet **27**: 271–276.

Melamed C., Kupiec M., 1992 Effect of donor copy number on the rate of gene conversion in the yeast
Saccharomyces cerevisiae. Mol Genet Genom **235**: 97–103.

Moran P., 1975 Wandering distributions and the electrophoretic profile. Theor Popul Biol **8**: 318–330.

Morgan A. P., Fu C.-P., Kao C.-Y., Welsh C. E., Didion J. P., Yadgary L., Hyacinth L., Ferris M. T., Bell T.
A., Miller D. R., Giusti-Rodriguez P., Nonneman R. J., Cook K. D., Whitmire J. K., Gralinski L. E., Keller
M., Attie A. D., Churchill G. A., Petkov P., Sullivan P. F., Brennan J. R., McMillan L., Pardo-Manuel de
Villena F., 2016 The Mouse Universal Genotyping Array: From substrains to subspecies. G3 **6**.

Muffato M., Louis A., Poisnel C. E., Crollius H. R., 2010 Genomicus: A database and a browser to study
gene synteny in modern and ancestral genomes. Bioinformatics **26**: 1119–1121.

Nachman M. W., Boyer S. N., Searle J. B., Aquadro C. F., 1994 Mitochondrial DNA variation and the
evolution of robertsonian chromosomal races of house mice, Mus domesticus. Genetics **136**: 1105–1120.

Nagylaki T., Petes T. D., 1982 Intrachromosomal gene conversion and the maintenance of sequence
homogeneity among repeated genes. Genetics **100**: 315–337.

Pamilo P., Nei M., 1988 Relationships between gene trees and species trees. Mol Biol Evol **5**: 568–583.

Pei B., Sisu C., Frankish A., Howald C., Habegger L., Mu X., Harte R., Balasubramanian S., Tanzer A.,
Diekhans M., Reymond A., Hubbard T. J., Harrow J., Gerstein M. B., 2012 The GENCODE pseudogene
resource. Genome Biol **13**: R51.

Pezer, Harr B., Teschke M., Babiker H., Tautz D., 2015 Divergence patterns of genic copy number
variation in natural populations of the house mouse (Mus musculus domesticus) reveal three conserved
genes with major population-specific expansions. Genome Res **25**: 1114–1124.

Phifer-Rixey M., Bomhoff M., Nachman M. W., 2014 Genome-wide patterns of differentiation among
house mouse subspecies. Genetics **198**: 283–297.

Rozen S., Skaletsky H., Marszalek J. D., Minx P. J., Cordum H. S., Waterston R. H., Wilson R. K., Page D.
C., 2003 Abundant gene conversion between arms of palindromes in human and ape Y chromosomes.
Nature **423**: 873–876.

Salcedo T., Geraldes A., Nachman M. W., 2007 Nucleotide variation in wild and inbred mice. Genetics
177: 2277–2291.

Sambrook J., Russell D. W. (Eds.), 2006 *Molecular cloning: A laboratory manual*. Cold Spring Harbor
Laboratory Press.

Scavetta R. J., Tautz D., 2010 Copy number changes of CNV regions in intersubspecific crosses of the
house mouse. Mol Biol Evol **27**: 1845–1856.

Schindelhauer D., 2002 Evidence for a fast, intrachromosomal conversion mechanism from mapping of
nucleotide variants within a homogeneous alpha-satellite DNA array. Genome Res **12**: 1815–1826.

She X., Cheng Z., Zöllner S., Church D. M., Eichler E. E., 2008 Mouse segmental duplication and copy
number variation. Nat Genet **40**: 909–914.

Shi J., Wolf S. E., Burke J. M., Presting G. G., Ross-Ibarra J., Dawe R. K., 2010 Widespread gene conversion
in centromere cores. PLoS Biol **8**: e1000327.

Soh Y., Alföldi J., Pyntikova T., Brown L., Graves T., Minx P., Fulton R., Kremitzki C., Koutseva N.,
Mueller J., Rozen S., Hughes J., Owens E., Womack J., Murphy W., Cao Q., de~Jong P., Warren W.,
Wilson R., Skaletsky H., Page D., 2014 Sequencing the mouse Y chromosome reveals convergent gene
acquisition and amplification on both sex chromosomes. Cell **159**: 800–813.

Stamatakis A., 2014 RAxML version 8: A tool for phylogenetic analysis and post-analysis of large
phylogenies. Bioinformatics **30**: 1312–1313.

Stankiewicz P., Lupski J. R. 2002 Genome architecture, rearrangements and genomic disorders. Trends
Genet **18**: 74-82.

Steckelberg A.-L., Boehm V., Gromadzka A., Gehring N., 2012 Cwc22 connects pre-mRNA splicing and
exon junction complex assembly. Cell Rep **2**: 454–461.

Suzuki H., Shimada T., Terashima M., Tsuchiya K., Aplin K., 2004 Temporal, spatial, and ecological
modes of evolution of Eurasian Mus based on mitochondrial and nuclear gene sequences. Mol
Phylogenet Evol **33**: 626–646.

Svenson K. L., Gatti D. M., Valdar W., Welsh C. E., Cheng R., Chesler E. J., Palmer A. A., McMillan L.,
Churchill G. A., 2012 High-resolution genetic mapping using the mouse Diversity Outbred population.
Genetics **190**: 437–447.

Swallow J. G., Carter P. A., Jr. T. G., 1998 Artificial selection for increased wheel-running behavior in
house mice. Behav Genet **28**: 227–237.

Treangen T. J., Salzberg S. L., 2011 Repetitive DNA and next-generation sequencing: Computational
challenges and solutions. Nat Rev Genet.

Turner J. M., Aprelikova O., Xu X., Wang R., Kim S., Chandramouli G. V., Barrett J., Burgoyne P. S., Deng
C.-X., 2004 BRCA1, histone H2AX phosphorylation, and male meiotic sex chromosome inactivation. Curr
Biol **14**: 2135–2142.

Turner D. J., Miretti M., Rajan D., Fiegler H., Carter N. P., Blayney M. L., Beck S., Hurles M. E., 2007 Germline rates of de novo meiotic deletions and duplications causing several genomic disorders. Nat Genet **40**: 90–95.

Waterston R. H., Chinwalla A. T., Cook L. L., Delehaunty K. D., Fewell G. A., Fulton L. A., Fulton R. S., Graves T. A., Hillier L. W., Mardis E. R., McPherson J. D., Miner T. L., Nash W. E., Nelson J. O., Nhan M. N., Pepin K. H., Pohl C. S., Ponce T. C., Schultz B., Thompson J., Trevaskis E., Waterston R. H., Wendl M. C., Wilson R. K., Yang S.-P., An P., Berry E., Birren B., Bloom T., Brown D. G., Butler J., Daly M., David R., Deri J., Dodge S., Foley K., Gage D., Gnerre S., Holzer T., Jaffe D. B., Kamal M., Karlsson E. K., Kells C., Kirby A., Kulbokas E. J., Lander E. S., Landers T., Leger J. P., Levine R., Lindblad-Toh K., Mauceli E., Mayer J. H., McCarthy M., Meldrim J., Meldrim J., Mesirov J. P., Nicol R., Nusbaum C., Seaman S., Sharpe T., Sheridan A., Singer J. B., Santos R., Spencer B., Stange-Thomann N., Vinson J. P., Wade C. M., Wierzbowski J., Wyman D., Zody M. C., Birney E., Goldman N., Kasprzyk A., Mongin E., Rust A. G., Slater G., Stabenau A., Ureta-Vidal A., Whelan S., Ainscough R., Attwood J., Bailey J., Barlow K., Beck S., Burton J., Clamp M., Clee C., Coulson A., Cuff J., Curwen V., Cutts T., Davies J., Eyras E., Grafham D., Gregory S., Hubbard T., Hunt A., Jones M., Joy A., Leonard S., Lloyd C., Matthews L., McLaren S., McLay K., Meredith B., Mullikin J. C., Ning Z., Oliver K., Overton-Larty E., Plumb R., Potter S., Quail M., Rogers J., Scott C., Searle S., Shownkeen R., Sims S., Wall M., West A. P., Willey D., Williams S., Abril J. F., Guigó R., Parra G., Agarwal P., Agarwala R., Church D. M., Hlavina W., Maglott D. R., Sapojnikov V., Alexandersson M., Pachter L., Antonarakis S. E., Dermitzakis E. T., Reymond A., Ucla C., Baertsch R., Diekhans M., Furey T. S., Hinrichs A., Hsu F., Karolchik D., Kent W. J., Roskin K. M., Schwartz M. S., Sugnet C., Weber R. J., Bork P., Letunic I., Suyama M., Torrents D., Zdobnov E. M., Botcherby M., Brown S. D., Campbell R. D., Jackson I., Bray N., Couronne O., Dubchak I., Poliakov A., Rubin E. M., Brent M. R., Flicek P., Keibler E., Korf I., Batalov S., Bult C., Frankel W. N., Carninci P., Hayashizaki Y., Kawai J., Okazaki Y., Cawley S., Kulp D., Wheeler R., Chiaromonte F., Collins F. S., Felsenfeld A., Guyer M., Peterson J., Wetterstrand K., Copley R. R., Mott R., Dewey C., Dickens N. J., Emes R. D., Goodstadt L., Ponting C. P., Winter E., Dunn D. M., Niederhausern A. C. von, Weiss R. B., Eddy S. R., Johnson L. S., Jones T. A., Elnitski L., Kolbe D. L., Eswara P., Miller W., O'Connor M. J., Schwartz S., Gibbs R. A., Muzny D. M., Glusman G., Smit A., Green E. D., Hardison R. C., Yang S., Haussler D., Hua A., Roe B. A., Kucherlapati R. S., Montgomery K. T., Li J., Li M., Lucas S., Ma B., McCombie W. R., Morgan M., Pevzner P., Tesler G., Schultz J., Smith D. R., Tromp J., Worley K. C., Lander E. S., Abril J. F., Agarwal P., Alexandersson M., Antonarakis S. E., Baertsch R., Berry E., Birney E., Bork P., Bray N., Brent M. R., Brown D. G., Butler J., Bult C., Chiaromonte F., Chinwalla A. T., Church D. M., Clamp M., Collins F. S., Copley R. R., Couronne O., Cawley S., Cuff J., Curwen V., Cutts T., Daly M., Dermitzakis E. T., Dewey C., Dickens N. J., Diekhans M., Dubchak I., Eddy S. R., Elnitski L., Emes R. D., Eswara P., Eyras E., Felsenfeld A., Flicek P., Frankel W. N., Fulton L. A., Furey T. S., Gnerre S., Glusman G., Goldman N., Goodstadt L., Green E. D., Gregory S., Guigó R., Hardison R. C., Haussler D., Hillier L. W., Hinrichs A., Hlavina W., Hsu F., Hubbard T., Jaffe D. B., Kamal M., Karolchik D., Karlsson E. K., Kasprzyk A., Keibler E., Kent W. J., Kirby A., Kolbe D. L., Korf I., Kulbokas E. J., Kulp D., Lander E. S., Letunic I., Li M., Lindblad-Toh K., Ma B., Maglott D. R., Mauceli E., Mesirov J. P., Miller W., Mott R., Mullikin J. C., Ning Z., Pachter L., Parra G., Pevzner P., Poliakov A., Ponting C. P., Potter S., Reymond A., Roskin K. M., Sapojnikov V., Schultz J., Schwartz M. S., Schwartz S., Searle S., Singer J. B., Slater G., Smit A., Stabenau A., Sugnet C., Suyama M., Tesler G., Torrents D., Tromp J., Ucla C., Vinson J. P., Wade C. M., Weber R. J., Wheeler R., Winter E., Yang S.-P., Zdobnov E. M., Whelan S., Worley K. C., Zody M. C., 2002 Initial sequencing and comparative analysis of the mouse genome. Nature **420**: 520–562.

Watterson G., 1975 On the number of segregating sites in genetical models without recombination. Theor Popul Biol **7**: 256–276.

White M. A., Ané C., Dewey C. N., Larget B. R., Payseur B. A., 2009 Fine-scale phylogenetic discordance
across the house mouse genome. PLoS Genet **5**: e1000729.

Yang H., Bell T. A., Churchill G. A., Pardo-Manuel de Villena F., 2007 On the subspecific origin of the
laboratory mouse. Nat Genet **39**: 1100–1107.

Yang H., Wang J. R., Didion J. P., Buus R. J., Bell T. A., Welsh C. E., Bonhomme F., Yu A. H.-T., Nachman
M. W., Pialek J., Tucker P., Boursot P., McMillan L., Churchill G. A., Pardo-Manuel de Villena F., 2011
Subspecific origin and haplotype diversity in the laboratory mouse. Nat Genet **43**: 648–655.

Yeh T.-C., Liu H.-L., Chung C.-S., Wu N.-Y., Liu Y.-C., Cheng S.-C., 2010 Splicing factor Cwc22 is required
for the function of Prp2 and for the spliceosome to escape from a futile pathway. Mol Cell Biol **31**: 43–53.

1065

1067

Figure 1. Origin and age of the *R2d2* duplication. (**A**) *R2d* copy number across the phylogeny of the
genus *Mus*. Each dot represents one individual; grey dots indicate diploid copy number 2 and black dots
copy number >2. The duplication event giving rise to *R2d1* and *R2d2* most likely occurred on the
highlighted branch. Approximate divergence times (REF: Suzuki 2004) are given in millions of years ago
(Mya) at internal nodes. (**B**) Schematic structure of the *R2d1-R2d2* locus. The mouse reference genome
(strain C57BL/6J, *M. m. domesticus*) contains a single copy of *R2d* at *R2d1*. Wild-derived inbred strains vary
in haploid copy number from 1 (PWK/PhJ, *M. m. musculus*) to 2 (CAST/EiJ, *M. m. castaneus*) to 33
(WSB/EiJ, *M. m. domesticus*). *R2d1* is located at approximately 77.9 Mbp and *R2d2* at 83.8 Mbp. (**C**)
Concatenated tree constructed from *R2d1* (reference genome) and *de novo* assembled *R2d2* and *M. caroli*
sequences assuming a strict molecular clock. The duplication node is indicated with a grey dot. (**D**)
Relationship between observed *R2d* copy-number states and inferred structure of the *R2d1-R2d2* locus.
The configuration of the *M. spretus – M. musculus* common ancestor (4 diploid copies) is boxed in black.
We have yet to identify samples with diploid copy number 2 but no *R2d1* (grey shaded box).

1081

1082

Figure 2. *Cwc22* paralogs in the mouse genome. (**A**) Location and organization of *Cwc22* gene copies present in mouse genomes. The intact coding sequence of *Cwc22* exists in in both *R2d1* (grey shapes) and *R2d2* (black shapes). Retrotransposed copies (empty shapes) exist in two loci on chrX and one locus on chr2, immediately adjacent *R2d2*. Among the retrotransposed copies, coding sequence is intact only in the copy on chr2. Exon numbers are shown in grey above transcript models. (**B**) Alternate transcript forms of *Cwc22*, using different 3' exons. Coding exons shown in blue and untranslated regions in black. All Ensembl annotated transcripts are shown in the lower panel (from UCSC Genome Browser.)

1090

1091

Figure 3. Copy-number variation of *R2d* in mouse populations worldwide. (**A**) Copy-number variation
as measured by quantitative PCR. The normalized deltaCt value is proportional to log2(copy number).
Samples are classified as having 2 diploid copies, >2 copies or >4 copies of *R2d* using linear discriminant
analysis (LDA). (**B**) Fine-mapping the location of *R2d2* in 83 samples genotyped on the Mouse Diversity
Array (MDA). Grey points give nominal p-values for association between *R2d* copy number and
genotype; red points show a smoothed fit through the underlying points. The candidate interval for *R2d2*
from Didion *et al.* (2015), shown as an orange shaded box, coincides with the association peak. (**C**) Local
phylogeny at chr2: 83-84 Mbp in 62 wild-caught mice and laboratory strains. Tips are colored by
subspecies of origin: *M. m. domesticus*, blue; *M. m. musculus*, red; *M. m. castaneus*, green; *M. m. molossinus*,
maroon; outgroup taxa, grey. Individuals with >4 diploid copies of *R2d* are shown as open circles. (**D**)
Haplotypes of laboratory strains and wild mice sharing a high-copy allele at *R2d2*. All samples share a
haplotype over the region shaded in orange.

1104

Figure 4. Expression of *Cwc22* isoforms. (**A**) Read coverage and splicing patterns in *Cwc22* in adult
mouse brain from three wild-derived inbred strains. Swoops below x-axis indicate splicing events
supported by 5 or more split-read alignments. Known transcripts of *Cwc22^R2d1* (grey, from Ensembl),
inferred transcripts from *Cwc22^R2d2* (black) and the sequence of retro-*Cwc22* mapped back to the parent
gene (blue) are shown in the lower panel. Red stars indicate retained introns; red arrow indicates
insertion site of an ERV in *R2d2*. (**B**) Estimated relative expression of *Cwc22* isoforms (y-axis) in adult
mouse brain and testis in wild-derived inbred strains (x-axis). TPM, transcripts per million, on log10
scale.

1113

Figure 5. Signatures of non-allelic gene conversion between *R2d1* and *R2d2*. (**A**) Phylogenetic trees for three representative intervals across *R2d*. Sequences are labeled according to their subspecies of origin using the same color scheme as in **Figure 1**; open circles are *R2d1*-like sequences and closed circles are *R2d2*-like**.** Trees are drawn so that *M. caroli*, the outgroup species used to root the trees, is always positioned at the top. The changing affinities of PWK/PhJ (red) and CAST/EiJ (green) along *R2d* are evidence of non-allelic gene conversion. (**B**) *R2d* sequences from 20 wild-caught mice and 5 laboratory inbred strains. Each track represents a single chromosome; grey regions are classified as *R2d1*-like based on manual inspection of sequence variants, and black-regions *R2d2*-like. Upper panel shows sequences from samples with a single copy of *R2d*, residing in *R2d1*. Lower panel shows representative *R2d2* sequences for comparison. Asterisks indicate samples for which read alignments are shown in panel C. (**C**) Upper panel, paire-end read alignments (visualized with IGV) across the proximal boundary (dashed line) of *R2d1* in a sample with a conversion tract extending to the boundary. Positions of derived variants shared with *R2d2* are indicated by black dots. Lower panel, read alignments across the boundary of a non-allelic gene conversion tract. *R2d1* sequence from a single chromosome is a mosaic of *R2d1*-like (grey) and *R2d2*-like (black) segments. A magnified view of read pairs in the 3.8 kbp surrounding the proximal boundary of the tract shows read pairs spanning the junction. Black dots indicate the position of derived alleles diagnostic for *R2d2*. The precise breakpoint lies somewhere in the yellow shaded region between the last *R2d1*-specific variant and the first *R2d2*-specific variant.

1133

Figure 6. Suppression of crossing-over around *R2d2*. (**A**) Cumulative number of unique recombination events in the middle region of chr2 in genomes of 4,640 Diversity Outbred mice. Recombination events involving the high-copy-number WSB/EiJ haplotype are shown in purple and all other events in grey. Dashed vertical lines indicate the position of *R2d1* (left) and *R2d2* (right). (**B**) Distribution of haplotype block sizes at *R2d2* in selected generations of the DO, for *R2d2*^{HC} (WSB/EiJ, purple) versus *R2d2*^{LC} (the other seven founder haplotypes, grey). Asterisks indicate generations in which the length distributions are significantly different by Wilcoxon rank-sum test.

1141

Figure 7. Rate of *de novo* copy-number changes at *R2d2*. Estimates of per-generation mutation rate for CNVs at *R2d2* (±1 bootstrap SE) in the Diversity Outbred population; among wild *M. m. domesticus;* and among wild *M. m. castaneus*. For comparison, mutation rates are shown for the CNV with the highest rate of recurrence in a C57BL/6J pedigree (Egan *et al.* 2007) and for a microsatellite whose mutation rate was estimated in the BXD panel (Dallas 1992).

1147

1148

Figure 8. Sequence and structural diversity around *R2d2*. (**A**) Geographic origin of wild mice used in
this study, color-coded by subspecies (blue, *M. m. domesticus*; red, *M. m. musculus*; green, *M. m. castaneus*).
Diploid copy number of the *R2d* unit is shown for wild samples for which integer copy-number estimates
are available: 26 *M. m. domesticus* and 10 *M. m. castaneus* with whole-genome sequencing data, and
representatives from *M. m. molossinus* and *M. m. musculus* for comparison. Schematic shows the
R2d1/R2d2 configurations corresponding to diploid copy numbers of 2 and 4. (**B**) Profiles of read depth
(first two panels), average sequence divergence to outgroup species *M. caroli* (d_{xy}, third panel), number of
segregating sites per base (S/L, fourth panel) and within-population average heterozygosity (π, fifth and
sixth panels). The region shown is 500 kbp in size; the insertion site of *R2d2* is indicated by the red dashed
line. Grey boxes along baseline show positions of repetitive elements (from UCSC RepeatMasker track);
black boxes show non-recombining haplotype blocks. Blue bars indicate the position of 7 tandem
duplications in the mm10 reference sequence with >99% mutual identity, each containing a copy of retro-
Cwc22. Grey shaded region indicates duplicate sequence absent from *M. caroli*. (**C**) Estimated per-site
nucleotide diversity within *M. m. domesticus R2d1*, *M. m. domesticus R2d2* and *M. m. castaneus R2d2*.

1163

1164

Figure 9. A region of excess sequence divergence on chromosome 4. (**A**) Genome-wide sequence
divergence estimates for representative samples from the sub-genus *Mus*. (**B**) Estimated sequence
divergence in 31 kbp windows across distal chr4 for the samples in panel A. Divergent regions identified
by the hidden Markov model (HMM) are indicated with black bars along the horizontal axis. (**C**)
Phylogenetic tree constructed from *Skint1* coding sequences reported in Boyden *et al.* (2008) (**D**)
Schematic representation of the process of gene duplication, followed by differential loss of paralogs
along independent lineages. (**E**) Dotplot of self-alignment of sequence from the region of distal chr4
containing the *Skint* gene family. Positions of *Skint* genes are indicated along the horizontal axis; *Skint1*
highlighted in red.

 SUPPLEMENTARY MATERIAL

Supplementary Figure 1. Conservation of synteny between mouse and four other mammals around
Cwc22^{R2d1} (upper panel) indicates that the *R2d1* sequence remains in its ancestral position. Chevrons
represent genes, alternating white and grey, and are oriented according to the strand on which the gene is
encoded. *Cwc22^{R2d2}* is novel in the mouse but its position relative to genes with conserved order is shown
in the lower panel. Note that synteny is disrupted in mouse and rat distal to *R2d2*.

Supplementary Figure 2. Pairwise alignment of *R2d2* contig (top) to the *R2d1* reference sequence
(bottom). Dark boxes show position of repetitive elements present in both sequences; syntenic positions
are connected by grey anchors, and blank space represents aligned bases in both sequences. Orange boxes
indicate position of repetitive elements present in the *R2d1* sequence but not detected in *R2d2*; blue boxes
indicate position of elements in *R2d2* but not *R2d1*. *Cwc22* transcripts are shown below the alignment.

Supplementary Figure 3. Phylogenetic tree constructed from amino acid sequences for mammalian
Cwc22 homologs (including all three mouse paralogs) with chicken as an outgroup. Node labels indicate
support in 100 bootstrap replicates.

Supplementary Figure 4. Alignment of amino acid sequences from mouse *Cwc22^{R2d1}*, *Cwc22^{R2d2}* and retro-
Cwc22, plus *Cwc22* orthologs from 19 other placental mammals plus opossum, platypus and chicken as
outgroups. Residues are colored according to biochemical properties and gaps are shown in grey.
Information content of each column in the alignment, measured as the Jenson-Shannon divergence, is
plotted in the lower panel.

Supplementary Figure 5. Diagnostic variants used to identify gene conversion tracts between *R2d2* and
R2d1. Each column represents a single variant (total of 1,411) between *R2d1* and *R2d2* for which *R2d2* has
the derived allele, and each row a single individual with whole-genome sequence data available (named
in **Supplementary Table 1**). Points are colored according to the "allelotype" of each variant detected in 8
or more reads in each sample: 0:0 (red), neither *R2d1* nor *R2d2* allele present; 1:0, *R2d1* but not *R2d2*
(orange); 1:1, both *R2d1* and *R2d2* (green); 0:1, *R2d2* but not *R2d1* (blue). Individuals are grouped
according to diploid *R2d* copy number.

Supplementary Figure 6. Partial loss of *R2d2* with structural rearrangement. (**A**) Inferred structure of the
R2d1-R2d2 region in IR:AHZ_STND:015, a wild *M. m. domesticus* individual from Iran. *R2d1* is present on
both chromosomes but only a fragment of *R2d2* remains on one chromosome, and it has been transposed
into the retro-*Cwc22* array. (**B**) Normalized depth of coverage (2 = normal diploid level) across *R2d*.
Regions in grey represent reads from *R2d1* alone, while region in black captures reads from *R2d1* and
R2d2, as shown by arrows from panel A. (**C**) Position of read pairs (red; not drawn to scale) with soft-
clipped alignments to *R2d1*. The proximal read aligns in the 3′ UTR of *Cwc22*, and the distal read across
an exon-intron boundary within the gene body. Note the "outward"-facing direction of the alignments.
(**D**) Positions of the mates of the reads in panel C. Note that the x-axis is reversed so that the exons of
retro-*Cwc22* (encoded on the plus strand) parallel those of *Cwc22* (encoded on the minus strand). The 3′
read maps across the boundary of th 3′ UTR of *Cwc22* and the ERV mediating the retrotransposition
event. The 5′ read maps across two exon-exon boundaries in retro-*Cwc22*, so there is no ambiguity
regarding its alignment to the retro-transposed copy. (**E**) Inferred structure of *Cwc22* paralogs in this
sample. Note that one of the copies of retro-*Cwc22* is now a mosaic of retrotransposed and *Cwc22^{R2d2}*-
derived sequence.

Supplementary Figure 7. Difference between expected and observed recombination fraction between
markers flanking *R2d2* in experimental crosses in which at least one parent is segregating for a high-copy

allele of *R2d2*. Thick and thin vertical bars show 90% and 95% confidence bounds, respectively, obtained
by non-parametric bootstrap.

Supplementary Figure 8. Targeted *de novo* assembly using the multi-string Burrows-Wheeler
Transform (msBWT). (**A**) The msBWT and its associated FM-index implicitly represent a suffix array of
sequencing reads, such that read suffixes sharing a k-mer prefix are adjacent in the data structure. This
allows rapid construction of a local de Bruijn graph starting from a k-mer seed (dark blue) and extending
by successive k-mers (light blue) containing the $(k − 1)$-length suffix of the previous k-mer. A $(k − 1)$-
length prefix with more than one possible suffix (red and orange) creates a branch point. Adjacent nodes
in the graph with in-degree and out-degree one can be collapsed into a single node, yielding a simplified
graph, which can then be traversed to obtain linear contig(s). (**B**) Paralogs of *R2d* can be disentangled
using the local de Bruijn graph by exploiting differences in copy number. Edges in the graph are
weighted by read count, and linear contigs for the *R2d1* and *R2d2* paralogs obtained by traversing the
graph in a manner that minimizes the variance in edge weights along possible paths. Phase-informative
reads (those overlapping multiple paralogous variants) provide a second source of evidence.

1231

Supplementary Table 1. List of mouse samples used in this study, with their taxonomic designation,
geographic origin, karyotype (STND, standard: 2n = 40; all others chromosomal races with Robertsonian
translocations) and *R2d2* copy-number classification.

Supplementary Table 2. Transposable-element insertions private to *R2d1* or *R2d2*. Coordinates are
offsets with respect to the start position of *R2d* (for *R2d1*: chr2: 77,869,657 in the reference genome; for
R2d2: the beginning of the *de novo* assembled contig in **Supplementary File 1**.)

Supplementary Table 3. Frequency table of copy-number status by geographic origin for wild-caught
and wild-derived Mus musculus individuals used in this study, stratified by subspecies.
"Europe/Mediterranean" includes continental Europe, the United Kingdom and countries in the
Mediterranean basin (Tunisia, Cyprus, Israel). "Asia" includes Asia, the Middle East and countries in the
Indian Ocean basin (Madagascar).

Supplementary Table 4. Individuals from the Diversity Outbred population carrying *de novo* copy-
number mutations at *R2d2*. Each was expected to be heterozygous for the WSB/EiJ allele (33 haploid
copies).

Supplementary Table 5. Regions of excess divergence between wild or wild-derived mice and the mouse
reference genome (GRCm38/mm10 build).

Supplementary Table 6. Regions of *R2d* targeted for *de novo* assembly in inbred strains.

Supplementary File 1. Compressed archive containing *R2d2* contig (from WSB/EiJ) and multiple
sequence alignments from selected regions in **Supplementary Table 6**.