Classification: Biological Sciences, Developmental Biology

Title: Divergence of gene regulatory network linkages during specification of ectoderm and

mesoderm in early development of sea urchins

Short Title: Divergence of GRNs in early echinoid embryos

Author Affiliations:

Eric M. Erkenbrack[a,1,3] and Eric H. Davidson[a,2]
10

[a]Division of Biology and Biological Engineering, California Institute of Technology, Pasadena,

California, 91125, United States of America

[1]Current address: Department of Ecology and Evolutionary Biology, Yale University, New Haven,

Connecticut, 06511, United States of America

[2]Deceased author

[3]Corresponding Author

Eric M. Erkenbrack, Yale University, Department of Ecology and Evolutionary Biology, New Haven,

CT, 06511, 203-737-3092, email: erkenbra@caltech.edu (EME), eric.erkenbrack@yale.edu (EME)

Keywords: gene regulatory network evolution, network topology, sea urchin embryogenesis,

embryonic axis specification, cidaroid, euechinoid

Abstract

Developmental gene regulatory networks (GRNs) are assemblages of gene regulatory interactions that direct ontogeny of animal body plans. Studies of GRNs operating in early development of euechinoid sea urchins has revealed that little appreciable change has occurred since their divergence approximately 90 million years ago (mya). These observations suggest that strong conservation of GRN architecture has been maintained in early development of the sea urchin lineage. To test whether this is true for all sea urchins, comparative analyses of echinoid taxa that diverged deeper in geological time must be conducted. Recent studies highlighted extensive divergence of skeletogenic mesoderm specification in the sister clade of euechinoids, the cidaroids, suggesting that comparative analyses of cidaroid GRN architecture may confer a greater understanding of the evolutionary dynamics of developmental GRNs. Here, we report spatiotemporal patterning of 55 regulatory genes and perturbation analyses of key regulatory genes involved in euechinoid oral-aboral patterning of non-skeletogenic mesodermal and ectodermal domains in early development of the cidaroid *Eucidaris tribuloides*. Our results indicate that developmental GRNs directing mesodermal and ectodermal specification have undergone marked alterations since the divergence of cidaroids and euechinoids. Notably, statistical and clustering analyses of echinoid temporal gene expression datasets indicate that regulation of mesodermal genes has diverged more markedly than regulation of ectodermal genes. Although research on indirect-developing euechinoid sea urchins suggests strong conservation of GRN circuitry during early embryogenesis, this study indicates that since the divergence of cidaroids and euechinoids developmental GRNs have undergone significant divergence.

Significance Statement

Sea urchins consist of two subclasses, cidaroid and euechinoids. Research on gene regulatory

networks (GRNs) in early development of three euechinoids indicates that little appreciable change

has occurred to their linkages since they diverged approximately 90 million years ago (mya). We

asked if this conservation extends to the all echinoids. Here we demonstrate extensive divergence

of GRN architecture in early embryonic specification of the oral-aboral axis in echinoids. We

systematically analyzed in a cidaroid spatiotemporal expression and function of regulatory genes

segregating euechinoid ectoderm and mesoderm. While we found evidence for diverged regulation

of both mesodermal and ectodermal genes, our comparative analyses indicated that, since these

two clades diverged 268.8 million years ago, mesodermal GRNs have undergone significantly

more alteration than ectodermal GRNs.

\body

55 Introduction

Integral to early development of a bilaterian is the development of the three embryonic

tissue-layer domains: endoderm, ectoderm, and mesoderm. Asymmetrically distributed RNA and

proteins in the egg provide the initial inputs into this process and thereby determine the spatial

coordinates of domain formation (1, 2). In the context of these maternal factors, zygotic

transcription of distinct sets of regulatory genes occurs in specific regions of the embryo, initiating

the genomically encoded regulatory program and its output of regulatory genes. In this way the

embryo becomes populated by distinct sets of transcription factors and cell signaling transcripts

called regulatory states (3). The spatial readout of developmental gene regulatory networks

(GRNs), regulatory states provide each cell with its molecularly distinct and functional identity (4,

5).

Sea urchins (class Echinodea) have long served as model systems to study mechanisms of

early development and, more recently, to study fundamental aspects of developmental GRNs.

Diversification of cell lineages and embryonic domains in sea urchin embryos depends on the

cleavage positions of their blastomeres (6, 7), facilitating interpretation of mechanisms of spatial

change. Sea urchin lineages have undergone multiple changes in life history strategies, providing

a convenient framework, with experimental replicates, to investigate evolution and mechanisms of

developmental programs (8). Importantly, a well-represented fossil record affords dating of

evolutionary events (9) and has established that the sister subclasses of sea urchins—cidaroids

and euechinoids—diverged from one another at least 268.8 million years ago (mya) (10).

Differences in the timing of developmental events in embryogenesis of cidaroids and euechinoids

have long been a topic of interest, but only recently have become the subject of molecular

research (11-18).

Research on early development of the euechinoid purple sea urchin *Strongylocentrotus*

purpuratus has brought into high resolution the players and molecular logic directing

developmental GRNs that specify its varied embryonic domains and subdomains (19-30).

Additionally, abundant comparative evidence exists for other euechinoid taxa, including *Lytechinus*

variegatus (31-37) and *Paracentrotus lividus* (38-42). Data from these three indirect-developing

euechinoids indicate that, although these taxa diverged from one another approximately 90 mya

(9, 43), very little appreciable change to developmental GRNs has been observed in their early

development (44-46). While there is evidence of a few heterochronic alterations to these GRNs,

e.g. a heterochronic shift in *snail* expression in *L. variegatus* versus *S. purpuratus* (22), numerous

studies have made clear the striking conservation of GRN linkages and spatiotemporal gene

expression in these lineages. Recently, studies of early development of the distantly-related,

indirect-developing cidaroid sea urchin *Eucidaris tribuloides* revealed that skeletogenic mesoderm

specification in this clade is markedly different from that observed in euechinoids (17, 18, 47).

These observations suggest that comparative analyses of GRN circuitry in early development of

cidaroids and euechinoids have the potential to illuminate the tempo and mode of evolution of

developmental GRNs.

Here, we coupled comparative spatiotemporal gene expression analyses with experimental

manipulations of regulatory genes specifying euechinoid non-skeletogenic mesodermal (NSM) and

ectodermal domains in the cidaroid *E. tribuloides* to reveal how these GRNs have changed since

the divergence of these two clades 268.8 mya. This study focused on oral-aboral (O-A; also called

dorsal-ventral) patterning, which has consequences for both ectoderm and mesoderm. O-A axis

specification is a well-documented process in euechinoids (46) and is a highly conserved

developmental mechanism in deuterostomes (48, 49). We present evidence that deployment and

regulatory interactions of regulatory genes specifying sea urchin ectoderm and mesoderm have

diverged substantially in indirect-developing echinoids. Importantly, our comparative data and

analyses suggest that regulatory interactions occurring in ectodermal GRNs have undergone less

divergence than those occurring in mesodermal domains. Our conclusions are supported by

comparative spatiotemporal data, statistical analyses of timecourse gene expression data in three

taxa of echinoids and perturbation analyses. Taken together regulatory genes expressed in

ectodermal domains exhibited stronger signals of conservation relative to those expressed in

mesodermal domains. These results suggest that embryonic domains and cell types in early

development of sea urchins have evolved at different rates since the divergence of the two sister

sub-classes of modern sea urchins. These alterations to GRN architecture have occured

throughout the network and frequently since their divergence. Additionally, these results offer an in

principle explanation for observations of rapid change to nearly all components of developmental

process in the development of direct-developing, non-feeding sea urchins (50-53).

Results

Ectodermal O-A regulatory states in the cidaroid *E. tribuloides* suggest striking conservation of regulatory gene deployment in echinoids

In euechinoids, ectoderm is segregated into a diverse set of regulatory states defined by the future location of the stomodeum (46, 54, 55). A critical factor in establishing oral-aboral (47) polarity in sea urchins is the signaling ligand *nodal* (35, 38). Nodal ligand activates the SMAD signaling pathway, which is directly upstream of *nodal* (itself), *not, lefty* and *chordin* in oral ectoderm (OE) (40, 56). In *E. tribuloides,* zygotic transcription of *nodal, not* and *lefty* begins by early blastula stage (Fig. 1*D*, 1*J*, and Fig. S1*D*). In contrast, transcriptional activation of *chordin* is delayed by at least 5 hours from this initial cohort, indicative of an intermediate regulator between *nodal* and *chordin* in *E. tribuloides* (Fig. 1D and 1E). From 17 hpf to 40 hpf, spatial expression of *nodal* is observed in a well-defined region in OE that expands slightly as gastrulation proceeds (Fig. 1D2-1D5 and Fig. S2). Unlike *nodal*, the spatial distribution of its targets is not solely restricted to a small field of cells in OE. *Lefty* (also known as Antivin), an antagonist of *nodal*, exhibits a broader pattern of expression that, by 50 hpf, expands into the oral side of the archenteron (Fig. S1D2-S1D5 and Fig. S2). Similarly, *chordin* transcripts are detected in OE throughout early *E. tribuloides* development (Fig. 1E1-1E5 and Fig. S2). The homeobox gene *not,* known to play a role directly downstream of *nodal* in euechinoid O-A ectodermal and mesodermal polarization (27, 57), was observed spatially in OE during gastrulation and later extends vegetally towards the perianal ectoderm and is observed in the archenteron (Fig. 1J1-1J5 and Fig. S2). In euechinoids, the *bmp2/4* ligand is a directly downstream of Nodal and is translocated across the embryo to the aboral side, where it stimulates via SMAD signaling in aboral ectoderm (AE) specification genes such as *tbx2/3* (38, 58, 59). In *E. tribuloides*, both *bmp2/4* and *tbx2/3* are

upregulated with the *nodal-not-lefty* cohort (Fig. S1G and Fig. 1I1-1I5). *Tbx2/3* exhibits spatial

expression from late blastula stage onwards that is complementary to OE genes in the lateral AE

(Fig. 1I2-1I5). By mid-gastrula stage, *tbx2/3* is also expressed in the archenteron and much later,

by 70 hpf, is expressed in the bilateral clusters of cells synthesizing the larval skeleton (Fig. S2),

which is similar to the spatial expression in two euechinoids with notably interesting heterochronic

differences (60, 61). Lastly, the Forkhead family transcription factor *foxq2* is sequentially restricted

to and specifically expressed in embryonic anterior neural ectoderm (ANE) territory in

deuterostomes (62). In euechinoids, *foxq2* restriction to ANE is a crucial component of O-A axis

specification, setting the anterior boundary of OE by restricting expression of *nodal* (23, 55). In *E.*

tribuloides, foxq2 exhibited an expression pattern consistent with observations in euechinoids and

other deuterostomes, suggesting conserved roles for this gene in ANE and O-A specification (Fig.

1B and Fig. S2).

Conserved spatiotemporal deployment of ciliary band regulatory genes

Free-feeding, indirect-developing sea urchins possess a single neurogenic ciliated band

(CB) early in development that circumnavigates the larval oral face and facilitates feeding and

locomotion (63). This structure has undergone frequent modification in the lineages leading to

modern sea urchins, viz. in planktotrophic larvae (64). In euechinoids *goosecoid (gsc), onecut* and

irxa contribute to the geometric patterning of CB formation (28, 40, 65). In euechinoids, *gsc* is

expressed in OE and is directly downstream of *nodal* signaling on the oral side of the embryo (40);

onecut (also known as *hnf6*) is a ubiquitous, maternally deposited factor that is restricted to a

region spanning the boundaries of OE and AE, at which lies progenitor CB territory; and *irxa* is

expressed exclusively in AE downstream of *tbx2/3* (40, 66). In the cidaroid *E. tribuloides*, *gsc* is

activated with the OE cohort and is also spatially restricted to OE (Fig. 1G1 and Fig. S3). As in

euechinoids, *onecut* is maternally deposited in *E. tribuloides* (Fig. 1H1). Although its early spatial

expression was not obtained, staining was observed in a restricted band of cells encircling the OE

by mid-gastrula stage. The spatial dynamics of *onecut* in *E. tribuloides* is notable insofar as its

spatial distribution in progenitor CB begins in the progenitor field of post oral CB and subsequently

extends in a narrow band of 4-8 cell diameters towards progenitor pre oral CB (Fig. 1H2-1H5 and

Fig. S3). This observation starkly contrasts that in euechinoids, in which *onecut* is observed

ubiquitously expressed early and later delimited as a whole to the CB territory by transcriptional

repressors in the OE and AE (28, 67). *Irxa* initiates zygotic expression at mid-blastula stage (~14

hpf) in *E. tribuloides*, and by 28 hpf is observed broadly in AE (Fig. 1C1-1C5). Unlike in

euechinoids, *irxa* is broadly distributed in *E. tribuloides* AE—much more so than *tbx2/3*—indicating

that it is likely broadly activated in the ectoderm and repressed in OE and ANE. These

spatiotemporal observations of key CB regulatory genes suggest that CB patterning mechanisms

are likely to be conserved in echinoids.

Non-skeletogenic mesodermal patterning and regulatory interactions

have diverged widely in echinoids

Non-skeletogenic mesoderm (NSM) in euechinoids arises at the vegetal plate from early

cleavage endomesodermal precursors and gives rise to four different cell types: blastocoelar cells,

pigment cells, circumesophageal cells and coelomic pouch cells (68). Experimental observations

indicate that euechinoids completely rely on presentation of Delta ligand in the adjacent SM to

upregulate NSM regulatory factors in veg2 endomesodermal cells (32, 69, 70). After Delta/Notch

signaling upregulates NSM regulatory genes in SM-adjacent cells, NSM is partitioned into aboral

NSM and oral NSM as a result of Nodal/SMAD signaling upregulating transcription factor *not* on

the oral side of the embryo (27, 71). The first evidence of NSM polarity in *E. tribuloides*, is seen in

the spatial distribution of *ese* and *gcm* transcripts shortly after gastrulation has commenced (Fig.

1K2-1K5 and 1L2-1L5). In euechinoids, *ese* is expressed in the oral NSM, and *gcm* is expressed in

the aboral NSM. In *E. tribuloides ese* is first in ANE and progenitor NSM just prior to the onset of

gastrulation, shortly after invagination of the archenteron it is subsequently restricted to one side in

NSM (Fig. 1K2-1K5). Similarly, *gcm* is expressed transiently in oral and aboral NSM, and by 28 hpf

is restricted to a cluster of cells just below the tip of the archenteron (Fig. 1L2-1L5). Later this

expression is seen solely on one side of the archenteron as *gcm*-positive cells ingress rapidly into

the blastocoel at 36 hpf (Fig. S4). Double fluorescent whole-mount *in situ* hybridization (WMISH)

revealed that *ese* and *gcm* in this cidaroid are restricted to opposite sides of the archenteron (Fig

S4G1 and S4G2), providing the first evidence of conserved polarity in NSM segregation in

echinoids.

197   To further clarify NSM regulatory states in *E. tribuloides* and their similarity to those in

euechinoids, we conducted additional single and double WMISH on NSM regulatory genes *ets1/2,*

gatac, gatae, prox, scl and *tbrain.* Directly downstream of *gcm* in euechinoids is *gatae* (27). In *S.*

purpuratus, *gatae* is observed in endomesoderm by blastula stage (72). In *E. tribuloides* NSM,

gatae is expressed throughout the endomesoderm at the time of SM ingression (~28 hpf) and later

is observed restricted to one side near the tip of the archenteron, as well as in the second wave of

ingressing mesenchyme (Fig. S1C2-S1C5 and Fig. S4). *Gatac (gata1/2/3), prox* and *scl*, all of

which are oral NSM genes in euechinoids (27), come off the baseline at similar times in *E.*

tribuloides and are detectable in a few cells at the base of the vegetal pole by 18 hpf by WMISH

(Fig. S4B1, S4E1, and S1F). Of these three genes, *scl* was the first to show O-A NSM polarity

followed by *gatac* (Fig. S1B5, S1E5, and S1F5). Surprisingly, by 36 hpf *prox* did not exhibit spatial

distribution of transcripts that unambiguously indicated O-A polarity (Fig. S1E2-S1E5 and Fig.

S4E6), suggesting that either *prox* is a general mesodermal regulatory factor in *E. tribuloides* or it

is spatially restricted later in gastrular development.

These data on spatial dynamics of NSM regulatory factors suggest that there exist

numerous regulatory states in the anterior archenteron. To provide some clarity, double fluorescent

WMISH (dfWMISH) indicated this was indeed the case. Previous observations suggested that *E.*

tribuloides mesodermal domains broadly express *ets1/2* and *tbrain* (17), and dfWMISH confirmed

this result (Fig. S4IS4I). Within this broad *ets1/2-tbrain* domain, three regulatory states are

identified (Fig. S4G-S4I): (1) orally localized *ets1/2, tbrain* and *ese*; (2) aborally localized *ets1/2,*

tbrain, and *gcm*; and (3) an anteriorly localized micromere-descendant regulatory state at the tip of

the archenteron of *ets1/2, tbrain, ese* and *alx1.* The only similarity between NSM regulatory states

of *E. tribuloides* and that of euechinoids is *ese* is restricted to oral NSM and *gcm* to aboral NSM

(27). The NSM regulatory states described here make it clear that since the divergence of

cidaroids and euechinoids there have been numerous changes to spatial and temporal gene

expression during the early patterning of echinoid NSM cell types.

Perturbation of Nodal signaling suggests highly conserved

regulatory interactions in patterning echinoid O-A ectoderm and

mesoderm

Next we sought to experimentally perturb O-A specification in *E. tribuloides* to determine the

extent to which this developmental program is conserved in echinoids. In euechinoids, perturbation

of animal-vegetal (A-V) axis polarity by disruption of nuclearization of β-catenin indirectly disrupts

O-A axis specification (31, 33). One mechanism underlying the crosstalk of these two

deuterostome specification events was found to be restriction of *foxq2* to ANE, as its presence in

OE blocked *nodal* transcription (23). In *E. tribuloides,* disrupting nuclearization of β-catenin at the

vegetal pole by overexpressing dn-Cadherin RNA led to upregulation of *foxq2* and strong downregulation of *nodal* and its euechinoid downstream components *bmp2/4, not,* and *tbx2/3* (Fig. S5A). This result suggests that the molecular crosstalk between β-catenin/TCF, *foxq2* and *nodal* are conserved in echinoids.

Next, we aimed to determine the spatiotemporal effects of perturbation of O-A specification by culturing *E. tribuloides* embryos in the presence of SB43152, a small molecule antagonist of the TGF-β (Nodal) receptor Alk4/5/7 (73). At four days post fertilization, these embryos exhibited strong aboralization, archenterons that failed to make contact with OE, and supernumerary skeletal elements (Fig. S5B). Quantitative PCR (qPCR) analysis at four different timepoints in *E. tribuloides* development showed strong downregulation of OE regulatory factors *chordin, gsc, lefty, nodal* and *not* (Fig. 2A). This result was confirmed spatially by WMISH for mRNA transcripts of *chordin, nodal* and *not* (Fig. 2B and S5C). Another critical OE regulatory gene is the secreted TGF-β ligand *bmp2/4.* This gene was clearly not affected to the same degree as the aforementioned cohort of OE factors (Fig. 2A). This result is strikingly different from the strong downregulation of *bmp2/4* observed in the euechinoid *P. lividus* when it was cultured in the presence of SB431542 or when injected with Nodal morpholino (MASO) (38, 40), indicating that regulation of *bmp2/4* may be under alternative control in *E. tribuloides*.

On the aboral side, a striking difference is the effect of SB431542 on *tbx2/3.* In euechinoids, *tbx2/3* is downstream of *bmp2/4* ligand, which diffuses from OE to AE (58, 59, 74). Treatment of *P. lividus* embryos with SB431542 inhibitor completely and specifically extinguishes *tbx2/3* in AE while not interfering with its SM expression (40). In *E. tribuloides,* qPCR data suggest SB431542 inhibitor has no effect on *tbx2/3* regulation (Fig. 5C). However, WMISH of *tbx2/3* in the presence of the inhibitor revealed its domain of expression had expanded into an equatorial ring (Fig. 2B and S5C). Similarly, whereas in *E. tribuloides* qPCR data indicate strong downregulation of the AE

regulatory factor *irxa* (Fig. 5C), its domain of expression expanded into OE in *P. lividus* embryos

upon SB431542 treatment (40). These results suggest GRN architecture immediately downstream

of the initial *nodal* and *bmp2/4* circuitry has diverged in echinoids and further indicate that

regulatory interactions immediately upstream of *bmp2/4, irxa* and *tbx2/3* are regulated differently in

E. tribuloides.

Between the OE and AE is the pro-neural ciliary band that expresses *onecut.* In

euechinoids, perturbation of Nodal signaling results in an expansion of *onecut* throughout the

ectoderm, leading to the suggestion that the default fate of ectoderm is a pro-neural CB territory

(40). When this perturbation was carried out in *E. tribuloides,* a very different result was obtained in

which we see *onecut* restricted to a single equatorial band of 6-10 cell diameters (Fig. 2C). A

previous study showed that disruption of *E. tribuloides* endomesoderm formation by treatment with

zinc results in embryos exhibiting a ring of highly concentrated proneural synaptotagmin-B positive

cells at the equator of the embryo (75). Together, these observations indicate that, in the absence

of Nodal signaling, there are anteriorly positioned repressors in *E. tribuloides* restricting CB fate to

the equator of the embryo and that CB restriction from the ANE is functioning differently in

cidaroids and euechinoids.

Lastly, in euechinoids studied thus far, polarity in NSM (47) lineages is downstream of

Nodal signaling via the transcription factor Not (27, 71). In *E. tribuloides,* qPCR data did not

indicate consistent differences in mRNA abundance for NSM regulatory genes (Fig. 2A). However,

WMISH assays revealed that embryos treated with SB431542 failed to restrict *gcm* to the aboral

side (Fig. 2B and S5C). This observation is consistent with the euechinoid GRN linkage

immediately downstream of Nodal signaling via the transcription factor *not,* which represses aboral

NSM in the oral-facing region of the archenteron (27). Indeed, in *E. tribuloides, not* is seen in the

archenteron throughout gastrulation (Fig. 1J3-1J5). These observations are consistent with a

conserved role for Nodal signaling via *not* in establishing polarity in NSM cell types in the

archenteron of *E. tribuloides.*

Regulatory interactions establishing ciliary band are conserved in echinoids

The spatial distributions of *gsc, onecut* and *irxa* are highly suggestive of a conserved

regulatory apparatus that spatially restricts CB to the boundary of OE and AE (Fig. 1 and Fig. S3).

Supporting this claim, microinjection of mutated or wild-type BACs encoding the *cis*-regulatory

region of *S. purpuratus onecut* and a GFP reporter were examined in *E. tribuloides*. In *S.*

purpuratus, onecut is restricted to the progenitor CB territory by *gsc* repression in OE and *irxa*

repression in AE (Fig. 3A). Recently, Barsi and Davidson (2016) experimentally validated four cis-

regulatory modules directing the activation of *onecut* by *soxb1* and repression of *onecut* by *gsc* in

OE and *irxa* in AE (Fig. 3B). Remarkably, these BACs faithfully recapitulated the regulatory activity

of these BACs in *E. tribuloides* (Fig. 3C-3H and Fig. S6). To test for this conservation, we assayed

a series of endogenous and site-directed mutagenesis *onecut* BACs from *S. purpuratus* by

microinjection (65). Remarkably, a BAC that has been shown to recapitulate the endogenous *S.*

purpuratus onecut expression pattern faithfully expressed reporter GFP in the CB of *E. tribuloides*

(Fig. 3H and S6A-S6B). Further, a BAC harboring mutated repressor sites for the oral repressor

gsc repeatedly exhibited ectopic expression in OE of *E. tribuloides* (Fig. 3E). When interpreted with

the spatial WMISH data of Figure 1, the early specification of CB regulatory factors suggests

divergence of initial activation of *onecut* and conserved GRN circuitry guiding the restriction of

onecut to CB territory by *irxa* and *gsc,* a striking example of a highly conserved developmental

program after 268 million years of evolution.

Comparative analyses of temporal gene expression in echinoids suggests ectodermal and mesodermal GRNs have diverged at different rates

The spatiotemporal data presented thus far are highly suggestive that O-A axis specification, as well as gastrular CB formation, in *E. tribuloides* is consistent with similar processes in euechinoids and that NSM specification has ostensibly diverged. Next we sought a comparative gene expression approach to reveal the extent of divergence between echinoid clades. We compared absolute mRNA transcript abundance of 18 regulatory genes in three species and 55 *S. purpuratus* regulatory genes (76) in early euechinoid developmental GRNs for 14 early developmental timepoints in *E. tribuloides* (Table S5 and S6). We scaled timepoints from *E. tribuloides* to the euechinoid *P. lividus* since both develop at a similar temperature and since scaling of developmental stages between *P. lividus* and *S. purpuratus* had been previously established (Table S5) (77). This allowed pairwise analyses of Spearmann's rank correlation coefficients, ρ, between the three species were conducted (Table S6). Plotting relative mRNA transcriptional dynamics for the three species were indicative of compelling correlation for ectodermal regulatory factors and supported the notion of poor correlation for regulatory factors driving mesoderm specification (Fig. 4 and S7). We next considered timecourse expression data of 55 orthologs of *E. tribuloides* and *S. purpuratus.* We binned the orthologs in ectoderm, endoderm and mesoderm based on their spatial distributions in *S. purpuratus* and compared them against the mean of all ρ values. Regulatory orthologs expressed in endoderm and ectoderm exhibited significantly higher ρ relative to the mean of all ρ values, suggesting strong conservation of transcriptional dynamics of these factors in echinoids (Fig. 5A). However, regulatory orthologs expressed in mesodermal cell lineages did not depart significantly from the mean ρ, suggesting

transcriptional dynamics of mesodermal regulatory factors have changed markedly since the cidaroid-euechinoid divergence (Fig. 5A and 5B). Furthermore, we conducted a comparative clustering analysis to identify statistical differences in the timecourse data. By clustering *S. purpuratus* regulatory orthologs into six distinct clusters and forcing assignment of each *E. tribuloides* ortholog into all clusters, we acquired cluster membership scores and clustering similarities for each ortholog (Table S7, S8, and S9). Considering *S. purpuratus* orthologs alone revealed regulatory genes clustering together based on whether they are expressed maternal, early or late in development (Fig. S8A). When *E. tribuloides* orthologs were forced into these *S. purpuratus* clusters and the highest membership value for each ortholog was averaged with other orthologs expressed in each germ layer, *E. tribuloides* orthologs expressed in ectodermal domains exhibited higher cluster membership scores than orthologs expressed in mesodermal domains (Fig. 5C, $p < 0.05$, Mann-Whitney U Test). Additionally, ectodermal orthologs jumped less frequently between clusters than mesodermal orthologs, suggesting timecourse data for ectodermal genes were more similar than those mesodermal genes (Fig. 5D and S8C). These analyses suggest that deployment of ectodermal regulatory genes is more similar between euechinoids and cidaroids than deployment of mesodermal regulatory genes. Further,

Discussion

Extensive divergence of ectodermal and mesodermal regulatory linkages has occurred in echinoid developmental GRNs

This study investigated embryonic pattern formation and regulatory interactions of the euechinoid ectodermal and mesodermal GRNs in a distantly related cidaroid. By contrasting our

findings with regulatory diagram of the *S. purpuratus* developmental GRN, we are able to better understand how the regulatory interactions patterning the early embryo have diverged since the cidaroids and euechinoids last shared a common ancestor in the Paleozoic 268.8 mya (Fig. 6). Numerous regulatory interactions were found to have diverged in these two clades, including (Fig 6A). Statistical analyses of comparative timecourse gene expression data revealed that gene expression profiles of regulatory gene orthologs expressed in ectodermal cell lineages are more similar to their euechinoid counterparts than those expressed in mesodermal cell lineages. This result is consistent the with the broad differences observed in cidaroids regarding SM specification, as well as that revealed here in NSM segregation and regulatory states. By contrasting these observations with those in other echinoderms, we can begin to appreciate the degree to which embryonic developmental GRNs are constrained or altered over vast evolutionary distances and can reconstruct the ancestral regulatory states that must have existed in the embryos of echinoderm ancestors (47). In the following we attempt to interpret the preceding results in an evolutionary context within the echinoderm phylum.

Divergent regulatory interactions in the evolution of echinoid ectodermal and mesodermal GRNs

Our analysis utilized the regulatory transactions of the well-known *S. purpuratus* GRN as a basis for comparison between the cidaroid *E. tribuloides* and euechinoids (Fig. 6A). The comparative approach combined with the well-documented fossil record of echinoids allowed us to reveal a first approximation of the magnitude of change incurred by these GRNs since these two clades last shared a common ancestor. Spatiotemporal expression patterns of ectodermal regulatory genes in *E. tribuloides* and euechinoids strongly suggest that alteration to this circuitry is nontrivial in early development relative to the circuitry directing development of mesodermal lineages. However, while numerous regulatory transactions are conserved in the ectodermal GRN,

deployment and rewiring of circuitry during the evolution of euechinoid lineages that possess

direct-developing, non-feeding larvae has occurred frequently (52, 78-80). And as we have seen

from this study, our data indicate that the ectodermal GRN has undergone at least one significant

alteration since the cidaroid-euechinoid divergence, viz. the *bmp2/4, tbx2/3* and *irxa.* These

observations support the idea that the constraints on alteration of GRN circuitry are not sufficient to

maintain these linkages over vast evolutionary timescales. For instance, perturbation of Nodal

signaling revealed that, while initial specification events are highly similar, alterations likely have

occurred to the regulation of *bmp2/4* and *tbx2/3.* In *E. tribuloides tbx2/3* is expressed in AE and

aboral NSM by mid-gastrula. By late gastrula, it is expressed in the lateral clusters of skeletogenic

synthesis, at the tip of the gut, in the gut endoderm, and residually in the ectoderm. This unfolding

pattern of *tbx2/3* expression in *E. tribuloides* has essentially been compressed into the early stages

of euechinoid development (60). In euechinoids, perturbation of Nodal signaling with SB431542

extinguishes aboral ectodermal *tbx2/3* specifically in *P. lividus*, while not affecting its expression in

SM (40). In *E. tribuloides* we observed the expression domain of *tbx2/3* expand into OE upon

perturbation with this inhibitor (Fig. 2B and S5C). This observation combined with the result that

bmp2/4 responds differently to Nodal perturbation suggests altered GRN circuitry downstream of

Nodal*.*

Evolution of ciliary band patterning mechanisms in echinoderms

Ciliary band formation and ANE patterning in *E. tribuloides* are evolutionarily interesting

given their position in the echinoderm phylogeny and given that cidaroids lack the pan-

deuterostome apical senory organ (12, 13, 81, 82). Previous studies indicated that ANE patterning

in *E. tribuloides* is more similar to outgroup echinoderms than it is in euechinoids; though

expression of CB and anterior regulatory factors, e.g. *onecut* and *nk2.1,* exhibited spatial

distributions similar to those seen in euechinoids (75). Here, we observed patterning and

regulation of CB that are consistent with the hypothesis that this process is conserved in echinoids

(Fig. 6B). Additionally, we observed the sequential spatial restriction of *foxq2* to ANE, a pan-

bilaterian observation driven by endomesodermal *wnt* factors (23, 83-86). These data suggest that

specification of the apical sensory organ in *E. tribuloides* is developmentally downstream of these

events and that the loss of this embryonic structure had little effect on conserved patterning of CB

and anterior localization of *foxq2*.

Regulatory states and polarity of NSM in *E. tribuloides*

Examining mesodermal polarity in euechinoids and outgroup echinoderms aids in the

establishment of a timeline of GRN evolution. In euechinoid mesodermal NSM, *gcm* is directly

downstream of Notch signaling and later is restricted prior to gastrulation to aboral NSM by Not,

which is directly downstream of Nodal/SMAD signaling (37, 71, 87, 88). In cidaroids, early

expression of *gcm* is not dependent on Delta/Notch [Erkenbrack, Davidson and Peter, forthcoming]

and its mesodermal polarity is only apparent 4-6 hours after gastrulation has begun. Thus, if *gcm* is

near the top of the NSM GRN in cidaroids (16), as is the case in euechinoids (88), then this

pregastrular NSM polarization can be viewed as a euechinoid synapomorphy. This hypothesis is

supported by the observation that no significant polarity occurs in mesodermal specification of

asteroids and holothuroids (89). Here we showed that the polarization of *gcm* and *ese* also seems

to be a derived feature of echinoids and the mechanism of NSM segregation via Nodal/SMAD

signaling is likely conserved (Fig. 6C). Two observations make *gcm* polarization via the

transcription factor *not* (27) likely to exist in *E. tribuloides*: (1) *not* expression is observed in the oral

side of the archenteron by early gastrula stage when *gcm* is spatially restricted (Fig. 1J4 and 1L3)

and (2) aboral localization of *gcm* does not obtain when O-A axis patterning, and thereby *not*

expression, is perturbed (Fig. 2B and S5C). Together these observations suggest a conserved role

for OE regulatory factors in patterning the NSM of echinoids and that, in the lineage leading to

modern euechinoids, deployment of GRN circuitry polarizing NSM underwent a heterochronic shift

in the lineage leading to euechinoids.

The data presented on O-A polarity in the NSM of *E. tribuloides* suggest that multiple

regulatory domains unfold at and around the tip of the archenteron as gastrulation proceeds. This

is also the case in euechinoids (90). However, this study determined that *ese* operates in the oral

NSM exclusive of *gcm* in the aboral NSM. While the regulatory states in *E. tribuloides* NSM need

further refinement by two-color WMISH, for our purposes the overt disorder in its formation relative

to the overt order of *S. purpuratus* NSM makes two salient points. First, early pregastrular or early

gastrular polarity of NSM regulatory states represents an echinoid novelty, as no evidence for early

mesodermal polarity exists in outgroup echinoderms (89, 91). Second, if we take *E. tribuloides* as

a proxy to the ancestral state for this character/regulatory state, then it is clear that the O-A polarity

observed in the euechinoid NSM was shifted to occur prior to gastrulation in the lineage leading to

modern euechinoids. On the other hand, an alternative evolutionary scenario is that NSM polarity

manifested in these two modern echinoids is the result of two independent evolutionary trajectories

with heterochronic and spatial differences, but both meeting a similar end in the diversification of

NSM cell types in early development. That at least two O-A regulatory states are common to these

embryos and that *gcm* is downstream of Nodal and much later is downstream of Notch signaling

provide support for the first scenario. Further investigation into the developmental timing and

regulatory states of NSM segregation in *E. tribuloides* and other cidaroids is required to parse out

the most likely evolutionary scenario of NSM evolution in echinoids.

442   **Evolution of global embryonic domains in early development of**

443   **echinoids**

Previous analyses of embryonic regulatory states in *E. tribuloides* surveyed SM regulatory

factors (17, 47) and anterior neural ectoderm specification (75). Additionally, two previous studies

investigated SM and early endomesodermal micromere regulatory factors in the Pacific-dwelling

cidaroid *Prionocidaris baculosa* (16, 92). Integrating these data into this study affords an analysis

of global embryonic regulatory states and GRN linkages over 268.8 mya of evolution in indirect-

developing sea urchins. From these studies, numerous alterations to deployment and GRN

circuitry at all levels of GRN topology can be enumerated. Here, we enumerate 19 changes in

spatiotemporal deployment or regulation of ectodermal and mesodermal embryonic regulatory

factors since the cidaroid-euechinoid divergence (Table S1). Prominent among alterations of

regulatory interactions are those that have occurred in establishing polarity in mesodermal

embryonic domains. Endodermal and ectodermal specification and regulatory states also have

undergone significant change, but to a lesser degree. One hypothesis that can accommodate

these observations is that endodermal and ectodermal developmental programs may be more

recalcitrant to change than mesodermal programs due to their more ancient evolutionary origin (5),

suggesting that accretion of process over evolutionary time is a mechanism of constraint in

developmental programs (5). Indeed, in euechinoids there have been additional layers of GRN

topology accrued in mesodermal specification, e.g. the *pmar1-hesc* double-negative gate novelty

(16, 17, 22), delta-dependent NSM specification (17, 34), etc., which cidaroids do not exhibit, and

which may explain the observation that little to no appreciable change has been observed in the

mesodermal developmental programs of *L. variegatus, P. lividus* and *S. purpuratus,*

representatives of modern euechinoid lineages that diverged approximately 90 mya.

Since the divergence of cidaroids and euechinoids, numerous alterations to developmental

GRNs have accrued in these lineages. This study revealed that changes to mesodermal GRN

architecture have occurred more frequently than alterations to ectoderm GRN architecture. These

results support the notion that GRN architecture evolves at different rates (93) and provide an in

principle explanation for the rapid evolution observed in both cidaroid and euechinoid sea urchin

lineages that have independently evolved direct-developing larval forms (53). It remains to be

determined by future research efforts whether the shared regulatory states between cidaroids and

euechinoids elucidated here are the product of conserved stretches of genomic DNA hardwired in

the *cis*-regulatory regions of the orthologous regulatory genes or whether they are the result of

diverged *cis*-regulatory modules producing similar developmental outcomes.

475 Materials and Methods

Animal and embryo culture, cloning, acquisition of spatial and temporal gene expression data and

microinjection perturbation data were as previously described (17, 47). Dosage of small molecule

inhibitors DAPT and SB431542 were determined by dilution series. Concentrations for DAPT and

SB431542 were 12 µM and 15 µM respectively. Additional details for previously described

methods, statistical analyses and all other experimental manipulations can be found in SI Materials

and Methods.

482 Acknowledgments

We appreciatively acknowledge numerous colleagues for discussing various aspects of this work
with us over the long course of its maturation: Dr. Julius C. Barsi kindly provided his knowledge
and the *Sp onecut* engineered BACs; Jennifer Wellman, M.S., for her insight on statistical
analyses; Stefan Materna provided raw data for *S. purpuratus* mRNA transcript abundance and
reviewed early versions of the manuscript; Matthias Futschik provided mfuzz analysis advice; Drs.
Oliver Griffith, David McClay, Tom Stewart and Jeffrey R. Thompson provided critical commentary
on figures and the manuscript. The commentary of two anonymous reviewers greatly improved
data analysis and the manuscript. This work was developed and written in the laboratory of
Professor Günter P. Wagner; his patience and support are fondly appreciated. This work was
funded by National Science Foundation CREATIV grant #1240626.
493

494 References

1.   Davidson EH (1986) *Gene activity in early development* (Academic Press, New York) 3rd
Ed.
2.   Wikramanayake AH*, et al.* (2003) An ancient role for nuclear beta-catenin in the evolution of
axial polarity and germ layer segregation. *Nature* 426(6965):446-450.
3.   Davidson EH (2006) *The regulatory genome: gene regulatory networks in development and
evolution* (Academic, Oxford).
4.   Peter IS & Davidson EH (2015) *Genomic Control Process, Development and Evolution*
(Academic Press, Oxford).
5.   Hashimshony T, Feder M, Levin M, Hall BK, & Yanai I (2015) Spatiotemporal
transcriptomics reveals the evolutionary history of the endoderm germ layer. *Nature*
519(7542):219-222.
6.   Davidson EH (1991) Spatial mechanisms of gene regulation in metazoan embryos.
Development 113(1):1-26.
7.   Davidson EH, Cameron RA, & Ransick A (1998) Specification of cell fate in the sea urchin
embryo: summary and some proposed mechanisms. *Development* 125(17):3269-3290.
8.   Wray GA & Bely AE (1994) The evolution of echinoderm development is driven by several
distinct factors. *Development* 1994(Supplement):97-106.
9.   Kroh A & Smith AB (2010) The phylogeny and classification of post-Palaeozoic echinoids. *J
Syst Palaeontol* 8(2):147-212.
10.   Thompson JR*, et al.* (2015) Reorganization of sea urchin gene regulatory networks at least
268 million years ago as revealed by oldest fossil cidaroid echinoid. *Sci Rep-Uk* 5.
11.   Tennent D (1914) The early influence of the spermatazoan upon the characters of echinoid
larvae. *Carn. Inst. Wash. Publ.* 182:129-138.
12.   Mortensen T (1938) Contributions to the study of the development and larval forms of
echinoderms. IV. *Danske Vid. Selsk. Ser.* 9(7(3)):1-59.
13.   Schroeder TE (1981) Development of a 'primitive' sea urchin (Eucidaris tribuloides):
irregularities in the hyaline layer, micromeres, and primary mesenchyme. *Biological Bulletin*
161(1):141-151.
14.   Wray GA & McClay DR (1988) The origin of spicule-forming cells in a 'primitive' sea urchin
(Eucidaris tribuloides) which appears to lack primary mesenchyme cells. *Development*
103(2):305-315.
15.   Wray GA & McClay DR (1989) Molecular heterochronies and heterotopies in early echinoid
development. *Evolution*:803-813.
16.   Yamazaki A, Kidachi Y, Yamaguchi M, & Minokawa T (2014) Larval mesenchyme cell
specification in the primitive echinoid occurs independently of the double-negative gate.
Development 141(13):2669-2679.
17.   Erkenbrack EM & Davidson EH (2015) Evolutionary Rewiring of Gene Regulatory Network
Linkages at Divergence of the Echinoid Subclasses. *PNAS*.
18.   Gao F*, et al.* (2015) Juvenile skeletogenesis in anciently diverged sea urchin clades. *Dev
Biol* 400(1):148-158.
19.   Angerer LM*, et al.* (2000) A BMP pathway regulates cell fate allocation along the sea urchin
animal-vegetal embryonic axis. *Development* 127(5):1105-1114.
20.   Davidson EH*, et al.* (2002) A genomic regulatory network for development. *Science*
295(5560):1669-1678.
21.   Revilla-i-Domingo R, Oliveri P, & Davidson EH (2007) A missing link in the sea urchin
embryo gene regulatory network: hesC and the double-negative specification of
micromeres. *Proc Natl Acad Sci U S A* 104(30):12383-12388.

22.   Oliveri P, Tu Q, & Davidson EH (2008) Global regulatory logic for specification of an
embryonic cell lineage. *Proc Natl Acad Sci U S A* 105(16):5955-5962.
23.   Yaguchi S, Yaguchi J, Angerer RC, & Angerer LM (2008) A Wnt-FoxQ2-nodal pathway links
primary and secondary axis specification in sea urchin embryos. *Developmental cell*
14(1):97-107.
24.   Su Y-H*, et al.* (2009) A perturbation model of the gene regulatory network for oral and
aboral ectoderm specification in the sea urchin embryo. *Developmental biology* 329(2):410-
421.
25.   Peter IS & Davidson EH (2010) The endoderm gene regulatory network in sea urchin
embryos up to mid-blastula stage. *Dev Biol* 340(2):188-199.
26.   Peter IS & Davidson EH (2011) A gene regulatory network controlling the embryonic
specification of endoderm. *Nature* 474(7353):635-639.
27.   Materna SC, Ransick A, Li E, & Davidson EH (2013) Diversification of oral and aboral
mesodermal regulatory states in pregastrular sea urchin embryos. *Dev Biol* 375(1):92-104.
28.   Barsi JC, Li EH, & Davidson EH (2015) Geometric control of ciliated band regulatory states
in the sea urchin embryo. *Development* 142(5):953-961.
29.   Nam JM*, et al.* (2008) Cis-regulatory control of the nodal gene, initiator of the sea urchin oral
ectoderm gene network. *Faseb J* 22.
30.   Cui M, Siriwon N, Li E, Davidson EH, & Peter IS (2014) Specific functions of the Wnt
signaling system in gene regulatory networks throughout the early sea urchin embryo.
Proceedings of the National Academy of Sciences 111(47):E5029-E5038.
31.   Wikramanayake AH, Huang L, & Klein WH (1998) beta-Catenin is essential for patterning
the maternally specified animal-vegetal axis in the sea urchin embryo. *P Natl Acad Sci USA*
95(16):9343-9348.
32.   Sherwood DR & McClay DR (1999) LvNotch signaling mediates secondary mesenchyme
specification in the sea urchin embryo. *Development* 126(8):1703-1713.
33.   Logan CY, Miller JR, Ferkowicz MJ, & McClay DR (1999) Nuclear beta-catenin is required
to specify vegetal cell fates in the sea urchin embryo. *Development* 126(2):345-357.
34.   Sweet HC, Gehring M, & Ettensohn CA (2002) LvDelta is a mesoderm-inducing signal in the
sea urchin embryo and can endow blastomeres with organizer-like properties. *Development*
129(8):1945-1955.
35.   Flowers VL, Courteau GR, Poustka AJ, Weng W, & Venuti JM (2004) Nodal/activin signaling
establishes oral–aboral polarity in the early sea urchin embryo. *Developmental dynamics*
231(4):727-740.
36.   Ettensohn CA, Kitazawa C, Cheers MS, Leonard JD, & Sharma T (2007) Gene regulatory
networks and developmental plasticity in the early sea urchin embryo: alternative
deployment of the skeletogenic gene regulatory network. *Development* 134(17):3077-3087.
37.   Croce JC & McClay DR (2010) Dynamics of Delta/Notch signaling on endomesoderm
segregation in the sea urchin embryo. *Development* 137(1):83-91.
38.   Duboc V, Rottinger E, Besnardeau L, & Lepage T (2004) Nodal and BMP2/4 signaling
organizes the oral-aboral axis of the sea urchin embryo. *Developmental Cell* 6(3):397-410.
39.   Duboc V, Lapraz F, Besnardeau L, & Lepage T (2008) Lefty acts as an essential modulator
of Nodal activity during sea urchin oral-aboral axis formation. *Developmental Biology*
320(1):49-59.
40.   Saudemont A*, et al.* (2010) Ancestral Regulatory Circuits Governing Ectoderm Patterning
Downstream of Nodal and BMP2/4 Revealed by Gene Regulatory Network Analysis in an
Echinoderm. *Plos Genet* 6(12).

41.   Lhomond G, McClay DR, Gache C, & Croce JC (2012) Frizzled1/2/7 signaling directs beta-
590   catenin nuclearisation and initiates endoderm specification in macromeres during sea urchin
591   embryogenesis. *Development* 139(4):816-825.
592   42.   Cavalieri V & Spinelli G (2014) Early asymmetric cues triggering the dorsal/ventral gene
593   regulatory network of the sea urchin embryo. *Elife* 3:e04664.
594   43.   Smith AB*, et al.* (2006) Testing the molecular clock: molecular and paleontological estimates
595   of divergence times in the Echinoidea (Echinodermata). *Mol Biol Evol* 23(10):1832-1851.
596   44.   Croce J*, et al.* (2011) Wnt6 activates endoderm in the sea urchin gene regulatory network.
597   *Development* 138(15):3297-3306.
598   45.   Ettensohn CA (2009) Lessons from a gene regulatory network: echinoderm skeletogenesis
599   provides insights into evolution, plasticity and morphogenesis. *Development* 136(1):11-21.
600   46.   Molina MD, de Croze N, Haillot E, & Lepage T (2013) Nodal: master and commander of the
601   dorsal-ventral and left-right axes in the sea urchin embryo. *Current Opinion in Genetics &*
602   *Development* 23(4):445-453.
603   47.   Erkenbrack EM*, et al.* (2016) Ancestral state reconstruction by comparative analysis of a
604   GRN kernel operating in echinoderms. *Dev Genes Evol* 226(1):37-45.
605   48.   Lapraz F, Haillot E, & Lepage T (2015) A deuterostome origin of the Spemann organiser
606   suggested by Nodal and ADMPs functions in Echinoderms. *Nat Commun* 6:8927.
607   49.   Duboc V & Lepage T (2008) A conserved role for the nodal signaling pathway in the
608   establishment of dorso-ventral and left-right axes in deuterostomes. *J Exp Zool Part B*
609   310b(1):41-53.
610   50.   Wray GA & Raff RA (1991) The evolution of developmental strategy in marine invertebrates.
611   *Trends Ecol Evol* 6(2):45-50.
612   51.   Smith MS, Collins S, & Raff RA (2009) Morphogenetic mechanisms of coelom formation in
613   the direct-developing sea urchin Heliocidaris erythrogramma. *Dev Genes Evol* 219(1):21-29.
614   52.   Smith MS, Turner FR, & Raff RA (2008) Nodal Expression and Heterochrony in the
615   Evolution of Dorsal-Ventral and Left-Right Axes Formation in the Direct-Developing Sea
616   Urchin Heliocidaris erythrogramma. *J Exp Zool Part B* 310b(8):609-622.
617   53.   Raff RA (2008) Origins of the other metazoan body plans: the evolution of larval forms.
618   *Philos T R Soc B* 363(1496):1473-1479.
619   54.   Su Y-H (2009) Gene regulatory networks for ectoderm specification in sea urchin embryos.
620   *Biochimica et Biophysica Acta (BBA)-Gene Regulatory Mechanisms* 1789(4):261-267.
621   55.   Li E, Cui M, Peter IS, & Davidson EH (2014) Encoding regulatory state boundaries in the
622   pregastrular oral ectoderm of the sea urchin embryo. *Proceedings of the National Academy*
623   *of Sciences* 111(10):E906-E913.
624   56.   Li E, Materna SC, & Davidson EH (2012) Direct and indirect control of oral ectoderm
625   regulatory gene expression by Nodal signaling in the sea urchin embryo. *Dev Biol*
626   369(2):377-385.
627   57.   Li E, Materna SC, & Davidson EH (2013) New regulatory circuit controlling spatial and
628   temporal gene expression in the sea urchin embryo oral ectoderm GRN. *Dev Biol*
629   382(1):268-279.
630   58.   Lapraz F, Besnardeau L, & Lepage T (2009) Patterning of the Dorsal-Ventral Axis in
631   Echinoderms: Insights into the Evolution of the BMP-Chordin Signaling Network. *Plos*
632   *Biology* 7(11).
633   59.   de-Leon SB-T, Su Y-H, Lin K-T, Li E, & Davidson EH (2013) Gene regulatory control in the
634   sea urchin aboral ectoderm: spatial initiation, signaling inputs, and cell fate lockdown.
635   *Developmental biology* 374(1):245-254.

60. Croce J, Lhomond G, & Gache C (2003) Coquillette, a sea urchin T-box gene of the Tbx2
subfamily, is expressed asymmetrically along the oral-aboral axis of the embryo and is
involved in skeletogenesis. *Mech Dev* 120(5):561-572.
61. Gross JM, Peterson RE, Wu SY, & McClay DR (2003) LvTbx2/3: a T-box family
transcription factor involved in formation of the oral/aboral axis of the sea urchin embryo.
Development 130(9):1989-1999.
62. Range R (2014) Specification and positioning of the anterior neuroectoderm in
deuterostome embryos. *genesis* 52(3):222-234.
63. Strathmann RR (1971) The feeding behavior of planktotrophic echinoderm larvae:
mechanisms, regulation, and rates of suspensionfeeding. *Journal of Experimental Marine*
Biology and Ecology 6(2):109-160.
64. Wray GA (1992) The Evolution of Larval Morphology during the Postpaleozoic Radiation of
Echinoids. *Paleobiology* 18(3):258-287.
65. Barsi JC & Davidson EH (2016) cis-Regulatory control of the initial neurogenic pattern of
onecut gene expression in the sea urchin embryo. *Developmental Biology* 409(1):310-318.
66. Poustka AJ*, et al.* (2007) A global view of gene expression in lithium and zinc treated sea
urchin embryos: new components of gene regulatory networks. *Genome Biol* 8(5):R85.
67. Otim O, Amore G, Minokawa T, McClay DR, & Davidson EH (2004) SpHnf6, a transcription
factor that executes multiple functions in sea urchin embryogenesis. *Dev Biol* 273(2):226-
243.
68. Cameron RA & Davidson EH (1991) Cell type specification during sea urchin development.
Trends Genet 7(7):212-218.
69. Sweet HC, Hodor PG, & Ettensohn CA (1999) The role of micromere signaling in Notch
activation and mesoderm specification during sea urchin embryogenesis. *Development*
126(23):5255-5265.
70. Materna SC & Davidson EH (2012) A comprehensive analysis of Delta signaling in pre-
gastrular sea urchin embryos. *Dev Biol* 364(1):77-87.
71. Duboc V*, et al.* (2010) Nodal and BMP2/4 pattern the mesoderm and endoderm during
development of the sea urchin embryo. *Development* 137(2):223-235.
72. Lee PY & Davidson EH (2004) Expression of Spgatae, the Strongylocentrotus purpuratus
ortholog of vertebrate GATA4/5/6 factors. *Gene Expression Patterns* 5(2):161-165.
73. Duboc V, Rottinger E, Lapraz F, Besnardeau L, & Lepage T (2005) Left-right asymmetry in
the sea urchin embryo is regulated by nodal signaling on the right side. *Developmental Cell*
9(1):147-158.
74. Bradham CA*, et al.* (2009) Chordin is required for neural but not axial development in sea
urchin embryos. *Dev Biol* 328(2):221-233.
75. Bishop CD, MacNeil KE, Patel D, Taylor VJ, & Burke RD (2013) Neural development in
Eucidaris tribuloides and the evolutionary history of the echinoid larval nervous system. *Dev*
Biol 377(1):236-244.
76. Materna SC, Nam J, & Davidson EH (2010) High accuracy, high-resolution prevalence
measurement for the majority of locally expressed regulatory genes in early sea urchin
development. *Gene Expr Patterns* 10(4-5):177-184.
77. Gildor T & de-Leon SB-T (2015) Comparative Study of Regulatory Circuits in Two Sea
Urchin Species Reveals Tight Control of Timing and High Conservation of Expression
Dynamics. *Plos Genet* 11(7):e1005435.
78. Wilson KA, Andrews ME, Turner FR, & Raff RA (2005) Major regulatory factors in the
evolution of development: the roles of goosecoid and Msx in the evolution of the direct-
developing sea urchin Heliocidaris erythrogramma. *Evolution & Development* 7(5):416-428.

79. Wilson KA, Andrews ME, & Raff RA (2005) Dissociation of expression patterns of
homeodomain transcription factors in the evolution of developmental mode in the sea
urchins Heliocidaris tuberculata and H-erythrogramma. *Evolution & Development* 7(5):401-
415.
80. Raff RA & Smith MS (2009) Axis Formation and the Rapid Evolutionary Transformation of
Larval Form. *Current Topics in Developmental Biology, Vol 86* 86:163-190.
81. Emlet RB (1988) Larval Form and Metamorphosis of a Primitive Sea-Urchin, Eucidaris-
Thouarsi (Echinodermata, Echinoidea, Cidaroida), with Implications for Developmental and
Phylogenetic Studies. *Biological Bulletin* 174(1):4-19.
82. Bennett KC, Young CM, & Emlet RB (2012) Larval Development and Metamorphosis of the
Deep-Sea Cidaroid Urchin Cidaris blakei. *Biological Bulletin* 222(2):105-117.
83. Tu Q, Brown CT, Davidson EH, & Oliveri P (2006) Sea urchin Forkhead gene family:
phylogeny and embryonic expression. *Dev Biol* 300(1):49-62.
84. Santagata S, Resh C, Hejnol A, Martindale MQ, & Passamaneck YJ (2012) Development of
the larval anterior neurogenic domains of Terebratalia transversa (Brachiopoda) provides
insights into the diversification of larval apical organs and the spiralian nervous system.
EvoDevo 3(1):1.
85. Yu JK*, et al.* (2008) The Fox genes of Branchiostoma floridae. *Dev Genes Evol* 218(11-
12):629-638.
86. Fritzenwanker JH, Gerhart J, Freeman RM, & Lowe CJ (2014) The Fox/Forkhead
transcription factor family of the hemichordate Saccoglossus kowalevskii. *Evodevo* 5.
87. Ransick A, Rast JP, Minokawa T, Calestani C, & Davidson EH (2002) New early zygotic
regulators expressed in endomesoderm of sea urchin embryos discovered by differential
array hybridization. *Dev Biol* 246(1):132-147.
88. Ransick A & Davidson EH (2006) cis-regulatory processing of Notch signaling input to the
sea urchin glial cells missing gene during mesoderm specification. *Dev Biol* 297(2):587-602.
89. McCauley BS, Wright EP, Exner C, Kitazawa C, & Hinman VF (2012) Development of an
embryonic skeletogenic mesenchyme lineage in a sea cucumber reveals the trajectory of
change for the evolution of novel structures in echinoderms. *Evodevo* 3(1):17.
90. Andrikou C, Pai CY, Su YH, & Arnone MI (2015) Logics and properties of a genetic
regulatory program that drives embryonic muscle development in an echinoderm. *Elife* 4.
91. Dylus DV*, et al.* (2016) Large-scale gene expression study in the ophiuroid Amphiura
filiformis provides insights into evolution of gene regulatory networks. *Evodevo* 7.
92. Yamazaki A, Kidachi Y, & Minokawa T (2012) "Micromere" formation and expression of
endomesoderm regulatory genes during embryogenesis of the primitive echinoid
Prionocidaris baculosa. *Dev Growth Differ* 54(5):566-578.
93. Davidson EH & Erwin DH (2006) Gene regulatory networks and the evolution of animal
body plans. *Science* 311(5762):796-800.
722

Figure Legends

Fig 1. Spatiotemporal dynamics of eleven euechinoid ectodermal and mesodermal

regulatory genes in the cidaroid *E. tribuloides*. Visualization of mRNA transcripts revealed by

whole mount in situ hybridization, and estimates of absolute mRNA transcript abundance

determined by qPCR during first 35 hours post fertilization (hpf). Individual data points are light

grey. Blue data points represent the mean at that particular timepoint. Red dashed lines indicate

the onset of zygotic transcription. Orange asterisks denote position of archenteron. (A1) Key to

embryonic domains of expression and orientation of embryo micrographs. The embryonic domain

of expression of each gene is indicated by colored boxes: ANE (anterior neural ectoderm), pink;

oral ectoderm (OE), yellow; ciliary band (CB), blue; aboral ectoderm (AE), green; NSM (non-

skeletogenic mesoderm), brown. (B1-B5) Spatiotemporal expression of *foxq2. Foxq2* spatial

distribution is detected prior to hatching and is spatially restricted to ANE by 17 hpf. (C1-C5)

Spatiotemporal expression of *irxa*. At 28 hpf, *irxa* is detected in AE and extends from the vegetal

endodermal domains to ANE. By 40 hpf *irxa* is seen extending anteriorly at the boundary of AE

and OE. (D1-D5) Spatiotemporal expression of *nodal*. *Nodal* spatial distribution is restricted to a

few cells in OE up to early-mid gastrula stage. (E1-E5) Spatiotemporal expression of *chordin.* First

observed in a few cells in OE at 17 hpf, *chordin* subsequently expands to extend from the perianal

ectoderm to ANE. (F1-F5) Spatiotemporal dynamics of *msx.* Detected exclusively in AE, *msx*

extends from the blastopore to ANE. (G1-G5) Spatiotemporal expression of *gsc.* By 22 hpf, *gsc* is

detected in a broad region surrounding OE and later is observed around the stomodeum. (H1-H5)

Spatiotemporal dynamics of *onecut.* By 40 hpf *onecut* is detected in the future post oral ciliary

band and is initiated in a band moving from the posterior to the anterior. (I1-I5) Spatiotemporal

expression of *tbx2/3.* Spatial distribution of *tbx2/3* at 17 hpf is detected broadly in aboral ectoderm

(AE) and later extends from the perianal ectoderm to lateral AE, but not past the embryonic

equator. (J1-J5) Spatiotemporal expression of *not. Not* spatial distribution is first detected in a

similar field of cells as *nodal* and *lefty;* however, the domain of *not* subsequently expands by 28 hpf

where it is seen in the oral side of the archenteron, where non-skeletogenic mesoderm (NSM) and

endoderm are being segregated. By 40 hpf, the spatial domain of *not* extends from anterior neural

ectoderm to the perianal ectoderm, is clearly seen in NSM, and was not detected in endodermal

lineages. (K1-K5) Spatiotemporal expression of *ese. Ese* is detected primarily in non-skeletogenic

mesoderm (NSM) but also in anterior neural ectoderm. By 28 hpf, *ese* is observed at the tip of the

archenteron and is asymmetrically polarized in NSM. (L1-L5) Spatiotemporal expression of *gcm.*

As gastrulation begins *gcm* is expressed broadly in NSM; however by 28 hpf it exhibits stark

polarity in a field of cells that resides basal to the tip of the archenteron. *Gcm* is also observed in a

few ectodermal cells at the time of primary mesenchymal ingression.

Fig 2. Perturbation of oral-aboral axis formation in *E. tribuloides* reveals alterations to

regulatory factor deployment in echinoids. (A1) Quantitative effect of SB431542 on expression

of 30 *E. tribuloides* regulatory genes as revealed by qPCR. Fold change in mRNA transcripts

(ddCt) is shown on the y-axis. Two timepoints from two independent replicates are shown.

Regulatory factors are listed on the x-axis and font color designates their embryonic domain: pink,

anterior neural ectoderm; green, aboral ectoderm; yellow, oral ectoderm; black, endoderm; brown,

mesoderm. (B) WMISH of select ectodermal and mesodermal regulatory genes in embryos

cultured in the presence of SB431542. At 28 hpf expression of *chordin, nodal* and *not* are

completely extinguished. Whereas *gcm* is regularly restricted to one side of the archenteron, in the

presence of SB431542 it exhibits expression throughout the archenteron. In the ectoderm,

expression *tbx2/3* expands from aboral ectoderm (AE) into oral ectoderm in the presence of the

inhibitor. (C) The ciliary band marker *onecut* is normally observed in a band of cells between the

boundaries of OE and AE. However, in the presence of SB431542, *onecut* is expressed in an

equatorial band that is 6-10 cell diameters across.

Fig 3. Expression dynamics in *E. tribuloides* of engineered reporter BACs harboring the

regulatory locus of *S. purpuratus onecut* suggest conservation of GRN circuitry positioning

ciliary band in echinoids . (A) Schematic of ciliary band (CB) restriction in the *S. purpuratus*

embryo. Red bars indicate repressor genes restrict CB genes from expression in those domains.

(B) Schematic showing the *S. purpuratus onecut* locus and the *cis*-regulatory modules (CRMs)

responsible for the geometric positioning of *onecut.* (C) Table showing reporter analysis of spatial

expression of five engineered *S. purpuratus onecut* BACs in *E. tribuloides.* The BACs were

previously utilized to analyze the *cis*-regulatory dynamics of *onecut* spatial restriction in *S.*

purpuratus (Barsi and Davidson, 2016)*.* (D) BAC reporter 1, which harbors mutations to all known

cis-regulatory modules (CRMs), showed a marked reduction of all reporter expression. (E) BAC

reporter 2 harbors mutations to a CRM driving zygotic activation. (F) BAC Reporter 4, which

harbors mutations to the aboral and oral ectodermal repression CRMs, increases the percent of

embryos that show reporter expression in the oral ectoderm, as well as normal reporter expression

in the ciliary band. (G) BAC reporter 3, which harbor mutated post-activation enhancer CRMs,

showed reduced reporter expression both in terms of percent embryos exhibiting reporter activity

and in the number of cells with reporter. (H) BAC Reporter 5, which harbors the unperturbed, wild-

type locus of *S. purpuratus onecut,* faithfully exhibits reporter GFP in the ciliary band domain of *E.*

tribuloides.

Fig 4. Comparative gene expression analysis of echinoid regulatory orthologs.

Strongylocentrotus purpuratus, purple line*; Paracentrotus lividus,* green dashed line; *Eucidaris*

tribuloides, black dashed line*.* Region of expression is indicated by a colored box. Transcripts per

embryo for each gene were normalized to their maximal expression over the first 30 hours of

development and are plotted against *E. tribuloides* development on the x-ordinate. Comparative

developmental staging for each species is listed in Table S5. Each analysis is accompanied by a

matrix of Spearman correlation coefficients (indicated as ρ).

Fig 5. Statistical analyses of comparative timecourse data in the euechinoid *S. purpuratus*

and the cidaroid *E. tribuloides.* (A-B) Distribution plots of Spearman's rank correlation

coefficients (ρ) for temporal dynamics in *E. tribuloides* and *S. purpuratus*. Genes were binned by

embryonic domain of *S. purpuratus* spatial expression. Boxplot boundaries show interquartile

range, means and standard deviation. Asterisks mark statistical significance as determined by a

two-tailed t-test. (A) Boxplots for statistical distribution of endodermal, ectodermal and mesodermal

regulatory factors in *E. tribuloides* and *S. purpuratus.* Mean ρ for endodermal and ectodermal

regulatory factors were significantly higher than the mean ρ, marked as a red dashed line.

Mesodermal regulatory factors did not significantly vary from the mean. (B) Boxplots for statistical

distribution of subdomains of ectodermal and mesodermal regulatory factors in *E. tribuloides* and

S. purpuratus. (C) Mean mfuzz membership scores derived from cluster analysis of regulatory

orthologs. Genes were binned by their *S. purpuratus* embryonic domain of expression.

Membership scores for *E. tribuloides* orthologs were binned based on their spatial distribution and

compiled for statistical analysis. *E. tribuloides* ectodermal orthologs exhibited higher membership

scores than mesodermal orthologs (Mann Whitney U Test, p<0.05). (D) Histogram showing both

the distribution of tissue-specific orthologs examined in this study and the percent of *S. purpuratus*

and *E. tribuloides* orthologs clustering together. Ectodermal and endodermal orthologs clustered

together more often than mesodermal orthologs

Fig 6. Modeling divergent and conserved regulatory transactions since the divergence of

cidaroids and euechinoids. (A) A truncated *S. purpuratus* GRN showing ectodermal and

mesodermal regulatory interactions investigated in this study. Solid black lines indicated conserved

interactions; dashed black lines indicate interactions that are likely conserved; solid red lines

indicate divergent regulatory interactions. Genes and their *cis*-regulatory regions are represented

as flat lines with arrows. Embryonic domains of the echinoid embryo are represented by large

boxes: pink, anterior neural ectoderm (ANE); yellow, oral ectoderm (OE); blue, ciliary band (CB);

green, aboral ectoderm (AE); brown, oral non-skeletogenic mesoderm (NSM); light brown, aboral

non-skeletogenic mesoderm (NSM); white, skeletogenic mesoderm (SM). (B) Regulatory states

and restriction of ectodermal CB in asteroids, cidaroids and euechinoids. Red bars indicate

repressor genes restrict CB in a particular embryonic domain. Color scheme is the same as in (A).

The top row of schematics represents the wild-type state in three echinoderm taxa. The bottom

row represents the effects of perturbation of critical ectodermal regulatory genes. (C) Mechanisms

of polarization of NSM in echinoderms. Mesodermal segregation is represented in three

echinoderm taxa: Asteroids, Cidaroids, and Euechinoids. In both cidaroids and euehinoids, Nodal

signaling via the transcription factor Not segregates oral-aboral NSM cell types by restricting

regulatory factors to the aboral side of the embryo. Whereas asteroids show no indication of NSM

oral-aboral polarity, cidaroids and euechinoids lineages have undergone dramatic heterochronic

shifts since the divergence of these two clades, resulting in the diversification of NSM regulatory

states.

Fig 1

 Fig 2

Fig 3

SpOnecut BAC Reporter*	BAC alteration	Embryos scored	Expressing reporter	Territory of reporter expression			
				CB	AE	OE	Gut/Meso
BAC 1 (D)	all CRMs	77	0.0%	0.0%	0.0%	0.0%	0.0%
BAC 2 (E)	zygotic activation	56	12.5%	7.1%	0.0%	0.0%	0.0%
BAC 3 (G)	post-activation	40	10.0%	7.5%	0.0%	0.0%	0.0%
BAC 4 (F)	AE & OE clearance	123	27.6%	12.2%	2.4%	10.4%	0.0%
BAC 5 (H)	none	102	17.6%	14.7%	1.0%	0.0%	0.0%

Fig 4

Fig 5