Drosophila Mutants that Are Motile but Respond Poorly to All Stimuli Tested

Mutants that have a large defect in interaction with stimuli or in executive function

Lar L. Vang and Julius Adler*

Departments of Biochemistry and Genetics, University of Wisconsin-Madison,
Madison, Wisconsin 53706, USA

*e-mail: adler@biochem.wisc.edu

KEY WORDS: *Drosophila*, motility, behavior, attractants and repellents, sensory
reception, decision making, executive function

ABSTRACT

     Adult *Drosophila melanogaster* fruit flies were placed into one end of a tube near
to repellents (benzaldehyde and heat) and away from the other end containing attractants
(light and a favored temperature). They escaped from the repellents and went to the
attractants. Motile mutants that failed to do that were isolated. They did not respond to
any external attractants tested or external repellents tested. In addition, they did not
respond well to internal stimuli like hunger, thirst, and sleep. Some possibilities for
explaining the deficiency: They may be considered defective in decision making that
results from faulty interaction with the sensing mechanism, or, alternatively, the mutants
may be defective in executive function. The mutants failed at both 34°C and, in one case
studied, at room temperature.

I. INTRODUCTION

     To study the mechanism of behavior, one can isolate mutants and explore their
properties. In this report we describe the isolation and study of *Drosophila* mutants that
are motile and able to reproduce but they are poorly responsive to all stimuli tested, both
external and internal stimuli. Some possibilities for explaining this deficiency (see Fig1):

Fig 1. A proposed schematic interaction. Decision making for stimuli of sensory
reception and decision making for relation to executive function [1, 2].

1) an abnormal relation between decision making and sensory reception or 2) between decision making and executive function [1, 2]. See Discussion. A preliminary report of some of the results has been presented [3, 4].

II. RESULTS

A. RESPONSES TO EXTERNAL STIMULI

1. RESPONSE TO STIMULI USED TOGETHER

In a 34°C dark room flies were started near two repellents (0.1M benzaldehyde and 37°C) at one end of a tube, away from two attractants (light at 1000 lux and 27°C) at the other end (Fig 2). The parent responded by going away from the repellents and to the

Fig 2. Apparatus for isolating and testing mutants in a 34°C room. At the left end were repulsive 0.1M benzaldehyde and repulsive 37°C (due to a hot plate at 150°C). At the right end were attractive light (1000 lux) and attractive 27°C (due to ice water). The middle was close to 34°C.

attractants (Fig 3A). Mutants that were not motile were rejected, only the motile mutants were studied. This consisted of five mutants, named 1 to 5. Fig 3B shows that such a mutant (Mutant 2) failed to respond when the four stimuli are together. S3 Fig shows that the other four mutants (Mutants 1, 3, 4, and 5) also failed to respond to the four stimuli together.

Fig 3. Response to stimuli used together. Repellents (0.1M benzaldehyde and high temperature (37°C) were at the origin end, attractants (light, 1000 lux, and a favored temperature (27°C) at the furthest end. (**A**) Parental response (n=7). (**B**) Mutant 2 (n=8). Flies were tested in a 34°C room with 10 to 20 flies used per trial. Data are mean±SEM.

2. RESPONSE TO INDIVIDUAL STIMULI

When only light was presented to flies that were derived from ones that had already experienced the four stimuli used together, the parent went to the light source (Fig 4A) while Mutant 1 did not (Fig 4B). Mutants 2, 3, 4, and 5 did not respond to only light either as shown in S4 Fig.

Fig 4. Response to light alone. Light (1000 lux) was placed at the right end as in Fig. 2. (**A**) Parental response (n=4). (**B**) Mutant 1 response (n=5). Flies were tested at 34°C with 10 to 20 flies used per replicate. Data are mean±SEM.

A similar result was found for benzaldehyde alone: see parent being repelled in Fig 5A and the mutant, Mutant 1, not repelled in Fig 5B. S5 Fig shows the same for another mutant tested, Mutant 2.

Fig 5. Response to benzaldehyde alone. Benzaldehyde (0.1M) was placed at the left end as in Fig 2. (**A**) Parental response (n=4). (**B**) Mutant 1 response (n=6). Flies were tested at

34°C with 10 to 20 flies used per trial. Data are mean±SEM. (The mutant largely remained near the benzaldehyde perhaps because its motility may have been inhibited by the benzaldehyde.)

Such was also found for heat alone: see results for parent (Fig 6A) and mutant (Fig 6B). S6 Fig shows the same for another mutant tested.

Fig 6. Response to heat gradient alone. The heat source was placed at the left end as in Fig. 2. (**A**) Parental response (n=4). (**B**) Mutant 1 response (n=5). Flies were tested at 34°C with 10 to 20 flies used per trial; the warm side measuring 37°C and the cool side at 27°C (as in Figure 2). Data are mean±SEM.

Thus the mutants were defective not only for the four stimuli used together but also for each stimulus used alone.

3. RESPONSE TO OTHER EXTERNAL STIMULI

These mutants were in addition tested with stimuli that were not among those four used to obtain the mutants.

In the case of the repellent quinine, flies were started in a 0.1M quinine half and then they had the opportunity to go into a non-quinine half (see ref. 4 for details of the method). The parent went into the non-quinine half (Fig 7A), but the mutant did not (Fig 7B). See S7 Fig for additional data.

Fig 7. Response to quinine. (**A**) Parental response (n=5). (**B**) Mutant 2 response (n=5). Flies were tested at 34°C with 10 to 20 flies used per trial using 0.1M quinine. Data are mean±SEM.

These mutants were also tested for response to sucrose [5]. Compared to the wild-type, both Mutants 1 and 2 consumed significantly less sucrose, about 30 to 40% as much as the wild-type. See Fig 8 for details.

Fig 8. Response to sucrose. Between 15 and 20 flies were starved per trial on agar for 17 to 20 hours and then transferred to a tube containing agar and food dye FD & C Blue No. 1 solution for 5 minutes. These flies were homogenized and an absorbance was obtained. This was then compared to flies that were likewise starved on agar for 17 to 20 hours but were then transferred to a tube containing agar, food dye FD & C Blue No. 1, and 0.1M sucrose. No sucrose: Mutant 1 (n=3), Mutant 2 (n=3), and CS (n=3). With sucrose: Mutant 1 (n=2), Mutant 2 (n=2), and CS (n=3). Data are mean±SEM.

To test response to gravity, these flies were placed into a vertical tube and pounded down, then at every minute the flies in each third of the tube were counted (see ref. 4 for details of the method). The parent responded by climbing up (Fig 9A) while the mutant climbed up less (Fig 9B). See S9 Fig for additional data.

Fig 9. Response to gravity. (**A**) Parental response (n=5). (**B**) Mutant 2 response (n=4). Flies were tested at 34°C with 10 to 20 flies used per trial. Data are mean±SEM.

Thus these mutants, isolated by use of the four stimuli, were defective even for stimuli that were not present during their isolation.

4. MOVEMENT WITHOUT ANY ADDED STIMULI

In the absence of any stimulus added by the experimenters, the parent (Fig 10A) and the mutant (Fig 10B) moved similarly, indicating that motility alone is about the

Fig 10. Response without added stimuli. (**A**) Parental response (n=4). (**B**) Mutant 1 response (n=6). Flies were tested at 34°C with 10 to 20 flies used per trial. Data are mean±SEM.

same in parent and mutant. This was found also for Mutant 4 (S10 Fig). Mutants 2, 3, and 5 showed a slightly slower motility (S10 Fig). These results tell that the mutants were motile.

5. EFFECT OF INCUBATION TEMPERATURE

All the work reported above was carried out in a 34°C room in order to allow, if necessary, isolation and study of conditional mutants, i.e. mutants defective at 34°C but not defective at room temperature. We found that at room temperature, when each stimulus was used together, there was substantial condensation along the interior of the tube, resulting in the flies becoming stuck when emptied into the test apparatus. Therefore we chose to measure response to light (1000 lux) at room temperature (21 to 23°C). The parent responded to light but all five of the mutants failed to respond to light or responded only 10% as well as the parent (S11 Fig), just as they did at 34°C. Thus the mutations are likely not conditional.

Being defective to all stimuli tested at the usual growth temperature, how could the mutants survive and grow? We don't know, but it seems that due to the crowded conditions and rich medium found in the growth tube at room temperature this defect is compensated for somehow.

B. RESPONSES TO INTERNAL STIMULI

1. HUNGER

Here we focus on hunger [6-11]. To measure hunger we used an apparatus (Fig 11), inspired by and modified from an earlier design [12], described in Supplemental

Fig 11. Apparatus for measuring hunger and for measuring thirst. For details see IV. METHODS, C. How to study response to internal stimuli. Tube 1 is called "origin". Flies were tested at room temperature (21-23°C).

Methods (Supplement: C. How to study response to internal stimuli, 1. Measuring hunger). Briefly, in a dark room at 21-23°C male flies – parent or mutants - were transferred into one end (tube 1) of a 5 x 140 cm apparatus containing throughout its length a 5 cm wide strip of wet paper to satisfy thirst but containing no food. Starvation

for food began once the flies were put in. Every 10 hours up to 40 hours the location of the flies was measured with light on for 30 seconds.

At 20 hours the parent had largely left the origin (tube 1) and had begun to accumulate at the end (tube 4) (Fig 12A, solid bars), while the mutant had moved

Fig 12. Movement of flies at 20 hours in search for food. Solid: water but no food (no sucrose). Open: water and food (0.1M sucrose). (**A**) Parental response with water only (n=5) and with water + sucrose (n=9). (**B**) Mutant 2 response with water only (n=5) and with water + sucrose (n=4). Data are mean±SEM. (See S13 Fig for Mutant 1.) Flies were tested at room temperature (21-23°C) with 40 to 60 flies used per trial.

towards the end very little (Fig 12B, solid bars). This is interpreted to mean that the parent is searching for food while the mutant is defective in searching for food. The flies at 40 hours had moved further than at 20 hours but this is not illustrated because by 40 hours the death of the flies, presumably due to starvation, had become a serious interference.

When food (0.1M sucrose) was added throughout the tube along with the wet strip of paper, the parent moved less far (rather than accumulating at the end) (Fig 12A, open bars), while the mutant remained mostly where placed (Fig 12B, open bars). Since sucrose inhibited the movement of the parent, it is supposed that movement without sucrose is due largely to hunger. From these results we conclude that the mutants are defective in hunger.

2. THIRST

To study thirst, flies were deprived of water by a procedure described in Supplemental Methods (Supplement: C. How to study response to internal stimuli, 2.

Measuring thirst). The procedure is the same as for hunger (Fig 11) except that water was omitted and solid sucrose was layered throughout.

By 30 hours the parent had moved out, presumably to search for water since addition of water inhibited this (Fig 13A). The mutant moved out less well than the

Fig 13. Movement of flies at 30 hours in search for water. Solid: food (sucrose crystals) but no water. Open: food (0.1M sucrose) and water. (**A**) Parental response with sucrose only (n=5) and with water + sucrose (n=6). (**B**) Mutant 1 response with sucrose only (n=3) and with water + sucrose (n=3). Flies were tested at room temperature (21-23°C) with 40 to 60 flies used per trial. Data are mean±SEM. (See S13 Fig for Mutant 2 data.)

parent (Fig 13B), so we conclude that the mutants are defective in thirst. Already by 30 hours dying, presumably from thirst, had become a problem; by 40 hours this had become severe, so those results are not illustrated.

3. SLEEP-WAKE

The parent and mutants isolated here were studied for sleep and wake according to the procedure of Pfeiffenberger et al. [13]. The parent was different from the mutants (Fig 14). The parent showed greatest activity at the start and end of the day but not in the

Fig 14. Circadian response. Individual flies are placed into a tube (5 x 20 mm) with an infrared light beam intersecting at the middle of the tube. Mutant 1 (n=24), Mutant 2 (n=24), and parental response (n=24) are recorded over a 24 hour period at 22° C. Data are mean±SEM. (S14 Fig shows smaller differences between parent and three other mutants.)

middle of the day. Mutant 2 showed high activity throughout the day. Mutant 1 was less active than the parent at the start of the day.

C. MAPPING OF THE MUTANTS

We used recombination and deficiency kits [14-16]. Mutants isolated were mated to attached-X females. When presented with the four stimuli (light, favorable and unfavorable temperature, and benzaldehyde) together, attached-X females behaved wild-type (S15 Fig), indicating that both Mutants 1 and 2 are X-linked. The mutants were then crossed to an X chromosome bearing *y cho cv v f.*

Mutant 1 was found to be between *v* and *f,* approximately in position 12 or 13. Of the deficiencies that can be found for this region, we found that *Df(1)BSC705* uncovered Mutant 1. We then tested two deficiencies that overlap *Df(1)BSC705, Df(1)BSC628* and *Df(1)ED7229. Df(1)BSC705* uncovered Mutant 1 but *Df(1)ED7229* did not. Therefore the mutation must lie within *Df(1)BSC705* but to the left of *Df(1)BSC628*. Of the genes that were found within this region, 12E3 to 12E5, and that we tested, all complimented. This would suggest that the gene may lie in a small gap between the deficiencies.

Mutant 2 was found to be between *cv* and *v*, approximately in position 8 or 9. Of the deficiencies tested in this region, *Df(1)BSC539* uncovered Mutant 2. Two deficiencies

that overlap with *Df(1)BSC539, Df(1)BSC754* and *Df(1)ED7005*, did not uncover Mutant 2 and the mutation must then be in the *Df(1)BSC539* segment to the left of *Df(1)BSC754*. Of the genes that were found within this region and that we tested, CG1791 failed to compliment, so we conclude that Mutant 2 lies in the *CG1791* gene, which has been shown to be related to fibrinogen [17, 18]. The *CG1791* gene has 3 transcripts: RA, RB, and RC [19] but we do not yet know how *CG1791* is connected to decision making or executive function.

This new information can be used to go on to learn what the next steps are. Ultimately all the steps between sensory reception and response can be studied in mutants in order to define the entire sequence.

D. RELATION TO REPRODUCTION

Although the mutants reported here are defective in the responses to all external stimuli tested and all internal stimuli tested, nevertheless they are able to reproduce, so it must be that the mechanism for mating is still active in these mutants. In a normal fly, mating uses many systems: hearing, vision, smell, taste, and contact; mating can, however, take place without some of these [20-23], but presumably contact is still required since the male must touch the female in mating. Contact may well be intact in the mutants isolated here and perhaps presence of the flies in crowded tubes might make some of the other requirements for mating not necessary.

In our previous work on isolating mutants such as these by G. Tang, V. Toniolo, Y. H. Ng, L. Vang, and J. Adler [24], the mutants isolated failed to reproduce. Possibly they lacked mating due to lacking this contact. The procedure for isolating mutants was different from the procedure here: instead of using the four stimuli together (as described in Fig 2), the stimuli were used one at a time.

III. DISCUSSION

Here we describe the isolation and some properties of *Drosophila* mutants that are motile but yet they each fail in response to all external attractants and repellents tested (Figs 3-9) and also they are deficient in response to internal stimuli tested (Figs 12 and 13). Thus, although the mutants are motile, they have

1. decreased responsiveness to light
2. decreased responsiveness to repulsive chemicals (like benzaldehyde)
3. decreased responsiveness to heat and to favorable temperature
4. decreased responsiveness to bitter tastants (like quinine)
5. decreased responsiveness to sweet tastants (like sucrose)
6. decreased responsiveness to gravity
7. decreased responsiveness to hunger
8. decreased responsiveness to thirst
9. abnormality in some sleep

Because all of these different behaviors are defective in each of these mutants (Figs. 3-9, 12 and 13), it seems reasonable to say that there is a central place which is responsible. This might possibly be decision making acting on sensory reception, or decision making acting on executive function (Fig. 1).

Decision making involves interaction with stimuli. The simple sensing-response relationships long assumed to underlie behaviors is actually more complex [25]. When there are no stimuli added there is a resting state, called "default" or "intrinsic", which yet allows many behaviors to occur, and when stimuli are presented there is an activated state, called "reflexive" or "responsive" [26-28]. Our mutants could be defective in making this relation between resting state and activated state so that responses to all stimuli are poor. Presumably some factor is needed for this coming together of the resting state and the activated state, and that factor may be deficient in the mutants.

An alternative is that the mutants are defective elsewhere, rather than in the interaction between decision-making and stimuli. What is defective may be the interaction between decision-making and executive function [1, 2]. "Executive functions are the high-level cognitive processes that facilitate new ways of behaving…The operation of executive processes accompanies a very wide range of behaviours…Similar progress has been made on work in animals" not just in humans [1]. It may be that executive function responds to a hypothetical control of the organism called here "The Boss", which is inside every organism in charge of it [3].

The place where the consequences of action of these mutants appears is not known. Possibilities are: 1) Outside of known bodies but rather in the surrounding fibers [29, p. 540]. 2) In the mushroom bodies [29, pp. 523, 548; 30], but this is not likely because when we tested mutants in the mushroom bodies they had a normal response (see S16 Fig). 3) In the central complex [29 pp. 521, 31].

In the case of insects, known parts of the central complex consist of the protocerebral bridge, the fan-shaped body, the ellipsoid body, and noduli (S1A Fig) and the central complex is involved in bringing about a behavioral response [32-36]. In the case of vertebrates, it is the prefrontal cortex [37] (S1B Fig) plus the basal ganglia [38] (S1C Fig) that determines a behavioral response.

Some 30 different mutants of *D. melanogaster* have been isolated that have defects in various parts of the central complex [33, 39-45]. These mutants have been kindly given to us by Roland Strauss, Burkhard Poeck, and Douglas Armstrong. Now we have shown here (S17 Fig) that several of these mutants can block the parental response away from benzaldehyde plus heat and towards light plus a favored temperature that is shown in Figure 3A; these several mutants are cbd^{ks96}, nob^{ks49}, and ccb^{ks145} [33] and *EB1* and *62y* [45]. This result may suggest that the mutants we isolated and report here cause defects in the central complex.

Neusser, Triphan, Mrontz, Poeck, and Strauss found that *D. melanogaster*

mutants, unlike their parent, failed to remember where they were located just before they were moved [46], see the diagram of this by Tomchik and Davis [47] (S2 Fig). The Strauss et al. work suggested to them that this is related to the prefrontal cortex of humans [46]. Our own work on decision making [48] also describes mutants of *D. melanogaster* that appear to lack steps like those handled by the prefrontal cortex of humans and other vertebrates. It would seem that the insect brain and the brain of higher-order animals share many qualities.

According to Straussfeld and Hirth [49], there is behavioral homology between the response by insects and the response by vertebrates: see Figure 2 of ref. 49 (S1D Fig) and they propose that this homology originated in very early times. Studies by Arendt's group indicate that the responses by insects and the by vertebrates are descended from a common ancient ancestor that represents an early stage of evolution of the nervous system and muscles [50].

Altogether, the mutants that we isolated and reported here may well have bearing for both insects and vertebrates.

RELATION TO CHARGE SYNDROME AND DEAF-BLIND CONDITION

The *Drosophila* mutants isolated in this report seem related to a rare human genetic condition that occurs in roughly 1 per 10,000 live births [51], called CHARGE syndrome, reviewed recently [52]. "Children with CHARGE syndrome are truly 'multi-sensory impaired', having difficulties not only in vision and hearing but also with the senses that perceive balance, touch, temperature, pain, pressure, and smell, as well as problems with breathing and swallowing, eating and drinking, digestion, and temperature control" [53]. This syndrome was described independently in 1979 by Hall [54] and Hittner et al. [55]. The term CHARGE was then coined for the disease to cover its various symptoms [56], although not all symptoms are always present. A human gene, *CHD7,* is known to be associated with some 65% of 379 published cases of human CHARGE syndrome [57-60].

CHARGE syndrome is now being studied also in mice [61], zebrafish [62], and *Drosophila* [63]. A number of genes have been identified that are closely related to the human gene. In *Drosophila*, kismet *(kis,* the *Drosophila* homolog of human *CHD7*) is located in chromosome 2 [64]. In our own work on *Drosophila* (this report) the mutants described are located in the X chromosome, so the research here might deal with a different, additional gene for CHARGE syndrome, or else it deals with a different function altogether.

While in CHARGE syndrome the deaf-blind symptoms occur already at birth, in a different inherited disease, Usher syndrome, the deaf syndrome occurs at birth but the blind syndrome occurs years later [65]. That defect is known to be caused by an abnormal autosomal gene as well as a number of still unidentified genes [66].

IV. METHODS

See Supplemental Methods for details. A. Isolation of mutants. B. How to study response to external stimuli. C. How to study response to internal stimuli.

ACKNOWLEDGEMENTS

We are most grateful to the Camille and Henry Dreyfus Foundation for six years of grants in support of this undergraduate research program. The following undergraduates all contributed greatly to this work: Laura B. Burbach, Darrius M. Chambers, Meghan M. Cohen, Sarah A. Cook, Sarah C. Drewes, Nicholas A. Dykstra, Leann M. Erlien, Aaron J. Feiner, Eleanor K. Ganz, Hannah A. Hecht, Alison M. Heydorn, Hertina E. Kan, Alex R. Kleven, Alexei V. Medvedev, Nathan J. Menninga, Yi Han Ng, Arti Y. Patel, Keegan M. Schlittler, Amelia Remiarez, Rutvi Shah, Zuoxiao Shi, Avi D. Stricker, Grace S. M. Tang, Victor A. Toniolo, Jarrett A. Warden, Sarah K. Warzon, Andrew M. Winter, Aleksander R. Wood, and Ke Xu. Lar Vang is currently associate research specialist in the Adler laboratory. Robert A. Kreber, a research specialist in Barry Ganetzky's laboratory, has helped us greatly in studies of the genetics of our mutants. We are grateful to Erin Gonzales and Jerry Yin for showing us how to use the *Drosophila* activity monitory system. Julius Adler thanks Barry Ganetzky for teaching him about fruit flies. Thanks to Millard Susman for criticism. We are thankful to Laura Vanderploeg for the beautiful art work.

BIBLIOGRAPHY

1. Gilbert SJ, Burgess PW. Executive function. Curr Biol 2008;18:R110-R114

2. Fuster JM. IV. Executive function. In: The Prefrontal Cortex, Fourth Edition. London: Academic Press; 2008. pp. 178-193.

3. Adler J. My life with nature. Annu Rev Biochem. 2011;80: 42-70.

4. Vang LL, Medvedev AV, Adler J. Simple ways to measure behavioral responses of *Drosophila* to stimuli and use of these methods to characterize a novel mutant. PLoS ONE. 2012;7: e37495.

5. Edgecomb RS, Harth CE, Schneiderman AM. Regulation of feeding behavior in adult *Drosophila melanogaster* varies with feeding regime and nutritional state. J Exp Biol. 1994;197: 215-235.

6. Melche C, Bader R, Pankratz MJ. Amino acids, taste circuits, and feeding behavior in *Drosophila*: towards understanding the psychology of feeding in flies and man. J Endocrinol. 2007;192: 467–472.

7. Fujikawa K, Takahashi A, Nishimura A, Itoh M, Takano-Shimizu T, et al. Characteristics of genes up-regulated and down-regulated after 24 h starvation in the head of *Drosophila.* Gene. 2009;446: 11-17.

8. Farhadian SF, Suárez-Fariñas M, Cho CE, Pellegrino M, Vosshall LB. Post-fasting olfactory, transcriptional, and feeding responses in *Drosophila*. Physiol Behav. 2012;105: 544-553.

9. Hong S-H, Lee K-S, Kwak S-J, Kim A-K, Bai H, et al. Minibrain/Dyrk1a regulates food intake through the Sir2-FOXO-sNPF/NPY pathway in *Drosophila* and mammals. PLoS Genet. 2012;8: e1002857.

10. Stafford JW, Lynd KM, Jung AY, Gordon MD. Integration of taste and calorie sensing in *Drosophila.* J Neurosci. 2012;32:14767-74.

11. Itskov PM, Ribeiro C. The dilemmas of the gourmet fly: the molecular and neuronal mechanisms of feeding and nutrient decision making in *Drosophila.* Front Neurosci. 2013;7: 1-7.

12. Browne LB, Evans DR. Locomotor activity of the blowfly as a function of feeding and starvation. J Insect Physiol. 1960;4: 27-37.

13. Pfeiffenberger C, Lear BC, Keagan KP, Allada R. Processing circadian data collected from the *Drosophila* activity monitoring (DAM) system. Cold Spring Harb Protoc. 2010; 2010: 1242-50.

14. Thibalt ST, Singer MA, Miyazaki WY, Milash B, Dompe NA, et al. A complementary transposon tool kit for *Drosophila melanogaster* using P and piggyback. Nat Genet. 2004;36: 283-87.

15. Cook RK, Christensen SJ, Deal JA, Coburn RA, Deal ME, et al. The generation of chromosomal deletions to provide extensive coverage and subdivision of the *Drosophila melanogaster* genome. Genome Biol. 2012;13: R21.

16. Roote J, Russell S. Toward a complete *Drosophila* deficiency kit. Genome Biol. 2012;13: 149.

17. Middha S, Wang X. Evolution and potential function of fibrinogen-like domains across twelve *Drosophila* species. BMC Genomics. 2008 May 30. doi: 10.1186/1471-2164-9-260

18. Wang X, Zhao Q, Christensen BM (2005) Identification and characterization of the fibrinogen-like domain of fibrinogen-related proteins in the mosquito, *Anopheles gambiae*, and the fruitfly, *Drosophila melanogaster*, genomes. BMC. 2005 Sep 8. doi:10.1186/1471-2164-6-114

19. Attrill H, Falls K, Goodman JL, Millburn GH, Antonazzo G, Rey AJ, Marygold SJ; the FlyBase Consortium. FlyBase: establishing a Gene Group resource for *Drosophila melanogaster*. Nucleic Acids Res. 2015 Oct 1. doi: 10.1093/nar/gkv1046

20. Villella A, Hall JC. Neurogenetics of courtship and mating in *Drosophila*. Adv Genetics. 2008;62: 67-184.

21. Krstic D, Boll W, Noll M. Sensory integration regulating male courtship behavior in *Drosophila*. PLoS ONE. 2009;4: e4457.

22. Pan Y, Meissner GW, Baker BS. Joint control of *Drosophila* male courtship behavior by motion cues and activation of male-specific P1 neurons. Proc Natl Acad Sci USA. 2012;109: 10065-10070.

23. Yamamoto D, Koganezawa M. Genes and circuits of courtship behavior in *Drosophila* males. Nat Rev Neurosci. 2013;14: 681-692.

24. Tang G. Isolation and characterization of *Drosophila melanogaster* central processing mutants. Biochemistry honors thesis. MINDS@UW-Madison. 2007. *http://minds.wisconsin.edu/handle/1793/7957*.

25. Gorostiza EA, Colomb J, Brembs B. A value-based behavioral choice underlies phototaxis in *Drosophila*. Anim Behav Cognit. 2016 submitted.

26. Raichle ME. Two views of brain function. Trends Cogn Sci. 2010; 14:189-190.

27. Bray D. Intrinsic activity in cells and the brain. Mol Bio Cell. 2014;25: 737-78.

28. Brembs B. What are brains good for, anyway? Lecture in "Brain, cognition, behavior, evolution: Polyglot to monoglot", Institute for Advanced Studies, Sao Paulo, 29 June 2015, organized by Hogan J. https://www.youtube.com/watch?v=rLdqEgNYsws

29. Power ME. The brain of *Drosophila*. J Morph. 1943;72: 517-59.

30. Zhang K, Guo JZ, Peng Y, Xi W, Guo A. Dopamine-mushroom body circuit regulates saliency-based decision-making in *Drosophila*. Science. 2007;316: 1901-1904.

31. Hanesch U, Fischbach KF, Heisenberg M. Neuronal architecture of the central complex in *Drosophila melanogaster*. Cell Tissue Res. 1989;257: 347-366.

32. Pfeiffer K, Homberg U. Organization and functional roles of the central cortex in the insect brain. Annu Re Entomol. 2014;59: 165-184.

33. Strauss R, Heisenberg M. A higher control center of locomotor behavior in the *Drosophila* brain. J Neurosci. 1993;13: 1852-1861.

34. Lin CY, Chuang CC, Hua TE, Chen CC, Dickson BJ, et al. A comprehensive wiring diagram of the protocerebral bridge for visual information processing in the *Drosophila* brain. Cell Rep. 2013;3: 1739-1753.

35. Seelig JD, Jayaraman V. Feature detection and orientation tuning in the *Drosophila* central complex. Nature. 2013;503: 262–266.

36. Wolff T, Iyer N A, Rubin GM. Neuroarchitecture and neuroanatomy of the *Drosophila* central complex: A GAL-based dissection of protocerebral bridge neurons and circuits. J Comp Neurol. 2015;523: 997-1037.

37. Fuster JM. The Prefrontal Cortex, Fourth Edition. London: Academic Press; 2008.

38. Graybiel AM. The basal ganglia. Curr Biol. 2000;10: R509-R511.

39. Renn SCP, Armstrong JD, Yang M, Wang Z, An X, et al. Genetic analysis of the *Drosophila* ellipsoid body neuropil: Organization and development of the central complex. J Neurobiol. 1999;41: 189-207.

40. Heisenberg M, Boehl K. Isolation of anatomical brain mutants of *Drosophila* by histological means. Z Naturforsch C. 1979;34: 143-147.

41. Martin J-R, Ernst R, Heisenberg M. Temporal pattern of locomotor activity in *Drosophila melanogaster*. J Comp Physiol A. 1999;184: 73-84.

42. Martin J-R, Raabe T, Heisenberg M. Central complex substructures are required for the maintenance of locomotor activity in *Drosophila melanogaster*. J Comp Physiol A. 1999;185: 277-288.

43. Strauss R. The central complex and the genetic dissection of locomotor behaviour. Curr Opin Neurobiol. 2002;12: 633-638.

44. Poeck B, Triphan T, Neuser K, Strauss R. Locomotor control by the central complex in *Drosophila* – an analysis of the *tay bridge* mutant. Develop Neurobiol. 2008;68: 1046-1058.

45. Young JM, Armstrong JD. Structure of the adult central complex in *Drosophila:* Organization of distinct neuronal subsets. J Comp Neurol. 2010;518: 1500-1524.

46. Neuser K, Triphan T, Mronz M, Poeck B, Strauss R. Analysis of a spatial orientation memory in *Drosophila*. Nature. 2008;453: 1244-1247.

47. Tomchik SM, Davis RL. Out of sight, but not out of mind. Nature. 2008;453: 1192-1194.

48. Adler J, Vang LL. Decision making by *Drosophila* flies. Submitted to bioRxiv 2016.

49. Strausfeld NJ, Hirth F. Deep homology of arthropod central complex and vertebrate basal ganglia. Science. 2013;340: 157-161.

50. Marlow H, Arendt D. Evolution: ctenophore genomes and the origin of neurons. Cur Biol. 2014;24: R757-R761.

51. Blake KD, Prasad C. CHARGE syndrome. Orphanet J of Rare Diseases. 2006 Sep 7. doi: 10.1186/1750-1172-1-34.

52. Hsu P, Alan M, Wilson M, Williams G, Curotta J, et al. CHARGE Syndrome: A review. J Paediat Child Health. 2014;50: 504-511.

53. Brown D. CHARGE syndrome "behaviors": Challenges or adaptations? Amer J Med Genet. 2005;133A: 268-272.

54. Hall BD. Choanal atresia and associated multiple anomalies. J Pediatr. 1997;95: 395-398.

55. Hittner HM, Hirsch NJ, Kreh GM, Rudolph AJ. Colobomatous, microphthalmia, heart disease, hearing loss, and mental retardation – a syndrome. J Pediatr Ophthalmol Strabismus. 1979;16: 122-128.

56. Pagon PA, Graham JM, Zonana J, Young SL. Coloboma, congenital heart disease, and choanal atresia with multiple anomalies: CHARGE association. J Pediatr. 1981;99: 223-227.

57. Vissers LE, van Ravenswaaij CM, Admiraal R, Hurst JA, de Vries BB, et al. Mutations in a new member of the chromodomain gene family cause CHARGE syndrome. Nature Genetics. 2004;36: 955-957.

58. Lalani SR, Safiullah AM, Fernbach SD, Harutyunyan KG, Thaller C, et al. Spectrum of CHD7 mutations in 110 individuals with CHARGE syndrome and genotype-phenotype correlation. Am J Hum Genet. 2006;78: 303-314.

59. Zentner GE, Layman WS, Martin DM, Scacheri PC. Molecular and phenotypic aspects of CHD7 mutation in CHARGE syndrome. Am J Med Genet A. 2010;152A: 674-686.

60. Janssen N, Bergman JE, Swertz MA, Tranebjaerg L, Lodahl M, et al. Mutation update on the *CHD7* gene involved in CHARGE syndrome. Human Mutation. 2012;33: 1149-1160.

61. Bosman EA, Penn AC, Ambrose JC, Kettleborough R, Stemple DL, et al. Multiple mutations in mouse *Chd7* provide models for CHARGE syndrome. Hum Mol Genet. 2005;14: 3463-3476.

62. Patton SA, Jacobs-McDaniels NL, Zaouter C, Drapeau P, Albertson RC, et al. Role of Chd7 in Zebrafish: a model for CHARGE syndrome. PLOS ONE. 2012;7: e31650.

63. Melicharek DJ, Ramirez LC, Singh S, Thompson R, Marenda DR. Kismet/CHD7 regulates axon morphology, memory and locomotion in a *Drosophila* model of CHARGE syndrome. Hum Mol Genet. 2010;19: 4253-4264.

64. Kennison JA, Tamkun JW (1988) Dosage-dependent modifiers of Polycomb and Antennapedia mutations in *Drosophila*. Proc Natl Acad Sci USA. 1988;85: 8136-8140.

65. Dammeyer J. Development and characteristics of children with Usher syndrome and CHARGE syndrome. Int J Pediatr Otorhi. 2012;76: 1292-1296.

66. Yan D, Liu XZ. Genetics and pathological mechanism of Usher syndrome. J Hum Genet. 2010;55: 327-355.

SUPPLEMENTAL FIGURES

CENTRAL COMPLEX

Strausfeld and Hirth, Science 340:157 (2013)
from U. Hanesch, K.-F. Fischach, M. Heisenberg, Neuronal architecture of the central
complex in *Drosophila melanogaster. Cell Tissue Res.* 257, 343 (1989).

S1A Fig. Central complex of the *Drosophila* brain.

The prefrontal cortex (blue)

Man

Chimp

Rhesus monkey

Dog

Cat

Modified from *The Prefrontal Cortex,*
J.M. Fuster, 2008, p.12

S1B Fig. The prefrontal cortex of various mammalian species.

The basal ganglia

The basic anatomy of the brain showing the major regions within the basal ganglia: the striatum (blue), which is made up of the caudate nucleus and the putamen; the pallidum (pink), which is made up of outer and inner segments; the subthalamic nucleus (green); and the substantia nigra (yellow).

A. M. Graybiel, Current Biology 10:R509 (2000)

S1C Fig. The basal ganglia of the human brain.

Proposed correspondences of the mammalian basal ganglia and insect central complex

Strausfeld and Hirth, Science 340:157 (2013)

S1D Fig. Comparison of the mammalian brain and the insect brain.

Fly remembers its location

a, A fly will walk back and forth between two opposing black stripes in a circular arena even if, midway, the target stripe disappears. Neuser et al.[1] find that if (**b**) a fly is distracted midway by a new black stripe to its side, the insect still remembers the position of the original stripe, and (**c**) when the distracter stripe subsequently disappears, resumes walking along its original course. The authors find that this behaviour depends on ring neurons of the ellipsoid body in the fly brain.
Tomchick & Davis, Nature 453 (2008) p. 1193

[1]Neuser, Triphan, Mronz, Poeck & Strauss, Nature 453 (2008) p.1244-1247

S2 Fig. Assay used to show that a fly remembers its original location.

S3 Fig. Response to four stimuli together. Repellents (0.1M benzaldehyde and high temperature (37°C)) were at the origin end, attractants (light and a favored temperature (27°C)) at the furthest end. (**C**) Mutant 1 response (n=5). (**D**) Mutant 3 response (n=5). (**E**) Mutant 4 response (n=5). (**F**) Mutant 5 response (n=4). Flies were tested in a 34°C room with 10 to 20 flies used per trial. Data are mean±SEM.

S4 Fig. Response to light alone. (**C**) Mutant 2 response (n=8). (**D**) Mutant 3 response (n=7). (**E**) Mutant 4 response (n=6). (*F*) Mutant 5 response (n=8). Flies were tested at 34°C with 10 to 20 flies used per trial. Data are mean±SEM.

S5 Fig. Response to benzaldehyde alone. (**C**) Mutant 2 response (n=4). Flies were tested at 34°C with 10 to 20 flies used per trial. Data are mean±SEM.

S6 Fig. Response to heat gradient alone. (**C**) Mutant 2 response (n=4). Flies were tested at 34°C with 10 to 20 flies used per trial; the warm side measured 37°C and the cool side 27°C. Data are mean±SEM.

S7 Fig. Response to quinine. (**C**) Mutant 1 response (n=10). Flies were tested at 34°C with 10 to 20 flies used per trial. Data are mean±SEM.

S9 Fig. Response to gravity. (**C**) Mutant 1 response (n=4). Flies were tested at 34°C with 10 to 20 flies used per trial. Data are mean±SEM.

S10 Fig. Response without added stimuli. (**C**) Mutant 2 response (n=6). (**D**) Mutant 3 response (n=5). (**E**) Mutant 4 response (n=6). (**F**) Mutant 5 response (n=6). Flies were tested at 34°C with 10 to 20 flies used per trial. Data are mean±SEM.

S11 Fig. Response to light at room temperature. (**A**) Parental response (n=4). (**B**) Mutant 1 response (n=4). (**C**) Mutant 2 response (n=6). (**D**) Mutant 3 response (n=3). (**E**) Mutant 4 respnose (n=5). (**F**) Mutant 5 response (n=4). Approximately 10 to 20 flies were used per trial. Data are mean±SEM.

S12 Fig. Movement of flies at 20 hours in search for food. Solid: water but no food (no sucrose). Open: water and food (sucrose). (**C**) Mutant 1 response with water only (n=4)

and with water+sucrose (n=4). Flies were tested at 22º C with 40 to 60 flies used per trial. Data are mean±SEM.

S13 Fig. Movement of flies at 30 hours in search for water. Solid: food but no water. Open: food and water. (**C**) Mutant 1 response with sucrose only (n=5) and with water+sucrose (n=4). Flies were tested at 22º C with 40 to 60 flies used per trial. Data are mean±SEM.

S14 Fig. Circadian response. Individual flies are placed into a tube (5 x 20 mm) with an infrared light beam intersecting at the middle of the tube. **(A)** Mutant 3 response (n=62), **(B)** Mutant 4 response (n=63), **(C)** Mutant 5 response (n=58), and parental fly response

(n=156). Flies were tested at 22º C. Graphs were obtained from traces of the original graphs as the original data were lost. Data are mean±SEM.

S15 Fig. Response of attached-X females from their respective mutant mate to four stimuli together. Repellents (0.1M benzaldehyde and high temperature (37°C)) were at the origin end, attractants (light and a favored temperature (27°C)) at the furthest end. (**A**) Female of parent response (n=4). (**B**) Female of Mutant 1 response (n=6). (**C**) Female of Mutant 2 response (n=4). Flies were tested at 34°C with 10 to 20 flies used per trial. Data are mean±SEM.

S16 Fig. Response of mutant strains with malformations of the mushroom bodies. Repellents (0.1M benzaldehyde and high temperature (37°C)) were at the origin end, attractants (light and a favored temperature (27°C)) at the furthest end. (**A**) Mutant *mbmc[1]* response (n=3) and (**B**) mutant *mbd[1]* response (n=3) compared to wild-type CS (n=7).). Flies were tested at 34°C with 10 to 20 flies used per trial. Data are mean±SEM. See [30] for more details.

S17 Fig. Response of GAL4 driver lines and mutant strains with malformations of the central complex. Repellents (0.1M benzaldehyde and high temperature (37°C)) were at the origin end, attractants (light and a favored temperature (27°C)) at the furthest end. (**A**) Response of cdb^{ks96} (n=3), (**B**) response of nob^{ks49} (n=6), (**C**) response of ccd^{ks145} (n=4), (**D**) response of $GAL4\text{-}EB1/+$ (n=3) and $GAL4\text{-}EB1/UAS\text{-}shi^{ts1}$ (n=10), and (**E**) response of $GAL4\text{-}62y/+$ (n=3) and $GAL4\text{-}62y/UAS\text{-}shi^{ts1}$ (n=4) compared to wild-type CS (n=7). Flies were tested at 34°C with 10 to 20 flies used per trial. Data are mean±SEM. See [33, 45] for more details.

SUPPLEMENTAL METHODS

A. ISOLATION OF MUTANTS

D. melanogaster (strain Canton-S) and attached-X flies were provided by Robert Kreber.

Flies were maintained on standard cornmeal-molasses agar medium at room temperature (21-23°C) in a room that was light for 12 hours and dark for 12 hours. Male flies were mutagenized overnight with 25mM ethyl methane sulfonate in 1% sucrose on Kimwipe paper (Kimberly-Clark, Roswell, GA). Then they were mated with virgin attached-X (*y w f* C(1)DX) females. At about six days the adult flies were removed. At around two weeks the F1 adults appeared and they were studied here.

At 2 to 5 days some 5300 of these F1 males plus females were screened about 500 at a time by a procedure shown in Fig 2. From 4-liter graduated cylinders (10.8 cm x 58 cm Nalgane polymethylpentene, Fisher, 08-572-5J, Hanover Park, IL) four 17.5 cm sections were cut (the end with the spout was not used) and labeled sections 1, 2, 3, and 4. (In Fig 2 sections 1, 2, 3, and 4 are called "origin", "next to origin", "next to furthest", and "furthest", respectively.) At room temperature the right end of section 4 was filled with 40 ml of 1.5% agar for control of the temperature-gradient described below. A 2-cm diameter hole was made in the middle of section 1 to be the point of entry for the flies. Then these four pieces were put into a 34°C room, where the experiments were to be carried out. Sections 1, 2, 3, and 4 were each fixed together with a transparent connector (a 15x20 cm piece cut from a transparency film for copiers, 3M Company, St. Paul, MN). A 45 cm fluorescent light (15 watts, cool white, lamp VTU15RTCPCO used without plastic cover, obtained from American Fluorescent Corp., Waukegan, IL or local hardware store such as Mennards, Madison, WI) was placed perpendicularly 15 cm from the right end of section 4 to produce a light gradient (stimulus 1). A temperature gradient (stimulus 2) was generated by placing a 2 L beaker of ice 1 cm from the right end of section 4 and by placing a hotplate (Barnstead Thermolyne, 1900, Dubuque, IA) on its side at a setting of 150°C 10 cm away from the left end of section 1; this was ready to use in about 20 minutes, when the temperature at the cool end was about 29° and at the warm end 38° (but not greater, so as to avoid a block in motility), as determined with a digital thermometer (63-1032, RadioShack, Fort Worth, TX) inside each end of the tube. At room temperature 0.1M benzaldehyde in 5 ml of melted 1.5% agar was pipetted onto the interior walls of a 5 cm-long cap (cut from a 17.5 cm section, see above) while rotating the cap until the agar hardened (cooler temperature may be needed). One end of the cap was then closed with Saran Wrap (S.C. Johnson Inc., Racine, WI). White tulle fabric or white organza fabric (Gifts International Inc., Ontario, CA, or local fabric store such as Hancock Fabrics, Madison, WI) was folded in half and taped to the other end of the benzaldehyde cap to provide a screen between section 1 and the cap in order to prevent the flies from making direct contact with this toxic benzaldehyde solution. Then the benzaldehyde cap was brought into the 34°C room and was affixed to the left end of section 1 with white label-tape (Fisher, 15-938). The benzaldehyde was allowed to diffuse for 30 minutes to provide a gradient of it (stimulus 4). (If a single stimulus had

been used instead of four stimuli together, the resulting mutants would more likely be defective in a gene for sensing that single stimulus instead of in a gene for the final pathway.)

A previous attempt to produce such mutants (2005-2007) by Grace Tang, Victor Toniolo, Yi Hang Ng, Lar Vang, and Julius Adler used an alternative yet similar procedure [23], but it failed. After initial mutagenization of males, about 33000 *Drosophila melanogaster* were tested, around 1500 males plus females at a time. Selected were those that failed in response to light, then those were tested in response to benzaldehyde, then those were tested in response to heat. After these three tests, finally 29 mutants resulted but they did not reproduce. Then we started over with the present procedure, which puts all the stimuli together for the isolation of mutants.

B. HOW TO STUDY RESPONSE TO EXTERNAL STIMULI

Flies were removed from their food vials into empty vials. The flies (about 500 flies for isolation of mutants, about 10 in later experiments) were then placed at 34°C in the dark for 30 minutes. They were added through the hole in section 1 by use of a funnel, and then the fluorescent lamp was turned on, which marked the beginning of the assay. Every minute for 20 minutes the number of flies in each quarter of the tube was counted, unless the number of flies was too large to easily count, in which case estimates were made. In cases where the flies were getting killed by the heat and/or benzaldehyde, the assays were run for a shorter time.

Parental flies went almost entirely to the attractant end (compartment 4). For isolation of mutants, those flies that failed to go to the attractant end (the ones in compartments 1, 2, and 3) were saved, placed in food vials, and retested in the same way one to two days later. Then each of those males that again failed was mated with virgin females, and the next generation (F2) of males was tested in the same way. This procedure yielded five mutants, called 1 to 5. Only males were used in all of the experiments presented in this article. (If a single stimulus had been used instead of four stimuli together, the resulting mutants would more likely be defective in a gene for sensing that single stimulus instead if in a gene for the final pathway.)

When light alone is used, the setup is the same as in the isolation of mutants but with only the light source used.

When heat alone is used, the setup is the same as in the isolation of mutants but with only the heat source used and with a light source placed parallel to the entire length of the tube.

When no stimuli was used, the setup is the same as in the isolation of mutants but with no stimuli present except for a parallel light source placed along the entire length of the tube.

See ref. 4 on testing for quinine and gravity. Each trial used about 10 male flies.

See ref. 5 on testing for sucrose. Each trial used about 10 male flies.

C. HOW TO STUDY RESPONSE TO INTERNAL STIMULI

1. Measuring hunger

The apparatus for measuring hunger consisted of four tubes as shown in Fig 10. Tube 1 is 20 cm long and 5 cm wide, tubes 2, 3, and 4 are 40 cm long and 5 cm wide (cut from clear acrylic tubes 6 feet long and 2 inches wide, TAP Plastics info@tapplastics.com). Each of the four tubes was drilled with four rows of 1 mm holes every 2.5 cm to provide exchange of air (possibly these holes are not needed when only 50 flies are used). Each of the four tubes was saturated with water by adding a strip of Kimwipe (15 in x 17 in, Kimberly-Clark) 5 cm wide and 20 or 40 cm long to which 5 or 10 ml of distilled water was added, and then the excess water was poured off. Between each of the four tubes was placed a funnel (45 mm wide, Kimble Kimax 28950-45) with its stem cut off (see Fig 10), to direct flies from one tube into the next and to minimize movement backward. Then as connector to connect the tubes, transparent packaging tape (5 cm width, SureTech Brands, Avon, OH) was placed all the way around between tubes 2 and 3 and between 3 and 4; between 1 and 2 it was placed over only the bottom half because the gate is in the way elsewhere. Then a gate (see Fig 10) (cut from transparency film for copiers, 3M, St. Paul, MN) was inserted between 1 and 2 (used only during the loading of the flies, then removed). The two open ends of the apparatus were capped with an acrylic cap we made, though Saran wrap (S. C. Johnson & Sons, Racine, WI) would be perhaps as effective.

The apparatus was put into a dark room at 21 to 23°C. About 50 male flies per trial were placed from their growth medium into tube 1. After about 5 minutes to allow the flies to become quiescent, the gate was removed and replaced by putting transparent packaging tape over the top half. The apparatus was then covered with black garbage bags to ensure darkness. Then the number of flies in each tube was counted at 0, 10, 20, 30, and 40 hours by having lights on for 30 seconds.

The effect of adding water plus sucrose was studied as above except that 0.1M sucrose was poured onto the strip of Kimwipe, 5 ml for tube 1 and 10 ml for tubes 2, 3, and 4, and then the excess was poured off at 10 sec.

2. Measuring thirst

The same apparatus (Fig 10) and the same method was used for studying thirst as for measuring hunger except there was no water added and sucrose crystals were placed on the Kimwipe, 250 mg in tube 1 and 500 mg in each of tubes 2, 3, and 4.