***Wolbachia* confers sex-specific resistance and tolerance to**

enteric but not systemic bacterial infection in *Drosophila*

Radhakrishnan B. Vasanthakrishnan [1,3§], Gupta Vanika [1§], Jonathon Siva-Jothy, Katy M.

Monteith[1], Sam P. Brown[4], Pedro F. Vale[1,2]*,

Affiliations

[1]Institute of Evolutionary Biology, School of Biological Sciences, University of Edinburgh.

Edinburgh EH9 3FL

[2] Centre for Immunity, Infection and Evolution, University of Edinburgh. Edinburgh EH9

3FL

[3] Current address: IGDR - CNRS UMR 6290, 2, Avenue Du Pr. Léon Bernard, 35043

Rennes, France.

[4] School of Biology, Georgia Institute of Technology, Atlanta, Georgia 30332–0230, USA.
18

19   [§] These authors contributed equally

23   *Corresponding author

24   E-mail: pedro.vale@ed.ac.uk

Abstract

Wolbachia-mediated protection against viral infection has been extensively demonstrated in *Drosophila* and in mosquitoes that are artificially inoculated *with D. melanogaster Wolbachia* (wMel), but to date no evidence for *Wolbachia*-mediated antibacterial protection has been demonstrated in *Drosophila*. Here we show that *D. melanogaster* carrying wMel shows reduced mortality during enteric – but not systemic - infection with the opportunist pathogen *Pseudomonas aeruginosa*, and that protection is more pronounced in male flies. *Wolbachia*-mediated protection is associated with increased early expression of the antimicrobial peptide *attacinA*, followed by increased expression of a ROS detoxification gene (*gstD8*), and other tissue damage repair genes which together contribute to greater host resistance and disease tolerance. These results highlight that the route of infection is important for symbiont-mediated protection from infection, that *Wolbachia* can protect hosts by eliciting a combination of resistance and disease tolerance mechanisms, and that these effects are sexually dimorphic.

Introduction

All organisms experience a combination of beneficial and detrimental colonisations by pathogens, commensals and symbionts, with profound effects on host physiology, behaviour, ecology and evolution (Bennett and Moran, 2015; Douglas, 2015; Gandon and Vale, 2014; Lewis and Lizé, 2015; Werren et al., 2008). Bacterial endosymbionts of insects, for example, are known to manipulate host reproduction (Engelstädter and Hurst, 2009; Werren et al., 2008), to alter the host's acquisition of essential nutrients(Douglas, 2015, 1998), and to provide protection from the deleterious effects of parasites and pathogens (Brownlie and Johnson, 2009; Hamilton and Perlman, 2013).

Wolbachia is a maternally-inherited, intracellular bacterium of arthropods and nematodes, and is one of the best studied microbial symbionts (Brownlie and Johnson, 2009; Hamilton and Perlman, 2013). Its host range is vast, with recent estimates that 48-57% of all terrestrial arthropods (Weinert et al., 2015), and at least 10% of all *Drosophila* species carry *Wolbachia* (Mateos et al., 2006). The ability of some *Wolbachia* strains to protect insect hosts from pathogenic infections make it particularly relevant for potential bio-control of insect vectored zoonotic infections, and more broadly, relevant as mediators of pathogen-mediated selection in insects (Brownlie and Johnson, 2009; Hamilton and Perlman, 2013; Karyn N. Johnson, 2015). *Aedes aegypti* and *Ae. albopictus* mosquitoes, for example, have been shown to become more resistant to Dengue and Chikungunya viruses, as well as malaria-causing *Plasmodium* when they are experimentally inoculated with *Wolbachia* (Bian et al., 2010; Kambris et al., 2010; Moreira et al., 2009). In *Drosophila*, there is also ample evidence that flies carrying *Wolbachia* are better able to survive infection by a number of naturally occurring RNA viruses (Hedges et al., 2008; Hedges and Johnson, 2008; Karyn N Johnson, 2015; Teixeira et al., 2008). This anti-viral protection is variable among strains of *Wolbachia* and correlates strongly with the reduction in viral titres within hosts (Martinez et al.,

2014), suggesting that *Wolbachia* generally enhances the ability to clear pathogens

(increasing host resistance) rather than the ability to repair damage independently of

pathogen clearance (disease tolerance) (Ayres and Schneider, 2012; Råberg et al., 2009).

In contrast to the strong evidence for *Wolbachia*-mediated antiviral protection,

its ability to protect its native fruit fly hosts from bacterial infections has not been

clearly demonstrated to date. In one study, carrying *Wolbachia* did not affect the

survival or immune activity of *D. simulans* or *D. melanogaster* during systemic infection

with *Pseudomonas aeruginosa*, *Serratia marcescens* or *Erwinia carotovora* (Wong et al.,

2011), while another study found that the presence of *Wolbachia* had no effect on the

ability to suppress pathogen growth during systemic infections by intracellular (*Listeria*

monocytogenes, *Salmonella typhimurium*) or extracellular bacterial pathogens

(*Providencia rettgeri*) (Rottschaefer and Lazzaro, 2012). Given that *Wolbachia* can

provide broad-spectrum protection against a range of pathogens, including bacteria, in

mosquitoes (Ye et al., 2013), the lack of evidence for antibacterial protection in flies is

puzzling. Some authors have proposed that antibacterial protection may only occur in

novel host-*Wolbachia* associations (like those of mosquitoes), although the exact

mechanism for such protection remains unclear (Wong et al., 2011; Zug and

Hammerstein, 2015).

One possibility is that the experimental conditions under which *Drosophila* are

commonly challenged with pathogens in the lab do not reflect the infections they are

likely to encounter in the wild. For example, experimental infections often focus on

systemic infection, introducing large quantities of bacterial pathogens via intra-thoracic

or abdominal injection (Neyen et al., 2014; Rottschaefer and Lazzaro, 2012; Wong et al.,

2011). The ecological context of fruit flies however, which consists mainly of foraging on

rotting organic matter, means that most pathogens in the wild are more likely to be

acquired orally, resulting in enteric, rather than systemic infections (Ferreira et al.,

2014; Stevanovic and Johnson, 2015). It is therefore conceivable that any form of

Wolbachia-mediated protection that could have evolved in the context of enteric infection may not be detected during systemic infection.

Here we show that the route of infection is indeed important for *Wolbachia*-mediated protection in *Drosophila*, which we find to occur during enteric - but not systemic - infection by the opportunist pathogen *Pseudomonas aeruginosa*. *P. aeruginosa* has an incredibly broad host range, infecting insects, nematodes, plants, and vertebrates, and is found in most environments (Apidianakis and Rahme, 2009; Neyen et al., 2014). Enteric infection of *Drosophila* by *P. aeruginosa* results in pathology to intestinal epithelia due to the formation of a bacterial biofilm in the crop, a food storage organ in the foregut (Mulcahy et al., 2011; Sibley et al., 2008). In the majority of enteric infections *P. aeruginosa* growth is restricted to the crop, and is sufficient to cause death (Chugani et al., 2001; Sibley et al., 2008). We exposed flies that were naturally infected with *Wolbachia*, and identical derived flies that were cured of *Wolbachia* infection, to *P. aeruginosa* either through intra-thoracic pricking (causing a systemic infection) or through the oral route of infection by feeding (causing an enteric infection). We then monitored how within-host microbe loads and survival varied throughout the course of an infection to assess if (1) *Wolbachia*-mediated protection occurred during systemic and enteric bacterial infection; (2) when protection was detected, if this was due to differences in the bacterial clearance rate (resistance) or if it aided host survival despite high microbe loads (tolerance); and (3) how these protective effects differed between male and female flies. We further characterized the expression of immune and damage repair genes previously shown to be involved in enteric bacterial infection in *Drosophila*.

Results

Wolbachia reduces mortality during enteric but not systemic bacterial infection

All flies infected systemically with PA14 via intra-thoracic pricking died within

24 hours (Fig 1a), and in line with previous work (Wong et al., 2011), we did not detect

any significant effect of *Wolbachia* status on the rate at which they died (Cox

Proportional Hazard Model, Likelihood Ratio X^2 = 0.003, DF=1, p=0.959), regardless of

sex ('Sex' effect, X^2= 0.860, DF=1, p=0.354). Flies that ingested and acquired an enteric

infection of PA14 died at a faster rate than control flies exposed only to a sucrose

solution (**Fig. 1b**; 'Infection status' effect, Likelihood Ratio X^2 = 64.27, DF=1, p<0.0001).

Fly mortality during enteric infection was significantly affected by their *Wolbachia*

status (X^2 = 6.32, DF=1, p=0.013). This protective effect was not substantial in female

flies: female flies without *Wolbachia* were 1.58 more likely to die than infected females

carrying *Wolbachia* (Cox risk ratio, X^2=1.27, p=0.2644). The protection in male flies was

more pronounced, where not carrying *Wolbachia* made PA14-infected males 2.26 times

more likely to die than their infected *Wolbachia*-positive counterparts (Cox risk ratio,

X^2= 4.22, p=0.0172)(Figure 1b). In order to understand the cause of the observed

protection during enteric but not systemic infection protection, the results below focus

only on flies that acquired infection orally.

Wolbachia does not affect the rate of bacterial clearance during enteric
infection
145
Following 12 hours of exposure to *P. aeruginosa*, bacterial loads decreased over

the course of the experiment in both male and female flies (**Fig. 2)** time effect

$F_{7,186}$=48.81, p<0.0001). However, the rate at which infection was cleared was not

affected by *Wolbachia* status ('Time × Wolbachia' interaction $F_{7,186}$=5.71, p=0.30), which

suggests that the presence of *Wolbachia* does not contribute to the clearance of this

bacterial gut infection. Regardless of *Wolbachia* status, we observed that males and

females showed different patterns of bacterial clearance over time (**Fig. 2**; 'Time × Sex'

interaction $F_{7,186}$=4.21, p=0.002). While males appeared to be able to clear the infection

almost entirely within a week (mean ± SEM 0.85 ±0.29 Log_{10} CFU per fly at 168 hours

post exposure) females appeared to stop clearing infection after 96h, maintaining a

relatively stable bacterial load of about 100 CFUs per fly until the end of the experiment

(**Fig. 2**). These sex differences in bacterial clearance were present regardless of the

Wolbachia status of the flies, suggesting they reflect sexual dimorphism in antibacterial

defence and not to sex-specific effects of *Wolbachia* (Sex × Wolbachia interaction

$F_{1,186}$=0.10, p=0.758).

Male flies with *Wolbachia* have lower bacterial loads in the early stages of enteric infection

162
163
164
The initial stages of exposure to pathogens can be crucial in determining

whether the host controls infection or if a pathogen grows to a point where hosts are

killed. Even though we detected no difference in the rate of clearance according to

Wolbachia status throughout the infection, at 12 and 24-hours post-infection male flies

harbouring *Wolbachia* showed significantly lower bacterial CFUs compared to those

without *Wolbachia* (Fig 2; Wol+: 3.86±0.22 Log_{10} CFU; Wol-: 4.56±0.22 Log_{10} CFU; $F_{1,20}$ =

5.27, p= 0.033). One explanation for the difference in initial microbe loads in males is

that *Wolbachia* could cause behavioural changes, such as reduced feeding rate, that

result in reduced infection. However, we did not find evidence that the lower CFU

numbers seen in *Wolbachia*-positive male flies resulted from lower feeding rates (Fig.

S1). Another potential explanation for the difference in initial microbe loads in males is a

Wolbachia-mediated antimicrobial response. Whole fly homogenate added to growing

cultures of PA14 (in both liquid and solid growth medium) showed greater

antimicrobial activity in homogenates of *Wolbachia* positive flies (Figure S2 and S3).

Wolbachia-positive flies show increased expression of IMD pathway genes

during the early stages of enteric infection

Given the preliminary evidence of increased antimicrobial activity in *Wolbachia* –

positive males (Figure S2 and S3), we decided to test for differences in the expression of

antimicrobial immune pathways. While previous work has found no effect of *Wolbachia*

on the expression of immune genes (Wong et al., 2011), or on bacterial

clearance(Rottschaefer and Lazzaro, 2012), this has not been tested in the context of

enteric bacterial infection. Other studies in *Wolbachia*-free flies have demonstrated that

the IMD pathway plays an active role in the response to enteric bacterial infection

(Buchon et al., 2014, 2009; Kuraishi et al., 2011). We therefore tested whether flies

carrying *Wolbachia* showed increased expression of genes involved in IMD-mediated

antimicrobial immunity.

192   In *Wolbachia*-positive females, we observed a significant increase in expression

(relative to uninfected females) of two IMD pathway receptor genes– *pgrp-lc* ($p =$

0.0002) and *pgrp–le* ($p = 0.004$) at 96-hours post-infection (Fig. 3). In PA14-infected

males, however, carrying *Wolbachia* resulted in a slight decrease in *pgrp–lc* expression

relative to uninfected males ($p = 0.03$), although this difference was transient and only

observed at 24-hours post-infection (Fig. 3). Overall there appears to be little effect of

Wolbachia on the expression of either receptor gene in male flies (Fig 3). We observed a

significant 3 to 4-fold increase in the expression of the antimicrobial peptide (AMP) gene

attacinA (*attA)* in *Wolbachia*-positive males at both 24 hours ($p = 0.002$) and 96 hours

($p < 0.001$) post-infection. We did not detect any difference in the expression of this

AMP gene in male flies that were free of *Wolbachia*, or in female flies, regardless of their

Wolbachia status (Fig. 3). These results suggest that the initial difference in clearance

between Wol+ and Wol- male flies could at least in part be due to *Wolbachia*-mediated

up-regulation of the AMP *attacinA*.

***Wolbachia* contributes to increased disease tolerance in male flies**

The data we describe above reveals interesting differences in the way male and

female flies fight enteric bacterial infection. Male flies are able to clear infection almost

completely, while female flies stop clearing infection after 96 hours and maintain a

stable bacterial load following that same time period, which suggests that male flies are

better than females at clearing enteric PA14 infection (**Fig. 2**). We could therefore

expect females to pay a survival cost due to higher bacterial loads but instead we find

that female flies have a similar survival probability to males, especially for flies that are

*Wolbachi*a negative (**Fig. 1**). This suggests that females are better able than males to

tolerate *P. aeruginosa* enteric infection because they are able to maintain a similar level

of health to females, while tolerating higher bacterial loads (Ayres and Schneider, 2012;

Medzhitov et al., 2012; Råberg et al., 2009). Male flies, however, showed a marked

increase in survival when they were *Wolbachia* positive compared to *Wolbachia*

negative males (**Fig. 1**), even though the rate at which both groups clear infection

appear identical (**Fig. 2**). This suggests increased tolerance in males mediated by the

presence of *Wolbachia*. In females, the survival benefit of *Wolbachia* appeared to be

minimal, suggesting that *Wolbachia*-mediated tolerance could be sex-specific.

To better assess these differences in disease tolerance mediated by sex and

Wolbachia status, we plotted the relationship between host health and microbe load for

matching time-points (Fig. 4). In all cases, these data were better described by a non-

linear 4-parameter logistic model than a linear model (Table S1). The 4-paraemter

logistic model is useful to compare how its maximum (reflecting health in the initial

stages of infection), baseline (reflecting the lowest survival reached during the

experiment), inflection point (the point at which fly survival reached halfway between

the baseline and maximum), and the growth rate (reflecting the rate at which fly

survival plummets) vary according to host sex and *Wolbachia* status. Each of these

parameters may reflect distinct mechanisms of damage repair involved in host infection

tolerance, so they are useful for further exploration of tolerance mechanisms (Ayres and

Schneider, 2012; Vale et al., 2016). In female flies, the logistic model explained about a

quarter of the variance ($R^2=0.24$), and a formal parallelism test found that the curves did

not show significantly different shapes ($F_{3,72}=0.886$, $p=0.452$). In male flies, the 4-

parameter logistic model explained over half the variance ($R^2=0.57$), and a formal

parallelism test revealed significant differences n the shapes of these two tolerance

curves between *Wolbachia*-positive and *Wolbachia*-negative males ($F_{3,72} = 2.98$,

$p=0.037$). These differences arise not only to the consistently lower maximum and

baseline survival in *Wolbachia*-negative males regardless of microbe load (Figure 4), but

also due to differences in the inflection point of each curve which occurs later in the

infectious period in *Wolbachia*-positive male flies (Figure 4).

***Wolbachia* is associated with higher expression of a ROS detoxification**

gene in males and epithelial repair genes in females during enteric

infection

250   Damage limitation mechanisms such as those involved in the response to

oxidative stress, epithelial renewal and damage repair improve host health during

infection while not acting directly to eliminate pathogens. They are therefore putative

mechanisms of disease tolerance (Ayres and Schneider, 2012; Vale et al., 2014) and have

previously been shown to be up-regulated during enteric bacterial infection (Buchon et

al., 2009). Given the differences we observed in the ability to tolerate enteric bacterial

infection (Fig. 4) we hypothesized that male and female flies may differ in the expression

of such genes according to their *Wolbachia* status.

258   In male flies, enteric infection with PA14, led to increased expression of *gstD8* -

involved in ROS detoxification (Buchon et al., 2009; Ha et al., 2005) – which was

significantly higher at 96-hours post-infection in those harbouring *Wolbachia*, while no

difference flies was observed in female *gstD8* expression according to *Wolbachia* status.

Since oral infection results in damage to insect guts (Buchon et al., 2014), we also

measured the expression of two genes involved in epithelial renewal and damage repair

(*gadd45* and CG32302) (Buchon et al., 2009). Both genes showed a significant increase

in expression in *Wolbachia*-positive females. *Gadd45* expression was marginally higher

in *Wolbachia*-positive females compared to those with out *Wolbachia* at 24-hours post-

infection (p = 0.045), but this differnce increased by 96-hours post-infection (p < 0.001).

CG32302 expression was only transiently differentially expressed in *Wolbachia*-positive

females at 24-hours post-infection. *Wolbachia*-negative males showed a significantly

higher expression relative to *Wolbachia*-positive males of both *gadd45* (p = 0.014) and

CG32302 (p = 0.004) at 24 hours post-infection, although this difference was no longer

observed by 96-hours post-infection.

Discussion

During the last decade, it has become well established that endosymbionts like

Wolbachia play a key role in conferring protection from pathogens in their insect hosts

(Brownlie and Johnson, 2009; Hamilton and Perlman, 2013; Karyn N. Johnson, 2015). In

its natural host *Drosophila*, *Wolbachia*-mediated protection is especially evident during

viral infections (Chrostek et al., 2013; Hedges et al., 2008; Martinez et al., 2014; Teixeira

et al., 2008), but protection from bacterial pathogens in *Drosophila* had not been

demonstrated to date (Rottschaefer and Lazzaro, 2012; Wong et al., 2011). Here we

show that the route of infection is important for *Wolbachia*-mediated protection from

bacterial infection. We find that *Wolbachia* can protect *Drosophila* from enteric bacterial

infection by eliciting a combination of resistance and disease tolerance mechanisms, and

that these protective effects are sexually dimorphic.

The route of infection matters

The role of *Wolbachia* in protecting hosts from infection, either by increasing

resistance or tolerance, is known in *Drosophila*-virus interactions, but previous work

testing for antibacterial protection in *Drosophila* did not find a significant effect of

Wolbachia (Rottschaefer and Lazzaro, 2012; Wong et al., 2011). Typically flies in

previous studies were inoculated by intra-thoracic pricking or injection, and therefore

experienced a systemic infection. In the wild however, infections are more likely to be

acquired through the fecal-oral route (during feeding on decomposing fruit), with most

pathogens colonising the gut before being shed through the faeces. *Drosophila*-

Wolbachia interactions would therefore have co-evolved mainly under selection by

pathogen infection in the gut, and any antibacterial protection that may have evolved as

a consequence would not be expected to manifest during a highly virulent systemic

infection (Liehl et al., 2006; Martins et al., 2013). Further, if *Wolbachia*-mediated

protection is especially efficient in the fly gut, the damage caused by a generalised

systemic infection could overwhelm any localised protection by *Wolbachia*, which could

explain the lack of observed protection in previous studies of systemic bacterial

infection in *Drosophila*. Future studies of host resistance and tolerance should therefore

favour natural routes of infection in order to gain a more realistic picture of the

mechanisms that hosts have evolved to fight infection.

***Wolbachia*-mediated protection is a combination of pathogen clearance**

and damage limitation

The mechanisms underlying *Wolbachia*-mediated protection are presently

unclear, especially given that the extent of the protection, and whether it acts to increase

resistance or tolerance appear to be pathogen specific (Chrostek et al., 2013; Martinez et

al., 2014; Teixeira et al., 2008). In mosquitos *Wolbachia* protection appears to be

involved in a combination of general immune priming (Rancès et al., 2012), resource

competition between *Wolbachia* and infectious agents (Caragata et al., 2013), and the

regulation of host genes involved in blocking pathogen replication (Zhang et al., 2013).

In the current experiment it is notable that bacterial numbers did not increase

throughout the course of the infection, but were cleared at a near exponential rate (Fig.

2). Despite this, flies still died from infection, although *Wolbachia* reduced the mortality

rate. One possibility is that most of the damage experienced by the host happens at the

early stages of infection, and the fact that *Wolbachia*-positive flies show lower bacterial

titres immediately after the exposure period (Fig. 2) may be the reason they also show

higher survival later during the infection. This may explain why the male *Wolbachia*

positive flies showed a slower rate of mortality, because *attacinA*-mediated clearance of

PA14 within the first 24 hours post infection (Fig. 3) would have minimized gut damage

caused by pathogen growth.

An alternative, although not mutually exclusive, possibility is that most of the

damage that causes fly death arises from immunopathology, as an indirect side effect of

mechanisms that clear pathogens. For example, a common and broad response to

infection in insects is the activation of pathways that result in the production of reactive

oxygen species (ROS)(Buchon et al., 2014; Ha et al., 2005; Zug and Hammerstein, 2015).

ROS production is tightly regulated (Buchon et al., 2013; Ha et al., 2005), and only

activated in response to pathogenic and not commensal bacteria (Lee et al., 2013).

Previous work has shown that both mosquitoes (Pan et al., 2012) and flies (Wong et al.,

2015) harbouring *Wolbachia* show higher ROS levels, but also that in some cases ROS

production can lead to oxidative stress, DNA damage (Brennan et al., 2012) and damage

to the fly gut epithelium (Buchon et al., 2014, 2013). We therefore hypothesised that

mechanisms involved in detoxifying ROS during enteric infection with PA14 may

underlie the differences in survival between flies with and without *Wolbachia* (Figure

1). We chose to measure the expression of *gstD8,* involved in ROS detoxification,

because it was previously shown to be up-regulated during enteric infection in

Drosophila with another bacterial pathogen, *Erwinia carotovora* (Buchon et al., 2009).

We found that the expression of *gstD8* was elevated in *Wolbachia*-positive males, but not female flies following 96-hours of oral exposure to *P. aeruginosa*. This pattern of expression is consistent with the increased survival observed in *Wolbachia*-positive males compared to males without the endosymbiont (Figure 1b).

In addition to this detoxification response, we also measured the expression of genes involved in tissue damage repair (*gadd45*) and a component of the peritrophic matrix (*CG32302*), a protective barrier in the fly gut (Lehane, 1997). In males, the presence of *Wolbachia* did not result in an increase in these genes within 96 hours of oral exposure to PA14, but females carrying the endosymbiont showed significantly higher expression than *Wolbachia*-negative flies of *gadd45*. This shows that *Wolbachia* induces different damage limitation mechanisms in males (ROS detoxification) and females (tissue damage repair). We also observed transient increases in the expression of CG32302, another component of gut renewal, in *Wolbachia*-positive females at 24-hours post-infection). There was also a transient increase in expression at 24-hours post-infection of *gadd45* and *CG32302* in *Wolbachia*-negative males (Figure 5). We interpret these increases as a response to increased damage to gut tissue cause by the 10-fold higher bacterial loads in these flies after 24 hours (Fig. 2), which was avoided in *Wolbachia*-positive males by *attacinA*-mediated clearance.

While previous work found no difference in genome-wide expression levels in adult *Drosophila* with or without *Wolbachia* (Teixeira, 2012), and only mild up-regulation of immune genes has been reported in *Drosophila* cell lines that are transiently infected (Xi et al., 2008), our gene expression results indicate that reducing immunopathology underlies *Wolbachia*-mediated protection from enteric bacterial infection. We are only beginning to understand the complex sequence of events that occur during gut infection in *Drosophila* (Lemaitre and Miguel-Aliaga, 2013), which not only consist of antimicrobial defence, but a multifaceted response that includes stress response, DNA damage repair, the renewal of damaged epithelial cells and gut structure,

369 and the maintenance of efficient metabolism (Buchon et al., 2010, 2009; Kuraishi et al.,

370 2011).

Sexual dimorphism in resistance and tolerance

The differences in gene expression we describe above reflect two major forms of

defence against infection: mechanisms that eliminate pathogens to reduce infection

loads, leading to resistance, and mechanisms that limit the damage caused by infection

without directly targeting the number of pathogens, leading to disease tolerance (Ayres

and Schneider, 2012; Medzhitov et al., 2012; Råberg et al., 2009; Vale et al., 2014). While

the majority of work on bacterial and viral infections in *Drosophila* (and most other

hosts) has historically focused on mechanisms that eliminate and clear pathogens

(Buchon et al., 2014; Obbard et al., 2009; Zambon et al., 2006), the role of mechanisms

that limit damage to increase host tolerance is increasingly recognised (Ayres and

Schneider, 2012; Medzhitov et al., 2012; Råberg et al., 2009; Soares et al., 2014; Vale et

al., 2014). For example recent work has highlighted how tissue damage repair

(Jamieson et al., 2013; Soares et al., 2014), immune regulation (Merkling et al., 2015;

Sears et al., 2011), and detoxification (Gozzelino et al., 2012; Pamplona et al., 2007) all

play a role in enhancing disease tolerance. Variation in disease tolerance is common

(Adelman et al., 2013; Howick and Lazzaro, 2014; Råberg et al., 2007; Vale and Little,

2012), and may arise from genetic differences in the physiological mechanisms that

promote greater tolerance(Råberg et al., 2007), variation in host nutritional states

(Howick and Lazzaro, 2014; Sternberg et al., 2012; Vale et al., 2011), and host gut

microbiota (Yilmaz et al., 2014). Here we also find evidence for variation in disease

tolerance associated with the presence of *Wolbachia*, and we find that these effects vary

according to host sex.

Wolbachia-mediated protection against viral infections, such as Drosophila C

Virus (DCV) acts by reducing viral replication (or increasing the host's ability to clear

infection), suggesting that *Wolbachia* increases resistance against DCV (Hedges et al.,

2008; Martinez et al., 2014; Teixeira et al., 2008). In other viral infections, for example

Flock House Virus (FHV), flies harbouring *Wolbachia* appear to become more tolerant to

infection, showing increased survival without any change in viral titres (Chrostek et al.,

2013; Teixeira et al., 2008). Our results show that *Wolbachia* affects both resistance and

tolerance to *P. aeruginosa* enteric infection and we found interesting differences

between sexes in these responses. Without *Wolbachia* females were more tolerant than

males, while males flies became more tolerant when carrying *Wolbachia*. While males

and females are generally susceptible to the same pathogens, sexual dimorphism in the

immune response is apparent in a wide range of species (Duneau and Ebert, 2012;

Marriott and Huet-Hudson, 2006; Zuk and McKean, 1996), and is documented for all

classes of viral, bacterial, fungal and parasitic infections [see (Cousineau and Alizon,

2014) for a review]. In invertebrate hosts, and especially in *Drosophila*, most studies

investigating the ability to resist or tolerate bacterial and viral infections have focused

primarily on the underlying immune mechanisms (Ayres and Schneider, 2012; Buchon

et al., 2014; Kemp and Imler, 2009; Neyen et al., 2014; Schneider et al., 2007), and

typically these studies have not focused on sexual differences in these mechanisms [but

see (Vincent and Sharp, 2014)]. Our results, together with a large body of work on

immune sexual dimorphism (Nunn et al., 2009), show that resistance and tolerance

mechanisms are likely to vary between males and females.

Evolutionary and epidemiological implications of sex-specific resistance and

tolerance

Variation in resistance and tolerance will directly impact the pathogen loads

within hosts (Gopinath et al., 2014; Lass et al., 2013; Susi et al., 2015; Vale et al., 2013),

and as a result, sexual dimorphism in these responses could generate potentially

important heterogeneity in pathogen spread and evolution (Cousineau and Alizon, 2014;

Duneau and Ebert, 2012; Gopinath et al., 2014; Vale et al., 2014). Given that *Wolbachia*

are estimated to be highly prevalent in insect populations (Weinert et al., 2015), it is

intriguing to consider the potential effects of sexual dimorphism in resistance and

tolerance in populations. Theoretical work shows that markedly different evolutionary

outcomes for the pathogen are expected when sexual dimorphism in resistance and

tolerance is present (Cousineau and Alizon, 2014). One reason is that the mortality rate

of males and females will vary due to dimorphism in resistance and tolerance, which in

itself will affect the evolutionary trajectories of pathogens (because it will bias infection

towards the one sex more than another). Further experimental work is currently needed

to test how pathogens are likely to evolve under different host sex ratios, especially

when sexual dimorphism in resistance and tolerance is present. Our work suggests that

bacterial oral infection in flies benefiting from sex-specific *Wolbachia*-mediated

tolerance would offer a useful model system to address these questions.

Materials And Methods

Fly stocks

Experiments were carried out using long-term lab stocks of *Drosophila melanogaster*
Oregon R (OreR). This line was originally infected with *Wolbachia* strain wMel,
(OreR[Wol+]). To obtain a *Wolbachia*-free line of the same genetic background (OreR[Wol-]),
OreR[Wol+] flies were cured of *Wolbachia* by rearing them on cornmeal Lewis medium
supplemented with 0.05 mg/ml tetracycline. This treatment was carried out at least 3
years before these experiments were conducted, and the *Wolbachia* status of both fly
lines was verified using PCR with primers specific to *Wolbachia* surface protein (wsp):
forward (5'-3'): GTCCAATAGCTGATGAAGAAAC; reverse (5'-3'):
CTGCACCAATAGCGCTATAAA. Both lines were kept as long-term lab stocks on a
standard diet of cornmeal Lewis medium, at a constant temperature of $18\pm1\,^{\circ}C$ with a
12-hour light/dark cycle. Flies were acclimatised at $25\,^{\circ}C$ for at least two generations
prior to experimental infections.

Bacterial cultures

Infections were carried out using the *P. aeruginosa* reference strain PA14, which has
been shown to have a very broad host range (He et al., 2004; Mikkelsen et al., 2011). To
obtain isogenic PA14 cultures, a frozen stock culture was streaked onto fresh LB agar
plates and single colonies were inoculated into 50 mL LB broth and incubated overnight
at $37\,^{\circ}C$ with shaking at 150 rpm. Overnight cultures were diluted 1:100 into 500 mL
fresh LB broth and incubated again at $37\,^{\circ}C$ with shaking at 150 rpm. At the mid log
phase ($OD_{600} = 1.0$) we harvested the bacterial cells by centrifugation at 8000 rpm for 10
min, washed the cells twice with 1xPBS and re-suspended the bacterial pellet in 5%
sucrose. The final inoculum was adjusted to $OD_{600} = 25$, and this was the bacterial
inoculum used for all flies inoculated orally (enteric infection).

Enteric and systemic *P. aeruginosa* infection

For systemic infection, flies were pricked at the pleural suture with a needle dipped in a

mid log phase (OD_{600} = 1.0) PA14 culture, grown as described above. Control flies were

pricked with a needle dipped in sterile LB broth. For the oral exposure (enteric

infection), the concentrated PA14 inoculum (OD_{600} = 25) was spotted onto a sterile filter

paper (80 µL/ filter paper), and placed onto a drop of solidified 5% sugar Agar inside

the lid of a 7ml Bijou tube. For the uninfected control treatment, filters received the

equivalent volume of 5% sucrose solution only. All filter papers were allowed to dry for

20 to 30 minutes at room temperature. We prepared one of these "inoculation lids" for

each individual fly. Two to four day-old flies were sex sorted and transferred

individually to empty plastic vials: 180 (90 male and 90 female) OreR^{Wol+}, and 180 (90

male and 90 female) OreR^{Wol-}. Following 2-4 hours of starvation, flies were transferred

individually to 7 ml Bijou tubes, and covered with previously prepared "inoculation lids"

containing a filter paper soaked in PA14 culture. Flies were left to feed on the bacterial

culture for approximately 12 hours at 25 ºC. Following this period, we sacrificed 6

exposed and 2 control flies and counted CFUs by plating the fly homogenate in

Pseudomonas isolating media (PAIM). The remaining flies were transferred to vials

containing 5% sugar agar and incubated at 25 ºC.

Quantification of within-host bacterial loads

Following the initial 12-hour exposure, every 24 hours we randomly sampled 5 to 7 live

flies per sex and *Wolbachia* status and quantified the microbe loads present inside the

flies. Briefly, a single fly was removed from the vial and transferred to 1.5 mL

microcentrifuge tubes. To guarantee we were only quantifying CFUs present inside the

fly, and not those possibly on its surface, each fly was surface sterilized by adding 75%

ethanol for 30-60 seconds (to kill the outer surface bacterial species). Ethanol was

discarded and flies were washed twice with distilled water. Plating 100 μL of the 2nd

wash in LB agar confirmed this method was efficient in cleaning the surface of the fly

(no viable CFUs were detected). Each washed whole fly was placed in 1 mL of 1X PBS in

a 1.5-mL screw-top microcentrifuge tube, centrifuged at 5000 rpm for 1 min and the

supernatant was discarded. 200 μL of LB broth was then added to each tube and the flies

were thoroughly homogenised using a motorised pestle for 1 min. A 100 μL aliquot of

homogenate was taken for serial dilution and different dilutions were plated on PAIM

agar plates, incubated at 37 ºC for 24 - 48 h and viable CFUs were counted.

Survival assays

We carried out separate experiments to measure how the presence of *Wolbachia*

affected fly mortality during either enteric or systemic infection, with identical fly

rearing and bacterial cultural conditions as those described above. For each survival

assay (enteric or systemic infection routes), two-to-four day-old flies were sexed and

exposed in groups of 10 flies to PA14, as described above. For each combination of male

or female OreR^{Wol+} and OreR^{Wol-} line, we set up fifty flies, with 10 flies per vial. The flies

that died from infection was recorded every approximately every hour until all flies had

died (systemic infection), or every 24 hours for up to 8 days (enteric infection).

Statistical analyses of host survival and microbe loads

Fly survival was analysed using a Cox proportional hazards model to compare survival

rates, with fly 'Sex', 'Infection status', and '*Wolbachia* status' and their interactions as

fixed effects. The significance of the effects was assessed using likelihood ratio tests

following a χ^2 distribution. For flies that were exposed orally to DCV, we compared

between pairs of treatments (control vs. infected or with and without *Wolbachia*) using

Cox risk ratios.

In orally infected flies, changes in the bacterial load within-hosts were analysed with a

linear model with Log CFU as the response variable, and fly 'Sex', '*Wolbachia* status', and

'Time (DPI)' as fixed effects. Differences in the mean bacterial load of flies immediately

following oral exposure (at 12 hours post-infection; Fig S1) were also analysed

separately using a linear model with Log CFU as the response variable and 'Sex' and

'*Wolbachia* status' as fixed effects. In all models all effects and their interactions were

tested in a fully factorial model, and models were simplified by removing non-significant

interaction terms. All analyses were conducted in JMP 12 (SAS).

Analysis of disease tolerance

Disease tolerance is defined as the ability to maintain health relative to changes in

microbe loads during an infection(Ayres and Schneider, 2012; Medzhitov et al., 2012;

Råberg et al., 2009). It is possible to analyse tolerance as the time-ordered health

trajectory of a host as microbe loads change (Doeschl-Wilson et al., 2012; Lough et al.,

2015; Schneider, 2011). To assess sex- and *Wolbachia*-mediated differences in how sick

a fly gets for a given pathogen load (tolerance) during the course of the infection, for

each time point we took the survival probability (as a measure host health) and PA14

CFUs present within the flies (as a measure of microbe load) for 5 replicate flies in each

sex/*Wolbachia* combination. While tolerance is commonly described as a linear reaction

norm (Lefèvre et al., 2011; Råberg et al., 2007; Simms, 2000), many other functional

forms are possible, including non-linear relationships (Vale et al., 2014). To assess the

form of the health/microbe relationship, we fit linear and non-linear 4-parameter

logistic model separately to the time-matched survival/microbe load plots. In all cases,

the 4 parameter logistic model – which is commonly use to asses dose-response curves

(Gottschalk and Dunn, 2005) - outperformed the linear fit (Table S1), and we present

only the logistic fit in the results section. To test if these tolerance curves differed with

Wolbachia status, we tested the parallelism of the models by comparing the error-sums

of square for a full model (where each group has different parameters in the logistic

model) to a reduced model (where models share all parameters except the inflection

point) (Gottschalk and Dunn, 2005).

Gene expression

We used qRT-PCR to test for differences in the expression of genes known to be involved

in either bacterial clearance or in the response to stress and gut damage during enteric

infection. Previous work has shown that the IMD pathway and some stress response and

damage repair genes are especially important during the fly's response to enteric

bacterial pathogens (Buchon et al., 2009). We therefore investigated the expression of

IMD pathway receptor genes (*pgrp – lc* and *pgrp – le*) and the antimicrobial peptide

effector gene *attacin A* (attA); *gstD8*, a gene that participates in the detoxification of

reactive oxygen species (ROS) produced during microbial immunity in the gut (Ha et al.,

2005); *gadd45*, a gene relevant for epithelial renewal, as it is involved in stress response

and wound healing in *Drosophila* (Stramer et al., 2008; Takekawa and Saito, 1998); and

CG32302, a gene identified as up-regulated during enteric bacterial infection(Buchon et

al., 2009). CG32302 has been annotated as a putative component of the peritrophic

matrix (Buchon et al., 2009), a protective barrier that separates the gut epithelium from

the invading bacteria (Lehane, 1997).

Our aim was to test if the expression of these genes varied in a sex- or

Wolbachia-specific manner in flies that were infected orally. Groups of 5 flies for each

sex / *Wolbachia* combination were exposed orally to *P. aeruginosa* infection in triplicate,

as described above, and then frozen in TRI reagent at 4, 24 and 96 hours post-infection.

Total RNA was extracted from flies homogenised in Tri Reagent (Ambion), reverse-

transcribed with M-MLV reverse transcriptase (Promega) and random hexamer

primers, and then diluted 1:10 with nuclease free water. The qRT-PCR was performed

on an Applied Biosystems StepOnePlus system using Fast SYBR Green Master Mix

(Applied Biosystems) with the following PCR cycle: 95°C for 2min followed by 40 cycles

of 95°C for 10 sec followed by 60°C for 30 sec. Three biological replicates and two qRT-

PCR reactions per replicate (technical replicates) were carried out per sample. Gene-

specific primers are reported in Table S1. Changes in gene expression were analysed

relative to the expression of *rp49,* an internal control gene. The relative fold-change

difference in expression between infected and health control flies was calculated as

described in (Livak and Schmittgen, 2001). Briefly:

$$\text{Fold change} = 2^{-\Delta\Delta Ct}$$

Where, $\Delta\Delta Ct$ = [(Ct of Gene A – Ct of Internal control) of Infected sample] –

[(Ct of Gene A – Ct of Internal control) of Control sample]

The fold change difference obtained was analysed using 3-way ANOVA with sex (Male,

Female), Time (2, 24 and 96 hours) and *Wolbachia* (Wol- and Wol+) as fixed factors.

Acknowledgements

591
592 We thank R. Popat for technical assistance, D. Obbard, F. Waldron, T.Little and S. Lewis

593 for helpful discussion and suggestions, and H. Cowan, H. Borthwick for media

594 preparation.

Funding information

597
598 This work was supported by a strategic award from the Wellcome Trust for the Centre

599 for Immunity, Infection and Evolution (http://ciie.bio.ed.ac.uk; grant reference no.

600   095831), and by a Society in Science – Branco Weiss fellowship (http://www.society-in-

601   science.org), both awarded to P. Vale. P. Vale is also supported by a Chancellor's

602   fellowship from the School of Biological Sciences

603
Author contributions

605   PFV and RBV conceived the study. RBV, GV, JSJ, KMM and PFV conducted experimental

606   work. GV and PFV analysed the data. PFV wrote the manuscript. SPB and PFV

607   contributed reagents and consumables. All authors commented on the manuscript.

References

Adelman, J.S., Kirkpatrick, L., Grodio, Hawley, D.M., 2013. House Finch Populations Differ in Early Inflammatory Signaling and Pathogen Tolerance at the Peak of Mycoplasma gallisepticum Infection. Am. Nat. 181, 674–689. doi:10.1086/670024

Apidianakis, Y., Rahme, L.G., 2009. Drosophila melanogaster as a model host for studying Pseudomonas aeruginosa infection. Nat Protoc. 4, 1285–1294. doi:10.1038/nprot.2009.124

Ayres, J.S., Schneider, D.S., 2012. Tolerance of Infections. Annu. Rev. Immunol. 30, 271–294. doi:10.1146/annurev-immunol-020711-075030

Bashir-Tanoli, S., Tinsley, M.C., 2014. Immune response costs are associated with changes in resource acquisition and not resource reallocation. Funct. Ecol. 28, 1011–1019. doi:10.1111/1365-2435.12236

Bennett, G.M., Moran, N.A., 2015. Heritable symbiosis: The advantages and perils of an evolutionary rabbit hole. Proc. Natl. Acad. Sci. U. S. A. 112, 10169–10176. doi:10.1073/pnas.1421388112

Bian, G., Xu, Y., Lu, P., Xie, Y., Xi, Z., 2010. The Endosymbiotic Bacterium Wolbachia Induces Resistance to Dengue Virus in Aedes aegypti. PLoS Pathog 6, e1000833. doi:10.1371/journal.ppat.1000833

Brennan, L.J., Haukedal, J.A., Earle, J.C., Keddie, B., Harris, H.L., 2012. Disruption of redox homeostasis leads to oxidative DNA damage in spermatocytes of Wolbachia-infected Drosophila simulans. Insect Mol. Biol. 21, 510–520. doi:10.1111/j.1365-2583.2012.01155.x

Brownlie, J.C., Johnson, K.N., 2009. Symbiont-mediated protection in insect hosts. Trends Microbiol. 17, 348–354. doi:10.1016/j.tim.2009.05.005

Buchon, N., Broderick, N.A., Kuraishi, T., Lemaitre, B., 2010. Drosophila EGFR pathway coordinates stem cell proliferation and gut remodeling following infection. BMC Biol. 8, 152.

Buchon, N., Broderick, N.A., Lemaitre, B., 2013. Gut homeostasis in a microbial world: insights from Drosophila melanogaster. Nat. Rev. Microbiol. 11, 615–626. doi:10.1038/nrmicro3074

Buchon, N., Broderick, N.A., Poidevin, M., Pradervand, S., Lemaitre, B., 2009. Drosophila Intestinal Response to Bacterial Infection: Activation of Host Defense and Stem Cell Proliferation. Cell Host Microbe 5, 200–211. doi:10.1016/j.chom.2009.01.003

Buchon, N., Silverman, N., Cherry, S., 2014. Immunity in Drosophila melanogaster — from microbial recognition to whole-organism physiology. Nat. Rev. Immunol. 14, 796–810. doi:10.1038/nri3763

Caragata, E.P., Rancès, E., Hedges, L.M., Gofton, A.W., Johnson, K.N., O'Neill, S.L., McGraw, E.A., 2013. Dietary Cholesterol Modulates Pathogen Blocking by Wolbachia. PLoS Pathog 9, e1003459. doi:10.1371/journal.ppat.1003459

Chrostek, E., Marialva, M.S.P., Esteves, S.S., Weinert, L.A., Martinez, J., Jiggins, F.M., Teixeira, L., 2013. Wolbachia Variants Induce Differential Protection to Viruses in Drosophila melanogaster: A Phenotypic and Phylogenomic Analysis. PLoS Genet. 9. doi:10.1371/journal.pgen.1003896

Chugani, S.A., Whiteley, M., Lee, K.M., D'Argenio, D., Manoil, C., Greenberg, E.P., 2001. QscR, a modulator of quorum-sensing signal synthesis and

virulence in Pseudomonas aeruginosa. Proc. Natl. Acad. Sci. 98, 2752–
2757. doi:10.1073/pnas.051624298
Cousineau, S.V., Alizon, S., 2014. Parasite evolution in response to sex-based host
heterogeneity in resistance and tolerance. J. Evol. Biol. 27, 2753–2766.
doi:10.1111/jeb.12541
Doeschl-Wilson, A.B., Bishop, S.C., Kyriazakis, I., Villanueva, B., 2012. Novel
methods for quantifying individual host response to infectious pathogens
for genetic analyses. Front. Livest. Genomics 3, 266.
doi:10.3389/fgene.2012.00266
Douglas, A.E., 2015. Multiorganismal Insects: Diversity and Function of Resident
Microorganisms. Annu. Rev. Entomol. 60, 17–34. doi:10.1146/annurev-
ento-010814-020822
Douglas, A.E., 1998. Nutritional Interactions in Insect-Microbial Symbioses:
Aphids and Their Symbiotic Bacteria Buchnera. Annu. Rev. Entomol. 43,
17–37. doi:10.1146/annurev.ento.43.1.17
Duneau, D., Ebert, D., 2012. Host sexual dimorphism and parasite adaptation.
PLoS Biol. 10, e1001271. doi:10.1371/journal.pbio.1001271
Engelstädter, J., Hurst, G.D.D., 2009. The Ecology and Evolution of Microbes that
Manipulate Host Reproduction. Annu. Rev. Ecol. Evol. Syst. 40, 127–149.
doi:10.1146/annurev.ecolsys.110308.120206
Ferreira, Á.G., Naylor, H., Esteves, S.S., Pais, I.S., Martins, N.E., Teixeira, L., 2014.
The Toll-Dorsal Pathway Is Required for Resistance to Viral Oral Infection
in Drosophila. PLoS Pathog. 10. doi:10.1371/journal.ppat.1004507
Gandon, S., Vale, P.F., 2014. The evolution of resistance against good and bad
infections. J. Evol. Biol. 27, 303–312. doi:10.1111/jeb.12291
Gopinath, S., Lichtman, J.S., Bouley, D.M., Elias, J.E., Monack, D.M., 2014. Role of
disease-associated tolerance in infectious superspreaders. Proc. Natl.
Acad. Sci. 111, 15780–15785. doi:10.1073/pnas.1409968111
Gottschalk, P.G., Dunn, J.R., 2005. Measuring parallelism, linearity, and relative
potency in bioassay and immunoassay data. J. Biopharm. Stat. 15, 437–
463. doi:10.1081/BIP-200056532
Gozzelino, R., Andrade, B.B., Larsen, R., Luz, N.F., Vanoaica, L., Seixas, E., Coutinho,
A., Cardoso, S., Rebelo, S., Poli, M., Barral-Netto, M., Darshan, D., Kühn, L.C.,
Soares, M.P., 2012. Metabolic Adaptation to Tissue Iron Overload Confers
Tolerance to Malaria. Cell Host Microbe 12, 693–704.
doi:10.1016/j.chom.2012.10.011
Ha, E.-M., Oh, C.-T., Bae, Y.S., Lee, W.-J., 2005. A Direct Role for Dual Oxidase in
Drosophila Gut Immunity. Science 310, 847–850.
doi:10.1126/science.1117311
Hamilton, P.T., Perlman, S.J., 2013. Host Defense via Symbiosis in Drosophila.
PLoS Pathog 9, e1003808. doi:10.1371/journal.ppat.1003808
Hedges, L.M., Brownlie, J.C., O'Neill, S.L., Johnson, K.N., 2008. Wolbachia and Virus
Protection in Insects. Science 322, 702. doi:10.1126/science.1162418
Hedges, L.M., Johnson, K.N., 2008. Induction of Host Defence Responses by
Drosophila C Virus. J. Gen. Virol. 89, 1497–1501. doi:10.1099/vir.0.83684-
0
He, J., Baldini, R.L., Déziel, E., Saucier, M., Zhang, Q., Liberati, N.T., Lee, D., Urbach,
J., Goodman, H.M., Rahme, L.G., 2004. The broad host range pathogen
Pseudomonas aeruginosa strain PA14 carries two pathogenicity islands

706   harboring plant and animal virulence genes. Proc. Natl. Acad. Sci. U. S. A.
707   101, 2530–2535.
708 Howick, V.M., Lazzaro, B.P., 2014. Genotype and diet shape resistance and
709   tolerance across distinct phases of bacterial infection. BMC Evol. Biol. 14,
710   56. doi:10.1186/1471-2148-14-56
711 Jamieson, A.M., Pasman, L., Yu, S., Gamradt, P., Homer, R.J., Decker, T., Medzhitov,
712   R., 2013. Role of Tissue Protection in Lethal Respiratory Viral-Bacterial
713   Coinfection. Science 340, 1230–1234. doi:10.1126/science.1233632
714 Johnson, K.N., 2015. Bacteria and antiviral immunity in insects. Curr. Opin. Insect
715   Sci. 8, 97–103.
716 Johnson, K.N., 2015. Bacteria and antiviral immunity in insects. Curr. Opin. Insect
717   Sci. doi:10.1016/j.cois.2015.01.008
718 Kambris, Z., Blagborough, A.M., Pinto, S.B., Blagrove, M.S.C., Godfray, H.C.J.,
719   Sinden, R.E., Sinkins, S.P., 2010. Wolbachia Stimulates Immune Gene
720   Expression and Inhibits Plasmodium Development in Anopheles gambiae.
721   PLoS Pathog 6, e1001143. doi:10.1371/journal.ppat.1001143
722 Kemp, C., Imler, J.-L., 2009. Antiviral immunity in drosophila. Curr. Opin.
723   Immunol. 21, 3–9. doi:10.1016/j.coi.2009.01.007
724 Kuraishi, T., Binggeli, O., Opota, O., Buchon, N., Lemaitre, B., 2011. Genetic
725   evidence for a protective role of the peritrophic matrix against intestinal
726   bacterial infection in Drosophila melanogaster. Proc. Natl. Acad. Sci. 108,
727   15966–15971. doi:10.1073/pnas.1105994108
728 Lass, S., Hudson, P.J., Thakar, J., Saric, J., Harvill, E., Albert, R., Perkins, S.E., 2013.
729   Generating super-shedders: co-infection increases bacterial load and egg
730   production of a gastrointestinal helminth. J. R. Soc. Interface 10,
731   20120588. doi:10.1098/rsif.2012.0588
732 Lee, K.-A., Kim, S.-H., Kim, E.-K., Ha, E.-M., You, H., Kim, B., Kim, M.-J., Kwon, Y.,
733   Ryu, J.-H., Lee, W.-J., 2013. Bacterial-Derived Uracil as a Modulator of
734   Mucosal Immunity and Gut-Microbe Homeostasis in Drosophila. Cell 153,
735   797–811. doi:10.1016/j.cell.2013.04.009
736 Lefèvre, T., Williams, A.J., de Roode, J.C., 2011. Genetic variation in resistance, but
737   not tolerance, to a protozoan parasite in the monarch butterfly. Proc. Biol.
738   Sci. 278, 751–759. doi:10.1098/rspb.2010.1479
739 Lehane, M.J., 1997. Peritrophic Matrix Structure and Function. Annu. Rev.
740   Entomol. 42, 525–550. doi:10.1146/annurev.ento.42.1.525
741 Lemaitre, B., Miguel-Aliaga, I., 2013. The digestive tract of Drosophila
742   melanogaster. Annu. Rev. Genet. 47, 377–404. doi:10.1146/annurev-
743   genet-111212-133343
744 Lewis, Z., Lizé, A., 2015. Insect behaviour and the microbiome. Curr. Opin. Insect
745   Sci.
746 Liehl, P., Blight, M., Vodovar, N., Boccard, F., Lemaitre, B., 2006. Prevalence of
747   Local Immune Response against Oral Infection in a
748   Drosophila/Pseudomonas Infection Model. PLoS Pathog 2, e56.
749   doi:10.1371/journal.ppat.0020056
750 Livak, K.J., Schmittgen, T.D., 2001. Analysis of relative gene expression data using
751   real-time quantitative PCR and the 2(-Delta Delta C(T)) Method. Methods
752   San Diego Calif 25, 402–408. doi:10.1006/meth.2001.1262
753 Lough, G., Kyriazakis, I., Bergmann, S., Lengeling, A., Doeschl-Wilson, A.B., 2015.
754   Health trajectories reveal the dynamic contributions of host genetic

resistance and tolerance to infection outcome. Proc R Soc B 282,
20152151. doi:10.1098/rspb.2015.2151

Marriott, I., Huet-Hudson, Y.M., 2006. Sexual dimorphism in innate immune
responses to infectious organisms. Immunol. Res. 34, 177–192.
doi:10.1385/IR:34:3:177

Martinez, J., Longdon, B., Bauer, S., Chan, Y.-S., Miller, W.J., Bourtzis, K., Teixeira,
L., Jiggins, F.M., 2014. Symbionts Commonly Provide Broad Spectrum
Resistance to Viruses in Insects: A Comparative Analysis of Wolbachia
Strains. PLoS Pathog 10, e1004369. doi:10.1371/journal.ppat.1004369

Martins, N.E., Faria, V.G., Teixeira, L., Magalhães, S., Sucena, É., 2013. Host
Adaptation Is Contingent upon the Infection Route Taken by Pathogens.
PLoS Pathog. 9. doi:10.1371/journal.ppat.1003601

Mateos, M., Castrezana, S.J., Nankivell, B.J., Estes, A.M., Markow, T.A., Moran, N.A.,
2006. Heritable Endosymbionts of Drosophila. Genetics 174, 363–376.
doi:10.1534/genetics.106.058818

Medzhitov, R., Schneider, D.S., Soares, M.P., 2012. Disease Tolerance as a Defense
Strategy. Science 335, 936–941. doi:10.1126/science.1214935

Merkling, S.H., Bronkhorst, A.W., Kramer, J.M., Overheul, G.J., Schenck, A., Van Rij,
R.P., 2015. The Epigenetic Regulator G9a Mediates Tolerance to RNA
Virus Infection in Drosophila. PLoS Pathog 11, e1004692.
doi:10.1371/journal.ppat.1004692

Mikkelsen, H., McMullan, R., Filloux, A., 2011. The Pseudomonas aeruginosa
Reference Strain PA14 Displays Increased Virulence Due to a Mutation in
ladS. PLoS ONE 6. doi:10.1371/journal.pone.0029113

Moreira, L.A., Iturbe-Ormaetxe, I., Jeffery, J.A., Lu, G., Pyke, A.T., Hedges, L.M.,
Rocha, B.C., Hall-Mendelin, S., Day, A., Riegler, M., Hugo, L.E., Johnson, K.N.,
Kay, B.H., McGraw, E.A., van den Hurk, A.F., Ryan, P.A., O'Neill, S.L., 2009. A
Wolbachia Symbiont in Aedes aegypti Limits Infection with Dengue,
Chikungunya, and Plasmodium. Cell 139, 1268–1278.
doi:10.1016/j.cell.2009.11.042

Mulcahy, H., Sibley, C.D., Surette, M.G., Lewenza, S., 2011. Drosophila
melanogaster as an Animal Model for the Study of Pseudomonas
aeruginosa Biofilm Infections In Vivo. PLoS Pathog 7, e1002299.
doi:10.1371/journal.ppat.1002299

Neyen, C., Bretscher, A.J., Binggeli, O., Lemaitre, B., 2014. Methods to study
Drosophila immunity. Methods San Diego Calif 68, 116–128.
doi:10.1016/j.ymeth.2014.02.023

Nunn, C.L., Lindenfors, P., Pursall, E.R., Rolff, J., 2009. On sexual dimorphism in
immune function. Philos. Trans. R. Soc. Lond. B Biol. Sci. 364, 61–69.
doi:10.1098/rstb.2008.0148

Obbard, D.J., Gordon, K.H.J., Buck, A.H., Jiggins, F.M., 2009. The evolution of RNAi
as a defence against viruses and transposable elements. Philos. Trans. R.
Soc. B Biol. Sci. 364, 99–115. doi:10.1098/rstb.2008.0168

Pamplona, A., Ferreira, A., Balla, J., Jeney, V., Balla, G., Epiphanio, S., Chora, Â.,
Rodrigues, C.D., Gregoire, I.P., Cunha-Rodrigues, M., Portugal, S., Soares,
M.P., Mota, M.M., 2007. Heme oxygenase-1 and carbon monoxide suppress
the pathogenesis of experimental cerebral malaria. Nat. Med. 13, 703–
710. doi:10.1038/nm1586

Pan, X., Zhou, G., Wu, J., Bian, G., Lu, P., Raikhel, A.S., Xi, Z., 2012. Wolbachia
induces reactive oxygen species (ROS)-dependent activation of the Toll
pathway to control dengue virus in the mosquito Aedes aegypti. Proc.
Natl. Acad. Sci. 109, E23–E31. doi:10.1073/pnas.1116932108
Råberg, L., Graham, A.L., Read, A.F., 2009. Decomposing health: tolerance and
resistance to parasites in animals. Philos. Trans. R. Soc. Lond. B. Biol. Sci.
364, 37–49. doi:10.1098/rstb.2008.0184
Råberg, L., Sim, D., Read, A.F., 2007. Disentangling genetic variation for resistance
and tolerance to infectious diseases in animals. Science 318, 812–814.
doi:10.1126/science.1148526
Rancès, E., Ye, Y.H., Woolfit, M., McGraw, E.A., O'Neill, S.L., 2012. The Relative
Importance of Innate Immune Priming in Wolbachia-Mediated Dengue
Interference. PLoS Pathog 8, e1002548.
doi:10.1371/journal.ppat.1002548
Rottschaefer, S.M., Lazzaro, B.P., 2012. No Effect of Wolbachia on Resistance to
Intracellular Infection by Pathogenic Bacteria in Drosophila melanogaster.
PLoS ONE 7, e40500. doi:10.1371/journal.pone.0040500
Schneider, D.S., 2011. Tracing Personalized Health Curves during Infections.
PLoS Biol. 9, e1001158.
Schneider, D.S., Ayres, J.S., Brandt, S.M., Costa, A., Dionne, M.S., Gordon, M.D.,
Mabery, E.M., Moule, M.G., Pham, L.N., Shirasu-Hiza, M.M., 2007.
Drosophila eiger Mutants Are Sensitive to Extracellular Pathogens. PLoS
Pathog 3, e41. doi:10.1371/journal.ppat.0030041
Sears, B.F., Rohr, J.R., Allen, J.E., Martin, L.B., 2011. The economy of inflammation:
when is less more? Trends Parasitol. 27, 382–387.
doi:10.1016/j.pt.2011.05.004
Sibley, C.D., Duan, K., Fischer, C., Parkins, M.D., Storey, D.G., Rabin, H.R., Surette,
M.G., 2008. Discerning the Complexity of Community Interactions Using a
Drosophila Model of Polymicrobial Infections. PLoS Pathog 4, e1000184.
doi:10.1371/journal.ppat.1000184
Simms, E.., 2000. Defining tolerance as a norm of reaction. Evol. Ecol. 14, 563–
570.
Soares, M.P., Gozzelino, R., Weis, S., 2014. Tissue damage control in disease
tolerance. Trends Immunol. 35, 483–494. doi:10.1016/j.it.2014.08.001
Sternberg, E.D., Lefèvre, T., Li, J., de Castillejo, C.L.F., Li, H., Hunter, M.D., de Roode,
J.C., 2012. Food plant derived disease tolerance and resistance in a natural
butterfly-plant-parasite interactions. Evol. Int. J. Org. Evol. 66, 3367–3376.
doi:10.1111/j.1558-5646.2012.01693.x
Stevanovic, A., Johnson, K.N., 2015. Infectivity of Drosophila C virus following
oral delivery in Drosophila larvae. J. Gen. Virol. doi:10.1099/vir.0.000068
Stramer, B., Winfield, M., Shaw, T., Millard, T.H., Woolner, S., Martin, P., 2008.
Gene induction following wounding of wild-type versus macrophage-
deficient Drosophila embryos. EMBO Rep. 9, 465–471.
doi:10.1038/embor.2008.34
Susi, H., Barrès, B., Vale, P.F., Laine, A.-L., 2015. Co-infection alters population
dynamics of infectious disease. Nat. Commun. 6.
doi:10.1038/ncomms6975

Takekawa, M., Saito, H., 1998. A family of stress-inducible GADD45-like proteins
mediate activation of the stress-responsive MTK1/MEKK4 MAPKKK. Cell
95, 521–530.
Teixeira, L., 2012. Whole-genome expression profile analysis of Drosophila
melanogaster immune responses. Brief. Funct. Genomics 11, 375–386.
doi:10.1093/bfgp/els043
Teixeira, L., Ferreira, Á., Ashburner, M., 2008. The Bacterial Symbiont Wolbachia
Induces Resistance to RNA Viral Infections in Drosophila melanogaster.
PLoS Biol 6, e1000002. doi:10.1371/journal.pbio.1000002
Vale, P.F., Choisy, M., Little, T.J., 2013. Host nutrition alters the variance in
parasite transmission potential. Biol. Lett. 9, 20121145.
doi:10.1098/rsbl.2012.1145
Vale, P.F., Fenton, A., Brown, S.P., 2014. Limiting Damage during Infection:
Lessons from Infection Tolerance for Novel Therapeutics. PLoS Biol 12,
e1001769. doi:10.1371/journal.pbio.1001769
Vale, P.F., Little, T.J., 2012. Fecundity compensation and tolerance to a sterilizing
pathogen in Daphnia. J. Evol. Biol. 25, 1888–1896. doi:10.1111/j.1420-
9101.2012.02579.x
Vale, P.F., McNally, L., Doeschl-Wilson, A.B., King, K.C., Domingo-Sananes, M.R.,
Popat, R., Allen, J.E., Soares, M.P., Kummerli, R., 2016. Beyond killing: can
we find new ways to manage infection? Evol. Med. Public Health in press.
Vale, P.F., Wilson, A.J., Best, A., Boots, M., Little, T.J., 2011. Epidemiological,
Evolutionary, and Coevolutionary Implications of Context-Dependent
Parasitism. Am. Nat. 177, 510–521.
Vincent, C.M., Sharp, N.P., 2014. Sexual antagonism for resistance and tolerance
to infection in Drosophila melanogaster. Proc. Biol. Sci. 281, 20140987.
doi:10.1098/rspb.2014.0987
Weinert, L.A., Araujo-Jnr, E.V., Ahmed, M.Z., Welch, J.J., 2015. The incidence of
bacterial endosymbionts in terrestrial arthropods. Proc. R. Soc. Lond. B
Biol. Sci. 282, 20150249. doi:10.1098/rspb.2015.0249
Werren, J.H., Baldo, L., Clark, M.E., 2008. Wolbachia: master manipulators of
invertebrate biology. Nat. Rev. Microbiol. 6, 741–751.
doi:10.1038/nrmicro1969
Wong, Z.S., Brownlie, J.C., Johnson, K.N., 2015. Oxidative Stress Correlates with
Wolbachia-Mediated Antiviral Protection in Wolbachia-Drosophila
Associations. Appl. Environ. Microbiol. 81, 3001–3005.
doi:10.1128/AEM.03847-14
Wong, Z.S., Hedges, L.M., Brownlie, J.C., Johnson, K.N., 2011. Wolbachia-Mediated
Antibacterial Protection and Immune Gene Regulation in Drosophila.
PLoS ONE 6, e25430. doi:10.1371/journal.pone.0025430
Xi, Z., Gavotte, L., Xie, Y., Dobson, S.L., 2008. Genome-wide analysis of the
interaction between the endosymbiotic bacterium Wolbachia and its
Drosophila host. BMC Genomics 9, 1. doi:10.1186/1471-2164-9-1
Ye, Y.H., Woolfit, M., Rancès, E., O'Neill, S.L., McGraw, E.A., 2013. Wolbachia-
Associated Bacterial Protection in the Mosquito Aedes aegypti. PLoS Negl
Trop Dis 7, e2362. doi:10.1371/journal.pntd.0002362
Yilmaz, B., Portugal, S., Tran, T.M., Gozzelino, R., Ramos, S., Gomes, J., Regalado, A.,
Cowan, P.J., d'Apice, A.J.F., Chong, A.S., Doumbo, O.K., Traore, B., Crompton,
P.D., Silveira, H., Soares, M.P., 2014. Gut Microbiota Elicits a Protective

Immune Response against Malaria Transmission. Cell 159, 1277–1289.
900   doi:10.1016/j.cell.2014.10.053

901   Zambon, R.A., Vakharia, V.N., Wu, L.P., 2006. RNAi is an antiviral immune
902   response against a dsRNA virus in Drosophila melanogaster. Cell.
903   Microbiol. 8, 880–889. doi:10.1111/j.1462-5822.2006.00688.x

904   Zhang, G., Hussain, M., O'Neill, S.L., Asgari, S., 2013. Wolbachia uses a host
905   microRNA to regulate transcripts of a methyltransferase, contributing to
906   dengue virus inhibition in Aedes aegypti. Proc. Natl. Acad. Sci. 110,
907   10276–10281. doi:10.1073/pnas.1303603110

908   Zug, R., Hammerstein, P., 2015. Wolbachia and the insect immune system: what
909   reactive oxygen species can tell us about the mechanisms of Wolbachia–
910   host interactions. Evol. Genomic Microbiol. 1201.
911   doi:10.3389/fmicb.2015.01201

912   Zuk, M., McKean, K.A., 1996. Sex differences in parasite infections: Patterns and
913   processes. Int. J. Parasitol. 26, 1009–1024. doi:10.1016/S0020-
914   7519(96)80001-4

Figure Legends

Figure 1. Fly survival after systemic (Fig. 1a) oral infection (Fig. 1b) with *P.*
***aerugino*sa PA14.** OreR[Wol-] (black line), OreR[Wol+] (grey line) were either pricked with a
needle dipped in PA14 culture (OD=1) , or left to feed on a PA14 culture (OD=23) or on a
control solution of 5% sugar for 12 hours. Survival was monitored for 24h (systemic
infection) or daily (oral infection). Data were analysed using a Cox Proportional Hazard
model.

Figure 2. Within-host microbe loads. The number of viable within-host CFUs was
quantified in 5-7 individual live flies following 12 hours of oral exposure, and then every
24 hours for a week. Males and females are plotted separately for OreR[Wol-] (black) and
OreR[Wol+] (grey) flies. Data shown are means ± SEM.

Figure 3. Gene expression of IMD pathway genes. Figures show gene expression

relative to *rp49* control gene in infected flies relative to uninfected flies. Mean of 3

biological replicates ± SE.

Figure 4. Disease tolerance. To measure tolerance we analysed the relationship

between host health and microbe loads. For each time point, we plot the survival

probability (as a measure of health) against the microbe load (number of CFU per fly)

for 5 biological replicates per sex and *Wolbachia* combination. Here we show the fit of a

4-parameter Logisitic model to the data (see Table S1 for model fits). The X axis is

reversed to read from beginning to the end of the infection (only clearance occurred.)

Figure 5. Gene expression of IMD pathway genes. Figures show gene expression

relative to *rp49* control gene in infected flies relative to uninfected flies. Mean of 3

biological replicates ± SE.

Supporting Information

Fig S1. Quantification of food intake. To quantify feeding, individual flies were fed on blue-dyed medium for 24 hours, homogenised and suspended in buffer. The absorbance of this suspension, which is proportional to the amount of food intake (Bashir-Tanoli and Tinsley, 2014), was measured at 520 nm. Data shown are means ± SEM.

Fig S2. Inhibition of PA14 growth in liquid culture. Single sex groups of *Wolbachia* positive or *Wolbachia* negative flies were homogenised, centrifuged and the supernatant was inoculated into PA14 cultures growing in 96-well plates. Absorbance was measured every 30 minutes.

Fig S3. Inhibition zone assay. Single sex groups of *Wolbachia* positive or *Wolbachia* negative flies were homogenised, centrifuged and the supernatant was spotted on a lawn of PA14 overnight. The figure shows the size of the zone where PA14 growth was inhibited. Data shown are means ± SEM.

S1 file. Materials and methods for feeding assay and PA14 growth inhibition assays.

Supplementary information

***Wolbachia* confers sex-specific resistance and tolerance to enteric but not**

systemic bacterial infection in *Drosophila*

Radhakrishnan B. Vasanthakrishnan [1,3§], Gupta Vanika [1§], Jonathon Siva-Jothy, Katy M.

Monteith[1], Sam P. Brown[4], Pedro F. Vale[1,2*],

Table S1: Primer list

Primer name	Gene function	Sequence (5'->3')
PGRP-LC Forward	IMD immune pathway-Extracellular receptor	TTGAACCAAAGTAAGATCAGAGAT
PGRP-LC Reverse		GTCCAGATATATTGTTGAATT
PGRP - LE Forward	IMD immune pathway-Intracellular receptor	GATGCCGACCAAAATACCAG
PGRP - LE Reverse		GTCTTCGAAATGTGTCGGAG
Attacin A Forward	IMD immune pathway-Antimicrobial peptide	GGCCCATGCCAATTTATTCA
Attacin A Reverse		CATTGCGCTGGAACTCGAA
Gst D8 Forward	Stress response	GGAATCCCGTGCCATTTTGA
Gst D8 Reverse		CCCATGTCGAAGTAGAGCCT
CG32302 Forward	Peritrophic matrix	CGATGGAGAACTGGAGGTGA
CG32302 Reverse		TATCAGTCACGCAGGTCAGG
Gadd45 Forward	Wounding stress	ACTGGACCTGGAGCTAGAGA
Gadd45 Reverse		CTTGGAGAGCACGTTGATGG

Feeding assay

All flies used were 0-72 hour old adult Oregon R flies, and Wol- and wol+ individuals

were raised separately in 6oz bottles on lewis medium at a constant temperature of

25ºC on a 12:12 light:dark cycle. Subjects were starved for 2 hours by placing them an

empty vial before CO_2 anaesthesia was used to separate sexes. While still anaesthetised

each fly was placed into an individual 23ml vial containing Blue-dyed Lewis medium

(Blue Dye number 1, 0.5g/litre). Between 22-24 individual flies per Sex/Wolbachia

combination were set up. All flies were left to feed on the blue dyed medium for 24hours

at 25 degrees under 12:12 light:dark cycle. After 24 hours flies were immediately frozen

to kill and then decapitated using a scalpel. Decapitation avoids inaccurate absorbance

readings due to eye pigments. Each individual fly was then placed in an Eppendorf tube

containing 100ul of ice-cold Ringer's solution, homogenised using a motorised pestle

and then centrifuged for 10mins at ~13300g at 20ºC. 80 µl of this supernatant were

loaded into a 96-well plate and the blue pigment was measured using a VersaMax

microplate reader, recording absorbance at 520nm.

1004

1005

1006   **Fig S1. Quantification of food intake.** To quantify feeding, individual flies were fed on

1007   blue-dyed medium for 24 hours, homogenised and suspended in buffer. The absorbance

1008   of this suspension, which is proportional to the amount of food intake (Bashir-Tanoli

1009   and Tinsley, 2014), was measured at 520 nm. Data shown are means ± SEM

1010

Pseudomonas aeruginosa PA14 inhibition assay

Wol+ and Wol- flies were grown in 6 oz plastic bottles on standard Lewis medium under standard laboratory conditions at 25°C , 12:12 Light:Dark cycle. Six days post eclosion, flies were sex separated and divided into cohorts of 10 flies each (n = 10 per sex per line). Flies were anaesthetized using cold anaesthesia. To each sample, 250 µl of LB was added and samples were crushed using an automated pestle and centrifuged at 11,000g for 5 minutes. Supernatant from these samples was used to assay antibacterial response. 100 µl from supernatant was transferred to a well in 96-well plate. PA14 cultures were grown as follows. A single colony was picked from the PA plate and 5 ml of LB medium was inoculated. The culture was grown overnight at 37°C at 200 rpm. 100 µl of the overnight culture was used to seed 5ml of LB. OD was monitored and culture was taken out after it reached OD_{600} = 1. The OD of the culture was adjusted to 0.02. Each well was seeded with 100 µl of this culture and was mixed well with fly homogenate by pipetting. For controls, LB without any bacteria was used. Bacterial growth in 96-well plate was monitored overnight using Thermo Scientific Varioskan Flash. Plate was incubated at 37°C and readings were collected every half an hour intervals for 16 hours. Data obtained was used to obtain lag phase time, growth rate and saturation time for each well using Growth Curve Analysis Tool (GCAT). These data were analysed using ANOVA with fly sex and Wolbachia status as factors.

1034

Fig S2. Inhibition of PA14 growth in liquid culture. Single sex groups of *Wolbachia*

1035

positive or *Wolbachia* negative flies were homogenised, centrifuged and the supernatant

1036

was inoculated into PA14 cultures growing in 96-well plates. Absorbance was measured

1037

every 30 minutes. This assay showed that PA14 cultures exhibited a significantly longer

1038

lag phase and lower total yield during saturation phase when exposed to homogenised

1039

Wolbachia-positive flies.

1040

1041

1042

We also carried out an inhibition zone assay on PA14 grown on a solid medium.

Bacterial cultures and fly homogenate were both prepared as described above. When

growing the PA14 culture, 100 µl of a 500-fold dilution was used to plate on LB agar

plates to obtain a uniform PA14 lawn. After 30 minutes, 100 µl of fly homogenate was

put on the plate. Plates were incubated at 37°C for overnight. Clearance zone formed by

fly homogenate was measured using Image J and area values were analysed using

ANOVA with fly sex and Wolbachia status as factors.

1050
1051   **Fig S3. Inhibition zone assay**. Single sex groups of *Wolbachia* positive or *Wolbachia*

1052   negative flies were homogenised, centrifuged and the supernatant was spotted on a

1053   lawn of PA14 overnight. The figure shows the size of the zone where PA14 growth was

1054   inhibited. Data shown are means ± SEM.

1055

1056

1057 **Table S1 Fits of non-linear tolerance curves**

Sex	Model	AICc	AICc Weight	R-Square
Female	Logistic 4P	59.83289049	0.80614163	0.239893273
	Linear	62.68315369	0.193858371	0.110112664
Male	Logistic 4P	18.97942649	0.511289458	0.572646097
	Linear	19.0697575	0.488710542	0.516645982

1058

1059 To assess the form of the health/microbe relationship, we fit linear and non-linear 4-

1060 parameter logistic model separately to the time-matched survival/microbe load plots. In

1061 all cases, the 4 parameter logistic model – which is commonly use to asses dose-

1062 response curves - outperformed the linear fit.

1063