Natural history of Asian lineage Zika virus infection in macaques

Dawn M. Dudley[1*], Matthew T. Aliota[2*], Emma L. Mohr[3*], Andrea M. Weiler[4], Gabrielle Lehrer-

Brey[4], Kim L. Weisgrau[4], Mariel S. Mohns[1], Meghan E. Breitbach[1], Mustafa N. Rasheed[1],

Christina M. Newman[1], Dane D. Gellerup[4], Louise H. Moncla[1,2], Jennifer Post[4], Nancy Schultz-

Darken[4], Josh A. Eudailey[5], M. Anthony Moody[5], Sallie R. Permar[5], Shelby L. O'Connor[1], Eva

G. Rakasz[4], Heather A. Simmons[4], Saverio Capuano III[4], Thaddeus G. Golos[4,6], Jorge E.

Osorio[2], Thomas C. Friedrich[2,4], and David H. O'Connor[1,4]

[1]Department of Pathology and Laboratory Medicine, University of Wisconsin-Madison;

[2]Department of Pathobiological Sciences, University of Wisconsin-Madison;[3]Department of

Pediatrics, School of Medicine and Public Health, University of Wisconsin-Madison; [4]Wisconsin

National Primate Research Center, University of Wisconsin-Madison; [5]Department of Pediatrics

and Human Vaccine Institute, Duke University Medical Center; [6]Department of Comparative

Biosciences and Obstetrics and Gynecology, University of Wisconsin-Madison.

*These authors contributed equally to this work.

Infection with Asian lineage Zika virus has been associated with Guillain-Barré

syndrome and fetal microcephaly[1-4]. Here we show that rhesus macaques are susceptible

to infection by an Asian lineage Zika virus isolate that shares more than 99% nucleotide

identity with strains currently circulating in the Americas. Following subcutaneous

inoculation, Zika virus RNA was detected in plasma one-day post infection (dpi) in all

animals (N = 3). Plasma viral loads peaked above 1×10^6 viral RNA copies/mL in two of

three animals. Viral RNA was also present in saliva, urine, and cerebrospinal fluid (CSF),

consistent with case reports from infected humans. Zika virus RNA persisted in both

plasma and urine at low levels for more than two weeks. Infection was associated with transient increases in proliferating natural killer cells, CD8+ and CD4+ T cells, and plasmablasts, suggesting pathogen sensing by the immune system. These data establish that Asian lineage Zika virus infection in rhesus macaques provides a relevant animal model for human infection. Furthermore, because fetal development is well characterized in rhesus macaques, infections in pregnant macaques will enable important studies of fetal defects associated with Zika virus.

Zika virus (ZIKV) is a mosquito-borne flavivirus first identified in 1947 that is divided into African and Asian lineages based on phylogeny and geographical origin[5]. The primary vector for ZIKV is *Aedes aegypti*, the same species that transmits dengue (DENV) and chikungunya (CHIKV) viruses[6]. Little was known about ZIKV when increased incidence of fetal microcephaly and Guillain-Barré syndrome occurred coincident with epidemic spread of Asian lineage ZIKV in South America. The virologic and immunologic parameters of ZIKV infection remain unknown, including tissue tropism, duration of infection, and the potential for persistence. To date, ZIKV RNA has been detected in plasma, urine, semen, CSF, breast milk and saliva of patients[4,7-10], but the extent to which this represents replication-competent, transmissible virus is unknown. Infectious virus has been isolated from saliva and a small number of case reports suggest that ZIKV may be transmitted by blood transfusion or sexual intercourse, but the likelihood of such non-vectored transmission remains undetermined[11-13]. These and many other questions cannot be resolved quickly by studying case reports of infected individuals alone.

We therefore sought to develop a relevant animal model to understand ZIKV pathogenesis, and to evaluate candidate vaccines and therapeutics. For decades, macaques have been used in both infectious disease and obstetric research. In fact, what is now known as the African lineage of ZIKV was initially discovered in a sentinel rhesus macaque kept in the Zika forest of Uganda for yellow fever surveillance[14]. But does the currently circulating Asian

lineage virus, which shares ~90% nucleotide identity with the African lineage, also infect

macaques[5]?

To determine this, we inoculated three Indian-origin rhesus macaques (*Macaca mulatta*)

subcutaneously with 1 x 10^6, 1 x 10^5, or 1 x 10^4 PFU ZIKV derived from a French Polynesian

virus isolate (Zika virus/H.sapiens-tc/FRA/2013/FrenchPolynesia-01_v1c1) (Fig. 1a). This range

of inocula is based on previous work in related flaviviruses such as West Nile virus (WNV) and

DENV, where it was estimated that mosquitoes delivered 1 x 10^4- 1x 10^6 PFU of virus[15]. DENV

infection of macaques also provides the precedence for using subcutaneous inoculations[16].

Blood was sampled daily for 11 days post-infection (dpi) and every 3-4 days thereafter. Viral

RNA (vRNA) was quantified by qRT-PCR[17]. Plasma viremia was detected in all three animals at

1 dpi and was proportional to the inoculum dose, underscoring the importance of selecting a

ZIKV challenge dose in macaques that approximates the dose encountered during either vector-

borne or sexual transmission (Fig. 1b)[8,11]. Peak plasma viremia occurred between 2 and 6 dpi,

and ranged from 8.2 x 10^4 to 2.5 x 10^6 vRNA copies/mL. The estimated doubling time for the

plasma viremia of individual animals ranged from 4.8 to 10.2 hours and appeared to be

independent of the infecting dose. By 10 dpi, plasma viral loads were undetectable (<100

copies/mL) in all three animals, although intermittent low-level detection (<550 copies/mL)

continued in two of three animals through 17 dpi.

To detect signs of morbidity, animals were evaluated daily for evidence of disease,

injury, or psychological abnormalities (e.g., inappetence, dehydration, diarrhea, depression,

inactivity, trauma, self-injurious or stereotypical behavior). All animals exhibited mild to

moderate inappetence, which led to mild weight loss in two animals. Two animals (912116 and

393422) also developed a very mild rash around the inoculation site one dpi that persisted for 4-

5 days. No other abnormal clinical signs were noted (e.g., increased body temperature, joint

pain, lymphadenopathy, lethargy).

To examine potential within-host evolution, we deep sequenced both the challenge stock

and plasma virus from each animal at 4 dpi. Like all flaviviruses, the ZIKV genome encodes a

single, large polyprotein that is co- and post-translationally processed into three structural

(envelope E, membrane precursor prM, capsid C) and seven non-structural (NS1, NS2A, NS2B,

NS3, NS4A, NS4B, and NS5) proteins[18]. The ZIKV challenge stock consensus sequence

matched the Genbank sequence (KJ776791) of the parental virus, but there were 8 sites where

between 5-40% of sequences contained variants (Fig. 2a). Maintenance of some of the

challenge stock variants, as well as de novo variants not detected in the challenge stock, were

noted in all three animals at 4 dpi (Fig. 2b). Variants were concentrated in the NS1, NS2A and

NS2B regions for the both the challenge stock and from the animals.

To measure how selection may be shaping the viral populations during replication in

macaques, we calculated measures of synonymous (πS) and nonsynonymous (πN) nucleotide

diversity in each coding region of the genome relative to the inoculum for each animal at 4 dpi.

The ratio of πS to πN can indicate how selection is acting on a gene. While most coding regions

revealed signatures of purifying selection, the regions encoding NS1 and NS2B exhibited

elevated levels of nonsynonymous diversity in all three animals (Fig. 2c). NS1 exists in multiple

forms, including a membrane--bound mNS1 form and a secreted sNS1 form. These forms are

implicated in a number of flavivirus--host interactions, including immune protection, disease

pathogenesis, and evasion of the complement system[19,20]. Thus, diversification in NS1 genes

could be the result of selective pressures imposed by the macaque host. Further study is

required to determine whether NS1 and NS2B are consistently under positive selection in

macaques, and to identify candidate selective pressures. However, our data suggest that they

may be good targets for future studies of ZIKV immunity and host adaptation.

Animals were monitored using complete blood counts and blood chemistry panels daily

for 11 dpi and then every 3 to 4 days thereafter (Extended Data Fig. 1). Similar to other viral

infections, a transient lymphopenia was observed beginning 1 dpi, reaching a nadir 4 dpi

(Extended Data Fig. 1f). The animal that received the highest dose of inoculum developed

leucocytosis at 7 dpi that persisted through 28 dpi and was characterized by a mature

neutrophilia. Each animal also exhibited evidence of a mild regenerative anemia characterized

by varying degrees of polychromasia and anisocytosis, but whether this was secondary to the

viral infection or simply a result of frequent blood draws could not be determined. All animals

developed an elevated serum creatine kinase (CK), which correlated directly with inoculum dose

and peaked 5 dpi (Extended Data Fig. 1). Increased CK is often noted in animals after repeated

sedation and venipuncture. Two of three animals also exhibited an approximate two-fold

increase in AST and ALT, which peaked at 5 dpi. These findings are not uncommon in acute

viral infections in humans. Both cellular dyscrasias and elevated transaminases have been

described in human ZIKV case reports; myositis has not been reported[21,22].

We next characterized the immune response to infection in lymphocyte subsets by

staining peripheral blood mononuclear cells (PBMC) for multiple lineage and activation markers.

Proliferating (Ki-67+) NK cells, CD8+ T cells, and CD4+ T cells expanded significantly above

baseline levels by 5 dpi; the most rapid increase occurred in the animal challenged with the

highest ZIKV dose (Fig. 3). We also enumerated circulating plasmablasts, defined as CD3-/20-

/14-/16-/11c-/123- and CD80+/HLA-DR+ cells, on 3, 7, 11 and 14 dpi (Fig. 3)[23]. The peak

plasmablast expansion was inoculum-dose-dependent by day 7 and the decline in plasmablast

cells was inversely related to inoculum dose. Together these data provide an estimate that peak

activation of the adaptive immune response and antibody production may occur 5-7 dpi and

depend on inoculum dose, informing future work to characterize these responses.

In addition to blood, several other body fluids were tested for ZIKV by qRT-PCR

including urine, saliva and CSF. Viruria was detected starting at 4 or 5 dpi and continued up to

14 dpi, in urine passively collected from cage pans (Fig. 1b). Despite possible degradation of

virus between the time of urination and sample collection and processing, 1×10^3-1×10^4 vRNA

copies/mL urine were detected at multiple timepoints. Virus was also detected in oral swabs

collected from all three animals, peaking at approximately 1 x 10^4 vRNA copies per sample in 2

of 3 animals (Fig. 1b). Notably, as with urine, the kinetics of virus detection in saliva occurred

after peak plasma viremia. Lumbar punctures were performed at 4 and 14 dpi to quantify viral

RNA in CSF; vRNA was detectable at 4 dpi in two of three animals (Fig. 1b).

Altogether, our study showed the persistence of ZIKV in the plasma of rhesus macaques

for approximately 10 days, similar to other vector-borne flaviviruses that cause acute, typically

self-limiting infections in humans. However, the prolonged detection of vRNA in urine and saliva

after apparent clearance from the blood, detection of virus in the CSF, and occasional plasma

"blips" after initial clearance, suggest that ZIKV could establish a more persistent infection.

Given reports of neurologic complications arising from ZIKV infection, the central nervous

system may be a reservoir site for persistent virus. Future work in rhesus macaques will seek to

determine whether virus from fluids other than blood can infect new hosts and what threshold of

viral load at these sites may be required for transmission.

While Asian lineage viruses are associated with fetal microcephaly or Guillain-Barré

syndrome, African lineage ZIKV is not, suggesting that the lineages may have important

biological differences. To begin to investigate this possibility, we also infected three macaques

with the African lineage Zika virus/R.macaque-tc/UGA/1947/MR766-3329, using the same three

doses as the Asian lineage infections (Extended Data Fig. 2). Although all three animals

became infected, viremia was only detected in two animals 1 dpi. The African lineage virus

achieved peak plasma levels at least one order of magnitude lower those observed in animals

infected with the Asian-lineage virus (Extended Data Fig. 2). This difference in *in vivo* viral

kinetics could be relevant for both virus transmission and pathogenesis. We are currently

working to complete the characterization of these infections.

Our study establishes the rhesus macaque as a relevant translational model for ZIKV

infection and pathogenesis in humans. Rhesus macaques support replication of both Asian

lineage and African lineage ZIKV, providing the possibility to conduct detailed comparative

pathogenesis studies. The large immunological toolset available for rhesus macaques will

enable investigations of immunity and potential vaccines. Pregnancy, the maternal-fetal

interface, and fetal development have been described in detail in rhesus macaques, so this

model will also enable assessments of the impact of maternal ZIKV infection on the developing

fetus.

Methods

Study design

This was a proof-of-concept study was designed to establish the infectivity and viral dynamics of

Asian lineage ZIKV. Because nothing is known about ZIKV dosing in macaques, one male

rhesus macaque of Indian ancestry was challenged with each of the following ZIKV doses:

1×10^6, 1×10^5, and 1×10^4 PFU ZIKV from each lineage. All macaques utilized in the study were

free of Macacine herpesvirus 1, Simian Retrovirus Type D, Simian T-lymphotropic virus Type 1,

and Simian Immunodeficiency Virus. In a follow-up study, two additional males and one female

macaque were infected with the African lineage ZIKV at the same three challenge doses as the

Asian lineage. Primary data from the study is available at goo.gl/rmNCqf.

Care and use of macaques at the Wisconsin National Primate Research Center

All macaque monkeys used in this study were cared for by the staff at the Wisconsin National

Primate Research Center (WNPRC) in accordance with the regulations and guidelines outlined

in the Animal Welfare Act and the Guide for the Care and Use of Laboratory Animals and the

recommendations of the Weatherall report. This study was approved by the University of

Wisconsin-Madison Graduate School Institutional Animal Care and Use Committee (Animal

Care and Use Protocol Number G005401). For all procedures (i.e., physical examination, virus

inoculation, blood and swab collection), animals were anesthetized with an intramuscular dose

of ketamine (10 mL/kg). Blood samples were obtained using a vacutainer system or needle and

syringe from the femoral or saphenous vein.

Inoculations

ZIKV strain H/PF/2013 (GenBank: KJ776791), originally isolated from a 51-year-old female in

France returning from French Polynesia with a single round of amplification on Vero cells, was

obtained from Xavier de Lamballerie (European Virus Archive, Marseille France). ZIKV strain

Zika virus/R.macaque-tc/UGA/1947/MR766-3329 (GenBank:LC002520), originally isolated from

a sentinel Rhesus monkey on 20 April 1947 in Zika Forest, Entebbe, Uganda with 149 suckling

mouse brain passages and two rounds of amplification on Vero cells, was obtained from Brandy

Russell (CDC, Ft. Collins, CO). Virus stocks were prepared by inoculation onto a confluent

monolayer of C6/36 mosquito cells. A single harvest of virus with a titer of 1.26×10^6 PFU/mL

for the Asian lineage (equivalent to 1.43×10^9 vRNA copies/mL) and 5.9×10^6 PFU/mL for the

African lineage was used for all three challenges of each lineage. The stock was thawed, diluted

in PBS to the appropriate concentration for each challenge, and loaded into a 1 mL syringe that

was kept on ice until challenge. Animals were anesthetized as described above, and 1 mL of

inocula was administered subcutaneously over the cranial dorsum. At the conclusion of the

procedure, animals were closely monitored by veterinary and animal care staff for adverse

reactions and signs of disease.

Viral RNA isolation from plasma

Fresh plasma and PBMC were isolated from EDTA-treated whole blood by Ficoll density

centrifugation at 1860 rcf for 30min. The plasma layer was collected and centrifuged for an

additional 8 min at 670 rcf to remove residual cells. RNA was extracted from 300 µl of plasma

using the Viral Total Nucleic Acid Purification Kit (Promega, Madison, WI) on a Maxwell 16 MDx

instrument. The RNA was then quantified by quantitative RT-PCR.

Viral RNA isolation from urine

Urine was collected from a pan beneath the animal's cage. Urine was centrifuged for 5 min at

500 rcf to remove cells and other debris. RNA was isolated from 300 µl urine using the Viral

Total Nucleic Acid Purification Kit (Promega, Madison, WI) on a Maxwell 16 MDx instrument.

Viral RNA isolation from oral swabs

Oral swab samples were collected from infected animals while anesthetized by gently running a

sterile swab under the animal's tongue. Swabs were placed immediately into either RNAlater or

viral transport medium (tissue culture medium 199 supplemented with 0.5% FBS and 1%

antibiotic/antimycotic) for 60-90 minutes. Samples were vortexed vigorously, then centrifuged

for 5 min at 500 rcf before removing the swabs. Samples were stored at either -20°C (RNAlater

samples) or -80°C (viral transport medium) until processing. Prior to extraction, virus was

pelleted by centrifugation for 1 hour at 4°C at 14000 rpm. Supernatant was removed, leaving the

virus in 200 µl media. Viral RNA was extracted from these samples using the Qiamp MinElute

Virus Spin kit (Qiagen, Germantown, Maryland) with all optional washes. Viral load data from

oral swabs are expressed as vRNA copies/mL eluate.

Quantitative reverse transcription PCR (qRT-PCR)

Viral RNA isolated from plasma, urine, or oral swabs was quantified by qRT-PCR using the

primers and probe designed by Lanciotti et al.[17] or using primers with a slight modification to

those described by Lanciotti et al. to accommodate the African lineage virus. The sequence of

these modified primers are as follows: forward-5'CGYTGCCCAACACAAGG-3', reverse 5'-

CCACYAAYGTTCTTTTGCABACAT-3', and probe 5'-6fam-

AGCCTACCTTGAYAAGCARTCAGACACYCAA-BHQ1-3'. The RT-PCR was performed using

the SuperScript III Platinum one-step quantitative RT-PCR system (Invitrogen, Carlsbad, CA) on

the LightCycler 480 instrument (Roche Diagnostics, Indianapolis, IN). Primers and probe were

used at final concentrations of 600 nm and 100 nm respectively, along with 150 ng random

primers (Promega, Madison, WI). Cycling conditions were as follows: 37°C for 15 min, 50°C for

30 min and 95°C for 2 min, followed by 50 cycles of 95°C for 15 sec and 60°C for 1 min. Virus

concentration was determined by interpolation onto an internal standard curve composed of

seven 10-fold serial dilutions of a synthetic ZIKV RNA fragment based on the Asian lineage. A

comparison of the crossing point detected by qRT-PCR found from the standard template, Asian

lineage virus and African lineage virus when using either the universal primer set developed by

our group, or the primers developed by Lanciotti et al., suggest that the efficiency of each primer

set is the same for both lineages of ZIKV (Extended Data Fig. 3).

Viral quantification by plaque assay

All ZIKV screens and titrations for virus quantification were completed by plaque assay on Vero

cell cultures. Duplicate wells were infected with 0.1 mL aliquots from serial 10-fold dilutions in

growth media and virus was adsorbed for one hour. Following incubation, the inoculum was

removed, and monolayers were overlaid with 3 ml containing a 1:1 mixture of 1.2% oxoid agar

and 2X DMEM (Gibco, Carlsbad, CA) with 10% (vol/vol) FBS and 2% (vol/vol)

penicillin/streptomycin. Cells were incubated at 37°C in 5% CO_2 for four days for plaque

development. Cell monolayers then were stained with 3 mL of overlay containing a 1:1 mixture

of 1.2% oxoid agar and 2X DMEM with 2% (vol/vol) FBS, 2% (vol/vol) penicillin/streptomycin,

and 0.33% neutral red (Gibco). Cells were incubated overnight at 37°C and plaques were

counted.

Immunophenotyping

The amount of activated/proliferating NK cells were quantified using a modified version of our

protocol detailed step-by step in OMIP-28[24]. Briefly 0.1 mL of EDTA-anticoagulated whole blood

samples were incubated for 15 min at room temperature in the presence of a mastermix of

antibodies against CD45 (clone D058-1283, Brilliant Violet 786 conjugate), CD3 (clone SP34-2

Alexa Fluor 700 conjugate), CD8 (clone SK2, Brilliant Violet 510), NKG2A/C (clone Z199, PE-

Cy7 conjugate), CD16 (clone 3G8, Pacific Blue conjugate), CD69 (clone TP1.55.3, ECD

conjugate), HLA-DR (clone 1D11, Brilliant Violet 650 conjugate), CD4 (clone SK3, Brilliant Violet

711 conjugate), CCR7 (clone 150503, Fluorescein conjugate), CD28 (clone CD28.2, PE

conjugate), and CD95 (clone DX2, PE-Cy5 conjugate) antigens. All antibodies were obtained

from BD BioSciences, except the NKG2A/C-specific antibody, which was purchased from

Beckman Coulter and the CCR7 antibody that was purchased from R&D Systems. Red blood

cells were lysed using BD Pharm Lyse, after which they were washed twice in media and fixed

with 0.125 ml of 2% paraformaldehyde for 15 min. After an additional wash the cells were

permeabilized using Life Technology's Bulk Permeabilization Reagent. The cells were stained

for 15 min. against Ki-67 (clone B56, Alexa Fluor 647 conjugate) while the permeabilizer was

present. The cells were then washed twice in media and resuspended in 0.125 ml of 2%

paraformaldehyde until they were run on a BD LSRII Flow Cytometer. Flow data were analyzed

using Flowjo version 9.8.2.

Plasmablast detection

Peripheral blood mononuclear cells (PBMCs) isolated from three ZIKV-infected rhesus monkeys

at 3, 7, 11, and 14 dpi were stained with the following panel of fluorescently labeled antibodies

(Abs) specific for the following surface markers: CD20 FITC (L27), CD80 PE(L307.4), CD123

PE-Cy7(7G3), CD3 APC-Cy7 (SP34-2), IgG BV605(G18-145) (all from BD Biosciences, San

Jose, CA), CD14 AF700 (M5E2), CD11c BV421 (3.9), CD16 BV570 (3G8), CD27 BV650(O323)

(all from BioLegend, San Diego, CA), IgD AF647 (polyclonal)(Southern Biotech, Birmingham,

AL), and HLA-DR PE-TxRed (TÜ36) (Invitrogen, Carlsbad, CA). LIVE/DEAD Fixable Aqua Dead

Cell Stain Kit (Invitrogen, Carlsbad, CA) was used to discriminate live cells. Briefly, cells were

resuspended in 1X PBS/1%BSA and stained with the full panel of surface Abs for 30 min in the

dark at 4°C, washed once with 1X PBS, stained for 30 min with LIVE/DEAD Fixable Aqua Dead

Cell Stain Kit in the dark at 4°C, washed once with 1X PBS, washed again with 1X

PBS/1%BSA, and resuspended in 2% PFA Solution. Stained PBMCs were acquired on a LSRII

Flow Analyzer (BD Biosciences, San Jose, CA) and the data was analyzed using FlowJo

software v9.7.6 (TreeStar, Ashland, OR). Plasmablasts were defined similarly to the method

previously described[23] excluding lineage cells (CD14+, CD16+, CD3+, CD20+, CD11c+,

CD123+), and selecting CD80+ and HLA-DR+ cells (known to be expressed on rhesus

plasmablasts and their human counterpart[25]).

Estimation of plasma viremia doubling time

The doubling time of plasma viremia was estimated in R version 3.2.3 (The R Foundation for

Statistical Computing, http://www.R-project.org). For each animal, the slope of the linear portion

of the line (between 1 and 2 dpi for the animals treated with 1×10^6 and 1×10^5 PFU and between

1, 2, and 3 dpi for the animal treated with 1×10^4 PFU) was generated by plotting the log of the

plasma viral loads. The linear portion represents the exponential growth phase and has been

used to estimate doubling time in other systems[26]. The slopes were then used in the equation:

log(2)/slope. Each result was then multiplied by 24 hours to produce a simple estimate of

doubling time in hours.

CBC and blood chemistry panels

CBCs were performed on EDTA-anticoagulated whole blood samples on a Sysmex XS-1000i

automated hematology analyzer (Sysmex Corporation, Kobe, Japan). Blood smears were

prepared and stained with Wright-Giemsa stain (Wescor Aerospray Hematology Slide Stainer;

Wescor Inc, Logan, UT). Manual slide evaluations were performed on samples as appropriate

when laboratory-defined criteria were met (including the presence of increased total white blood

cell counts, increased monocyte, eosinophil, and basophil percentages, decreased hemoglobin,

hematocrit, and platelet values, and unreported automated differential values). Individuals

performing manual slide evaluations screened both white blood cells (WBC) and red blood cells

(RBC) for cellular maturity, toxic change, and morphologic abnormalities.

Whole blood was collected into serum separator tubes (Becton, Dickinson and

Company, Franklin Lakes, NJ) for blood chemistry analysis and processed per manufacturer's

instructions. Blood chemistry panels were performed on the serum using a Cobas 6000

analyzer (Roche Diagnostics, Risch-Rotkreuz, Switzerland). Results from CBC and blood

chemistry panels were reported with species, age, and sex-specific reference ranges.

Zika virus deep sequencing from plasma

A vial of the same ZIKV strain H/PF/2013 virus stock that infected macaques was deep

sequenced by preparing libraries of fragmented double-stranded cDNA using methods similar to

those previously described[27]. Briefly, the sample was centrifuged at 5000 rcf for 5 min. The

supernatant was then filtered through a 0.45-um filter. The Qiagen QiAmp Minelute viral RNA

isolation kit (omitting carrier RNA) was used to isolate vRNA. The eluted RNA was then treated

with DNAse I. Double stranded DNA was prepared with the Superscript double stranded cDNA

synthesis kit (Invitrogen) and priming with random hexamers. Agencourt Ampure XP beads

were used to purify double stranded DNA. The purified DNA was fragmented with the Nextera

XT kit (Illumina), tagged with Illumina-compatible primers, and then purified with Agencourt

Ampure XP beads. Purified libraries were then sequenced with 2 x 300 bp kits on an Illumina

MiSeq.

Zika virus samples isolated from animals were PCR amplified to create a series of five

overlapping amplicons that span the entire viral polyprotein, as described previously for other

RNA viruses[28]. Briefly, vRNA was isolated using the Qiagen QiAmp Minelute viral RNA isolation

kit, according to manufacturer's protocol (Qiagen). The following primer pairs were used to

generate viral cDNA amplicons using the Superscript III One-step RT-PCR kit (Invitrogen) and

including MgSO4:

ZFP-1F: GCGACAGTTCGAGTTTGAAGCG; ZFP-1R: ATCCAAAGTCCCAGGCTGTG;

ZFP-2F: AGATCCCGGCTGAAACACTG; ZFP-2R: CCCATGTGATGTCACCTGCT;

ZFP-3F: TACTCACAGCTGTTGGCCTG; ZFP-3R: CACCTCGGTTTGAGCACTCT;

ZFP-4F: TGTTTGGCTGGCCTATCAGG; ZFP-4R: CTGCGGATCCTTTCAATGCG;

ZFP-5F: TATGGGGGAGGACTGGTCAG; ZFP-5R: ACTAGCAGGCCTGACAACAC.

The following cycling conditions were used: 55°C for 30 min; 94°C 2 min; 35 cycles of the

following: 94°C 15 sec, 56°C 30 sec, and 68°C 3.5 min; 68°C 10 min. Viral cDNA amplicons

were size selected by agarose gel electrophoresis and then purified using the Qiagen MinElute

Gel extraction kit. Purified PCR products were pooled and then ~1ng of DNA was fragmented

using the Nextera XT kit (Illumina), tagged with Illumina-compatible primers, and then purified

with Agencourt Ampure XP beads. Purified libraries were then sequenced with 2 x 300bp kits on

an Illumina MiSeq.

Sequences were analyzed using a workflow (https://bitbucket.org/dhoconno/zequencer)

implemented in Geneious Pro (Biomatters, Ltd.). Briefly, duplicate reads were initially removed

with the BBMap v35.82 dedupe tool (https://sourceforge.net/projects/bbmap/) using a kmer seed

length of 31 bp and allowing zero edits or substitutions. Non-duplicate reads were quality

trimmed with BBDuk v35.82 to remove Truseq, Nextera, and PhiX adapters from both the left

and right ends of sequences. Regions with low quality sequences (average quality < 20) were

removed from both ends of remaining sequence reads, and only reads longer than than 150 bp

are retained. BBMap v35.82 was used to map these reads to the ZIKV strain H/PF/2013

Genbank reference sequence (KJ776791) in local alignment mode using the normal sensitivity

preset. The Geneious Pro 9.1.2 variant caller is used to detect and annotate variants. Variants

are called that fit the following conditions: present in >=5% of the mapped reads, have a

minimum p-value of 10e-60, and a minimum strand bias of 10e-5. Areas with low coverage

(<20x) are annotated.

Calculating nucleotide diversity

Synonymous (πS) and Nonsynonymous (πN) nucleotide diversity were calculated for each

coding region using Popoolation version 1.2.2[29]reference-based mappings were exported from

Geneious as bam files, which were converted to sorted pileup files using SAMtools[30]. πN and

πS estimates were calculated for each coding region in the genome using the Syn-

nonsyn-at-position.pl script. πS > πN suggests that purifying selection is acting to remove new

variation, πN > πS suggests that diversifying selection is favoring new mutations, and πN = πS

indicates no strong selection. To reduce the effects of uneven coverage across sites, pileup files

were subsampled such that every position was covered by exactly 1000x using the subsample-

pileup.pl script in Popoolation. Regions in which coverage was <1000x were excluded from the

analysis. πN and πS estimates were then calculated for each coding region in the genome

using the Syn-nonsyn-at-position.pl script. Corrections were disabled.

References

1.   Brasil, P. et al. Zika Virus Infection in Pregnant Women in Rio de Janeiro - Preliminary

Report. *N Engl J Med* (2016).

2.   Cao-Lormeau, V. M. et al. Guillain-Barré Syndrome outbreak associated with Zika virus

infection in French Polynesia: a case-control study. *Lancet* (2016).

3.   Calvet, G. et al. Detection and sequencing of Zika virus from amniotic fluid of fetuses with

microcephaly in Brazil: a case study. *Lancet Infect Dis* (2016).

4.   Carteaux, G. et al. Zika Virus Associated with Meningoencephalitis. *N Engl J Med* (2016).

5.   Haddow, A. D. et al. Genetic characterization of Zika virus strains: geographic expansion

of the Asian lineage. *PLoS Negl Trop Dis* **6**, e1477 (2012).

6.   Marchette, N. J., Garcia, R. & Rudnick, A. Isolation of Zika virus from Aedes aegypti

mosquitoes in Malaysia. *Am J Trop Med Hyg* **18**, 411-415 (1969).

7.   de M Campos, R. et al. Prolonged detection of Zika virus RNA in urine samples during the

ongoing Zika virus epidemic in Brazil. *J Clin Virol* **77**, 69-70 (2016).

8.   Mansuy, J. M. et al. Zika virus: high infectious viral load in semen, a new sexually

transmitted pathogen. *Lancet Infect Dis* (2016).

9.   Musso, D. et al. Detection of Zika virus in saliva. *J Clin Virol* **68**, 53-55 (2015).

10. Dupont-Rouzeyrol, M., Biron, A., O'Connor, O., Huguon, E. & Descloux, E. Infectious Zika

viral particles in breastmilk. *Lancet* (2016).

11. Musso, D. et al. Potential sexual transmission of Zika virus. *Emerg Infect Dis* **21**, 359-361

(2015).

12. Foy, B. D. et al. Probable non-vector-borne transmission of Zika virus, Colorado, USA.

Emerg Infect Dis **17**, 880-882 (2011).

13. Cunha, M. S. et al. First Complete Genome Sequence of Zika Virus (Flaviviridae,

Flavivirus) from an Autochthonous Transmission in Brazil. *Genome Announc* **4**, (2016).

14. DICK, G. W., KITCHEN, S. F. & HADDOW, A. J. Zika virus. I. Isolations and serological

specificity. *Trans R Soc Trop Med Hyg* **46**, 509-520 (1952).

15. Styer, L. M. et al. Mosquitoes inoculate high doses of West Nile virus as they probe and

feed on live hosts. *PLoS Pathog* **3**, 1262-1270 (2007).

16. Osorio, J. E. et al. Efficacy of a tetravalent chimeric dengue vaccine (DENVax) in

Cynomolgus macaques. *Am J Trop Med Hyg* **84**, 978-987 (2011).

17.   Lanciotti, R. S. et al. Genetic and serologic properties of Zika virus associated with an

408   epidemic, Yap State, Micronesia, 2007. *Emerg Infect Dis* **14**, 1232-1239 (2008).

409   18.   Lindenbach, B. D. & Rice, C. M. Molecular biology of flaviviruses. *Adv Virus Res* **59**, 23-61

410   (2003).

411   19.   Muller, D. A. & Young, P. R. The flavivirus NS1 protein: molecular and structural biology,

412   immunology, role in pathogenesis and application as a diagnostic biomarker. *Antiviral Res*

413   **98**, 192-208 (2013).

414   20.   Avirutnan, P. et al. Antagonism of the complement component C4 by flavivirus

415   nonstructural protein NS1. *J Exp Med* **207**, 793-806 (2010).

416   21.   Tappe, D. et al. Acute Zika virus infection after travel to Malaysian Borneo, September

417   2014. *Emerg Infect Dis* **21**, 911-913 (2015).

418   22.   Zammarchi, L. et al. Zika virus infections imported to Italy: clinical, immunological and

419   virological findings, and public health implications. *J Clin Virol* **63**, 32-35 (2015).

420   23.   Silveira, E. L. et al. Vaccine-induced plasmablast responses in rhesus macaques:

421   phenotypic characterization and a source for generating antigen-specific monoclonal

422   antibodies. *J Immunol Methods* **416**, 69-83 (2015).

424   **References from methods section only**

425   24.   Pomplun, N., Weisgrau, K. L., Evans, D. T. & Rakasz, E. G. OMIP-028: activation panel

426   for Rhesus macaque NK cell subsets. *Cytometry A* **87**, 890-893 (2015).

427   25.   Wrammert, J. et al. Rapid cloning of high-affinity human monoclonal antibodies against

428   influenza virus. *Nature* **453**, 667-671 (2008).

429   26.   Staprans, S. I. et al. Simian immunodeficiency virus disease course is predicted by the

430   extent of virus replication during primary infection. *J Virol* **73**, 4829-4839 (1999).

431   27.   Lauck, M. et al. Discovery and full genome characterization of two highly divergent simian

432   immunodeficiency viruses infecting black-and-white colobus monkeys (Colobus guereza)

in Kibale National Park, Uganda. *Retrovirology* **10**, 107 (2013).

28.   Gellerup, D. D. et al. Conditional Immune Escape during Chronic Simian

Immunodeficiency Virus Infection. *J Virol* **90**, 545-552 (2015).

29.   Kofler, R. et al. PoPoolation: a toolbox for population genetic analysis of next generation

sequencing data from pooled individuals. *PLoS One* **6**, e15925 (2011).

30.   Li, H. et al. The Sequence Alignment/Map format and SAMtools. *Bioinformatics* **25**, 2078-

2079 (2009).

Supplementary Information is linked to the online version of the paper at

www.nature.com/nature.

Acknowledgements We thank the Veterinary, animal care, Scientific Protocol Implementation,

and the Pathology staff at the Wisconsin National Primate Research Center (WNPRC) for their

contribution to this study. We thank the DHHS/PHS/NIH (R01AI116382-01A1) and

(DP2HD075699) for funding. We also thank the P51OD011106 awarded to the WNPRC,

Madison-Wisconsin. This research was conducted in part at a facility constructed with support

from Research Facilities Improvement Program grants RR15459-01 and RR020141-01. The

publication's contents are solely the responsibility of the authors and do not necessarily

represent the official views of NCRR or NIH.

Author contributions D.H.O., T.C.F., J.E.O., M.T.A., E.M., T.G.G. and D.M.D. designed the

experiments. D.H.O., D.M.D., M.T.A., E. M., T.C.F., and L.H.M. drafted the manuscript. M.T.A.,

and J.E.O. provided and prepared viral stocks and performed plaque assays. A.M.W., G.L-B.,

and T.C.F. developed and performed viral load assays. K.L.W. and E.G.R. performed

immunophenotyping assays. M.S.M., M.E.B., M.N.R., C.M.N., and D.M.D. coordinated and

processed macaque samples for distribution. D.D.G., S.L.O., and D.M.D. designed and

performed the sequencing experiments. L.H.M. and T.C.F. performed nucleotide diversity

calculations. J.P., N.S-D., H.A.S., S.C., coordinated the macaque infections, sampling, and

performed blood chemistries and CBC analysis. J.A.E., M.A.M., and S.R.P. performed the

plasmablast experiments.

Author information: Sequences are pending deposition into the Sequence Read Archive at

NCBI. The bioSample accession number is SAMN04589872. All data from these studies are

available at zika.labkey.com. Reprints and permissions information is available at

www.nature.com/reprints. The authors declare no competing interests. Correspondence and

requests for materials should be addressed to dhoconno@wisc.edu.

Figure Legends

Figure 1. Study design and ZIKV viral load from rhesus macaque fluids. a. ZIKV, strain

H/FP2013 from the European virus archive (Genbank: KJ776791) was inoculated onto C6/36

mosquito cells to generate the stock Zika virus/H.sapiens-tc/FRA/2013/FrenchPolynesia-

01_v1c1. Doses administered subcutaneously per animal are shown. b. Viral loads were

measured in plasma, urine, saliva and cerebral spinal fluid for each animal through 28 dpi. CSF

was collected on days 4 and 14. All animals were negative on day 14.

Figure 2. Genetic diversity of the challenge stock and virus isolated from animals. a. The

ZIKV challenge stock was deep sequenced from all three animals. Nucleotide sites where at

least 5% of sequences obtained from the challenge stock are different from the Genbank

sequence are shown. b. The plasma virus of each animal from 4 dpi was deep sequenced.

Non-synonymous variants present in the challenge stock relative to the Genbank consensus

sequence are shown in black in all three plots; synonymous variants are shown in grey. Variants

detected relative to the challenge stock in each animal are colored based on animal.

Non-synonymous variants are shown with dark colored markers, while synonymous variants are

shown in a lighter shade. Changes found in both the stock and the animals are connected with a

line. c. Deep sequence data from all 3 animals on day 4 post-infection were used to calculate

measures of synonymous (πS) and nonsynonymous (πN) nucleotide diversity for each coding

region in the genome. The $\pi N/\pi S$ ratio for each gene is shown with color corresponding to

animal. The capsid, propeptide and membrane coding regions in animal 393422 were excluded

from the analysis due to coverage <1000x.

Figure 3. Immune cell expansion following ZIKV infection. Expansion of Ki-67+ (activated)

CD4+, CD8+ and NK cells were measured daily for 11 days and then every 3 days for each

animal. Total number of activated cells per microliter of blood are plotted for each animal for

each cell type over time. The inset of each graph shows the total number of plasmablast cells

found in PBMCs collected at 3, 7, 11 and 14 dpi for each animal. Viral load is graphed over the

columns as a reference.

Extended Data Figure 1. Complete blood counts and chemistries for macaques infected

with ZIKV. Animals were infected with different doses of ZIKV as described in panel a.

Chemistries (AST in b; ALT in c; CK in d) were measured prior to infection and on 2, 5, and 11

dpi. Complete blood counts (WBC in e; % lymphocytes in f; RBC in g; platelets in h) were

measured prior to infection, daily for 11 days after infection and then every 3 days until 28 dpi.

Extended Data Figure 2. Infection with African lineage ZIKV. ZIKV strain MR766 (GenBank:

LC002520), originally isolated from a sentinel rhesus macaque on 20 April 1947 in Zika Forest,

Entebbe, Uganda with 149 suckling mouse brain passages and two rounds of amplification on

Vero cells, was obtained from Brandy Russell (CDC, Ft. Collins, CO). Virus stocks were

prepared by inoculation onto a confluent monolayer of C6/36 mosquito cells to generate the

stock Zika virus/R.macaque-tc/UGA/1947/MR766-3329. The viral loads for the first seven dpi

are shown relative to the viral loads of animals receiving the same dose of Asian lineage ZIKV.

Extended Data Figure 3. Efficiency of qRT-PCR on different templates with different

primers. The region of NS1 targeted by the Lanciotti et al. (2008) QRT-PCR assay is not

completely conserved between Asian and African lineage Zika viruses. We therefore introduced

ambiguous bases at sites in the primers and probe that varied between lineages. To determine

whether the modified oligos amplify and detect both African and Asian lineage viruses with

equal efficiency we compared tenfold dilution series of a Ugandan Zika virus, French Polynesian

Zika virus and our in-vitro transcribed RNA standard in this assay (which we term "universal")

with the same samples amplified with the sequence-specific (Lanciotti) primers and probe. Both

assays amplified samples with essentially equivalent efficiencies. (The slopes of the lines are

1.011, 1.012 and 1.020 respectively). Not surprisingly, the concentration of the Ugandan virus

is roughly 2-fold higher when measured with the universal assay than when measured with the

original Lanciotti assay. The computed concentration of the French Polynesian virus is the same

regardless of assay used to quantify the virus.

Figure 1. Study design and ZIKV viral load from rhesus macaque fluids. a. ZIKV, strain H/FP2013 from the European virus archive (Genbank: KJ776791) was inoculated onto C6/36 mosquito cells to generate the stock Zika virus/H.sapiens-tc/FRA/2013/FrenchPolynesia-01_v1c1. Doses administered subcutaneously per animal are shown. b. Viral loads were measured from plasma, urine, saliva or cerebral spinal fluid for each animal through 28 dpi. CSF was collected on days 4 and 14. All animals were negative on day 14.

a.

Site relative to KJ776791	Change	Amino Acid Change	Codon Change	Protein Effect	Variant Frequency
801	T -> C	F->L	UUC->CUC	Substitution	17.9%
2318	C -> T		CUC->CUT	None	18.7%
2863	T -> C	L->P	CUC->CCC	Substitution	7.7%
3223	A -> T	K->I	AAA->ATA	Substitution	10.5%
3475	T -> C	M->T	AUG->ACG	Substitution	7.9%
3740	C -> T		GUC->GUT	None	11.1%
3835	C -> T	A->V	GCG->GTG	Substitution	15.8%
4242	G -> C	E->Q	GAG->CAG	Substitution	40.9%

b.

c.

Figure 2. Genetic diversity of the challenge stock and virus isolated from animals. a. The ZIKV challenge stock was deep sequenced from all three animals. Nucleotide sites where at least 5% of sequences obtained from the challenge stock are different from the Genbank sequence are shown. b. The plasma virus of each animal from 4 dpi was deep sequenced. Nonsynonymous variants present in the challenge stock relative to the Genbank consensus sequence are shown in black in all three plots; synonymous variants are shown in grey.

Variants detected relative to the challenge stock in each animal are colored based on animal. Nonsynonymous variants are shown with dark colored markers, while synonymous variants are shown in a lighter shade. Changes found in both the stock and the animals are connected with a line. c. Deep sequence data from all 3 animals on day 4 postinfection were used to calculate measures of synonymous (πS) and nonsynonymous (πN) nucleotide diversity for each coding region in the genome. The $\pi N/\pi S$ ratio for each gene is shown with color corresponding to animal. The capsid, propeptide and membrane coding regions in animal 393422 were excluded from the analysis due to coverage <1000x.

Figure 3. Immune cell expansion following ZIKV infection. Expansion of Ki-67+ (activated) CD4+, CD8+ and NK cells were measured daily for 11 days and then every 3 days for each animal. Total numbers of activated cells per microliter of blood are plotted for each animal for each cell type over time. The inset of each graph shows the total number of plasmablast cells found in PBMCs collected at days 3, 7, 11 and 14 dpi for each animal. Viral load is graphed over the columns as a reference.

a.

826226
10⁶ PFU

393422
10⁵ PFU

912116
10⁴ PFU

b.

c.

d.

e.

f.

g.

h.

Extended Data Figure 1. Complete blood counts and chemistries for macaques infected with ZIKV. Animals were infected with different doses of ZIKV as described in panel a. Chemistries (AST in b; ALT in c; CK in d) were measured prior to infection and on 2, 5, and 11 dpi. Complete blood counts (WBC in e; % lymphocytes in f; RBC in g; platelets in h) were measured prior to infection, daily for 11 days after infection and then every 3 days until 28 dpi.

Extended Data Figure 2. Infection with African lineage ZIKV. ZIKV strain MR766 (GenBank: LC002520), originally isolated from a sentinel rhesus macaque on 20 April 1947 in Zika Forest, Entebbe, Uganda with 149 suckling mouse brain passages and two rounds of amplification on Vero cells, was obtained from Brandy Russell (CDC, Ft. Collins, CO). Virus stocks were prepared by inoculation onto a confluent monolayer of C6/36 mosquito cells to generate the stock Zika virus/R.macaque-tc/UGA/1947/MR766-3329. The viral loads for the first seven dpi are shown relative to the viral loads of animals receiving the same dose of Asian lineage ZIKV.

Extended Data Figure 3. Efficiency of qRT-PCR on different templates with different primers. The region of NS1 targeted by the Lanciotti et al. (2008) QRT-PCR assay is not completely conserved between Asian and African lineage Zika viruses. We therefore introduced ambiguous bases at sites in the primers and probe that varied between lineages. To determine whether the modified oligos amplify and detect both African and Asian lineage viruses with equal efficiency we compared tenfold dilution series of a Ugandan Zika virus, French Polynesian Zika virus and our in-vitro transcribed RNA standard in this assay (which we term "universal") with the same samples amplified with the sequence-specific (Lanciotti) primers and probe. Both assays amplified samples with essentially equivalent efficiencies. (The slopes of the lines are 1.011, 1.012 and 1.020 respectively). Not surprisingly, the concentration of the Ugandan virus is roughly 2-fold higher when measured with the universal assay than when measured with the original Lanciotti assay. The computed concentration of the French Polynesian virus is the same regardless of assay used to quantify the virus.