A rhesus macaque model of Asian lineage Zika virus infection

Dawn M. Dudley[1*], Matthew T. Aliota[2*], Emma Mohr[3*], Andrea M. Weiler[4], Gabrielle Lehrer-

Brey[4], Kim L. Weisgrau[4], Mariel S. Mohns[1], Meghan E. Breitbach[1], Mustafa N. Rasheed[1],

Christina M. Newman[1], Dane D. Gellerup[4], Louise H. Moncla[1,2], Jennifer Post[4], Nancy Schultz-

Darken[4], Michele L. Schotkzo[4], Jennifer M. Hayes[4], Josh A. Eudailey[5], M. Anthony Moody[5],

Sallie R. Permar[5], Shelby L. O'Connor[1], Eva G. Rakasz[4], Heather A. Simmons[4], Saverio

Capuano III[4], Thaddeus G. Golos[4,6], Jorge E. Osorio[2], Thomas C. Friedrich[2,4], and David H.

O'Connor[1,4]

[1]Department of Pathology and Laboratory Medicine, University of Wisconsin-Madison;

[2]Department of Pathobiological Sciences, University of Wisconsin-Madison;[3]Department of

Pediatrics, School of Medicine and Public Health, University of Wisconsin-Madison; [4]Wisconsin

National Primate Research Center, University of Wisconsin-Madison; [5]Department of Pediatrics

and Human Vaccine Institute, Duke University Medical Center; [6]Department of Comparative

Biosciences and Obstetrics and Gynecology, University of Wisconsin-Madison.

*These authors contributed equally to this work.

Infection with Asian lineage Zika virus has been associated with Guillain-Barré

syndrome and fetal abnormalities [1-4], but the mechanisms and risk factors for these

outcomes remain unknown. Here we show that rhesus macaques are susceptible to

infection by an Asian-lineage virus closely related to strains currently circulating in the

Americas. Following subcutaneous inoculation, Zika virus RNA was detected in plasma

one day post-infection (dpi) in all animals (N = 8, including 2 animals infected during the

first trimester of pregnancy). Plasma viral loads peaked above 1×10^5 viral RNA

copies/mL in seven of eight animals. Viral RNA was also present in saliva, urine, and

cerebrospinal fluid (CSF), consistent with case reports from infected humans. Viral RNA

was cleared from plasma and urine by 21 dpi in non-pregnant animals. In contrast, both

pregnant animals remained viremic longer, up to 57 days. In all animals, infection was

associated with transient increases in proliferating natural killer cells, CD8+ T cells, CD4+

T cells, and plasmablasts. Neutralizing antibodies were detected in all animals by 21 dpi.

Rechallenge of three non-pregnant animals with the Asian-lineage Zika virus 10 weeks

after the initial challenge resulted in no detectable virus replication, suggesting that

primary Zika virus infection elicits protective immunity against homologous virus strains.

These data establish that Asian-lineage Zika virus infection of rhesus macaques provides

a relevant animal model for studying pathogenesis in pregnant and non-pregnant

individuals and evaluating potential interventions against human infection, including

during pregnancy.

Zika virus (ZIKV) is a mosquito-borne flavivirus first identified in 1947[5]. Little was known

about ZIKV when fetal abnormalities and Guillain-Barré syndrome were reported coincident with

epidemic spread of Asian-lineage ZIKV in South America[6,7,1]. Animal models are essential for

quickly understanding ZIKV transmission and pathogenesis, as well as for evaluating candidate

vaccines and therapeutics. ZIKV infects immunocompromised mice[8] but such mice do not mimic

key attributes of human infection and fetal development.

In contrast, immunocompetent macaque monkeys are widely used in both infectious

disease and obstetric research. To determine whether a phyisiologically relevant dose and route

of Asian lineage ZIKV infects immunocompetent pregnant and non-pregnant macaques, we

inoculated eight Indian-origin rhesus macaques (*Macaca mulatta*) subcutaneously with ZIKV

derived from a French Polynesian virus isolate (Zika virus/H.sapiens-

tc/FRA/2013/FrenchPolynesia-01_v1c1). The eight animals were divided into three cohorts as

shown in Extended Data Fig. 1.

To define the minimal dosage necessary to establish infection, two macaques per group

were infected with 1×10^6, 1×10^5, or 1×10^4 PFU ZIKV (cohorts 1 and 2) (Fig. 1a). This dose

range of inocula is based on previous work in related flaviviruses such as West Nile virus (WNV)

and DENV, where it was estimated that mosquitoes delivered 1×10^4- 1×10^6 PFU of virus[9,10].

This is also the range found in mosquito saliva in a recent publication specifically for Brazilian

Zika virus [11]. None of the animals developed significant clinical disease (see Extended Data Fig.

2 and SI Data)

Blood was sampled daily for 10-11 days post-infection (dpi) and every 3-7 days

thereafter. Viral RNA (vRNA) was quantified by qRT-PCR from plasma[12] and was detected in all

six animals at 1 dpi (Fig. 1b). Peak plasma viremia occurred between 2 and 6 dpi, and ranged

from 8.2×10^4 to 2.5×10^6 vRNA copies/mL. These results resemble findings in humans in

Colombia where the mean serum viral load was 2.6×10^5 copies/ml ± 10 copies/ml in acutely

infected individuals (n=10, range=537 - 6.9×10^5 copies/ml) (manuscript submitted[13]). Infectious

titers, measured from serum in cohort 2 animals, were 500-1000-fold less than copies of vRNA

detected from plasma at the same time points (Fig. 1c). Copies of vRNA detected in the serum

and plasma were very similar as shown in Extended Data Fig. 3. The estimated doubling time

for plasma viremia averaged 7.7 hours (range = 4.8-10.2 hours) and was independent of the

infecting dose and sex of the macaque. By 10 dpi, plasma viral loads were undetectable (<100

vRNA copies/mL) in all six animals, although intermittent low-level detection (<550 vRNA

copies/mL) continued sporadically through 17 dpi. Thereafter viral RNA remained undetectable

in all fluids throughout follow-up (longest follow-up 70 dpi; Fig. 1b insets).

We also measured ZIKV vRNA by qRT-PCR in other body fluids including urine, saliva,

CSF and vaginal fluid. Viruria was detected starting at 2-5 dpi and as late as 17 dpi, in urine

passively collected from cage pans (Fig. 1b). Despite possible degradation of virus between the

time of urination and sample collection and processing, 1×10^3-1×10^4 vRNA copies/mL urine

were detected at multiple timepoints. Virus was also detected in oral swabs collected from all six

animals, peaking at over 1 x 10^3 vRNA copies per sample in 3 of 6 animals (Fig. 1b). Notably,

as with urine, the kinetics of virus detection in saliva occurred after peak plasma viremia.

Cisterna magna punctures were performed at 4 and 14 dpi to quantify viral RNA in CSF; vRNA

was detectable at 4 dpi in 3 out of 5 animals from which CSF could be obtained (Fig. 1d).

Vaginal swabs collected from the 3 female animals in cohort 2 had detectable vRNA starting at

1 and/or 7 dpi, but were undetectable at 14, 21, and 28 dpi (Fig. 1e).

We next characterized the immune response to infection by staining peripheral blood

mononuclear cells (PBMC) for multiple lineage and activation markers. Proliferating (Ki-67+) NK

cells, CD8+ T cells, and CD4+ T cells expanded significantly above baseline levels by 6 dpi

(Fig. 2a,b). NK and CD8+ T cell expansion increased as plasma vRNA loads decreased starting

at 6 dpi. We also enumerated circulating plasmablasts, defined as CD3-/20-/14-/16-/11c-/123-

and CD80+/HLA-DR+ cells, on 0, 3, 7, 11 and 14 dpi (Fig. 2c)[14]. The peak plasmablast

expansion occurred between 7 and 10 dpi in 5/6 animals. Serum neutralizing antibody

responses were also measured by plaque reduction neutralization tests (PRNT$_{90}$). All animals

exhibited high neutralizing antibody (nAb) titers as early as 14 dpi (Fig. 2d), the earliest time

point tested. Cohort 1 animals were tested at 64 dpi and cohort 2 animals were tested at 14 and

28 dpi. Together these data suggest that peak activation of the adaptive immune response and

antibody production occurs 5-7 dpi and may both be important to control viral replication as

evidenced by reducing vRNA loads in the plasma at these time points.

To determine whether activation of T cells correlated with the appearance of Zika virus-

specific responses, we performed interferon-gamma ELISPOT on PBMCs collected at 4, 10 and

14 dpi for cohort 2 animals. Cells were stimulated with pools of 15mer peptides collectively

representing the amino acid sequence of the Asian-lineage NS5 protein (GenBank: KU321639).

We detected specific IFN-gamma secretion in response to 12 of 16 peptide pools in at least one

animal (Extended Data Fig. 4a). Overall, this data supports that there are ZIKV-specific T cell

responses in all animals tested.

To determine whether the immune responses that we detected following primary

challenge were protective against homotypic rechallenge, we rechallenged the three animals in

cohort 1 ten weeks after primary infection with 1×10^4 PFU of a homologous virus (Fig. 1b inset

and Extended Data Fig. 1). Plasma, urine and saliva vRNA loads remain negative to at least 9

dpi (as of 5/4/2016), indicating complete protection against ZIKV re-infection.

We also challenged two time-mated rhesus macaques at approximately gestation day 31

and 38 (mid-first trimester) with 1×10^4 PFU of ZIKV (cohort 3; see Extended Data Fig. 1 and

Fig. 3a). Both animals were viremic by 1 dpi and exhibited peak plasma viral loads of $> 4 \times 10^5$

vRNA copies/ml by 3 or 6 dpi (Fig. 3b). Infectious virus was also quantitated by plaque assay

from the serum of 660875 (Extended Data Fig. 5). In contrast to their non-pregnant

counterparts, both animals maintained persistent plasma viremia (vRNA copies/ml) to 57+ and

29 dpi (Fig. 3b). This is similar to a case described by Driggers et. al., where a pregnant mother

had persistent ZIKV vRNA detected from 35 to 70 dpi that did not resolve until termination of

pregnancy[7]. The fetus was found to have 2×10^8 copies/ml of virus in brain tissue and it is

speculated that the fetus may have been the source of the prolonged maternal plasma viremia.

We will continue to monitor these pregnant animals for vRNA in the blood and amniotic fluid and

will determine the infection status of the fetus upon termination of the pregnancy either at full

term or earlier if necessary for the health and safety of the mother. Amniocentesis using

ultrasound guidance was performed at 43 dpi for 827577 and 36 dpi for 660875 and both were

negative for ZIKV RNA.

Both animals generated similar activation of NK, CD8+ T cell and CD4+ T cell responses

above baseline as non-pregnant animals (Fig. 3c). Expansion of plasmablast cells was also

observed by 10-21 dpi with one animal expanding more than, and one animal expanding less

than, the average non-pregnant animal (Fig. 3d). Neutralizing antibodies were detected by 21

dpi for 827577 and 10 dpi for 660875 and were similar to the cohort 2 non-pregnant animals at

28 dpi (Fig. 3e). One pregnant animal exhibited an increase in CK above RI and above levels

expected with repeated ketamine sedation and blood collection. The same pregnant animal also

developed persistent regenerative anemia characterized by circulating nucleated erythrocytes.

Updates to the cohort 3 experiments are available in real-time at goo.gl/rmNCqf.

Altogether, our study shows the persistence of ZIKV RNA in the plasma of rhesus

macaques for approximately 10 days, similar to other vector-borne flaviviruses that cause acute,

typically self-limiting infections in humans. This work also shows that natural ZIKV infection

elicits a robust immune response including ZIKV-specific T cell response and neutralizing

antibody responses that confers protection against reinfection. However, the prolonged

detection of vRNA in urine and saliva after apparent clearance from the blood, detection of virus

in the CSF, and occasional plasma "blips" after initial clearance, suggest that ZIKV may persist

longer, at low levels, in certain tissues. Future work in rhesus macaques will seek to determine if

and where these reservoirs may exist and whether they seed virus into fluids that might allow for

human-to-human transmission.

Our study establishes immunocompetent rhesus macaques infected with physiologically

relevant ZIKV as a relevant translational model for infection and pathogenesis. The large

immunological toolset available for rhesus macaques will enable investigations of immunity and

potential vaccines. Pregnancy, the maternal-fetal interface, and fetal development have been

described in detail in rhesus macaques, so this model will also enable assessments of the

impact of maternal ZIKV infection on the developing fetus. We have established persistent

viremia in pregnant macaques despite activation of NK cells and T cells as well as development

of a neutralizing antibody response. We continue to follow these pregnant animals and will

establish whether fetal infection and/or abnormalities have occurred through serial ultrasound

assessments of the fetus and placenta as well as tissue analysis at pregnancy termination.

Methods

Study design

This was a proof-of-concept study designed to establish the infectivity and viral dynamics of

Asian lineage ZIKV. Because nothing is known about ZIKV dosing in macaques, one male and

one female rhesus macaque of Indian ancestry were each challenged with the following ZIKV

doses: 1×10^6, 1×10^5, and 1×10^4 PFU ZIKV. Two pregnant macaques at 31 and 38 days of

gestation were infected with 1×10^4 PFU ZIKV. We selected 2 animals per inoculum dose and 2

pregnant animals as a minimum number of animals for this pilot study to provide proof-of-

concept and design larger studies necessary to place statistical significance on the findings. All

macaques utilized in the study were free of Macacine herpesvirus 1, Simian Retrovirus Type D,

Simian T-lymphotropic virus Type 1, and Simian Immunodeficiency Virus. Primary data from the

study is available at goo.gl/rmNCqf.

Care and use of macaques at the Wisconsin National Primate Research Center

All macaque monkeys used in this study were cared for by the staff at the Wisconsin National

Primate Research Center (WNPRC) in accordance with the regulations and guidelines outlined

in the Animal Welfare Act and the Guide for the Care and Use of Laboratory Animals and the

recommendations of the Weatherall report. This study was approved by the University of

Wisconsin-Madison Graduate School Institutional Animal Care and Use Committee (Animal

Care and Use Protocol Number G005401). For all procedures (i.e., physical examinations, virus

inoculations, ultrasound examinations, blood and swab collection), animals were anesthetized

with an intramuscular dose of ketamine (10 mL/kg). Blood samples were obtained using a

vacutainer system or needle and syringe from the femoral or saphenous vein.

Inoculations

ZIKV strain H/PF/2013 (GenBank: KJ776791), originally isolated from a 51-year-old female in

France returning from French Polynesia with a single round of amplification on Vero cells, was

obtained from Xavier de Lamballerie (European Virus Archive, Marseille France). We deep

sequenced the challenge stock to verify the expected origin. The ZIKV challenge stock

consensus sequence matched the Genbank sequence (KJ776791) of the parental virus, but

there were 8 sites where between 5-40% of sequences contained variants that appear to be

authentic (6/8 were non-synonymous changes) (Extended Data Fig. 6). Sequences have been

deposited in the Sequence Read Archive (SRA) under accession number SRP072852 .

186
Virus stocks were prepared by inoculation onto a confluent monolayer of C6/36 mosquito

cells. These cell lines were obtained from ATCC, were not further authenticated and were not

specifically tested for mycoplasma. A single harvest of virus with a titer of 1.26 x 10^6 PFU/mL

(equivalent to 1.43 x 10^9 vRNA copies/mL) was used for all 8 challenges. The stock was

thawed, diluted in PBS to the appropriate concentration for each challenge, and loaded into a 1

mL syringe that was kept on ice until challenge. Animals were anesthetized as described above,

and 1 mL of inocula was administered subcutaneously over the cranial dorsum. Post-

inoculation, animals were closely monitored by veterinary and animal care staff for adverse

reactions and signs of disease.

Viral RNA isolation from plasma

Fresh plasma and PBMC were isolated from EDTA-treated whole blood by Ficoll density

centrifugation at 1860 rcf for 30min. The plasma layer was collected and centrifuged for an

additional 8 min at 670 rcf to remove residual cells. RNA was extracted from 300 µl of plasma

using the Viral Total Nucleic Acid Purification Kit (Promega, Madison, WI) on a Maxwell 16 MDx

instrument. The RNA was then quantified by quantitative RT-PCR.

Viral RNA isolation from urine

Urine was collected from a pan beneath the animal's cage. Urine was centrifuged for 5 min at

500 rcf to remove cells and other debris. RNA was isolated from 300 µl urine using the Viral

Total Nucleic Acid Purification Kit (Promega, Madison, WI) on a Maxwell 16 MDx instrument.

Viral RNA isolation from oral swabs

Oral swab samples were collected from infected animals while anesthetized by gently running a

sterile swab under the animal's tongue. Swabs were placed immediately into either RNAlater or

viral transport medium (tissue culture medium 199 supplemented with 0.5% FBS and 1%

antibiotic/antimycotic) for 60-90 minutes. Samples were vortexed vigorously, then centrifuged

for 5 min at 500 rcf before removing the swabs. Samples were stored at either -20°C (RNAlater

samples) or -80°C (viral transport medium) until processing. Prior to extraction, virus was

pelleted by centrifugation for 1 hour at 4°C at 14000 rpm. Supernatant was removed, leaving the

virus in 200 µl media. Viral RNA was extracted from these samples using the Qiamp MinElute

Virus Spin kit (Qiagen, Germantown, Maryland) with all optional washes. Viral load data from

oral swabs are expressed as vRNA copies/mL eluate.

Quantitative reverse transcription PCR (qRT-PCR)

Viral RNA isolated from plasma, urine, or oral swabs was quantified by qRT-PCR using the

primers and probe designed by Lanciotti et al. [12].The RT-PCR was performed using the

SuperScript III Platinum one-step quantitative RT-PCR system (Invitrogen, Carlsbad, CA) on the

LightCycler 480 instrument (Roche Diagnostics, Indianapolis, IN). Primers and probe were used

at final concentrations of 600 nm and 100 nm respectively, along with 150 ng random primers

(Promega, Madison, WI). Cycling conditions were as follows: 37°C for 15 min, 50°C for 30 min

and 95°C for 2 min, followed by 50 cycles of 95°C for 15 sec and 60°C for 1 min. Virus

concentration was determined by interpolation onto an internal standard curve composed of

seven 10-fold serial dilutions of a synthetic ZIKV RNA fragment based on the Asian-lineage

(ZIKV strain H/PF/2013).

Viral quantification by plaque assay

Titrations for replication competent virus quantification of the challenge stock as well as from

serum collected at multiple time points from cohort 2 animals were completed by plaque assay

on Vero cell cultures. Vero cells were obtained from ATCC, were not further authenticated and

were not specifically tested for mycoplasma. Duplicate wells were infected with 0.1 mL aliquots

from serial 10-fold dilutions in growth media and virus was adsorbed for one hour. Following

incubation, the inoculum was removed, and monolayers were overlaid with 3 ml containing a 1:1

mixture of 1.2% oxoid agar and 2X DMEM (Gibco, Carlsbad, CA) with 10% (vol/vol) FBS and

2% (vol/vol) penicillin/streptomycin. Cells were incubated at 37°C in 5% CO_2 for four days for

plaque development. Cell monolayers then were stained with 3 mL of overlay containing a 1:1

mixture of 1.2% oxoid agar and 2X DMEM with 2% (vol/vol) FBS, 2% (vol/vol)

penicillin/streptomycin, and 0.33% neutral red (Gibco). Cells were incubated overnight at 37°C

and plaques were counted. Titers of virus detected from the serum of cohort 2 animals were

compared to plasma and serum viral load assays. For both the challenge stock and the virus

isolated from macaque serum, the level of infectious virus detected by plaque assay was ~500-

1000-fold less than the number of viral RNA particles detected by qRT-PCR in either the plasma

or serum. This was true throughout the duration of viremia where plaque assay titers were

detectable.

Plaque reduction neutralization test (PRNT$_{90}$)

Macaque serum samples were screened for ZIKV neutralizing antibody utilizing a plaque

reduction neutralization test (PRNT). Endpoint titrations of reactive sera, utilizing a 90% cutoff

(PRNT90) were performed as described[15] against ZIKV strain H/PF/2013.

Immunophenotyping

The amount of activated/proliferating NK cells were quantified using a modified version of our

protocol detailed step-by step in OMIP-28[16]. Briefly 0.1 mL of EDTA-anticoagulated whole blood

samples were incubated for 15 min at room temperature in the presence of a mastermix of

antibodies against CD45 (clone D058-1283, Brilliant Violet 786 conjugate), CD3 (clone SP34-2

Alexa Fluor 700 conjugate), CD8 (clone SK2, Brilliant Violet 510), NKG2A/C (clone Z199, PE-

Cy7 conjugate), CD16 (clone 3G8, Pacific Blue conjugate), CD69 (clone TP1.55.3, ECD

conjugate), HLA-DR (clone 1D11, Brilliant Violet 650 conjugate), CD4 (clone SK3, Brilliant Violet

711 conjugate), CCR7 (clone 150503, Fluorescein conjugate), CD28 (clone CD28.2, PE

conjugate), and CD95 (clone DX2, PE-Cy5 conjugate) antigens. All antibodies were obtained

from BD BioSciences, except the NKG2A/C-specific antibody, which was purchased from

Beckman Coulter, and the CCR7 antibody that was purchased from R&D Systems. Red blood

cells were lysed using BD Pharm Lyse, after which they were washed twice in media and fixed

with 0.125 ml of 2% paraformaldehyde for 15 min. After an additional wash the cells were

permeabilized using Life Technology's Bulk Permeabilization Reagent. The cells were stained

for 15 min. with Ki-67 (clone B56, Alexa Fluor 647 conjugate) while the permeabilizer was

present. The cells were then washed twice in media and resuspended in 0.125 ml of 2%

paraformaldehyde until they were run on a BD LSRII Flow Cytometer. Flow data were analyzed

using Flowjo version 9.8.2.

Interferon-gamma ELISPOT assay

Peripheral blood mononuclear cells (PBMCs) were isolated from EDTA-treated whole blood by

using Ficoll-Paque Plus (GE Health Sciences) density centrifugation. Enzyme-linked

immunosorbent spot (ELISPOT) assays were conducted according to the manufacturer's

protocol. Briefly, 1 x 10^5 cells in 100ul of R10 medium were added to pre-coated monkey

gamma interferon (IFNγ) ELISpot-PLUS plates (Mabtech Inc., Mariemont, OH) with peptide at a

final concentration of 1uM. Full proteome peptides derived from the ZIKV NS5 sequence

(GenBank: KU321639.1) used in this study were synthesized by GenScript (Piscataway, NJ).

Pools were created using 10 overlapping 15mer peptides, each at a working concentration of

1mM. Concanavalin A (10uM) was used as a positive control. Assays of all samples were

repeated in duplicate or triplicate. Cells alone in the absence of stimulant were used as a

negative control. Wells were imaged by using an AID ELISPOT reader, and spots were counted

by using an automated program with parameters including size, intensity, and gradient. The limit

of detection was set at 100 spot-forming cells per million PBMCs.

Plasmablast detection

Peripheral blood mononuclear cells (PBMCs) isolated from three ZIKV-infected rhesus monkeys

at 3, 7, 11, and 14 dpi were stained with the following panel of fluorescently labeled antibodies

(Abs) specific for the following surface markers: CD20 FITC (L27), CD80 PE(L307.4), CD123

PE-Cy7(7G3), CD3 APC-Cy7 (SP34-2), IgG BV605(G18-145) (all from BD Biosciences, San

Jose, CA), CD14 AF700 (M5E2), CD11c BV421 (3.9), CD16 BV570 (3G8), CD27 BV650(O323)

(all from BioLegend, San Diego, CA), IgD AF647 (polyclonal)(Southern Biotech, Birmingham,

AL), and HLA-DR PE-TxRed (TÜ36) (Invitrogen, Carlsbad, CA). LIVE/DEAD Fixable Aqua Dead

Cell Stain Kit (Invitrogen, Carlsbad, CA) was used to discriminate live cells. Briefly, cells were

resuspended in 1X PBS/1%BSA and stained with the full panel of surface Abs for 30 min in the

dark at 4°C, washed once with 1X PBS, stained for 30 min with LIVE/DEAD Fixable Aqua Dead

Cell Stain Kit in the dark at 4°C, washed once with 1X PBS, washed again with 1X

PBS/1%BSA, and resuspended in 2% PFA Solution. Stained PBMCs were acquired on a LSRII

Flow Analyzer (BD Biosciences, San Jose, CA) and the data was analyzed using FlowJo

software v9.7.6 (TreeStar, Ashland, OR). Plasmablasts were defined similarly to the method

previously described[14] excluding lineage cells (CD14+, CD16+, CD3+, CD20+, CD11c+,

CD123+), and selecting CD80+ and HLA-DR+ cells (known to be expressed on rhesus

plasmablasts and their human counterpart[17]).

Estimation of plasma viremia doubling time

The doubling time of plasma viremia was estimated in R version 3.2.3 (The R Foundation for

Statistical Computing, http://www.R-project.org). For each animal, the slope of the linear portion

of the line (between 1 and 2 dpi for the animals treated with $1x10^6$ and $1x10^5$ PFU and between

1, 2, and 3 dpi for the animal treated with $1x10^4$ PFU) was generated by plotting the log of the

plasma viral loads. The linear portion represents the exponential growth phase and has been

used to estimate doubling time in other systems[18]. The slopes were then used in the equation:

log(2)/slope. Each result was then multiplied by 24 hours to produce a simple estimate of

doubling time in hours.

CBC and blood chemistry panels

CBCs were performed on EDTA-anticoagulated whole blood samples on a Sysmex XS-1000i

automated hematology analyzer (Sysmex Corporation, Kobe, Japan). Blood smears were

prepared and stained with Wright-Giemsa stain (Wescor Aerospray Hematology Slide Stainer;

Wescor Inc, Logan, UT). Manual slide evaluations were performed on samples as appropriate

when laboratory-defined criteria were met (including the presence of increased total white blood

cell counts, increased monocyte, eosinophil, and basophil percentages, decreased hemoglobin,

hematocrit, and platelet values, and unreported automated differential values). Individuals

performing manual slide evaluations screened both white blood cells (WBC) and red blood cells

(RBC) for cellular maturity, toxic change, and morphologic abnormalities.

Whole blood was collected into serum separator tubes (Becton, Dickinson and

Company, Franklin Lakes, NJ) for blood chemistry analysis and processed per manufacturer's

instructions. Blood chemistry panels were performed on the serum using a Cobas 6000

analyzer (Roche Diagnostics, Risch-Rotkreuz, Switzerland). Results from CBC and blood

chemistry panels were reported with species, age, and sex-specific reference ranges.

Zika virus deep sequencing of the challenge stock

A vial of the same ZIKV strain H/PF/2013 virus stock that infected macaques was deep

sequenced by preparing libraries of fragmented double-stranded cDNA using methods similar to

those previously described[19]. Briefly, the sample was centrifuged at 5000 rcf for 5 min. The

supernatant was then filtered through a 0.45-μm filter. The Qiagen QiAmp Minelute viral RNA

isolation kit (omitting carrier RNA) was used to isolate vRNA. The eluted RNA was then treated

with DNAse I. Double stranded DNA was prepared with the Superscript double stranded cDNA

synthesis kit (Invitrogen) and priming with random hexamers. Agencourt Ampure XP beads

were used to purify double stranded DNA. The purified DNA was fragmented with the Nextera

XT kit (Illumina), tagged with Illumina-compatible primers, and then purified with Agencourt

Ampure XP beads. Purified libraries were then sequenced with 2 x 300 bp kits on an Illumina

MiSeq. Of note, challenge stock viral loads were 1.43×10^9 vRNA copies/ml. This results in an

input of 7.15×10^8 RNA copies into the sequencing reactions. This far exceeds the average

depth of coverage of 11,877 (+/- 4658) sequences per nucleotide site indicating little resampling

effects in our data analysis.

References

1.   Brasil, P. et al. Zika Virus Infection in Pregnant Women in Rio de Janeiro - Preliminary

354   Report. *N Engl J Med* (2016).

2. Cao-Lormeau, V. M. et al. Guillain-Barré Syndrome outbreak associated with Zika virus

infection in French Polynesia: a case-control study. *Lancet* (2016).

3. Calvet, G. et al. Detection and sequencing of Zika virus from amniotic fluid of fetuses with

microcephaly in Brazil: a case study. *Lancet Infect Dis* (2016).

4. Carteaux, G. et al. Zika Virus Associated with Meningoencephalitis. *N Engl J Med* (2016).

5. Musso, D. & Gubler, D. J. Zika Virus. *Clin Microbiol Rev* **29**, 487-524 (2016).

6. Sarno, M. et al. Zika Virus Infection and Stillbirths: A Case of Hydrops Fetalis,

Hydranencephaly and Fetal Demise. *PLoS Negl Trop Dis* **10**, e0004517 (2016).

7. Driggers, R. W. et al. Zika Virus Infection with Prolonged Maternal Viremia and Fetal Brain

Abnormalities. *N Engl J Med* (2016).

8. Rossi, S. L. et al. Characterization of a Novel Murine Model to Study Zika Virus. *Am J Trop*

Med Hyg (2016).

9. Styer, L. M. et al. Mosquitoes inoculate high doses of West Nile virus as they probe and

feed on live hosts. *PLoS Pathog* **3**, 1262-1270 (2007).

10. Cox, J., Mota, J., Sukupolvi-Petty, S., Diamond, M. S. & Rico-Hesse, R. Mosquito bite

delivery of dengue virus enhances immunogenicity and pathogenesis in humanized mice.

J Virol **86**, 7637-7649 (2012).

11. Dutra, H. L. et al. Wolbachia Blocks Currently Circulating Zika Virus Isolates in Brazilian

Aedes aegypti Mosquitoes. *Cell Host Microbe* (2016).

12. Lanciotti, R. S. et al. Genetic and serologic properties of Zika virus associated with an

epidemic, Yap State, Micronesia, 2007. *Emerg Infect Dis* **14**, 1232-1239 (2008).

13. Aliota MT, P. S. A., Dario Velez I, and Osorio JE. The wMel strain of Wolbachia reduces

transmission of Zika virus by Aedes aegypti. *Scientific reports, in review* (2016).

14. Silveira, E. L. et al. Vaccine-induced plasmablast responses in rhesus macaques:

phenotypic characterization and a source for generating antigen-specific monoclonal

antibodies. *J Immunol Methods* **416**, 69-83 (2015).

15.   Lindsey, H. S., Calisher, C. H. & Mathews, J. H. Serum dilution neutralization test for

382   California group virus identification and serology. *J Clin Microbiol* **4**, 503-510 (1976).

383   16.   Pomplun, N., Weisgrau, K. L., Evans, D. T. & Rakasz, E. G. OMIP-028: activation panel

384   for Rhesus macaque NK cell subsets. *Cytometry A* **87**, 890-893 (2015).

385   17.   Wrammert, J. et al. Rapid cloning of high-affinity human monoclonal antibodies against

386   influenza virus. *Nature* **453**, 667-671 (2008).

387   18.   Staprans, S. I. et al. Simian immunodeficiency virus disease course is predicted by the

388   extent of virus replication during primary infection. *J Virol* **73**, 4829-4839 (1999).

389   19.   Lauck, M. et al. Discovery and full genome characterization of two highly divergent simian

390   immunodeficiency viruses infecting black-and-white colobus monkeys (Colobus guereza)

391   in Kibale National Park, Uganda. *Retrovirology* **10**, 107 (2013).

Supplementary Information is linked to the online version of the paper at

www.nature.com/nature.

Acknowledgements We thank the Veterinary, Animal Care, Scientific Protocol Implementation,

and the Pathology staff at the Wisconsin National Primate Research Center (WNPRC) for their

contribution to this study. We thank the DHHS/PHS/NIH (R01AI116382-01A1 to D.H.O.),

(R01AI107157-01A1 to T.G.G.) and (DP2HD075699 to S.R.P.) for funding. We also thank the

P51OD011106 awarded to the WNPRC, Madison-Wisconsin. This research was conducted in

part at a facility constructed with support from Research Facilities Improvement Program grants

RR15459-01 and RR020141-01. The publication's contents are solely the responsibility of the

authors and do not necessarily represent the official views of NCRR or NIH.

Author contributions D.H.O., T.C.F., J.E.O., M.T.A., E.M., T.G.G. and D.M.D. designed the

experiments. D.H.O., D.M.D., M.T.A., E. M., T.C.F., and L.H.M. drafted the manuscript. M.T.A.,

and J.E.O. provided and prepared viral stocks and performed plaque assays. A.M.W., G.L-B.,

and T.C.F. developed and performed viral load assays. K.L.W. and E.G.R. performed

immunophenotyping assays. M.S.M., M.E.B., M.N.R., C.M.N., and D.M.D. coordinated and

processed macaque samples for distribution. D.D.G., S.L.O., and D.M.D. designed and

performed the sequencing experiments. L.H.M. and T.C.F. performed nucleotide diversity

calculations. J.P., N.S-D., H.A.S., S.C., and J.M.H. coordinated the macaque infections,

sampling, and performed blood chemistries and CBC analysis. M.L.S. coordinated experiments

and helped perform ultrasounds on the pregnant macaques. J.A.E., M.A.M., and S.R.P.

performed the plasmablast experiments.

Author information: Sequences have been deposited in the Sequence Read Archive (SRA)

under accession number SRP072852. All data from these studies are available at

zika.labkey.com. Reprints and permissions information is available at www.nature.com/reprints.

The authors declare no competing interests. Correspondence and requests for materials should

be addressed to dhoconno@wisc.edu.

Figure Legends

Figure 1. Animal cohort definitions and ZIKV viral load from rhesus macaque fluids. a.

Animals included in this study and the ZIKV doses used to infect them. Solid lines and bars

throughout the figure represent cohort 1 animals by color while stripped bars and dotted lines

represent cohort 2 animals by color. **b**. Viral RNA loads measured in plasma, urine, and saliva

for the two animals challenged with each dose of virus through 28 dpi. Cohort 1 animals are

represented by a solid line while cohort 2 animals are represented by a dotted line for each fluid.

Inset: vRNA loads from cohort 1 animals measured before and after rechallenge with homotypic

Zika virus as indicated by an arrow. **c.** Number of plaque forming units per ml of serum for

cohort 2 animals. **d.** Viral RNA load per ml of CSF collected on 4 and 14 dpi. e. Viral RNA load

per vaginal swab collected on 0, 7, 14, 21 and 28 dpi. NA: sample not available.

Figure 2. Immune cell expansion and neutralizing antibody titers following ZIKV infection.

a. Solid dots, lines and bars with corresponding color represent cohort 1 animals and open

circles, dotted lines or stripped bars represent cohort 2 animals throughout the figure. **b**.

Expansion of Ki-67+ (activated) NK cells, CD8+ T cells and CD4+ T cells were measured daily

for 10 days and then on days 14, 21 and 28 post-infection. Absolute numbers of activated

cells/µl of blood are presented relative to the baseline value set to 100%. **c.** Total number of

plasmablast cells found in PBMCs collected at 0 (cohort 2 only), 3, 7, 11 and 14 dpi for each

animal. **d.** $PRNT_{90}$ titers for cohort 1 and cohort 2.

Figure 3. Characterization of Zika virus infection in animals infected during the first

trimester of pregnancy. a. Schematic of animals presented in this figure as cohort 3. Dots,

lines and bars representing each animal match in color throughout the figure. **b**. vRNA copies of

the plasma, urine, and saliva from each pregnant animal. Oral swabs could not be obtained

from 660875. **c.** Absolute numbers of Ki-67+ NK, CD8+ T cell and CD4+ T cell populations

presented as a percentage relative to baseline (x100) over time in each animal. **d.** Plasmablast

expansion over time from each pregnant animal. The average plasmablast expansion of cohort

1 and cohort 2 animals infected with the 10^4 PFU is presented by the black line. Error bars

represent standard deviation. **e.** $PRNT_{90}$ titers over time for each animal. Lines representing the

titers from cohort 2 animals are overlayed at 28 dpi for reference (top to bottom: 610107,

181856, and 411359).

Extended Data Figure 1. Schematic representation of the timeline of infection and

sampling for each animal in the presented studies. Cohort 1 received the first ZIKV

challenges and were then rested for 6 weeks before a rechallenge. For all studies, samples

were collected daily for 10 days and then on 14, 21, and 28 dpi as indicated by hashes in the

timelines. Cohort 3 represents the two pregnant animals that were challenged on 2 different

days. Both animals are currently in the once weekly sampling phase until the pregnancies come

to term (~165 gestational days). Cohort 2 was a repeat experiment of cohort 1 that allowed for

additional experiments and sample collection (e.g., serum plaque infectivity) that were not

feasible when we initiated cohort 1 studies. These animals are currently in a 6-week rest period

and will be rechallenged on June 6, 2016. Ages of all animals are indicated under each

macaque identification number.

Extended Data Figure 2. Complete blood counts and serum chemistries for macaques

infected with ZIKV. a. Animals were infected with different doses of ZIKV. Cohort 1 animals

are represented by solid lines and cohort 2 animals are represented by dotted lines. All non-

pregnant animals had serum chemistry analysis performed at -7, 0, 1, 2, 3, 4, 6, and 14 dpi or at

-6, 2, 5 and 11 dpi. **b.** AST blood chemistries **c.** ALT serum chemistries. **d.** CK serum

chemistries. Complete blood counts were measured prior to infection, daily for 10-11 days after

infection and then every 3-7 days until 28 dpi. **e.** white blood cell counts. **f.** % lymphocytes. **g.**

red blood cell counts.

476
Extended Data Figure 3. qRT-PCR detection of ZIKV RNA is equally sensitive from serum

and plasma. vRNA copies/ml of plasma or serum were quantitated by qRT-PCR over multiple

time points from cohort 2 animals. Sufficient baseline samples were not available from serum.

Extended Data Figure 4. Antigen-specific T cell responses by IFNγ-ELISPOT. a. Average

spot forming cell counts for PBMC collected from each animal at 4, 10 and 14 dpi. Data were

baseline corrected by subtracting the average negative control values from each response. A

threshold of 10.0 SFC/100,000 cells was set as the minimum value to be considered a positive

T cell response, as indicated by the dashed line. **b.** Each pool was comprised of 10 overlapping

15mer peptides offset by 4 amino acids. **c.** Peptide pools eliciting T cell responses at 4, 10 and

14 dpi for each animal. The region of the NS5 protein that is represented by each pool of

overlapping 15mers is provided. MHC class I haplotypes of each cohort 2 animal are also

presented. All three animals shared the A004 and B012b major histocompatibility complex

haplotypes and two animals shared the A023 haplotype. Therefore, it was not surprising that 3

pools were recognized by 2 different animals likely sharing the MHC class I allele that is

presenting one of the peptides in those pools. Grayed pools were positive in more than one

animal and bolded pools were positive at more than one time point in the same animal.

Extended Data Figure 5. Plaque assay titers in a pregnant animal. Log_{10} PFU/ml serum

(dotted line) is plotted relative to vRNA copies/ml plasma (solid line) for 660875.

Extended Data Figure 6. Genetic diversity of the ZIKV challenge stock. The ZIKV challenge

stock was deep sequenced from all three animals. Nucleotide sites where at least 5% of

sequences obtained from the challenge stock are different from the Genbank sequence are

shown.

Figure 1| Animal cohort definitions and ZIKV viral load from rhesus macaque fluids. a. Animals included in this study and the ZIKV doses used to infect them. Solid lines and bars throughout the figure represent cohort 1 animals by color while stripped bars and dotted lines represent cohort 2 animals by color. **b.** Viral RNA loads measured in plasma, urine, and saliva for the two animals challenged with each dose of virus through 28 dpi. Cohort 1 animals are represented by a solid line while cohort 2 animals are represented by a dotted line for each fluid Inset: vRNA loads from cohort 1 animalsmeasured before and after rechallenge with homotypic Zika virus as indicated by an arrow. **c.** Number of plaque forming units per ml of serum for cohort 2 animals. **d.** Viral RNA load per ml of CSF collected on 4 and 14 dpi. e. Viral RNA load per vaginal swab collected on 0, 7, 14, 21 and 28 dpi. NA: sample not available.

Figure 2| Immune cell expansion and neutralizing antibody titers following ZIKV infection. a. Solid dots, lines and bars with corresponding color represent cohort 1 animals and open circles, dotted lines or stripped bars represent cohort 2 animals throughout the figure. **b.** Expansion of Ki-67+ (activated) NK cells, CD8+ T cells and CD4+ T cells were measured daily for 10 days and then on days 14, 21 and 28 post-infection. Absolute numbers of activated cells/μl of blood are presented relative to the baseline value set to 100%. **c.** Total number of plasmablast cells found in PBMCs collected at 0 (cohort 2 only), 3, 7, 11 and 14 dpi for each animal. **d.** PRNT90 titers for cohort 1 and cohort 2 .

Figure 3| Characterization of Zika virus infection in animals infected during the first trimester of pregnancy. a. Schematic of animals presented in this figure as cohort 3. Dots, lines and bars representing each animal match in color throughout the figure. **b.** vRNA copies of the plasma, urine, and saliva from each pregnant animal. Oral swabs could not be obtained from 660875. **c.** Absolute numbers of Ki-67+ NK, CD8+ T cell and CD4+ T cell populations presented as a percentage relative to baseline (x100) over time in each animal. **d.** Plasmablast expansion over time from each pregnant animal. The average plasmablast expansion of cohort 1 and cohort 2 animals infected with 10^4 PFU ZIKV is presented by the black line. Error bars represent standard deviation. **e.** PRNT90 titers over time for each animal. Lines representing the titers from cohort 2 animals are overlaid at 28 dpi for reference (top to bottom: 610107, 181856, and 411359).

Extended Data Figure 1| Schematic representation of the timeline of infection and sampling for each animal in the presented studies. Cohort 1 received the first ZIKV challenges and were then rested for 6 weeks before a rechallenge. For all studies, samples were collected daily for 10 days and then on 14, 21, and 28 dpi as indicated by hashes in the timelines. Cohort 3 represents the two pregnant animals that were challenged on 2 different days. Both animals are currently in the once weekly sampling phase until the pregnancies come to term (~165 gestational days). Cohort 2 was a repeat experiment of cohort 1 that allowed for additional experiments and sample collection (e.g., serum plaque infectivity) that were not feasible when we initiated cohort 1 studies. These animals are currently in a 6-week rest period and will be rechallenged on June 6, 2016. Ages of all animals are indicated under each macaque identification number.

Extended Data Figure 2| Complete blood counts and serum chemistries for macaques infected with ZIKV.
a. Animals were infected with different doses of ZIKV. Cohort 1 animals are represented by solid lines and cohort 2 animals are represented by dotted lines. All non-pregnant animals had serum chemistry analysis performed at -7, 0, 1, 2, 3, 4, 6, and 14 dpi or at -6, 2, 5 and 11 dpi. **b.** AST blood chemistries **c.** ALT serum chemistries. **d.** CK serum chemistries. Complete blood counts were measured prior to infection, daily for 10-11 days after infection and then every 3-7 days until 28 dpi. **e.** white blood cell counts. **f.** % lymphocytes. **g.** red blood cell counts.

Extended Data Figure 3| qRT-PCR detection of ZIKV RNA is equally sensitive from serum and plasma. vRNA copies/ml of plasma or serum were quantitated by qRT-PCR over multiple time points from cohort 2 animals. Sufficient baseline samples were not available from serum.

b. Asian lineage ZIKV peptide pools

Pool ID	Target sequence
A	ZIKV Asian lineage NS5 1-51
B	ZIKV Asian lineage NS5 41-91
C	ZIKV Asian lineage NS5 81-131
D	ZIKV Asian lineage NS5 121-171
E	ZIKV Asian lineage NS5 161-211
F	ZIKV Asian lineage NS5 201-251
G	ZIKV Asian lineage NS5 241-291
H	ZIKV Asian lineage NS5 281-331
I	ZIKV Asian lineage NS5 321-371
J	ZIKV Asian lineage NS5 361-411
K	ZIKV Asian lineage NS5 401-451
L	ZIKV Asian lineage NS5 441-491
M	ZIKV Asian lineage NS5 481-531
N	ZIKV Asian lineage NS5 521-571
O	ZIKV Asian lineage NS5 561-611
P	ZIKV Asian lineage NS5 601-651

c. Summary of NS5-specifc T cell responses

Animal ID	MHC Haplotype	DPI	Pool ID	Target Sequence
411359	A004, A023, B012b, B024a	10	D	ZIKV Asian NS5 121-171
			H	**ZIKV Asian NS5 281-331**
			O	ZIKV Asian NS5 561-575
		14	**H**	**ZIKV Asian NS5 281-331**
			J	ZIKV Asian NS5 361-411
181856	A001, A004, B012b, B055	4	F	ZIKV Asian NS5 201-251
		10	P	ZIKV Asian NS5 601-651
		14	O	ZIKV Asian NS5 561-575
610107	A004, A023, B012b, B048	4	**K**	**ZIKV Asian NS5 401-451**
		10	A	ZIKV Asian NS5 1-51
			B	**ZIKV Asian NS5 41-91**
			C	ZIKV Asian NS5 81-131
			D	**ZIKV Asian NS5 121-171**
			F	**ZIKV Asian NS5 201-251**
			K	**ZIKV Asian NS5 401-451**
		14	**B**	**ZIKV Asian NS5 41-91**
			D	**ZIKV Asian NS5 121-171**
			F	**ZIKV Asian NS5 201-251**
			I	ZIKV Asian NS5 321-371
			N	ZIKV Asian NS5 521-571

Extended Data Figure 4| Antigen-specific T cell responses by IFNγ-ELISPOT. a. Average spot forming cell counts for PBMC collected from each animal at 4, 10 and 14 dpi. Data were baseline corrected by subtracting the average negative control values from each response. A threshold of 10.0 SFC/100,000 cells was set as the minimum value to be considered a positive T cell response, as indicated by the dashed line. **b.** Each pool was comprised of 10 overlapping 15mer peptides offset by 4 amino acids. **c.** Peptide pools eliciting T cell responses at 4, 10 and 14 dpi for each animal. The region of the NS5 protein that is represented by each pool of overlapping 15mers is provided. MHC class I haplotypes of each cohort 2 animal are also present-ed. All three animals shared the A004 and B012b major histocompatibility complex haplotypes and two animals shared the A023 haplotype. Therefore, it was not surprising that 3 pools were recognized by 2 different animals likely sharing the MHC class I allele that is presenting one of the peptides in those pools. Grayed pools were positive in more than one animal and bolded pools were positive at more than one time point in the same animal.

Extended Data Figure 5| Plaque assay titers in a pregnant animal. Log10 PFU/ml serum (dotted line) is plotted relative to vRNA copies/ml plasma (solid line) for 660875.

Site relative to KJ776791	Change	Amino Acid Change	Codon Change	Protein Effect	Variant Frequency
801	T -> C	F->L	UUC->CUC	Substitution	17.9%
2318	C -> T		CUC->CUT	None	18.7%
2863	T -> C	L->P	CUC->CCC	Substitution	7.7%
3223	A -> T	K->I	AAA->ATA	Substitution	10.5%
3475	T -> C	M->T	AUG->ACG	Substitution	7.9%
3740	C -> T		GUC->GUT	None	11.1%
3835	C -> T	A->V	GCG->GTG	Substitution	15.8%
4242	G -> C	E->Q	GAG->CAG	Substitution	40.9%

Extended Data Figure 6| Genetic diversity of the ZIKV challenge stock. The ZIKV challenge stock was deep sequenced from all three animals. Nucleotide sites where at least 5% of sequences obtained from the challenge stock are different from the Genbank sequence are shown.

SI Guide

Supplementary Information Data

Morbidity, complete blood counts and serum blood chemistry findings in nonpregnant rhesus macaques exposed to ZIKV. Data is shown in Extended Data Fig. 2.

To detect signs of morbidity, animals were evaluated daily for evidence of

disease, injury, or psychological abnormalities (e.g., inappetence, dehydration, diarrhea,

depression, inactivity, trauma, self-injurious or stereotypical repetitive behaviors (e.g.,

pacing) often seen in captive animals). Five of six animals exhibited mild to moderate

inappetence, which resulted in mild weight loss in four animals. Two animals (912116

and 393422) also developed a very mild rash around the inoculation site at 1 dpi that

persisted for 4-5 days. No other abnormal clinical signs were noted (e.g., increased body

temperature, joint pain, lymphadenopathy, lethargy).

Daily complete blood counts (CBCs) were evaluated for all six non-pregnant

animals for 10 dpi and then every 3 to 7 days thereafter and serum chemistry analyses

were performed intermittently post-infection as per protocol (Extended Data Fig. 2).

Reference intervals (RI) developed for the WNPRC colony for species, gender, and age

were used to evaluate results. All six animals developed elevated serum creatine kinase

(CK), which peaked by 5 dpi (Extended Data Fig. 2d). Increases in serum CK are

strongly associated with muscle damage and myositis (skeletal, smooth, and cardiac),

but can also be caused by repeated sedation, hemolysis, and endocrine abnormalities

[20,21]. Predictable increases in CK have been noted in nonhuman primates undergoing

repeated sedation and venipuncture [21]. However, CK levels did not remain elevated

during the 10-day period of daily sedation and blood collection, suggesting other causes

for the noted increase in values. Future studies are planned to determine if CK increases

may be due to viral myositis. ALT values in three of six animals exceeded the maximum

WNPRC RI (Extended Data Fig. 2c) and the expected increase associated with repeated

ketamine sedation [21]. Although AST values exceeded upper RI values for all six non-

pregnant animals after infection, they did not increase above the expected levels

previously associated with repeated ketamine sedation described by Lugo-Roman et al.

(Extended Data Fig. 2b) [21].

All the animals displayed decreased total WBC numbers following infection, but

only one animal fell below the RI with a value of 2.88 ths/ul (3.70-15.70). WBC numbers

rebounded almost completely to pre-infection levels around 10 dpi in all six animals

(Extended Data Fig. 1e). Both animals that received the highest dose of inoculum

developed persistent mature neutrophilia around 7-14 dpi that lasted through 28 dpi.

Five of 6 non-pregnant animals had mild regenerative anemia characterized by varying

degrees of polychromasia and anisocytosis, but whether this was secondary to the viral

infection or simply a result of frequent blood collections could not be determined. Platelet

values for all six animals remained within RI. Both cellular dyscrasias and elevated

transaminases have been described in human ZIKV case reports; myositis has not been

reported [22,23].

References

20.   Stockham, S. L. *Fundamentals of veterinary clinical pathology.* (Iowa State Press,

Ames, Iowa, 2002).

21.   Lugo-Roman, L. A., Rico, P. J., Sturdivant, R., Burks, R. & Settle, T. L. Effects of

serial anesthesia using ketamine or ketamine/medetomidine on hematology and

serum biochemistry values in rhesus macaques (Macaca mulatta). *J Med Primatol*

39, 41-49 (2010).

22.   Tappe, D. et al. Acute Zika virus infection after travel to Malaysian Borneo,

September 2014. *Emerg Infect Dis* **21**, 911-913 (2015).

23.   Zammarchi, L. et al. Zika virus infections imported to Italy: clinical, immunological

and virological findings, and public health implications. *J Clin Virol* **63**, 32-35

(2015).