Title: Repeated duplication of Argonaute2 is associated with strong selection and testis specialization

in *Drosophila*

Authors: Samuel H. Lewis[1,2]*, Claire L. Webster[1,3], Heli Salmela[4] & Darren J. Obbard[1,5]

Affiliations:

[1]Institute of Evolutionary Biology, University of Edinburgh, Kings Buildings, EH9 3JT, United Kingdom

[2]Present Address: Department of Genetics, University of Cambridge, Downing Street, Cambridge,

CB2 3EH

[3]Present Address: Life Sciences, University of Sussex, United Kingdom

[4]Department of Biosciences, Centre of Excellence in Biological Interactions, University of Helsinki,

Helsinki, Finland

[5]Centre for Immunity, Infection and Evolution, University of Edinburgh, Kings Buildings, EH9 3JT,

United Kingdom

*Author for correspondence: sam.lewis@gen.cam.ac.uk

Abstract

Argonaute2 (Ago2) is a rapidly evolving nuclease in the *Drosophila melanogaster* RNAi pathway that

targets viruses and transposable elements in somatic tissues. Here we reconstruct the history of Ago2

duplications across the *Drosophila obscura* group, and use patterns of gene expression to infer new

functional specialization. We show that some duplications are old, shared by the entire species group,

and that losses may be common, including previously undetected losses in the lineage leading to *D.*

pseudoobscura. We find that while the original (syntenic) gene copy has generally retained the

ancestral ubiquitous expression pattern, most of the novel Ago2 paralogues have independently

specialised to testis-specific expression. Using population genetic analyses, we show that most testis-

specific paralogues have significantly lower genetic diversity than the genome-wide average. This

suggests recent positive selection in three different species, and model-based analyses provide

strong evidence of recent hard selective sweeps in or near four of the six *D. pseudoobscura* Ago2

paralogues. We speculate that the repeated evolution of testis-specificity in *obscura* group Ago2

genes, combined with their dynamic turnover and strong signatures of adaptive evolution, may be

associated with highly derived roles in the suppression of transposable elements or meiotic drive. Our

study highlights the lability of RNAi pathways, even within well-studied groups such as *Drosophila*,

and suggests that strong selection may act quickly after duplication in RNAi pathways, potentially

giving rise to new and unknown RNAi functions in non-model species.

Introduction

Argonaute genes are found in almost all eukaryotes, where they play a key role in antiviral immune defence, gene regulation and genome stability. They carry out this diverse range of functions through their role in RNA interference (RNAi) mechanisms, an ancient system of nucleic acid manipulation in which small RNA (sRNA) molecules guide Argonaute proteins to nucleic acid targets through base complementarity (reviewed in [1]). Gene duplication has occurred throughout the evolution of the Argonaute gene family, with ancient duplication events characteristic of some lineages – such as three duplications early in plant evolution [2], and multiple expansions and losses throughout the evolution of nematodes (reviewed in [3]) and the Diptera [4]. After duplication, Argonautes have often undergone functional divergence, involving changes in expression patterns and altered small RNA (sRNA) binding partners [5–7]. Duplication early in eukaryotic evolution produced two distinct Argonaute subfamilies, Ago and Piwi, which have since been retained in the vast majority of Metazoa [8]. Members of the Ago subfamily are expressed in both somatic and germline tissue, and variously bind sRNAs derived from host transcripts (miRNAs, endo-siRNAs) or transposable elements (TE endo-siRNAs) and viruses (viRNAs). In contrast, in most vertebrates and arthropods, the Piwi subfamily members are expressed only in association with the germline (reviewed in [9]), and bind sRNAs from TEs and host loci (piRNAs), suggesting that the Piwi subfamily specialised to a germline-specific role on the lineages leading to vertebrates and arthropods.

After the early divergence of the Ago and Piwi subfamilies, subsequent duplications gave rise to three Piwi subfamily members (Ago3, Aubergine (Aub) and Piwi) and two Ago subfamily members (Ago1 & Ago2) in *Drosophila melanogaster*. All three Piwi subfamily genes are associated with the germline and bind Piwi-interacting RNAs (piRNAs) derived from TEs and other repetitive genomic elements: Ago3 and Aub amplify the piRNA signal through the "Ping-Pong" cycle (reviewed in [10]), and Piwi suppresses transposition by directing heterochromatin formation [11]. These functional differences are associated with contrasting selective regimes, with Aub evolving under positive selection [12] and more rapidly than Ago3 and Piwi [13]. In contrast, Ago1 binds microRNAs (miRNAs), and regulates gene expression by inhibiting translation and marking transcripts for degradation (reviewed in [14]). This function imposes strong selective constraint on Ago1, resulting in slow evolution and very few adaptive substitutions [12,13,15]. Finally, Ago2 binds small interfering RNAs (siRNAs) from viruses (viRNAs) and TEs (endo-siRNAs), and functions in gene regulation [16], dosage compensation [17],

and the ubiquitous suppression of viruses [18,19] and TEs [20,21]. Ago2 also evolves under strong

positive selection, with frequent selective sweeps [12,13,15,22,23], possibly driven by an arms race

with virus-encoded suppressors of RNAi (VSRs) [15,24,25].

In contrast to *D. melanogaster*, from which most functional knowledge of Ago2 in arthropods is

derived, an expansion of Ago2 has been reported in *D. pseudoobscura* [26], providing us with an ideal

opportunity to study how the RNAi pathway evolves after duplication. Given the roles of *D.*

melanogaster Ago2 in antiviral defence [18,19], TE suppression [20,21], dosage compensation [17],

and gene regulation [16], we hypothesized that duplication in *D. pseudoobscura* may have led to

subfunctionalization of Ago2 to a subset of these roles, or even to the evolution of entirely new

functions. To elucidate the evolution and function of Ago2 paralogues in *D. pseudoobscura* and its

relatives, we identified and dated Ago2 duplication events across available *Drosophila* genomes and

transcriptomes, tested for divergence in expression patterns between the Ago2 paralogues in *D.*

subobscura, *D. obscura* and *D. pseudoobscura*, and quantified the evolutionary rate and positive

selection acting on each of these paralogues. We find that testis-specificity of Ago2 paralogues has

evolved repeatedly in the *obscura* group, and that the majority of paralogues show evidence of recent

positive selection.

<u>Results</u>

<u>Ago2 has undergone numerous ancient and recent duplications in the *obscura* group</u>

Ago2 duplications had previously been noted in *D. pseudoobscura* [26], but their age and distribution

in other species was unknown. We used BLAST [27] and PCR to identify 65 Ago2 homologues in 39

species sampled across the Drosophilidae, including 30 homologues in 9 *obscura* group species. To

characterize the relationships between Ago2 homologues in the *obscura* group and the other

Drosophilidae, and estimate the date of the duplication events that produced them, we carried out a

strict clock Bayesian phylogenetic analysis (Figure 1). This showed that there are early diverging

Ago2 clades in the *obscura* group: the Ago2e subclade that diverged from other Ago2 paralogues

around 21mya (±10 My), and the Ago2a and Ago2f subclades that were produced by a gene

duplication event around 16mya (±7 My). Subsequently there have been a series of more recent

duplications in the *D. pseudoobscura* subgroup Ago2a-d lineage. Using published genomes,

Figure 1: An approximately time-scaled Bayesian gene tree of Ago2 in the Drosophilidae. Duplication events are marked by yellow diamonds, Bayesian posterior support is shown for nodes for which it is less than 100%, and the genes and species that are the focus of the present study are marked in bold. Ago2 has duplicated at least twelve times in the Drosophilidae: seven times in the *obscura* group, twice early in the *melanogaster* group, and once each in the lineages leading to *D. willistoni*, *S. deflexa* and *D. kikkawai*. There has also been a potentially recent duplication of Ago2a on the *D. affinis* / *D. azteca* lineage (~5mya), although the low support for this node may suggest that these paralogues could also nest within the *D. pseudoobscura* / *D. persimilis* expansion, with one paralogue sister to the Ago2a1-Ago2b subclade and the other sister to the Ago2c-Ago2d subclade. After duplication, Ago2 paralogues in the *obscura* group have specialised to the testis three times independently (marked with ♂), and have been retained for an extended period of time (>10 My in the case of Ago2e), suggesting an adaptive basis for testis-specificity. The labelling a-e of paralogous clades corresponds to reference [26], while clade f is newly reported here.

transcriptomes and PCR we were unable to identify Ago2e in *D. subobscura*, Ago2e or Ago2f in *D.*

lowei, or Ago2f in *D. pseudoobscura*, *D. persimilis* and *D. azteca*. While some of these losses may

reflect incomplete genome assemblies or unexpressed genes in transcriptome surveys, we attempted

to validate the losses in *D. pseudoobscura* and *D. subobscura* by extensive PCR, and were again

unable to recover these genes.

In release 3.03 of the *D. pseudoobscura* genome Ago2b-Ago2e have confirmed locations, but Ago2a1

and Ago2a3 (the very recent paralogues newly identified here) lie in tandem on an unplaced contig

with a third incomplete copy (Ago2a2) between them. We used PCR to confirm the existence,

orientation, and relative positioning of these genes, and to identify the location of this contig, which

lies in reverse orientation on chromosome XL-group1a (predicted coordinates 3,463,701-3,489,689).

We then combined this information with our phylogenetic analysis to reconstruct the positional

evolution of *D. pseudoobscura* Ago2 paralogues (Figure 2).

Figure 2: The course of duplications and translocations of Ago2 paralogues in *D. pseudoobscura.* A complex series of duplications and translocations has produced six Ago2 paralogues in *D. pseudoobscura*, located on four different chromosome arms. Chromosome arms correspond to Muller Elements A (X/XL), B (2L), C (2R), D (3L), E (3R) & F (4) (adapted from [91] Fig. 1). Firstly, the Ago2a1-e ancestor duplicated ~21mya to form Ago2a1-d and Ago2e, the latter of which moved onto chromosome 2L. Next, the 3L arm fused with the X chromosome, moving Ago2a1-d onto the X: this happened 7-15mya, after the divergence of the *obscura* group into Palearctic (e.g. *D. subobscura*) and Nearctic (e.g. *D. pseudoobscura*) clades [92]. Ago2a1-d then duplicated ~6mya, forming Ago2c-d and Ago2a1-b, the latter of which moved onto chromosome 2. After this, Ago2a1-b duplicated ~5mya, producing Ago2b and Ago2a1-3, the latter of which moved onto the left arm of the X chromosome. This was followed by a duplication of Ago2c-d ~2mya, forming Ago2d and Ago2c, the latter of which moved onto chromosome 2. Finally, Ago2a1-3 duplicated ~27kya, producing Ago2a1 and Ago2a3 in tandem. Note that due to differences in evolutionary rate between branches, the timings of these events should be treated with caution.

<u>Ago2 paralogues in *D. subobscura*, *D. obscura* and *D. pseudoobscura* are probably functional</u>

Our phylogenetic analysis (Figure 1) revealed that the Ago2 paralogues in the *obscura* group have

retained coding sequences for millions of generations, showing that they have remained functional for

this period. They have also retained PAZ and PIWI domains and a bilobal structure (characteristic of

Argonaute proteins across the tree of life), suggesting that they are part of a functional RNAi pathway.

In *D. melanogaster*, Ago2 plays a key role in antiviral immunity, but is ubiquitously and highly

expressed in both males and females, and is not strongly induced by viral challenge (Figure 3a, [28]).

To test whether this expression pattern has been conserved after Ago2 duplication, or whether any

Ago2 paralogues have become inducible by viral challenge, we measured the expression of each

Ago2 paralogue in female and male *D. subobscura*, *D. obscura* and *D. pseudoobscura* after infection

with Drosophila C Virus (DCV). These species are separated by ~10My of evolution, and represent

the three major clades within the *obscura* group. Members of the *obscura* group are known to be

highly susceptible to DCV, supporting high viral titres and displaying rapid mortality [29]. We found

that only one paralogue is expressed in both sexes at a high level in *D. subobscura* (Ago2a), *D.*

obscura (Ago2a) and *D. pseudoobscura* (Ago2c). Unexpectedly, and with only one exception, the

other Ago2 paralogues in all species were only expressed in males (Figure 3b-d), raising the

possibility that they have specialised to a sex-specific role. The one exception was *D. pseudoobscura*

Ago2d, which is the ancestral paralogue in this species (inferred by synteny), and for which we could

not detect any expression.

Figure 3: Expression patterns of Ago2 paralogues under challenge with Drosophila C Virus. In each *obscura* group species, only one Ago2 paralogue has retained the ancestral pattern of ubiquitous stable expression in each sex (illustrated by *D. melanogaster*). In contrast, all other paralogues are expressed in males only (apart from *D. pseudoobscura* Ago2d, which is unexpressed in either sex). The high degree of sequence similarity between Ago2a1 and Ago2a3 prevented us from amplifying these genes separately in qPCR, and here they are combined as "Ago2a". Error bars indicate 1 standard error estimated from 2 technical replicates in each of three different genetic backgrounds. Apparent differences in expression between sexes and species should be interpreted with caution, as these may be driven by differences in expression levels of the reference gene (RpL32).

<u>Ago2 paralogues have repeatedly specialised to the testis</u>

To determine whether the strongly male-biased expression pattern is associated with a testis-specific role, we quantified the tissue-specific expression patterns of Ago2 paralogues in *D. subobscura*, *D. obscura* and *D. pseudoobscura*. In *D. melanogaster* the single copy of Ago2 was expressed in all adult tissues (Figure 4a), and transcripts were present in the embryo (S1 Figure). In *D. subobscura*, *D. obscura* and *D. pseudoobscura*, we found that the Ago2 paralogues exhibited striking differences in their tissue-specific patterns of expression (Figure 4b-d). In each species, one paralogue has retained the ancestral ubiquitous expression pattern in adult tissues. In contrast, every other paralogue was expressed only in the testis, except for the non-expressed *D. pseudoobscura* Ago2d. None of the testis-specific paralogues in *D. pseudoobscura* was detectable in embryos (S1 Figure).

Interestingly, the ubiquitously expressed paralogue in *D. subobscura* and *D. obscura* is the ancestral gene (Ago2a in both cases, as inferred by synteny with *D. melanogaster*), but in *D. pseudoobscura* another paralogue (Ago2c) has evolved the ubiquitous expression pattern, and the ancestral gene (Ago2d) was not expressed at a detectable level in any tissue. When interpreted in the context of the phylogenetic relationships between these paralogues, the most parsimonious explanation is that testis-specificity evolved at least three times: firstly at the base of the Ago2e clade, secondly at the base of the Ago2f clade, and thirdly at the base of the *D. pseudoobscura-D. persimilis* Ago2a-Ago2b subclade (Figure 1).

Figure 4: Tissue-specific expression patterns of Ago2 paralogues.
In each of the three *obscura* group species tested, one paralogue has retained the ancestral ubiquitous expression pattern, while the others have specialised to the testis (with the exception of *D. pseudoobscura* Ago2d). The high degree of sequence similarity between Ago2a1 and Ago2a3 prevented us from amplifying these genes separately in qPCR, and here they are combined as "Ago2a". Error bars indicate 1 standard error estimated from 2 technical replicates in each of five different genetic backgrounds. *D. melanogaster* expression levels were taken from a single RNA-seq experiment [71].

<u>Testis-specificity is associated with faster protein evolution</u>

To test for differences in evolutionary rate between testis-specific and ubiquitously expressed Ago2

paralogues, we fitted sequence evolution models to the set of drosophilid Ago2 sequences depicted in

Figure 1 using codeml (PAML, Yang 1997). These tests estimate separate dN/dS ratios (ω) and

likelihoods for different subclades in the gene tree, providing a test for differential rates of evolution.

We found that most support (Akaike weight = 0.99) falls behind a model specifying a different ω for

each obscura group Ago2 subclade, and another separate ω for the *D. pseudoobscura-D. persimilis*

Ago2a-Ago2b subclade. Under this model, the testis-specific *D. pseudoobscura-D. persimilis* Ago2a-

Ago2b subclade has the highest rate of protein evolution (ω=0.32±0.047 SE), followed by the testis-

specific Ago2f subclade (ω=0.21±0.014), the ubiquitous Ago2a subclade (ω=0.19±0.012), the testis-

specific Ago2e subclade (ω=0.16±0.010), and finally the other Drosophilid Ago2 sequences

(ω=0.12±0.002). This shows that the evolution of testis-specificity is generally accompanied by an

increase in the rate of protein evolution. We also used the Bayes Empirical Bayes sites test in codeml

to identify codons evolving under positive selection across the entire gene tree, and the branch-sites

test to identify codons under positive selection in the *obscura* group Ago2 subclade. While we found

no positively-selected codons with the sites test, we identified three codons under positive selection

(297, 338 & 360) in the *obscura* group Ago2 subclade with the branch-sites test (likelihood ratio test

M8 vs M8a, p<0.005).

<u>McDonald-Kreitman tests identify strong positive selection on *D. pseudoobscura* Ago2e</u>

This increase in evolutionary rate after the evolution of testis-specificity may have occurred as a result

of positive selection, or the relaxation of selective constraint. However, unless there are multiple

substitutions within single codons, this will be hard to detect using methods such as codeml.

Therefore, as a second test for positive selection on Ago2 paralogues in *D. subobscura*, *D. obscura*

and *D. pseudoobscura*, we gathered intraspecies polymorphism data for each Ago2 paralogue in

these species (S4 Appendix), and performed McDonald-Kreitman (MK) tests (S1 Table). The MK test

uses a comparison of the numbers of fixed differences between species at nonsynonymous (Dn) and

synonymous (Ds) sites, and polymorphisms within a species at nonsynonymous (Pn) and

synonymous (Ps) sites to infer the action of positive selection. If all mutations are either neutral or

strongly deleterious, the Dn/Ds ratio should be approximately equal to the Pn/Ps ratio; however, if

there is positive selection, an excess of nonsynonymous differences is expected [31]. The majority of

MK tests were non-significant (Fisher's exact test, p>0.1), despite often displaying relatively high

K_A/K_S ratios e.g. *D. pseudoobscura* Ago2a1 (K_A/K_S =0.34), Ago2b (K_A/K_S =0.43) & Ago2d (K_A/K_S

=0.36). However, the low diversity at these loci (<10 polymorphic sites in most cases; see below) will

mean that the MK approach has little power, and that estimates of the proportion of substitutions that

are adaptive (α) are likely to be poor. In contrast to the other loci, we identified strong positive

selection acting on *D. pseudoobscura* Ago2e – which has relatively high genetic diversity – with α at

100% (α=1.00; Fisher's exact test, p=0.0004). This result is driven by the extreme skew in the

proportion of nonsynonymous to synonymous polymorphisms (0 Pn to 17 Ps), despite substantial

numbers of fixed differences (77 Dn to 120 Ds), and is robust to the choice of outgroup (S2 Table).

<u>The majority of Ago2 paralogues have extremely low levels of sequence diversity</u>

When strong selection acts to reduce genetic diversity at a locus, it can also reduce diversity at linked

loci before recombination can break up linkage [32]. Recent positive selection can therefore be

inferred from a reduction in synonymous site diversity compared with other genes. Because MK tests

can only detect multiple long-term substitutions, and are hampered by low diversity, diversity-based

approaches offer a complementary way to detect very recent strong selection. We therefore

compared the synonymous site diversity at each Ago2 paralogue in *D. pseudoobscura* with the

distribution of genome-wide synonymous site diversity. We found that all paralogues have unusually

low diversity relative to other loci: Ago2a1, Ago2b and Ago2c fall into the lowest percentile, Ago2a3

and Ago2d into the 2nd lowest percentile and Ago2e into the 8th lowest percentile (S3 Figure). A

multi-locus extension of the HKA test (ML-HKA [33]) confirmed that the diversity of Ago2a1-Ago2e is

significantly lower than the *D. pseudoobscura* genome as a whole (Akaike weight = 0.98).

Unfortunately, population-genomic data are not available for *D. subobscura* and *D. obscura*,

preventing a similar analysis. However, we found similar results for Ago2a and Ago2e when

comparing the diversity of *D. subobscura* and *D. obscura* Ago2 paralogues to levels of diversity

inferred from transcriptome data (data from [34]), suggesting that this effect is not limited to *D.*

pseudoobscura and these genes may therefore have been recent targets of selection in multiple

species. In *D. obscura*, Ago2a and Ago2e fall into the 2nd and 4th lowest diversity percentile

respectively, whereas Ago2f falls into the 19th percentile (S3 Figure). In *D. subobscura*, Ago2a falls

into the 7th percentile, whereas Ago2f falls into the 16th percentile (S3 Figure). The prevalence of low

intraspecific diversity for testis-specific paralogues is consistent with recent selective sweeps,

suggesting that positive selection, not merely relaxation of constraint, has contributed to the increased

evolutionary rate seen after specialization to the testis.

<u>Four out of six *D. pseudoobscura* Ago2 show a strong signature of recent hard selective sweeps</u>

The impact of selection on linked diversity (a selective sweep) is expected to leave a characteristic

footprint in local genetic diversity around the site of selection, and this forms the basis of explicit

model-based approaches to detect the recent action of positive selection [35]. For *D. pseudoobscura*,

population genomic data for 11 haplotypes is available from [36], permitting an explicit model-based

test for recent hard selective sweeps near to Ago2 paralogues. We therefore combined our Ago2 data

with 111kb haplotypes from [36] to analyse the neighbouring region around each paralogue. Ago2a1

and Ago2a3 form a tandem repeat, and were therefore analysed together as a single potential sweep.

We found strong evidence for recent selective sweeps at or very close to Ago2a1/3, Ago2b and

Ago2c, which display sharp troughs in their diversity levels, and large peaks in the composite

likelihood of a sweep, which far exceed a significance threshold derived from coalescent simulation

($p < 0.01$) (Figure 5). These localised reductions in diversity remain when our own Ago2 haplotype data

is removed, showing the results are robust to the fact that our Ago2 sequence data is derived from a

different population to the genome-wide data of [36] (S5 Figure; note that sequence data for Ago2

paralogues cannot be derived from the data of reference [36], because of their extreme similarity). In

addition, there is ambiguous evidence for a sweep at Ago2d, in the form of one significant ($p < 0.01$)

likelihood peak just upstream of the paralogue, but two other peaks ~1kb and ~3kb further upstream.

There is no evidence for a hard sweep at Ago2e, which has no diversity trough or likelihood peak.

Figure 5: Selective sweeps at *D. pseudoobscura* Ago2 paralogues.

For each paralogue, diversity at all sites (Watterson's θ) is displayed in red, and the likelihood of a sweep centred at that site (composite likelihood ratio, CLR) is displayed in blue. The significance threshold for the CLR is displayed by the horizontal dotted line (p<0.01, derived from the 10th-highest CLR out of 1000 coalescent simulations, assuming constant recombination rate and N_e). There is strong evidence for sweeps at Ago2a, Ago2b and Ago2c, indicated by troughs in their diversity levels and peaks in the likelihood of a sweep.

Discussion

Testis-specificity may indicate a loss of antiviral function

We have found that Ago2 paralogues in the *obscura* group have repeatedly evolved divergent

expression patterns after duplication, with the majority of paralogues specializing to the testis. This is

the first report of testis-specificity for any arthropod Ago2, which is ubiquitously expressed in *D.*

melanogaster [37], and provides a strong indication that these paralogues have diverged in function.

This testis-specificity (Figure 4), combined with a lack of upregulation on viral challenge (in contrast to

virus-responsive genes in the Toll [38] and Jak-STAT [39] signalling pathways in *D. melanogaster*),

suggests that these Argonautes are likely to have lost their ancestral ubiquitous antiviral role. In

contrast, one paralogue in each species has retained this ubiquitous expression pattern (*D.*

subobscura Ago2a, *D. obscura* Ago2a & *D. pseudoobscura* Ago2c, Figure 4), suggesting that these

paralogues have retained roles in antiviral defence [18,19], dosage compensation [17] and/or somatic

TE suppression [20,21].

Both ubiquitous and testis-specific Ago2 paralogues show evidence of recent positive selection

We identified selective sweeps at the ubiquitously expressed Ago2 paralogue in *D. pseudoobscura*

Ago2c, and very low diversity in the ubiquitously expressed Ago2 paralogues of *D. subobscura* and *D.*

obscura (Ago2a), suggesting that all of these genes may have recently experienced strong positive

selection. This is consistent with previous findings of strong selection and rapid evolution of Ago2 in

D. melanogaster [15,22,23] which has also experienced recent sweeps in *D. melanogaster*, *D.*

simulans, and *D. yakuba* [23], and across the *Drosophila* more broadly [12]. It has previously been

suggested that this is driven by arms-race coevolution with viruses [12,13], some of which encode

viral suppressors of RNAi (VSRs) that block Ago2 function [40]. The presence of VSR-encoding

viruses, such as Nora virus, in natural *obscura* group populations [34], combined with the host-

specificity that VSRs can display [25], suggest that arms-race dynamics may also be driving the rapid

evolution of ubiquitously expressed Ago2 paralogues in the *obscura* group.

Potential testis-specific functions

In contrast to their ancestral ubiquitous expression pattern, the dominant fate for Ago2 paralogues in

the *obscura* group appears to have been specialization to the testis. Paralogues often undergo a brief

period of testis-specificity soon after duplication [41,42], and this has given rise to the 'out-of-the-

testis' hypothesis, in which new paralogues are initially testis-specific before evolving functions in

other tissues [43]. However, two lines of evidence suggest an adaptive basis for the testis-specificity

observed for the *obscura* group Ago2 paralogues. First, testis-specificity has been retained for more

than 10 million years in Ago2e and Ago2f, in contrast to the broadening of expression over time

expected under the out-of-the-testis hypothesis [41,43]. Second, all testis-specific Ago2 paralogues

in *D. pseudoobscura* show evidence either of long-term positive selection (MK test for the high-

diversity Ago2e) or of recent selective sweeps (in low-diversity Ago2a1/3 and Ago2b), and the testis-

specific *D. obscura* Ago2e displays a reduction in diversity, potentially driven by selection.

Under a subfunctionalization model for Ago2 testis-specialization, four candidate selective pressures

seem likely: testis-specific dosage compensation, antiviral defence, TE suppression, and/or the

suppression of meiotic drive. Of these, testis-specific dosage compensation seems the least likely to

drive testis-specificity because the male-specific lethal (MSL) complex, which Ago2 directs to X-linked

genes to carry out dosage compensation in the soma of *D. melanogaster*, is absent from testis [44].

Testis-specific antiviral defence seems similarly unlikely, as the only known paternally-transmitted

Drosophila viruses (Sigmaviruses; Rhabdoviridae) pass through both the male and female gametes

[45], and so the potential benefits of testis-specificity seem unclear. In contrast, the suppression of

TEs or meiotic drive seem more promising candidate selective forces. First, numerous TEs transpose

preferentially in the testis, such as *Penelope* in *D. virilis* [46] and *copia* in *D. melanogaster* [47,48],

which could impose a selection pressure on Ago2 paralogues to provide a testis-specific TE

suppression mechanism. Nevertheless, it should be noted that all members of the canonical anti-TE

Piwi subfamily (Ago3, Aub and Piwi) are also expressed in *obscura* group testis (S2 Figure),

suggesting that if Ago2 paralogues have specialised to suppress TEs, they are doing so alongside the

existing TE suppression mechanism. Second, testis-specificity could have evolved to suppress

meiotic drive, which is prevalent (in the form of sex-ratio distortion) in the *obscura* group [49–53], and

which is suppressed by RNAi-based mechanisms in other species [54–56]. A high level of meiotic

drive in the *obscura* group could therefore impose selection for the evolution of novel suppression

mechanisms, leading to the repeated specialization of Ago2 paralogues to the testis.

<u>Prospects for novel functions during the evolution of RNAi</u>

The functional specialization that we observe for *obscura* group Ago2 paralogues raises the prospect

of undiscovered derived functions following Argonaute expansions in other lineages. Ago2 has

duplicated frequently across the arthropods, with expansions present in insects (*Drosophila willistoni*

(Figure 1) & *Musca domestica* [57]), crustaceans (*Penaeus monodon* [6]) and chelicerates

(*Tetranychus urticae*, *Ixodes scapularis*, *Mesobuthus martensii* & *Parasteatoda tepidariorum* [58]).

The prevalence of testis-specificity in *obscura* group Ago2 paralogues raises the possibility that

specialization to the germline may be more widespread following Argonaute duplication. The

expression of Ago2 paralogues has previously been characterized in *P. monodon*, and shows that

one paralogue has indeed specialised to the germline of both males and females, but not the testis

alone [6]. Publicly available RNAseq data from the head, gonad and carcass of male and female

Musca domestica [59] suggests that neither Ago2 paralogue has specialised to the testis (S6 Figure).

However, public data from the head, thorax and abdomen of male and female *D. willistoni* [60] shows

that one Ago2 paralogue (FBgn0212615) is expressed ubiquitously, while the other (FBgn0226485) is

expressed only in the male abdomen (S6 Figure), consistent with the evolution of testis-specificity

after duplication. This raises the possibility that a testis-specific selection pressure may be driving the

retention and specialization of Ago2 paralogues across the arthropods.

In conclusion, we have identified rapid and repeated evolution of testis-specificity after the duplication

of Ago2 in the *obscura* group, associated with low genetic diversity and signatures of strong selection.

Ago2 and other RNAi genes have undergone frequent expansions in different eukaryotic lineages

[4,61], and have been shown to switch between ubiquitous and germline- or ovary-specific functions

in isolated species. This study provides evidence for the evolution of a new testis-specific RNAi

function, and suggests that positive selection may act on young paralogues to drive the rapid

evolution of novel RNAi mechanisms across the eukaryotes.

Materials and Methods

Identification of Ago2 homologues in the Drosophilidae

We used tBLASTx to identify Ago2 homologues in transcriptomes and genomes of 39 species of the

Drosophilidae, using previously-characterised Ago2 from the closest possible relative to provide the

query for each species. If blast returned partial hits, we aligned all hits from the target species to all

Argonautes from the query species, and assigned hits to the appropriate Ago lineage based on a

neighbour-joining tree. For each query sequence, we then manually curated partial blast hits into

complete genes using Geneious v5.6.2 (http://www.geneious.com [62]) (see Supplementary Materials

for sequence accessions).

Additionally, we used degenerate PCR to identify Ago2 paralogues in *D. azteca* and *D. affinis*, and

paralogue-specific PCR with a touchdown amplification cycle to validate the Ago2 paralogues

identified in *D. subobscura*, *D. obscura* and *D. pseudoobscura*. For each reaction, unincorporated

primers were removed with ExonucleaseI (New England Biolabs) and 5' phosphates were removed

with Antarctic Phosphatase (NEB), the PCR products were sequenced by Edinburgh Genomics using

BigDye V3 reagents on a capillary sequencer (Applied Biosystems), and Sanger sequence reads

were trimmed and assembled using Geneious v.5.6.2 (http://www.geneious.com [62]). We also used

a combination of PCR and blast searches to locate *D. pseudoobscura* Ago2a1 & Ago2a3, which lie on

the unplaced "Unknown_contig_265" in release 3.03 of the *D. pseudoobscura* genome (all PCR

primers are detailed in S4 Table)*.

<u>Phylogenetic analysis of drosophilid Ago2 paralogues</u>

To characterise the evolutionary relationships between Ago2 homologues in the Drosophilidae, we

aligned sequences using translational MAFFT [63] with default parameters. We noted that there is a

high degree of codon usage bias in *D. pseudoobscura* Ago2e (effective number of codons

(ENC)=34.24) and *D. obscura* Ago2e (ENC=40.36), and a lesser degree in *D. subobscura* Ago2f

(ENC=45.63) and *D. obscura* Ago2f (ENC=48.39). To reduce the impact of codon usage bias, which

disproportionately affects synonymous sites, we stripped all third positions [64]. We then inferred a

gene tree using the Bayesian approach implemented in BEAST v1.8.1 [65] under a nucleotide model,

assuming a GTR substitution model, variation between sites modelled by a gamma distribution with

four categories, and base frequencies estimated from the data. We used the default priors for all

parameters, except tree shape (for which we specified a birth-death speciation model) and the date of

the *Drosophila-Sophophora* split. To estimate a timescale for the tree, we specified a normal

distribution for the date of this node using values based on mutation rate estimates in [66], with a

mean value of 32mya, standard deviation of 7mya, and lower and upper bounds of 15mya and 50mya

respectively. We ran the analysis for 50 million steps, recording samples from the posterior every

1,000 steps, and inferred a maximum clade credibility tree with TreeAnnotator v1.8.1 [65]. Note that

precise date estimates are not a primary focus of this study, but that other calibrations [67,68] would

lead to more ancient estimates of divergence, and thus stronger evidence for selective maintenance.

 <u>Domain architecture and structural modelling of Ago2 paralogues in the *obscura* group</u>

To infer the location of each domain in each paralogue identified in *D. subobscura*, *D. obscura* and *D.*

pseudoobscura, we searched the Pfam database [69]. To test for structural differences between the

D. pseudoobscura paralogues, we built structural models of each paralogue based on the published

X-ray crystallographic structure of human Ago2 [70]. We used the MODELER software in the

Discovery Studio 4.0 Modeling Environment (Accelrys Software Inc., San Diego, 2013) to calculate

ten models, selected the most energetically favourable for each protein, and assessed model quality

with the 3D-profile option in the software. To assess variation in selective pressure across the

structure of each paralogue, we mapped polymorphic residues onto each structure using PyMol

v.1.7.4.1 (Schrödinger, LLC).

<u>Quantification of virus-induced expression of Ago2 paralogues</u>

We exposed 48-96hr post-eclosion virgin males and females of *D. melanogaster*, *D. subobscura*, *D.*

obscura and *D. pseudoobscura* to Drosophila C virus (DCV), by puncturing the thorax with a pin

contaminated with DCV at a dose of approximately 4×10^7 $TCID^{50}$ per ml. Infection with DCV using this

method has previously been shown to lead to a rapid and ultimately fatal increase in DCV titre in *D.*

melanogaster and *obscura* group species [29]. All flies were incubated at 18C on a 12L:12D light

cycle, with *D. melanogaster* on Lewis medium and *D. subobscura*, *D. obscura* and *D. pseudoobscura*

on banana medium. We sampled 4-7 individuals per species at 0, 8, 16, 24, 48 and 72 hours post

infection. At each time-point we extracted RNA using TRIzol reagent (Ambion) and a

chloroform/isopropanol extraction, treated twice with TURBO DNase (Ambion), and reverse-

transcribed using M-MLV reverse transcriptase (Promega) primed with random hexamers. We then

quantified the expression of Ago2 paralogues in these samples by qPCR, using Fast Sybr Green

(Applied Biosystems) and custom-designed paralogue-specific qPCR primer pairs (see Table S6 for

primer sequences). Due to their high level of sequence similarity (99.9% identity), no primer pair could

distinguish between *D. pseudoobscura* Ago2a1 and Ago2a3, so these two genes are presented

together as "Ago2a". All qPCR reactions for each sample were run in duplicate, and scaled to the

internal reference gene Ribosomal Protein L32 (RpL32). To capture the widest possible biological

variation, the three biological replicates for each species each used a different wild-type genetic

background (see S3 Table for backgrounds used).

 Quantification of Ago2 paralogue expression in different tissues and life stages

For *D. subobscura*, *D. obscura* and *D. pseudoobscura*, we extracted RNA from the head,

testis/ovaries and carcass of 48-96hr post-eclosion virgin adults, with males and females extracted

separately. Each sample consisted of 8-15 individuals in *D. subobscura*, 10 individuals in *D. obscura*

and 15 individuals in *D. pseudoobscura*. We then used qPCR to quantify the expression of each Ago2

paralogue in each tissue, with two technical replicates per sample (reagents, primers and cycling

conditions as above). We carried out five replicates per species, each using a different wild-type

background (see S3 Table for details of backgrounds used). To provide an informal comparison with

the expression pattern of Ago2 before duplication (an "ancestral" expression pattern), we used the

BPKM (bases per kilobase of gene model per million mapped bases) values for Ago2 calculated from

RNA-seq data from the body (carcass and digestive system), head, ovary and testis of 4 day old *D.*

melanogaster adults by [71], scaling each BPKM value to the value for RpL32 in each tissue. Due to

the design of that experiment, the body data are derived from pooled samples of males and females

[71].

To quantify expression of Ago2 paralogues in *D. pseudoobscura* embryos, we collected eggs within

30 minutes of laying, and used qPCR to measure the expression of each Ago2 paralogue (reagents

and primers as above) in two separate wild-type genetic backgrounds (MV8 and MV10). As above,

we estimated an ancestral expression pattern of Ago2 before duplication from the BPKM values for

Ago2 in 0-2hr old *D. melanogaster* embryos according to [71], scaled to the BPKM value for RpL32 in

embryos. To determine any changes in the expression of other *D. pseudoobscura* Argonautes (Ago1,

Ago3, Aub & Piwi) that are associated with Ago2 duplication, we measured their expression in adult

tissues and embryos as detailed above, and compared this with the expression of the Argonautes in

D. melanogaster as measured by [71].

Testing for evolutionary rate changes associated with tissue-specificity of Ago2

We used codeml (PAML, Yang 1997) to fit variants of the M0 model (a single dn/ds ratio, ω) to the 65

drosophilid Ago2 homologues shown in Figure 1. In contrast to the tree topology, which was based on

1st and 2nd positions only, the alignment for the codeml analysis included all positions. To compare the

evolutionary rates of ubiquitously expressed and testis-specific Ago2 paralogues, we fitted a model

specifying one ω for the Ago2 paralogues that were shown to be testis-specific by qPCR, and another

ω for the rest of the tree. We also fitted two models to account for rate variation between the *obscura*

group Ago2 subclades. The first model specified a separate ω for the Ago2a subclade, the Ago2e

subclade, the Ago2f subclade and the rest of the tree. The second model additionally incorporated an

extra ω specified for the *D. pseudoobscura-D. persimilis* Ago2a-Ago2b subclade (which is testis-

specific, in contrast with the rest of the *obscura* group Ago2a subclade). We used Akaike weights to

assess which model provided the best fit to the data, given the number of parameters.

<u>Sequencing of Ago2 paralogue haplotypes from *D. subobscura*, *D. obscura* and *D. pseudoobscura*</u>

To gain genotype data for the Ago2 paralogues in *D. subobscura*, *D. obscura* and *D. pseudoobscura*,

we sequenced the Ago2 paralogues from six males and six females of each species, each from a

different wild-collected line (detailed in S3 Table, sequence polymorphism data in S4 Appendix). We

extracted genomic DNA from each individual using the DNeasy Blood and Tissue kit (Qiagen), and

amplified and Sanger sequenced each Ago2 paralogue from each individual (reagents and PCR

primers as above, sequencing primers detailed in S5 Table). We trimmed and assembled Sanger

sequence reads using Geneious v.5.6.2 (http://www.geneious.com [62]), and identified polymorphic

sites by eye. After sequencing Ago2a (annotated as a single gene in the *D. pseudoobscura* genome),

we discovered two very recent Ago2a paralogues (Ago2a1 & Ago2a3), both of which had been cross-

amplified. For each *D. pseudoobscura* individual we therefore re-sequenced Ago2a3 using one primer

targeted to its neighbouring locus GA22965, and used this sequence to resolve polymorphic sites in

the Ago2a1/Ago2a3 composite sequence, thereby gaining both genotypes for each individual. For

each Ago2 paralogue, we inferred haplotypes from these sequence data using PHASE [72], apart

from the X-linked paralogues (Ago2a1, Ago2a3 & Ago2d) in *D. pseudoobscura* males, for which

phase was obtained directly from the sequence data. The hemizygous haploid X-linked sequenced

were used in phase inference, and should substantially improve the inferred phasing of female

genotypes.

To quantify differences between paralogues in their population genetic characteristics, we aligned

haplotypes using translational MAFFT [63], and used DnaSP v.5.10.01 [73] to calculate the following

summary statistics for each Ago2 paralogue: π (pairwise diversity, with Jukes-Cantor correction as

described in [74]) at nonsynonymous (π_a) and synonymous (π_s) sites, Tajima's *D* [75] and the

effective number of codons (ENC) [76]. To compare the ENC for each gene with the genome as a

whole, we used codonW v1.4.2 [77] to calculate the ENC for the longest ORF from each gene or

transcript in the genomes of *D. subobscura*, *D. obscura* and *D. pseudoobscura* (ORF sets detailed

below). In each species, we then compared the ENC values of each Ago2 paralogue with this

genome-wide ENC distribution.

<u>Testing for positive selection on Ago2 paralogues in the *obscura* group</u>

We used McDonald-Kreitman (MK) tests [31] to test for positive selection on each Ago2 paralogue.

For each paralogue, we chose an outgroup with divergence at synonymous sites (K_S) in the range

0.1-0.2 where possible. However, the prevalence of duplications and losses of Ago2 paralogues in the

obscura group meant that for some tests a suitably divergent extant outgroup sequence did not exist.

In these cases, we reconstructed hypothetical ancestral sequences using the M0 model in PAML [30].

To assess the effect of these outgroup choices on our results, we repeated each test with another

outgroup, and found no effect of outgroup choice on the significance of any tests, and only marginal

differences in estimates of α and ω_α (results of tests using primary and alternative outgroups are

detailed in S1 & S2 Tables).

A complementary approach to identifying positive selection is to test for reduced diversity at a locus

compared with the genome as a whole. To compare the diversity of each *D. pseudoobscura* Ago2

paralogue with the genome-wide distribution of synonymous site diversity, we used genomic data for

12 lines generated by [36]. We mapped short reads to the longest ORF for each gene in the R3.2

gene set using Bowtie2 v2.1.0 [78], and estimated synonymous site diversity (θ_W based on fourfold

synonymous sites) at each ORF using PoPoolation [79]. We then plotted the distribution of

synonymous site diversity, limited to genes in the size range of 0.75kb - 3kb for comparability with the

Ago2 paralogues, and compared the fourfold synonymous site diversity levels of each *D.*

pseudoobscura Ago2 paralogue with this distribution. Some *D. pseudoobscura* paralogues are

located on autosomes (Ago2b, Ago2c & Ago2e) and some on the X chromosome (Ago2a1, Ago2a3 &

Ago2d). Therefore, because of the different population genetic expectations for autosomal and X-

linked genes [80], we examined separate distributions for autosomal and X-linked genes. To provide

an additional test for reduced diversity at *D. pseudoobscura* Ago2 paralogues, we performed

maximum-likelihood Hudson-Kreitman-Aguadé tests [33], using divergence from *D. affinis* and

intraspecific polymorphism data for 84 *D. pseudoobscura* loci generated by [81]. We performed 63

tests to encompass all one, two, three, four, five and six-way combinations of the paralogues, and

calculated Akaike weights from the resulting likelihood estimates to provide an estimate of the level of

support for each combination.

To infer a genome-wide distribution of synonymous site diversity for *D. obscura* and *D. subobscura*,

for which genomic data are unavailable, we used pooled transcriptome data from wild-collected adult

male flies that had previously been generated for surveys of RNA viruses [25,34]. To generate a *de*

novo transcriptome for each species, we assembled short reads with Trinity r20140717 [82]. For each

species, we mapped short reads from the pooled sample to the longest ORF for each transcript,

estimated synonymous site diversity at each locus using PoPoolation [79], and plotted the distribution

of diversity (as described above for *D. pseudoobscura*). The presence of heterozygous sites in males

(identified by Sanger sequencing) confirmed that all Ago2 paralogues in *D. subobscura* and *D.*

obscura are autosomal: we therefore compared the synonymous site diversity for these paralogues

with the autosomal distribution, and do not show the distributions for putatively X-linked genes. Our

use of transcriptome data for *D. obscura* and *D. subobscura* will bias the resulting diversity

distributions in three ways. First, variation in expression level will cause individuals displaying high

levels of expression to be over-represented among reads, downwardly biasing diversity. Second,

highly expressed genes are easier to assemble, and highly expressed genes tend to display lower

genetic diversity [83,84]. Third, high-diversity genes are harder to assemble, *per se*. However, as all

three biases will tend to artefactually reduce diversity in the genome-wide dataset relative to Ago2,

this makes our finding that Ago2 paralogues display unusually low diversity conservative.

Identifying selective sweeps in Ago2 paralogues of *D. pseudoobscura*

To test whether the unusually low diversity seen in the *D. pseudoobscura* Ago2 paralogues is due to

recent selection or generally reduced diversity in that region of the genome, we compared diversity at

each paralogue to diversity in their neighbouring regions. We obtained sequence data for the 50kb

either side of each of these paralogues from the 11 whole genomes detailed in [36]. Note that the very

high similarity of these Ago2 paralogues means that they cannot be accurately assembled from short

read data, and are not present in the data from [36]. For each genome, we therefore replaced the

poorly-assembled region corresponding to the paralogue with one of our own Sanger-sequenced

haplotypes, making a set of 11 ca. 102kb sequences for each paralogue. We aligned these

sequences using PRANK [85] with default settings, and calculated Watterson's θ at all sites in a

sliding window across each alignment, with a window size of 5kb and a step of 1kb. For Ago2a1 and

Ago2a3, which are located in tandem, we analysed the same genomic region. Since our Ago2

haplotypes were sampled from a different North American population of *D. pseudoobscura* to those of

[36], an apparent reduction in local diversity might result from differences in diversity between the two

populations. Therefore, we also repeated these analyses on a dataset in which our Sanger

sequenced haplotypes were removed, leaving missing data.

To test explicitly for selective sweeps at each region, we used Sweepfinder [86] to calculate the

likelihood and location of a sweep in or near each Ago2 paralogue. We specified a grid size of 20,000,

a folded frequency spectrum for all sites, and included invariant sites. To infer the significance of any

observed peaks in the composite likelihood ratio, we used ms [87] to generate 1000 samples of 11

sequences under a neutral coalescent model. We generated separate samples for each region

surrounding an Ago2 paralogue, conditioning on the number of polymorphic sites observed in that

region, the sequence length equal to the alignment length, and an effective population size at 10^6

(based on a previous estimate for *D. melanogaster* by [88]). We specified the recombination rate at

5cM/Mb, a conservative value based on previous estimates for *D. pseudoobscura* [36], which will lead

to larger segregating linkage groups and therefore a more stringent significance threshold.

<u>Acknowledgements</u>

This work was supported by a Natural Environment Research Council Doctoral Training Grant (NERC

DG NE/J500021/1 to SHL), the Academy of Finland (265971 to HS), a University of Edinburgh

Chancellor's Fellowship and a Wellcome Trust Research Career Development Fellowship (WT085064

to DJO), and a Wellcome Trust strategic award to the Centre for Immunity, Infection and Evolution

(WT095831 to the CIIE). We thank Ben Longdon and Brian Charlesworth for providing us with strains

of *D. obscura* and *D. pseudoobscura* respectively, and Francis Jiggins for providing us with DCV.

References

1. Meister G. Argonaute proteins: functional insights and emerging roles. Nat Rev Genet.
2013;14: 447–59.

2. Singh RK, Gase K, Baldwin IT, Pandey SP. Molecular evolution and diversification of the
Argonaute family of proteins in plants. BMC Plant Biol. 2015;15: 1–16.

3. Buck AH, Blaxter M. Functional diversification of Argonautes in nematodes: an expanding
universe. 2013;41: 881–6.

4. Lewis SH, Salmela H, Obbard DJ. Duplication and diversification of Dipteran Argonaute genes,
and the evolutionary divergence of Piwi and Aubergine. Genome Biol Evol. 2016; Advance Ac:
1–30.

5. Lu H-L, Tanguy S, Rispe C, Gauthier J-P, Walsh T, Gordon K, et al. Expansion of genes
encoding piRNA-associated argonaute proteins in the pea aphid: diversification of expression
profiles in different plastic morphs. PLoS One. 2011;6: e28051.

6. Leebonoi W, Sukthaworn S, Panyim S, Udomkit A. A novel gonad-specific Argonaute 4 serves
as a defense against transposons in the black tiger shrimp Penaeus monodon. Fish Shellfish
Immunol. 2015;42: 280–288.

7. Miesen P, Girardi E, van Rij RP. Distinct sets of PIWI proteins produce arbovirus and
transposon-derived piRNAs in Aedes aegypti mosquito cells. Nucleic Acids Res. 2015;43:
6545–56.

8. Cerutti H, Casas-Mollano JA. On the origin and functions of RNA-mediated silencing: from
protists to man. Curr Genet. 2006;50: 81–99.

9. Ross RJ, Weiner MM, Lin H. PIWI proteins and PIWI-interacting RNAs in the soma. Nature.
2014;505: 353–9.

10. Luteijn MJ, Ketting RF. PIWI-interacting RNAs: from generation to transgenerational
epigenetics. Nat Rev Genet. 2013;14: 523–34.

11. Sienski G, Dönertas D, Brennecke J. Transcriptional silencing of transposons by Piwi and
maelstrom and its impact on chromatin state and gene expression. Cell. 2012;151: 964–980.

12. Kolaczkowski B, Hupalo DN, Kern AD. Recurrent adaptation in RNA interference genes across

the Drosophila phylogeny. Mol Biol Evol. 2011;28: 1033–1042.

13.   Obbard DJ, Gordon KHJ, Buck AH, Jiggins FM. The evolution of RNAi as a defence against

viruses and transposable elements. Philos Trans R Soc London Biol Sci. 2009;364: 99–115.

14.   Eulalio A, Huntzinger E, Izaurralde E. Getting to the Root of miRNA-Mediated Gene Silencing.

Cell. 2008;132: 9–14.

15.   Obbard DJ, Jiggins FM, Halligan DL, Little TJ. Natural selection drives extremely rapid

evolution in antiviral RNAi genes. Curr Biol. 2006;16: 580–5.

16.   Wen J, Duan H, Bejarano F, Okamura K, Fabian L, Brill JA, et al. Adaptive Regulation of

Testis Gene Expression and Control of Male Fertility by the Drosophila Harpin RNA Pathway.

Mol Cell. 2015;57: 165–178.

17.   Menon DU, Meller VH. A role for siRNA in X-chromosome dosage compensation in Drosophila

melanogaster. Genetics. 2012;191: 1023–8.

18.   Li H, Li WX, Ding SW. Induction and suppression of RNA silencing by an animal virus.

Science. 2002;296: 1319–1321.

19.   van Rij RP, Saleh M-C, Berry B, Foo C, Houk A, Antoniewski C, et al. The RNA silencing

endonuclease Argonaute 2 mediates specific antiviral immunity in Drosophila melanogaster.

Genes Dev. 2006;20: 2985–95.

20.   Czech B, Malone CD, Zhou R, Stark A, Schlingeheyde C, Dus M, et al. An endogenous small

interfering RNA pathway in Drosophila. Nature. 2008;453: 798–802.

21.   Chung W-J, Okamura K, Martin R, Lai EC. Endogenous RNA interference provides a somatic

defense against Drosophila transposons. Curr Biol. 2008;18: 795–802.

22.   Obbard DJ, Welch JJ, Kim K-W, Jiggins FM. Quantifying adaptive evolution in the Drosophila

immune system. PLoS Genet. 2009;5: e1000698.

23.   Obbard DJ, Jiggins FM, Bradshaw NJ, Little TJ. Recent and recurrent selective sweeps of the

antiviral RNAi gene Argonaute-2 in three species of Drosophila. Mol Biol Evol. 2011;28: 1043–

56.

24.   Marques JT, Carthew RW. A call to arms: coevolution of animal viruses and host innate

immune responses. Trends Genet. 2007;23: 359–364.

25.   van Mierlo JT, Overheul GJ, Obadia B, van Cleef KWR, Webster CL, Saleh M-C, et al. Novel

582   Drosophila Viruses Encode Host-Specific Suppressors of RNAi. PLoS Pathog. 2014;10:

583   e1004256.

584   26.   Hain D, Bettencourt BR, Okamura K, Csorba T, Meyer W, Jin Z, et al. Natural variation of the

585   amino-terminal glutamine-rich domain in Drosophila argonaute2 is not associated with

586   developmental defects. PLoS One. 2010;5: e15264.

587   27.   Altschul SF, Madden TL, Schaffer AA, Zhang J, Zhang Z, Miller W, et al. Gapped BLAST and

588   PSI-BLAST: a new generation of protein database search programs. Nucleic Acids Res.

589   1997;25: 3389–3402.

590   28.   Aliyari R, Wu Q, Li H-W, Wang X-H, Li F, Green LD, et al. Mechanism of induction and

591   suppression of antiviral immunity directed by virus-derived small RNAs in Drosophila. Cell Host

592   Microbe. 2008;4: 387–97.

593   29.   Longdon B, Hadfield JD, Day JP, Smith SCL, McGonigle JE, Cogni R, et al. The causes and

594   consequences of changes in virulence following pathogen host shifts. PLoS Pathog. 2015;11:

595   e1004728.

596   30.   Yang Z. PAML: a program package for phylogenetic analysis by maximum likelihood. Comput

597   Appl Biosci. 1997;13: 555–6.

598   31.   McDonald JH, Kreitman M. Adaptive protein evolution at the Adh locus in Drosophila. Nature.

599   1991;351: 652–654.

600   32.   Maynard Smith J, Haigh J. The hitch-hiking effect of a favourable gene. Genet Res. 1974;23:

601   23–35.

602   33.   Wright SI, Charlesworth B. The HKA test revisited: a maximum-likelihood-ratio test of the

603   standard neutral model. Genetics. 2004;168: 1071–6.

604   34.   Webster CL, Longdon B, Lewis SH, Obbard DJ. Twenty five new viruses associated with the

605   Drosophilidae (Diptera); 2016. Preprint. Available: http://dx.doi.org/10.1101/041665. Accessed

606   21 March 2016.

607   35.   Nielsen R, Bustamante C, Clark AG, Glanowski S, Sackton TB, Hubisz MJ, et al. A scan for

608   positively selected genes in the genomes of humans and chimpanzees. PLoS Biol. 2005;3:

e170.

36.   McGaugh SE, Heil CSS, Manzano-Winkler B, Loewe L, Goldstein S, Himmel TL, et al.

Recombination modulates how selection affects linked sites in Drosophila. PLoS Biol. 2012;10:

e1001422.

37.   Celniker SE, Dillon LAL, Gerstein MB, Gunsalus KC, Henikoff S, Karpen GH, et al. Unlocking

the secrets of the genome. Nature. 2009;459: 927–930.

38.   Zambon RA, Nandakumar M, Vakharia VN, Wu LP. The Toll pathway is important for an

antiviral response in Drosophila. Proc Natl Acad Sci. 2005;102: 7257–62.

39.   Dostert C, Jouanguy E, Irving P, Troxler L, Galiana-Arnoux D, Hetru C, et al. The Jak-STAT

signaling pathway is required but not sufficient for the antiviral response of drosophila. Nat

Immunol. 2005;6: 946–953.

40.   Bronkhorst AW, van Rij RP. The long and short of antiviral defense: small RNA-based

immunity in insects. Curr Opin Virol. 2014;7C: 19–28.

41.   Assis R, Bachtrog D. Neofunctionalization of young duplicate genes in Drosophila. Proc Natl

Acad Sci. 2013;110: 17409–14.

42.   Assis R, Bachtrog D. Rapid divergence and diversification of mammalian duplicate gene

functions. BMC Evol Biol. 2015;15: 138.

43.   Kaessmann H. Origins, evolution, and phenotypic impact of new genes. Genome Res.

2010;20: 1313–26.

44.   Conrad T, Akhtar A. Dosage compensation in Drosophila melanogaster: epigenetic fine-tuning

of chromosome-wide transcription. Nat Rev Genet. 2012;13: 123–134.

45.   Longdon B, Jiggins FM. Vertically transmitted viral endosymbionts of insects: do sigma viruses

walk alone? Proc R Soc B. 2012;279: 3889–3898.

46.   Rozhkov N V, Aravin AA, Zelentsova ES, Schostak NG, Sachidanandam R, McCombie WR, et

al. Small RNA-based silencing strategies for transposons in the process of invading Drosophila

species. RNA. 2010;16: 1634–45.

47.   Pasyukova E, Nuzhdin S, Li W, Flavell AJ. Germ line transposition of the copia

retrotransposon in Drosophila melanogaster is restricted to males by tissue-specific control of

copia RNA levels. Mol Gen Genet. 1997;255: 115–124.

48.   Morozova T V, Tsybulko EA, Pasyukova EG. Regularory elements of the copia

retrotransposon determine different levels of expression in different organs of males and

females of Drosophila melanogaster. Genetika. 2009;45: 169–177.

49.   Gershenson S. A New Sex-Ratio Abnormality in Drosophila obscura. Genetics. 1928;13: 488–

507.

50.   Sturtevant AH, Dobzhansky T. Geographical Distribution and Cytology of "Sex Ratio" in

Drosophila Pseudoobscura and Related Species. Genetics. 1936;21: 473–490.

51.   Wu CI, Beckenbach AT. Evidence for extensive genetic differentiation between the sex-ratio

and the standard arrangement of Drosophila pseudobscura and D. persimilis and identification

of hybrid sterility factors. Genetics. 1983;105: 71–86.

52.   Jaenike J. Sex chromosome meiotic drive. Annu Rev Ecol Syst. 2001;32: 25–49.

53.   Unckless RL, Larracuente AM, Clark AG. Sex-ratio meiotic drive and Y-linked resistance in

Drosophila affinis. Genetics. 2015;199: 831–40.

54.   Tao Y, Araripe L, Kingan SB, Ke Y, Xiao H, Hartl DL. A sex-ratio meiotic drive system in

Drosophila simulans. II: An X-linked distorter. PLoS Biol. 2007;5: 2576–2588.

55.   Kotelnikov RN, Klenov MS, Rozovsky YM, Olenina L V., Kibanov M V., Gvozdev V a.

Peculiarities of piRNA-mediated post-transcriptional silencing of Stellate repeats in testes of

Drosophila melanogaster. Nucleic Acids Res. 2009;37: 3254–3263.

56.   Gell SL, Reenan RA. Mutations to the piRNA pathway component aubergine enhance meiotic

drive of segregation distorter in Drosophila melanogaster. Genetics. 2013;193: 771–784.

57.   Scott JG, Warren WC, Beukeboom LW, Bopp D, Clark AG, Giers SD, et al. Genome of the

house fly, Musca domestica L., a global vector of diseases with adaptations to a septic

environment. Genome Biol. 2014;15: 466–482.

58.   Palmer WJ, Jiggins FM. Comparative Genomics Reveals the Origins and Diversity of

Arthropod Immune Systems. Mol Biol Evol. 2015;32: 2111–2129.

59.   Meisel RP, Scott JG, Clark AG. Transcriptome Differences between Alternative Sex

Determining Genotypes in the House Fly, Musca domestica. Genome Biol Evol. 2015;7: 2051–

2061.

60.   Meisel RP, Malone JH, Clark AG. Disentangling the relationship between sex-biased gene
667   expression and X-linkage. Genome Res. 2012;22: 1255–1265.

668   61.   Mukherjee K, Campos H, Kolaczkowski B. Evolution of animal and plant dicers: early parallel
669   duplications and recurrent adaptation of antiviral RNA binding in plants. Mol Biol Evol.
670   2013;30: 627–41.

671   62.   Kearse M, Moir R, Wilson A, Stones-Havas S, Cheung M, Sturrock S, et al. Geneious Basic:
672   An integrated and extendable desktop software platform for the organization and analysis of
673   sequence data. Bioinformatics. 2012;28: 1647–1649.

674   63.   Katoh K, Misawa K, Kuma K, Miyata T. MAFFT : a novel method for rapid multiple sequence
675   alignment based on fast Fourier transform. Nucleic Acids Res. 2002;30: 3059–3066.

676   64.   Behura SK, Severson DW. Codon usage bias: causative factors, quantification methods and
677   genome-wide patterns: with emphasis on insect genomes. Biol Rev. 2013;88: 49–61.

678   65.   Drummond AJ, Suchard MA, Xie D, Rambaut A. Bayesian phylogenetics with BEAUti and the
679   BEAST 1.7. Mol Biol Evol. 2012;29: 1969–1973.

680   66.   Obbard DJ, MacLennan J, Kim KW, Rambaut A, O'Grady PM, Jiggins FM. Estimating
681   divergence dates and substitution rates in the Drosophila phylogeny. Mol Biol Evol. 2012;29:
682   3459–3473.

683   67.   Russo C a, Takezaki N, Nei M. Molecular phylogeny and divergence times of Drosophilid
684   species. Mol Biol Evol. 1995;12: 391–404.

685   68.   Tamura K. Temporal Patterns of Fruit Fly (Drosophila) Evolution Revealed by Mutation Clocks.
686   Mol Biol Evol. 2004;21: 36–44.

687   69.   Finn RD, Mistry J, Tate J, Coggill P, Heger a., Pollington JE, et al. The Pfam protein families
688   database. Nucleic Acids Res. 2009;38: D211–D222.

689   70.   Schirle NT, Macrae IJ. The Crystal Structure of Human Argonaute2. Science. 2012;336: 1037–
690   1040.

691   71.   Brown JB, Boley N, Eisman R, May GE, Stoiber MH, Duff MO, et al. Diversity and dynamics of
692   the Drosophila transcriptome. Nature. 2014;512: 393–399.

72.   Stephens M, Smith NJ, Donnelly P. A new statistical method for haplotype reconstruction from
population data. Am J Hum Genet. 2001;68: 978–989.

73.   Librado P, Rozas J. DnaSP v5: a software for comprehensive analysis of DNA polymorphism
data. Bioinformatics. 2009;25: 1451–2.

74.   Lynch M, Crease TJ. The analysis of population survey data on DNA sequence variation. Mol
Biol Evol. 1990;7: 377–394.

75.   Tajima F. Statistical method for testing the neutral mutation hypothesis by DNA polymorphism.
Genetics. 1989;123: 585–595.

76.   Wright F. The "effective number of codons" used in a gene. Gene. 1990;87: 23–29.

77.   Peden J. Analysis of codon usage bias. PhD Thesis, The University of Nottingham. 1995.
Available:
http://citeseerx.ist.psu.edu/viewdoc/download?doi=10.1.1.186.1796&rep=rep1&type=pdf

78.   Langmead B, Trapnell C, Pop M, Salzberg SL. Ultrafast and memory-efficient alignment of
short DNA sequences to the human genome. Genome Biol. 2009;10: R25.

79.   Kofler R, Orozco-terWengel P, De Maio N, Pandey RV, Nolte V, Futschik A, et al. PoPoolation:
a toolbox for population genetic analysis of next generation sequencing data from pooled
individuals. PLoS One. 2011;6: e15925.

80.   Vicoso B, Charlesworth B. Evolution on the X chromosome: unusual patterns and processes.
Nat Rev Genet. 2006;7: 645–653.

81.   Haddrill PR, Loewe L, Charlesworth B. Estimating the parameters of selection on
nonsynonymous mutations in Drosophila pseudoobscura and D. miranda. Genetics. 2010;185:
1381–96.

82.   Grabherr MG, Haas BJ, Yassour M, Levin JZ, Thompson DA, Amit I, et al. Full-length
transcriptome assembly from RNA-Seq data without a reference genome. Nat Biotechnol.
2011;29: 644–52.

83.   Pal C, Papp B, Hurst LD. Highly expressed genes in yeast evolve slowly. Genetics. 2001;158:
927–931.

84.   Lemos B, Bettencourt BR, Meiklejohn CD, Hartl DL. Evolution of proteins and gene expression

levels are coupled in Drosophila and are independently associated with mRNA abundance,

protein length, and number of protein-protein interactions. Mol Biol Evol. 2005;22: 1345–1354.

85. Löytynoja A, Goldman N. An algorithm for progressive multiple alignment of sequences with

insertions. Proc Natl Acad Sci. 2005;102: 10557–10562.

86. Nielsen R, Williamson S, Kim Y, Hubisz MJ, Clark AG, Bustamante C. Genomic scans for

selective sweeps using SNP data. Genome Res. 2005;15: 1566–75.

87. Hudson RR. Generating samples under a Wright-Fisher neutral model of genetic variation.

Bioinformatics. 2002;18: 337–338.

88. Li H, Stephan W. Inferring the demographic history and rate of adaptive substitution in

Drosophila. PLoS Genet. 2006;2: 1580–1589.

89. Anders S, Pyl PT, Huber W. HTSeq - A Python framework to work with high-throughput

sequencing data. Bioinformatics. 2015;2: 166–169.

90. Mortazavi A, Williams BA, McCue K, Schaeffer L, Wold B. Mapping and quantifying

mammalian transcriptomes by RNA-Seq. Nat Methods. 2008;5: 621–628.

91. Schaeffer SW, Bhutkar A, McAllister BF, Matsuda M, Matzkin LM, O'Grady PM, et al. Polytene

chromosomal maps of 11 Drosophila species: the order of genomic scaffolds inferred from

genetic and physical maps. Genetics. 2008;179: 1601–55.

92. Segarra C, Aguadé M. Molecular organization of the X chromosome in different species of the

obscura group of Drosophila. Genetics. 1992;130: 513–521.

 Supporting Information Captions

S1 Figure: The expression of *D. pseudoobscura* Ago2 paralogues in embryos. Error bars

indicate 1 standard error estimated from 2 technical replicates in each of two different genetic

backgrounds. *D. melanogaster* expression levels were taken from a single publicly-available RNA-seq

experiment [71]. Ago2c is highly expressed in embryos, but none of the testis-specific Ago2

paralogues (Ago2a, Ago2b & Ago2e) are expressed.

S2 Figure: The tissue-specific expression patterns of other members of the Argonaute gene

family (Ago1, Ago3, Aub & Piwi) in *D. melanogaster* and *D. pseudoobscura*. For *D.*

pseudoobscura embryo, error bars indicate 1 standard error estimated from 2 technical replicates in

each of two different genetic backgrounds. For all other *D. pseudoobscura* tissues, error bars indicate

1 standard error estimated from 2 technical replicates in each of five different genetic backgrounds. *D.*

melanogaster expression levels were taken from a single RNA-seq experiment [71]. In *D.*

pseudoobscura, Ago1 is expressed in all tissues, but the other genes are only expressed in the

embryo and germline.

S3 Figure: The distribution of synonymous site diversity across genes, derived from genome

(*D. pseudoobscura*) or transcriptome (*D. subobscura* & *D. obscura*) data. The percentile of the

distribution into which each paralogue falls is indicated in brackets under the paralogue name. In each

species, members of the Ago2a and Ago2e subclades have very low diversity compared with the

genome as a whole.

S4 Figure: The distribution of codon usage bias, derived from genome (*D. pseudoobscura*) or

transcriptome (*D. subobscura* & *D. obscura*) data. The percentile of the distribution into which

each paralogue falls is indicated in brackets under the paralogue name. Ago2e has a very low

effective number of codons (ENC) compared with the genome as a whole, indicating a high degree of

codon usage bias.

S5 Figure: Genetic diversity in the regions surrounding each *D. pseudoobscura* Ago2

paralogue, with Ago2 paralogue haplotype sequences removed. After specifying Ago2 paralogue

sequence data as missing information, sharp troughs in diversity remain at Ago2a, Ago2b and Ago2c,

indicating a selective sweep.

S6 Figure: The tissue-specific expression patterns of the Argonaute gene family in *D. willistoni*

and *M domestica*. Transcriptome data for *D. willistoni* were taken from [60], and transcriptome data

for *M. domestica* were taken from [59]. For both species, we mapped reads to coding sequences

using Bowtie 2.1 [78], counted reads mapping to each coding sequence using HTSeq [89], and

converted counts to reads per kilobase per million reads (RPKM [90]) to account for coding sequence

length and sequencing depth. For *M. domestica*, error bars indicate two biological replicates, each in

a different genetic background.

S1 Table: McDonald-Kreitman test results. Pn & Ps are the number of within-species

polymorphisms after singletons have been removed. All values are displayed to 2dp, except ω_a which

is displayed to 4dp.

S2 Table: McDonald-Kreitman test results with alternative outgroups. Pn & Ps are the number of

within-species polymorphisms after singletons have been removed. All values are displayed to 2dp,

except ω_a which is displayed to 4dp.

S3 Table: Genetic backgrounds used in each experiment. Line refers to an individual isofemale

line, and Origin refers to the geographic location where the female who founded that line was caught.

S4 Table: Primers used for PCR and qPCR amplification of Ago2 paralogues. All primers are

displayed in the 5' to 3' direction.

S5 Table: Primers used for Sanger sequencing of Ago2 paralogue haplotypes. All primers are

displayed in the 5' to 3' direction.

S1 Appendix: Sequence alignment of drosophilid Ago2 homologues. This alignment has had all

3[rd] positions stripped, and was used for time-scaled phylogenetic analysis of drosophilid Ago2

evolution.

S2 Appendix: Sequence alignment of drosophilid Ago2 homologues. This alignment has had all

3[rd] positions stripped, and was used for model-based analysis of differential evolutionary rate and

codon-specific positive selection.

S3 Appendix: Sequence metadata for drosophilid Ago2 homologues.

S4 Appendix: Sequence polymorphism data for *D. subobscura*, *D. obscura* and *D.*

***pseudoobscura* Ago2 paralogues**

S5 Appendix: Raw data used to plot Figures 3, 4, 5, S1, S2, S3, S4 & S6.