Modification of plant regeneration medium decreases the time for recovery of *Solanum*

lycopersicum cultivar M82 stable transgenic lines

Sarika Gupta and Joyce Van Eck*

The Boyce Thompson Institute, 533 Tower Road, Ithaca, NY 14853 USA

*Corresponding author, e-mail: jv27@cornell.edu; phone: 607-254-1686; fax: 607-254-1242

Sarika Gupta, e-mail: sarikagupt9@gmail.com

Abstract

Background: Tomato (*Solanum lycopersicum*) has rapidly become a valuable model as a result

of the availability of a high quality reference genome, extensive genetic resources, and efficient

gene transfer methodology. A high-throughput method to obtain transgenic lines sooner than

standard methods has the potential to greatly advance gene function studies.

Results: The goal of this study was to optimize our current gene transfer (transformation)

method for tomato by investigating medium components that would result in a decreased time

for recovery of transgenic lines. The methodology reported here is based on infection of 6-day-

old cotyledon explants from *Solanum lycopersicum* cultivar M82 *in vitro*-grown seedlings with

Agrobacterium tumefaciens strain LBA4404 containing the binary vector pBI121. This vector

contains the β-glucuronidase reporter gene and the neomycin phosphotransferase II selectable

marker gene that confers resistance to kanamycin. Modification of our standard plant

regeneration medium by the addition of indole-3-acetic acid (IAA) at concentrations of either

0.05 mg/l or 0.1 mg/l greatly decreased the time for recovery of transgenic lines by 6 weeks as

compared to our standard medium that contains zeatin alone. Moreover, addition of 1 mg/l

IAA to the root induction medium resulted in faster root development. We observed 50% and

54% transformation efficiency when plant regeneration medium containing 0.05 mg/l and 0.1

mg/l IAA, respectively, was used for transformation experiments. Transgenic lines recovered

with the optimized method were similar in development to our standard method. No negative

effects were observed as a result of the addition of IAA to the medium.

Conclusions: The results showed that addition of IAA at concentrations of either 0.05 or 0.1

mg/l to the plant regeneration medium reduced the total time for recovery of stable transgenic

lines by 6 weeks. The ability to recover transgenic lines in a shorter time results in higher

throughput for the introduction of gene constructs and has the potential to decrease the time

needed to complete investigations of gene function. The ability to recover transgenic lines for

each gene construct in a shorter time saves labor and resource expenses because culture

maintenance is also reduced.

Keywords: *Agrobacterium tumefaciens*, indole-3-acetic acid, *Solanaceae*, *Solanum*

pimpinellifolium, tomato

Background

Tomato, *Solanum lycopersicum*, is a member of the *Solanaceae* family, which contains

approximately 3,000 plant species and includes some of the most economically important food

crops. It is native to South America and was brought to Europe in the 1500s and then to North

America in the 1800s [1]. Tomato is a perennial plant that has two different growth habits,

determinate and indeterminate. There are two different market types of tomatoes, fresh

market and processing. According to the Agricultural Marketing Resource Center, in 2014 the

US dollar value for fresh market tomatoes was 1.14 billion and 1.325 billion for processing

types, which are used to make products such as juice, sauces, and ketchup [2]. In addition to

being an economically important food crop, tomato is an excellent source of health beneficial

nutrients including beta-carotene and lycopene.

Over the years, utilization of tomato as a model plant species has increased because of

readily available resources such as mutant populations [3], bioinformatics tools [4], and a high

quality reference genome [5]. In addition, since the very first report of *Agrobacterium*-

mediated transformation of tomato by McCormick et al. [6], there have been other reports of

successful transformations of different genotypes [7-12]. A key aspect for the adoption of a

model plant species is the availability of efficient transformation methodology. This was

certainly the case for *Arabidopsis*, which is by far the most widely used model for plant research

programs [13].

While there are several methods available for plant transformation, A*grobacterium*

tumefaciens-mediated transformation has become the most extensively used method [14, 15].

Despite its effectiveness for gene transfer in tomato, there is still need for improvement.

Improving methodology to decrease the time from introduction of a gene construct of interest

to recovery of stable transgenics would improve the throughput and shorten the timeframe for

studies that utilize tomato transgenic lines.

We were interested in finding an approach to decrease the time to obtain transgenic lines

of the processing type tomato M82 because this genotype is used for gene function studies in

our lab as well as others [16, 17]. We chose to start by investigating supplementation of our

standard plant regeneration and rooting media with a growth regulator that had the potential

to speed up plant development [12].

Cytokinins and auxins are important hormones that influence growth and developmental

processes in plants. Interactions between cytokinins and auxins have been shown to be

necessary for the shoot apex growth [18, 19]. Auxin has also been shown to play a role in the

specification of the root apical meristem [18, 20, 21]. The hormonal interactions can be utilized

in the area of tissue culture to leverage the presence of the hormones in the medium. In this

study, we report the effects of the addition of the auxin, indole-3-acetic acid (IAA) on the

recovery time of M82 transgenic lines.

Methods

Plant material

Seeds of *Solanum lycopersicum* cv M82 were surface sterilized in 20% (v/v) bleach solution

containing Tween-20 for 20 min followed by 3 rinses in sterile water. Seeds were germinated in

Magenta GA7 boxes (Caisson Labs, Logan, UT) that contained 50 ml of Murashige and Skoog

(MS) [22] (Caisson Labs) based medium containing 2.15 g/l MS salts, 100 mg/l myo-inositol, 2

mg/l thiamine, 0.5 mg/l pyridoxine, 0.5 mg/l nicotinic acid, 10 g/l sucrose and 8 g/l Sigma agar

(Sigma-Aldrich, St. Louis, MO). Cultures were maintained at 24°C under a 16h light/8h dark

photoperiod at 57 – 65 uE m^{-2} s^{-1}.

One day prior to infection with *Agrobacterium*, cotyledon explants and feeder layer plates

were prepared. Feeder layers were prepared before cutting the explants by dispensing 2 ml of

a 1-week-old NT1 suspension culture onto KCMS medium (4.3 g/l MS salts, 100 mg/l myo-

inositol, 1.3 mg/l thiamine, 0.2 mg/l 2,4-dichlorophenoxy acetic acid, 200 mg/l KH$_2$PO$_4$, 0.1 mg/l

kinetin, 30 g/l sucrose, 5.2 g/l Agargel (Sigma Aldrich), pH 6.0.   The suspension was covered

with a sterile 7 cm Whatman filter paper. Explants were excised from 6-day-old seedlings

before the first true leaves emerged. To prepare the explants, seedlings were placed on a

sterile paper towel moistened with sterile water. Cotyledons were excised at the petioles, cut

into approximately 1 cm sections, placed adaxial side down on the KCMS feeder layer plates,

and maintained at 24°C under a 16 h light/8 h dark photoperiod at 57 – 65 uE m^{-2} s^{-1}.

Bacterial strain and binary vector

Electroporation was used to introduce the pBI121 vector into the *Agrobacterium tumefaciens*

strain LBA4404. A single, well-formed colony from the selection plate was transferred to 50 ml

of YEP selective medium that contained 50 mg/l kanamycin and maintained in a shaking

incubator at 28°C for 18 – 24 hrs or the length of time needed to reach an OD$_{600}$ of 0.6 - 0.7.

The *Agrobacterium* suspension was centrifuged at 8000 rpm for 10 min at 20°C. The pellet was

resuspended in 50 ml of 2% MSO medium (4.3 g/l MS salts, 100 mg/l myo-inositol, 0.4 mg/l

thiamine, and 20 g/l sucrose) by vortexing.

Agrobacterium-mediated transformation

Cotyledon explants were incubated in the *Agrobacterium*/2% MSO suspension for 5 min,

transferred to a sterile paper towel to allow excess suspension to briefly drain, placed back

onto the feeder plates with the adaxial sides down, and co-cultivated in the dark at 19°C for 48

hrs. Explants, adaxial side up, were transferred to our standard plant regeneration selective

medium designated 2ZK that contained 4.3 g/l MS salts, 100 mg/l myo-inositol, 1 ml/l Nitsch

vitamins (1000x), 20 g/l sucrose, 2 mg/l trans-zeatin, 75 mg/l kanamycin, 300 mg/l timentin,

and 5.2 g/l Agargel. One week later, the explants were transferred onto 2ZK medium

containing IAA at either 0 mg/l, 0.01 mg/l, 0.05 mg/l, 0.1 mg/l, or 0.5 mg/l IAA.

After two weeks, explants were transferred onto 1ZK medium that contained 4.3 g/l MS

salts, 100 mg/l myo-inositol, 1 ml/l Nitsch vitamins (1000x), 20 g/l sucrose, 1 mg/l trans-zeatin,

75 mg/l kanamycin, 300 mg/l timentin, 5.2 g/l Agargel, and IAA at either 0 mg/l, 0.01 mg/l, 0.05

mg/l, or 0.1 mg/l IAA, or 0.5 mg/l in plates or Magenta GA7 boxes depending upon the size of

the shoots regenerating from the cotyledon explants.

Shoots were excised and transferred to selective rooting medium designated RMK (4.3 g/l

MS salts, 1 ml/l Nitsch vitamins (1000x), 30 g/l sucrose, pH 6.0, 8 g/l Difco Bacto agar (Becton,

Dickinson and Company, Franklin Lakes, NJ), 75 mg/l kanamycin, 300 mg/l timentin, and IAA at

either 0 mg/l or 1 mg/l in Magenta GA7 boxes.

Unless otherwise noted, the pH of all media was adjusted to 5.8 before autoclaving. For all

media, the trans-zeatin, IAA, kanamycin, and timentin were dispensed from filter sterilized

stock solutions into autoclaved medium that was allowed to cool to 55°C. Cotyledon explant

cultures were transferred to freshly prepared medium every two weeks.

GUS histochemical assay

Histochemical assay of β-glucuronidase (GUS) activity was performed on leaves from putative

transgenic and control (non-transformed) plants. Leaves were vacuum infiltrated for 20 – 30

min in buffer (0.8g/l 5-bromo-4-chloro-3-indolyl- β −D-glucuronide (X-Gluc), 0.1 M Na$_2$HPO$_4$,

0.1 M NaH$_2$PO$_4$ phosphate, 10 mM ethylenediamine tetraacetic acid (EDTA), 1.6 mM

potassium-ferricyanide and 1.6 mM potassium −ferrocyanide, 5% v/v Triton X-100, and 20% v/v

methanol) before incubation at 37°C overnight. The chlorophyll was removed from the leaves

by 3 - 4 washes with 70% ethanol at room temperature. The leaves were examined with a Leica

S8APO stereomicroscope outfitted with a digital camera.

Polymerase chain reaction analysis

To confirm the presence of the neomycin phosphotransferase II selectable marker gene (*nptII*),

DNA was extracted from leaves of putative transgenic lines and controls (non-transformed)

with the Qiagen DNeasy plant mini kit (Hilden, Germany) as per the manufacturer's

instructions. Primers used to detect *nptII* were forward 5'-GGC TGG AGA GGC TAT TC-3' and

reverse 5'-GGA GGC GAT AGA AGG CG-3'. The diagnostic amplicon size expected with these

primers is approximately 700 bp. The PCR program started with a one-step cycle of 2 min at

95^{o}C, followed by 29 cycles of 30 s at 94^{o}C, 45 s at 57^{o}C, 50 s at 72^{o}C, and a 10 min final

extension at 72^{o}C. DNA was separated and visualized by electrophoresis through a 1%

agaraose, ethidium bromide-stained gel.

Experimental Design

A total of 5 different experiments were performed. Three biological replicates were used for

each IAA concentration in each experiment. A total of 750 cotyledon explants were used per

IAA concentration investigated. The standard error was calculated.

Results

Optimization of IAA concentration for recovery of stable transgenic lines

Following the co-cultivation period, cotyledon explants were transferred to our standard

selective plant regeneration medium designated 2ZK that contains 2 mg/l trans-zeatin as the

only plant growth regulator. One week later, the explants were transferred to 2ZK

supplemented with different IAA concentrations (0 mg/l, 0.01 mg/l, 0.05 mg/l, 0.1 mg/l, 0.5

mg/l) to determine if the addition of IAA would decrease the time from infection with

Agrobacterium to recovery of stable transgenic lines. We continued to use this same series of

IAA concentrations in the subsequent selective plant regeneration medium designated 1ZK.

Medium supplemented with IAA resulted in shoots that were more fully developed earlier in

the culture process as compared to medium without IAA (Fig. 1A). In Figure 1A, cotyledon

cultures shown in a – e represent controls that were not infected with *Agrobacterium*. We

observed that as the IAA concentration increased, the level of plant regeneration from the

controls decreased (Fig. 1A, a – d). For our standard method without IAA (Fig. 1A, e), the level

of plant regeneration is significantly less in comparison with medium that contained IAA.

Cotyledon explants infected with *Agrobacterium* and cultured on medium containing IAA

exhibited the same pattern of shoot development as the cotyledons not infected, in that we

observed more well-developed shoots at an early stage of culture post infection (Fig. 11 f – i).

In general, earlier emergence of well-developed shoots from *Agrobacterium*-infected

cotyledon explants on medium containing IAA translated to the recovery of whole rooted plants

in less time as compared with medium that did not contain IAA (Table 1). Medium containing

either 0.05 mg/l or 0.1 mg/l IAA resulted in the shortest time, 11 wks, for recovery of stable

transgenic lines. There appeared to be a threshold of IAA concentration and effect on recovery

time because at 0.5 mg/l IAA the time was similar to our standard method. We observed a

similar decrease in time when transformations of other tomato genotypes were performed by

different lab members who tested plant regeneration medium that contained 0.1 mg/l IAA

(data unpublished).

Effect of IAA on transformation efficiency and rooting

The formula below was used to calculate transformation efficiency (TE).

Transformation efficiency (%):
194   Total number of rooted shoots   x 100
195   Total number of cotyledon explants infected with *Agrobacterium*
196

Overall, the TE was lower when medium containing IAA was used as compared to the TE of 88%

when medium containing trans-zeatin as the only growth regulator was used (Table 1).

When putative transgenic lines were approximately 3 - 4 cm tall, they were removed from

the cotyledon explants and transferred to either our standard selective rooting medium (RMK)

without IAA or RMK supplemented with 1 mg/l IAA.   We chose this concentration based on

previous work with tomato transgenic lines recovered from a few genotypes that did not root

as well as M82 on our standard rooting medium (data not published). We observed that shoots

cultured on RMK supplemented with IAA resulted in the emergence of roots after 6 - 7 days as

compared to 11 - 14 days on medium without IAA. The addition of IAA to the medium did not

result in any phenotypic differences of the plants as compared to medium that did not contain

IAA.

Characterization of putative transgenic lines

The first level of analysis to confirm the recovered plants from *Agrobacterium*-infected

cotyledons were transgenic was a histochemical assay for the GUS reporter protein. Whole

leaves from plants rooted on RMK were used for the analysis. All leaves exhibited GUS activity,

although we observed variation in the level of intensity with some leaves exhibiting a darker

coloration than others (Fig 1B). GUS activity was not observed in leaves from non-transformed

control plants.

To further confirm the recovered plants were indeed stable transgenic lines, we did PCR

analysis for the presence of the *nptII* selectable marker gene in plants found to be positive for

GUS activity. Total genomic DNA was isolated from the leaves of the GUS-positive lines and

non-transformed control plants. PCR amplification of the *nptII* gene was detected in plants that

were also GUS positive. No amplified product was detected in DNA from the control, (non-

transgenic) plants (Fig. 1C).

Modified protocol

Based on our findings, we now follow a modified protocol as outlined in Figure 2 for all

Agrobacterium-mediated tomato transformations. IAA concentrations of 0.5 and 0.1 mg/l IAA

both resulted in a 6-week decrease for recovery of stable transformants, however, we chose to

use 0.1 mg/l IAA in our modified protocol because of the 54% transformation efficiency (Table

1). In addition to M82, we have applied this protocol to other tomato genotypes and also

observed a decrease in time for recovery of transgenic lines as compared to our previous

tomato transformation methodology (data not published).

Discussion

For development of stable transformation methodology, the foremost factors to be

considered are transformation efficiency and the time from infection with *Agrobacterium*

tumefaciens until the recovery of transgenic lines. Methods that provide both high efficiency

and the shortest time to recovery of transgenic lines lead to a high-throughput pipeline that

allows earlier evaluation of gene function. In turn, a high-throughput pipeline decreases the

amount of labor and resources needed, which can translate into significant financial savings.

The focus of our study was to investigate medium components that had the potential to

decrease the time for recovery of stable tomato transgenic lines. Our standard method, which

has a high transformation efficiency at approximately 90%, takes 17 weeks for recovery of

transformants. The interest in optimization of our methods stemmed from an increased need

for transgenic lines because tomato has become the model species of choice for many studies

that include ripening, abiotic and biotic tolerance, and nutritional content [23-26]. In addition,

with the recent demonstration of successful genome editing by CRISPR/Cas9 in tomato, the

interest in applying this technology for the study of gene function will increase [16, 17, 27].

Therefore, a transformation methodology that can deliver modified lines in a shorter time

frame will help to advance these studies.

Our standard protocol is a modified version of methods reported by Fillatti et al. [8] in

which zeatin is the only growth regulator incorporated into the plant regeneration medium

[12]. We chose to start our investigation by examining additional growth regulators that, in

combination with zeatin, would greatly reduce the time for recovery of stable transgenic lines

but not have a significant negative effect on transformation efficiency. In a literature search,

we found several reports that demonstrated a positive effect on tomato plant regeneration and

transformation efficiency when indole-3-acetic acid (IAA) was incorporated into zeatin-

containing plant regeneration medium [10, 28, 29]. However, they did not report any effects

observed on the time required to recover transgenic plants.

We found that addition of either 0.05 or 0.1 mg/l IAA to our standard plant regeneration

medium that contains trans-zeatin as the only growth regulator decreased the time for

recovery of stable transgenic lines from 17 to 11 weeks. Previous reports have demonstrated

that shoot apical meristem development involves interactions among cytokinin signaling

pathway components, auxin, and several families of transcription factors [18]. It is possible that

the addition of IAA to our standard plant regeneration medium facilitates interactions among

the cytokinin signaling and auxin regulated genes, which results in faster shoot development

from the cotyledon explants.

Although there was a reduction in transformation efficiency with the addition of IAA from

approximately 90% to about 50%, this level is acceptable considering transgenic lines can be

evaluated significantly earlier than when our standard method was used. This decrease in time

allows researchers to test their material earlier and make changes to their approaches sooner if

results are unsatisfactory for their genes of interest.

In addition to supplementation of the standard plant regeneration medium with IAA, we

also investigated effects of adding IAA to the rooting medium, which was not a component in

our standard rooting medium. Inclusion of IAA in *in vitro* rooting medium has been reported

for tomato, however, it is not routinely added because tomato readily develops roots in culture

medium without growth regulators [9]. Our interest was to determine if supplementation

decreased the time to rooting, which we did observe. Auxin is produced in both shoots and

roots and the auxin produced in the roots helps in root development [30-32]. It is possible that

IAA, when exogenously added, increases the levels of auxin in the plants, hence resulting in the

cells differentiating earlier to form roots. However, research needs to be conducted to confirm

this hypothesis.

Conclusions

Interest in tomato as a model has increased over the years and we have seen a rise in the

number of research groups that require stable transgenic lines for various studies.

Modification of our standard plant regeneration medium through the addition of either 0.05 or

0.1 mg/l AA shortened the recovery of transgenic lines by 6 weeks for the M82 tomato cultivar.

Application of this modification for transformation of other tomato genotypes in our lab also

resulted in a decreased time for recovery of stable transgenic lines.

A shorter recovery time for stable transgenic lines is highly desirable for functional studies to

allow earlier determination of the genes and networks involved in phenotypes of interest. A

decrease in recovery time would also provide a higher throughput process, which has the

potential for cost savings related to labor and resources. Optimization studies of standard

transformation methodologies for different plant species should always be considered in order

to alleviate bottlenecks for generation of stable transgenic lines [33]. Availability of efficient

transformation methods is especially critical with the rapid development of genome editing

technologies, which will result in an increased demand for generation of transgenic lines for

basic research studies that can lead to crop improvement.

Competing interests

The authors declare that they have no competing interests.

Authors' contributions

SG and JVE designed the experiments, SG performed the experiments, SG and JVE wrote the

manuscript. Both authors read and approved the manuscript.

Acknowledgements

We thank Cynthia Du for assistance with part of the experimental process. We thank Cynthia

Du, Patricia Keen, and Michelle Tjahjadi for their critical review of the manuscript. Support for

this work was through a grant from the National Science Foundation Plant Genome Research

Program (IOS-1237880).

References

1.   Jones JBJ: **Tomato Plant Culture: In the Field, Greenhouse, and Home Garden**. Boca
Raton, FL: CRC Press LLC; 1998.
2.   *Agricultural Marketing Resource Center*.
http://www.agmrc.org/commodities__products/vegetables/tomatoes.
3.   Emmanuel E, Levy AA: **Tomato mutants as tools for functional genomics**. *Curr Opin*
Plant Biol 2002, **5**:112-117.
4.   Bombarely A, Menda N, Tecle IY, Buels RM, Strickler S, Fischer-York T, Pujar A, Leto J,
Gosselin J, Mueller LA: **The Sol Genomics Network (solgenomics.net): growing**
tomatoes using Perl. *Nucleic Acids Res* 2011, **39**:1149-1155.
5.   Consortium TTG: **The tomato genome sequence provides insights into fleshy fruit**
evolution. *Nature* 2012, **485**:635-641.
6.   Mccormick S, Niedermeyer J, Fry J, Barnason A, Horsch R, Fraley R: **Leaf disk**
transformation of cultivated tomato (*L. esculentum*) using *Agrobacterium*
tumefaciens. *Plant Cell Rep* 1986, **5**:81-84.
7.   Chyi Y-S, Phillips GC: **High efficiency *Agrobacterium*-mediated transformation of**
***Lycopersicon* based on conditions favorable for regeneration** *Plant Cell Rep* 1987,
6:105-108.
8.   Fillatti JJ, Kiser J, Rose R, Comai L: **Efficient transfer of a glyphosate tolerance gene into**
tomato using a binary *Agrobacterium tumefaciens* vector. *Bio-Technol* 1987, **5**:726-
730.
9.   Frary A, Earle ED: **An examination of factors affecting the efficiency of *Agrobacterium*-**
mediated transformation of tomato. *Plant Cell Rep* 1996, **16**:235-240.
10. Park SH, Morris JL, Park JE, Hirschi KD, Smith RH: **Efficient and genotype-independent**
***Agrobacterium* - mediated tomato transformation**. *J Plant Physiol* 2003, **160**:1253-
1257.
11. Sun HJ, Uchii S, Watanabe S, Ezura H: **A highly efficient transformation protocol for**
Micro-Tom, a model cultivar for tomato functional genomics. *Plant Cell Physiol* 2006,
47:426-431.
12. Van Eck J, Kirk DD, Walmsley AM: **Tomato (*Lycopersicum esculentum*)**. In: *Methods in*
Molecular Biology, Agrobacterium Protocols. Edited by Wang K. Totowa, NJ: Humana
Press Inc.; 2006:459-473.
13. Somerville C, Koornneef M: **A fortunate choice: the history of *Arabidopsis* as a model**
plant. *Nature reviews Genetics* 2002, **3**:883-889.
14. Gelvin SB: ***Agrobacterium*-mediated plant transformation: the biology behind the**
"gene-jockeying" tool. *Microbiol Mol Biol Rev* 2003, **67**:16-37.
15. Pitzschke A, Hirt H: **New insights into an old story: *Agrobacterium*-induced tumour**
formation in plants by plant transformation. *EMBO J* 2010, **29**:1021-1032.
16. Brooks C, Nekrasov V, Lippman ZB, Van Eck J: **Efficient gene editing in tomato in the**
first generation using the Clustered Regularly Interspaced Short Palindromic
Repeats/CRISPR-Associated9 system. *Plant Physiol* 2014, **166**:1292-1297.

17.   Xu C, Liberatore KL, MacAlister CA, Huang Z, Chu YH, Jiang K, Brooks C, Ogawa-Ohnishi
352   M, Xiong G, Pauly M *et al*: **A cascade of arabinosyltransferases controls shoot meristem**
353   **size in tomato**. *Nat Genet* 2015, **47**:784-792.
354   18.   Gupta S, Rashotte AM: **Down-stream components of cytokinin signaling and the role of**
355   **cytokinin throughout the plant**. *Plant Cell Rep* 2012, **31**:801-812.
356   19.   Shimizu-Sato S, Tanaka M, Mori H: **Auxin-cytokinin interactions in the control of shoot**
357   **branching**. *Plant Mol Biol* 2009, **69**:429-435.
358   20.   Friml J, Vieten A, Sauer M, Weijers D, Schwarz H, Hamann T, Offringa R, Jurgens G:
359   **Efflux-dependent auxin gradients establish the apical-basal axis of *Arabidopsis***. *Nature*
360   2003, **426**:147-153.
361   21.   Sabatini S, Beis D, Wolkenfelt H, Murfett J, Guilfoyle T, Malamy J, Benfey P, Leyser O,
362   Bechtold N, Weisbeek P *et al*: **An auxin-dependent distal organizer of pattern and**
363   **polarity in the *Arabidopsis* root**. *Cell* 1999, **99**:463-472.
364   22.   Murashige T, Skoog F: **A revised medium for rapid growth and bio assays with tobacco**
365   **tssue cultures**. *Physiol Plantarum* 1962, **15**:473-497.
366   23.   Martel C, Vrebalov J, Tafelmeyer P, Giovannoni JJ: **The tomato MADS-box transcription**
367   **factor RIPENING INHIBITOR interacts with promoters involved in numerous ripening**
368   **processes in a COLORLESS NONRIPENING-dependent manner**. *Plant Physiol* 2011,
369   **157**:1568-1579.
370   24.   Sun W, Xu X, Zhu H, Liu A, Liu L, Li J, Hua X: **Comparative transcriptomic profiling of a**
371   **salt-tolerant wild tomato species and a salt-sensitive tomato cultivar**. *Plant Cell Physiol*
372   2010, **51**:997-1006.
373   25.   Nguyen HP, Chakravarthy S, Velasquez AC, McLane HL, Zeng L, Nakayashiki H, Park DH,
374   Collmer A, Martin GB: **Methods to study PAMP-triggered immunity using tomato and**
375   ***Nicotiana benthamiana***. *Mol Plant Microbe Interact* 2010, **23**:991-999.
376   26.   Gonzali S, Mazzucato A, Perata P: **Purple as a tomato: towards high anthocyanin**
377   **tomatoes**. *Trends Plant Sci* 2009, **14**:237-241.
378   27.   Ito Y, Nishizawa-Yokoi A, Endo M, Mikami M, Toki S: **CRISPR/Cas9-mediated**
379   **mutagenesis of the RIN locus that regulates tomato fruit ripening**. *Biochem Biophys*
380   *Res Commun* 2015, **467**:76-82.
381   28.   Gubis J, Lajchova Z, Farago J, Jurekova Z: **Effect of growth regulators on shoot induction**
382   **and plant regeneration in tomato (*Lycopersicon esculentum* Mill.)**. *Biologia* 2004,
383   **59**:405-408.
384   29.   Yasmeen A: **An improved protocol for the regeneration and transformation of tomato**
385   **(cv Rio Grande)**. *Acta Physiol Plant* 2009, **31**:1271-1277.
386   30.   Petersson SV, Johansson AI, Kowalczyk M, Makoveychuk A, Wang JY, Moritz T, Grebe M,
387   Benfey PN, Sandberg G, Ljung K: **An auxin gradient and maximum in the *Arabidopsis***
388   **root apex shown by high-resolution cell-specific analysis of IAA distribution and**
389   **synthesis**. *Plant Cell* 2009, **21**:1659-1668.
390   31.   Stepanova AN, Robertson-Hoyt J, Yun J, Benavente LM, Xie DY, Dolezal K, Schlereth A,
391   Jurgens G, Alonso JM: **TAA1-mediated auxin biosynthesis is essential for hormone**
392   **crosstalk and plant development**. *Cell* 2008, **133**:177-191.
393   32.   Overvoorde P, Fukaki H, Beeckman T: **Auxin Control of Root Development**. *Cold Spring*
394   *Harb Perspect Biol* 2010, **2**:1-16.

395   33.   Altpeter F, Springer NM, Bartley LE, Blechl AE, Brutnell TP, Citovsky V, Conrad LJ, Gelvin
396   SB, Jackson DP, Kausch AP *et al*: **Advancing crop transformation in the era of genome**
397   **editing**. *Plant Cell* 2016, **in press**.
398

Figure Legends

Figure 1

Results for the recovery of *Solanum lycopersicum* cv M82 stable transgenics from

***Agrobacterium tumefaciens*-infected cotyledon explants cultured on plant regeneration**

medium supplemented with different concentrations of indole-3-acetic acid (IAA). (A)

Agrobacterium tumefaciens-infected cotyledon explants (approximately 5 weeks post infection)

cultured on selective plant regeneration medium containing the following amounts of IAA in

mg/l (f) 0.01, (g) 0.05, (h) 0.1, (i) 0.5, and (j) 0. Images a – e represent the corresponding non-

infected controls for each IAA concentration, respectively.

(B) Histochemical analysis for GUS expression in leaves taken from independent transgenic lines

designated 1 - 9 recovered from selective plant regeneration medium that contained 0.1 mg/l

IAA. GUS expression was not observed in the non-transformed controls.

(C) Agarose gel of PCR products showing the expected ~700 bp product amplified from the *nptII*

selectable marker gene in 10 independent transgenic lines (lanes 1 – 10). These lines were

recovered from selective plant regeneration medium that contained 0.1 mg/l IAA. C = the

control

Figure 2

Schematic representation of the optimized *Agrobacterium tumefaciens*-mediated

transformation methodology for *Solanum lycopersicum* cv M82. See the Materials and

Methods for details on seed sterilization and all media compositions.

<u>Tables</u>

Table 1: Results for recovery of stable transgenic lines of *Solanum lycopersicum* cv M82 from

***Agrobacterium tumefaciens*-infected cotyledon explants cultured on selective plant**

regeneration medium supplemented with different indole-3-acetic acid (IAA) concentrations.

IAA (mg/l)	Total number explants	Total number rooted plants	Average transformation efficiency (+) SE*	Total time for recovery of transgenic lines (wks)
0	750	660	88 + 2.2	17
0.01	750	390	52 + 1.0	15
0.05	750	375	50 + 1.5	11
0.1	750	405	54 + 1.2	11
0.5	750	360	48 + 2.0	16

433   *Average transformation efficiency was calculated as percent of stable transgenic lines

434   recovered from the total number of cotyledon explants infected with *Agrobacterium*

435   *tumefaciens*. Transformation efficiency values shown are the average from 5 experiments ± the

436   standard error (SE) calculated from 3 biological replicates.

A

B

C

Gupta and Van Eck_Figure 2

Germination of sterilized seeds *in vitro*

6 days ↓

Preparation of explants from cotyledons

24 hrs ↓

Incubation in *Agrobacterium* suspension

5 min ↓

Co-cultivation of explants

2 days ↓

Transfer explants to 2ZK selective plant regeneration medium

1 wk ↓

Transfer explants to 2ZIK selective plant regeneration medium

2 wks ↓

Transfer explants to 1ZIK selective medium

~4 wks ↓

Remove shoots from explants transfer to RMIK

~3 wks ↓

Well-rooted plants