***Zfp423* regulates Sonic hedgehog signaling via primary cilium function**

Short title: *Zfp423* regulates cilium function in Shh signaling

Chen-Jei Hong[1,#a] and Bruce A. Hamilton1,*

[1]Department of Cellular & Molecular Medicine, Department of Medicine,

Moores UCSD Cancer Center, and Institute for Genomic Medicine

University of California, San Diego

La Jolla, CA 92093-0644

[#a] Current address: Neural Regeneration Laboratory, Neurological Institute, Taipei

Veterans General Hospital, Taipei, Taiwan.

*Corresponding author

email: bah@ucsd.edu

ABSTRACT

Zfp423 encodes a 30-zinc finger transcription factor that intersects several canonical

signaling pathways. *Zfp423* mutations result in ciliopathy-related phenotypes, including

agenesis of the cerebellar vermis in mice and Joubert syndrome (JBTS19) and

nephronophthisis (NPHP14) in humans. Unlike most ciliopathy genes, *Zfp423* encodes a

nuclear protein and its developmental expression is complex, leading to alternative

proposals for cellular mechanisms. Here we show that Zfp423 is expressed by cerebellar

granule cell precursors, that loss of *Zfp423* in these precursors leads to cell-intrinsic

reduction in proliferation, loss of response to Shh, and primary cilia abnormalities that

include diminished translocation of both Smoothened and IFT88. Loss of Zfp423 alters

expression of several genes encoding key cilium components, including increased

expression of *Tulp3*. *Tulp3* is a direct binding target of Zfp423 and reducing the

overexpression of Tulp3 in *Zfp423*-deficient cells suppresses Smoothened translocation

defects. These results define *Zfp423* deficiency as a bona fide ciliopathy, acting

upstream of Shh signaling, and indicate a mechanism intrinsic to granule cell precursors

for the resulting cerebellar hypoplasia.

AUTHOR SUMMARY

Ciliopathies are a broad group of individually rare genetic disorders that share

overlapping phenotypes and mutations in genes that make components of the primary

cilium. Mutations in *ZNF423* are an exception. Patients and mouse models show

characteristic hypoplasia of the cerebellar midline (Joubert syndrome), but the gene

encodes a nuclear transcription factor. The mouse gene, *Zfp423*, is expressed in a

dynamic developmental pattern, leaving the cellular mechanism for this brain

malformation unresolved. One report suggested reduced Purkinje cell expression of

Shh, a key mitogen for cerebellar granule cell precursors (GCPs) whose signal

transduction occurs at the primary cilium, as the key event. We show that *Zfp423*

mutants expressed normal Shh levels, but that *Zfp423*-depleted GCPs were unable to

respond. Primary cilia on *Zfp423*-mutant GCPs in situ typically had a wider base and

longer extension. ZNF423-depletion in a human cell culture model resulted in defective

translocation of Smoothened, a key event in Shh signaling, and of the intraflagellar

transport protein IFT88. RNA-Seq and RT-qPCR experiments identified known

ciliopathy genes as potential conserved targets of *ZNF423* and *Zfp423*. One of these,

TULP3, is both up-regulated in *ZNF423*/*Zfp423*-deficient cells and directly bound by

Zfp423 in granule cell precursors. Reversing the overexpression of TULP3 in ZNF423-

depleted human cell culture model reversed the defect in Smoothened translocation.

INTRODUCTION

Cerebellar granule cell precursors (GCPs) are both an important model for neuronal development and a site of clinically important developmental abnormalities. GCP proliferation is highly responsive to Purkinje cell-derived sonic hedgehog (Shh) over a wide dynamic range [1-3]. Increasing or decreasing developmental Shh signaling can alter GCP dynamics sufficiently to reshape the cerebellum [4]. Correspondingly, ligand-independent signaling through the Shh pathway is a common feature in medulloblastoma, the most common pediatric brain tumor [5-7], and the Shh signaling pathway is a focal point of therapeutic development [8-10]. Shh signaling at primary cilia, where multiple components are trafficked to create a focused signaling module [11-14], is required for GCP expansion [15]. How Shh signaling is integrated with other signals that impact GCP proliferation, migration, and differentiation is not fully understood.

Through alternative interactions with multiple signaling and transcriptional pathways, Zfp423 (and its human orthologue, ZNF423) is well positioned to integrate extracellular signals into a coherent developmental response. Zfp423 was first described (as Roaz) through inhibitory interaction with EBF (Olf1) helix–loop–helix transcription factors [16]. Zfp423 is also a coactivator for BMP-activated SMADs [17, 18], retinoic acid receptors [19, 20], and Notch intracellular domain [21]. Intriguingly, Zfp423 activities on EBF and ligand-activated factors appear to be mutually inhibitory, further suggesting an integrative network function. *ZNF423* also appears to be a target of some cancers and low expression in neuroblastoma [20] or epigenetic silencing by Polycomb repressive complex 2 in glioma [22] is associated with poor prognosis.

Zfp423-deficient mice have a variety of developmental defects, including fully penetrant loss of cerebellar vermis [23-25] and variable loss of cerebellar hemispheres dependent on modifier genes and other factors [26]. Zfp423-deficient animals are also

defective in forebrain development–including hypoplasia of the hippocampus and

incomplete corpus callosum [23, 24], in olfactory neurogenesis [27], and in induction of

adipose tissue [28, 29]. Mechanistically, literature to date has focused on physical

interactions between Zfp423 and other transcription factors. Alternative levels of

integration, such as alterations to cellular signaling centers, have not been well explored.

Hildebrandt and co-workers identified mutations in human *ZNF423* among

patients with ciliopathy diagnoses [30]. Patients from all three *ZNF423* families had

cerebellar vermis hypoplasia or Joubert Syndrome, while two also had nephronophthisis

and other clinical features. Cellular assays with patient mutations showed effects on

proliferation and DNA damage response, presenting a new pathogenic mechanism in

ciliopathy disorders, but did not assess cilium structure or function. This raises the

question of whether *ZNF423* and *Zfp423* mutations phenocopy ciliopathies by acting on

downstream signaling events or represent bona fide ciliopathies by affecting cilium

function upstream of signaling.

Results here provide new insights into Zfp423-dependent developmental

mechanisms. Distinctly different models have been proposed for the cerebellar

hypoplasia in *Zfp423* mice and, by extension, human patients. Based on *Zfp423* gene-

trap expression in postnatal Purkinje cells, one group proposed a non-autonomous

mechanism mediated by diminished Shh production [25], as seen in some Purkinje cell-

selective mutations [31]. In situ hybridization, however, showed *Zfp423* expression in

both the ventricular zone and the external germinal layer (EGL) in developing

cerebellum, equally consistent with a GCP-intrinsic mechanism [23, 24]. We show that

GCPs express Zfp423 protein in situ and in primary culture, that loss of Zfp423 blocked

their ability to respond to Shh, altered cilium morphologies, decreased Smoothened

translocation, and increased expression of several cilium-related genes, including *Tulp3*.

Reversing the Tulp3 overexpression restored Smoothened translocation. Our results

demonstrate a GCP-intrinsic role for Zfp423 upstream of Shh signaling and suggest

excess Tulp3 as a potential target to restore function in ZNF423/Zfp423-deficient cells.

RESULTS

Zfp423 **is expressed in granule cell precursors.**

In situ hybridization had previously shown *Zfp423* RNA expression in ventricular

zone, external germinal layer and rhombic lip [23, 24], while lacZ reporter expression in

a gene trap line suggested expression restricted to Purkinje cells [25], leading to

different proposals for developmental defects in mutant embryos. To resolve which cells

express Zfp423 in developing cerebellum, we examined Zfp423 protein expression and

RNA in isolated cell populations (Fig 1). Zfp423 immunoreactivity showed strong,

nuclear-limited signal in postmitotic Purkinje cells and most migrating GCPs in the

rhombic lip and external germinal layer (EGL), with somewhat less intense staining in a

fraction of ventricular zone cells (Fig 1A). Comparing sections from control (+/+) and

Zfp423 null mutant (*nur12*) congenic littermates that were processed in parallel on single

slides demonstrated both specificity and sensitivity of the signal. Similar results have

been obtained with two additional, independent antibodies developed against Zfp423 or

its human ortholog (L. Flores-García and B.A.H.). As a further test of cell type

distribution, we monitored by reverse transcription and quantitative polymerase chain

reaction (RT-qPCR) the expression of *Zfp423* and several cell-type selective markers in

cerebellar cells isolated by Percoll gradient centrifugation at E18.5 (Fig 1B,C). These

data indicated that highly enriched GCP cultures retain *Zfp423* expression while losing

Purkinje cell selective markers such as *Rora* and *Shh*. Furthermore, RT-qPCR

experiments for expression of *Shh*, the primary mitogen for GCPs, was not significantly

different between *Zfp423*-mutant and control cerebellum among several independent

samples (Fig 1D). These experiments demonstrate nuclear Zfp423 expression in GCPs

in situ, with continued expression in primary culture, and support the potential for a cell-

intrinsic role for Zfp423 in the granule cell deficits seen in $Zfp423^{-/-}$ animals.

Fig 1. Zfp423 is expressed in cerebellar granule cell precursors. (**A**) Antibody

staining at E16.5 shows nuclear expression of Zfp423 in cells of the rhombic lip (rl) and

external germinal layer (egl), as well as ventricular zone (VZ) derivatives in control, but

not in $Zfp423^{nur12}$ mutant cerebellum. (**B**) E18.5 granule cell precursors purified by

Percoll gradient centrifugation retain Zfp423 expression. Quantitative RT-PCR shows co-

expression of Zfp423 with granule-specific Atoh1 and Barhl1 RNAs, with loss of Purkinje

cell marker Rora. Histogram shows average expression ratio among technical replicates

from a single purification, normalized to Gapdh and expressed relative to whole

cerebellum. Cb, whole cerebellum; GC, granule cell/precursor fraction; P/G, Purkinje cell

and Glial fraction. Rora level expressed relative to P/G fraction; Cb fraction not run. (**C**)

Quantitative RT-PCR with hydrolysis probes (TaqMan) assays shows enrichment of

Zfp423 in the granule cell precursor fraction of a second independent purification. In

contrast, no amplification from the granule cell preparation is detected for Shh, a GCP

mitogen expressed by Purkinje cells. (**D**) RT-qPCR for Shh expression in whole

cerebellum from control and nur12 mutant animals dissected on P3. Mean values from

technical replicates were normalized to Gapdh and expressed as a fraction of a control

sample for 7 control and 5 mutant embryos. Similar results were obtained when

normalized to *Ppia* cyclophilin. Group means were not significantly different (p=0.55, t-

test).

***Zfp423* is required for ex vivo proliferation and Shh responsiveness.**

To test whether GCP-intrinsic *Zfp423* expression is relevant to their proliferation

phenotype, we transfected purified GCPs with shRNA vectors (Fig 2) that we previously

validated for reducing Zfp423 levels [32]. Transfected GCPs were identified by

fluorescence of an enhanced green fluorescent protein (EGFP) reporter in the vector.

Cells entering S phase were marked by BrdU incorporation (A). The proportion of

EGFP[+] cells that were also BrdU[+] was taken as a mitotic index specific for transfected

cells. As predicted from *Zfp423* mutant animals [23], shRNA significantly reduced this

index relative to the corresponding index of non-transfected (EGFP[-]) cells in the same

cultures (p = 0.040-0.0014, paired *t* tests) and to approximately one-third the level of

cells transfected by empty vector, irrelevant target (luciferase), or scrambled sequence

shRNA controls (Fig 2B; p<10[−7], one-factor ANOVA with post-hoc Tukey HSD test).

Because the transfection efficiency in primary GCPs is modest (~15%), most transfected

cells were surrounded by non-transfected cells. This minimizes the likelihood of any

confounding effects secondary to disrupted cell interactions. These data show that

Zfp423 has a cell-intrinsic effect on GCP proliferation ex vivo and suggest that the effect

might be cell autonomous within a population of GCPs, as proliferation phenotype does

not appear to depend on the status of adjacent cells in the culture.

Because previous reports have shown that Shh is the principle mitogen for GCP

proliferation in vivo and in culture, with a wide dynamic range, we next tested whether

Zfp423-depletion in GCPs might alter the dose-response curve for exogenous Shh

treatment (Fig 2C). We exposed Zfp423-knockdown and control GCPs to a range of 0-

10 µg/ml recombinant Shh concentrations and measured the resulting mitotic index for

replicate cultures at each concentration. Two-factor ANOVA on resulting cell count data

showed significant impacts of Shh dose ($p = 2.0 \times 10^{-4}$), Zfp423 vs. control shRNA ($p =$

9.3×10^{-10}), and the interaction between dose and shRNA ($p = 6.4 \times 10^{-6}$). Remarkably,

Zfp423-shRNA cells were refractory to Shh levels more than an order of magnitude

greater than that required to stimulate proliferation of control-shRNA cells. These results

indicate a Zfp423-dependent step critical to Shh signal transduction in GCPs.

Fig 2. *Zfp423* shRNA blocks proliferation and Shh response of purified granule cell

precursors. (*A*) Examples of GCPs transfected with scrambled control or *Zfp423*-

targeted shRNA constructs, in the presence of 3 µg/ml Shh. Transfected cells express

EGFP from the shRNA vector (green). Proliferating cells are labeled by BrdU

incorporation for 6 hr, 48 hr after transfection (red). Control panels show dividing

transfected cells (closed arrowheads) and a non-dividing, non-transfected, cell (open

arrowhead). *Zfp423* (shRNA mZ1) panels show a dividing non-transfected cell (closed

arrowhead) and a non-diving transfected cell (open arrowhead). At least two

transfections for each of three independent DNA preparations were assayed for each

construct. Channel levels were adjusted for merged images. (*B*) The proportion of

EGFP$^+$ cells labeled by BrdU is plotted for each replicate culture. shRNA sequence was

highly significant ($p < 2.2 \times 10^{-16}$, ANOVA), with each *Zfp423* shRNA construct different

from every control ($p < 10^{-7}$ for each pair-wise comparison, Tukey HSD). Controls were

not significantly different from each other (p > 0.4), nor were the two *Zfp423* hairpin

sequences (p > 0.4). Empty, pLKO vector with no hairpin sequence; luc, luciferase; scr,

scrambled sequence control; mZ1 and mZ2 are non-overlapping shRNA sequences

designed against *Zfp423* from the RNAi consortium collection. (**C**) GCP cultures were

transfected with control (scr) or *Zfp423*-targeted (mZ1) shRNA constructs and treated

with 0-10 μg/ml Shh. Transfected cells were recognized by EGFP fluorescence,

replicating cells by BrdU incorporation. Plotted points indicate mitotic index of

transfected cells for each of four replicate experiments at each of the indicated Shh

concentrations. A minimum of three independent DNA preparations was used for

transfection in each condition. The effect of *Zfp423*-targeting shRNA, Shh

concentration, and the interaction between hairpin identity and Shh concentration were

each strongly supported statistically (p= 9.3×10^{-10}, 2.0×10^{-4}, and 6.4×10^{-6}

respectively, two-factor ANOVA).

213   Zfp423-deficient precursors have abnormal cilium morphology

Because Shh signaling is transduced through components localized to the

primary cilium, we next asked whether Zfp423-deficient GCPs made cilia and basal

bodies in normal frequency and quality (Fig 3). Cerebellum sections from E18.5 mice

showed both structures intact among cells in the EGL and with comparable frequency

per cell between *Zfp423^{nur12}*-homozygous and littermate controls by double label

immunofluorescence for acetylated α-tubulin (Ac-αTub) in the cilium and γ-tubulin (γTub)

in the basal body (Fig 3A). Volume measurements extracted from optical sections,

however, showed a significant increase in cilium volume in *Zfp423^{nur12}* EGL compared to

littermate controls (Fig 3B). As replicate measurements did not follow a normal

distribution within or between biological samples, we made several assessments using

non-parametric tests. Among co-processed sections from eight littermate pairs, average

volumes were larger in the mutant than in the control animal for all eight comparisons (p

= 0.0078, Wilcoxon signed rank test, two tailed). The distribution of volumes between

genotypes for all 499 discrete measurements irrespective of pairing was also highly

significant (p = 8.7 x 10^{-6}, Kolmogorov-Smirnov test). Mutant cilia also had substantially

higher variance than those in control littermates (0.89 vs. 0.33; p = 1.3 x 10^{-4}, Ansari-

Bradley test). Similar analysis of basal bodies (Fig 3C) did not support a difference in

paired sample averages (p = 1), but did support a slightly lower distribution of basal body

volumes over 662 individual measurements irrespective of pairs (p = 0.031) without

significant difference in variances (p = 0.19). Mean voxel values for acetylated α-tubulin

were 1.02 in control, 1.37 in mutant sections. Mean voxel values for γ-tubulin were 1.35

in control, 1.11 in mutant sections.

Fig 3. Confocal images show altered primary cilium volumes in *Zfp423^nur12* EGL.

(***A***) Confocal micrographs show representative projection images showing ciliary

axoneme (anti-acetylated α-tubulin, Ac-αTub) and basal bodies (anti-γ-tubulin, γTub) in

sections of E18.5 cerebellum from animals of the indicated *Zfp423* genotypes. All

images are from sagittal sections, 0.7-0.9 mm from the midline. Scale bar, 2 μm. (***B,C***)

Measured volumes for each of 499 axonemes (***B***) and 662 basal bodies (***C***) identified by

the Velocity program from reconstructed optical sections. Measurements from paired

mutant and control animals are color coded in the scatter plot. The differences in overall

distributions between mutant and control are significant for both axonemes (p = 8.7 x

10^{-6}) and basal bodies (p = 0.031, Kolmogorov-Smirnov test) in pooled data from eight

pairs of animals.

To understand better the structural basis for these volume differences, we

measured discrete parameters by structured illumination "super-resolution" microscopy

of cilia from mutant and control littermate animals (Fig 4). We obtained similar results

using either Ac-αTub (Fig 4A-C) or Arl13b (Fig 4D-F) as a marker protein. In Ac-αTub

measurements, mutant cilia tended to be longer (Fig 4B), though this effect did not reach

conventional significance (p = 0.079, Wilcoxon rank sum test, two tails), and mutant cilia

had a significantly wider base compared with control (Fig 4C, p = 0.0091). Structured

illumination measurements for Arl13b, associated with the ciliary membrane, showed

significant increases in both length (Fig 4E, p=0.028) and width (Fig 4F, p=0.013) of

GCP cilia in an independent set of animals. Taken together, these observations indicate

a cellular basis for ciliopathy-related phenotypes of *Zfp423*-deficient mice.

Fig 4. Structured illumination microscopy shows altered cilium dimensions in *Zfp423^{nur12}* **EGL.** (*A*) Super-resolution images show axoneme (Ac-αTub) and basal body (γTub). Scale bar, 2 μm. (*B,C*) Box and scatter plots show distributions of cilium length (*B*) and base width (*C*) measurements. Significance was assessed from paired samples. Length, p = 0.079, base width p = 0.0091, Wilcoxon rank sum test with two tails. (*D*) Structured illumination micrographs of cilia in the EGL. Arl13b and γ-tubulin in green, acetylated α-tubulin in red. (*E*) Length measurements from Arl13b signal in consecutively imaged cilia showed a shift toward longer axonemes in *nur12* (p = 0.028, Wilcoxon rank sum test). (*F*) Width of the axoneme base was also increased in *nur12* (p = 0.013, Wilcoxon rank sum test).

ZNF423 **is required for quantitative Smoothened translocation during Shh signaling**

Because Zfp423 is required for Shh response and affected the distribution of cilium morphologies, we asked whether it also affected trafficking of Smoothened, using a cell culture model (Fig 5). DAOY is a human medulloblastoma-derived cell line that expresses markers consistent with a GCP lineage [33, 34] and *ZNF423*, the human homologue of *Zfp423* (Fig 5A). DAOY cells were infected with a pseudotyped lentivirus expressing both a Smoothened-enhanced green fluorescent protein (EGFP) fusion protein and shRNA to either human *ZNF423* or a non-targeting control (Fig 5B). Cells were examined after serum starvation to produce a high percentage of cells with a definitive primary cilium. Two non-overlapping *ZNF423* shRNA were each effective in reducing its level of expression relative to multiple controls, a representative image is shown in Fig 5C. Quantification of fluorescence Western blots showed that, relative to control shRNAs, *ZNF423* knockdown cultures expressed ~30% ZNF423, ~110%

acetylated α-tubulin, and ~95% IFT88 levels scaled to either GAPDH or β-actin as

internal loading controls. We then tested the effect of ZNF423 knockdown on

Smoothened translocation. Infected DAOY cells were treated with 0.1 µg/ml Shh as a

low dose sufficient for signaling [1]. Cells were fixed 6 hr after stimulation and processed

for immunofluorescence to visualize localization of the EGFP tag on Smoothened and

acetylated α-tubulin as a marker of cilium location. Only EGFP+ cells were included to

avoid measurements from uninfected cells. To compare localization of Smoothened-

EGFP relative to acetylated α-tubulin, immunofluorescence micrographs (Fig 5D)

presented in randomized order were assessed categorically by five observers blinded to

experimental conditions, for >800 cilia across five independent experiments. Summary

results (Fig 5E) showed significantly less Smoothened-EGFP translocation in *ZNF423*-

depleted cells (1.2×10^{-13}, chi-square test). Results showed significant differences in each

independent experiment ($p = 0.0058$ to 2.9×10^{-5}).

Mean fluorescence intensities along the length of each cilium were also

calculated from the raw image files as a second analytical approach to the same

experiments. Cilium annotations and measurements were performed blind to treatment

group on images in randomized order. ZNF423-depleted cells showed ~30% reduction

in both mean and median Smoothened-EGFP intensities in cilia, with a highly significant

population shift between shRNA groups ($p=7.0 \times 10^{-11}$, Wilcoxon rank sum test; each of

five replicates was independently significant, $p<0.025$). Ratios of Smoothened-EGFP to

acetylated α-tubulin showed even stronger difference between groups ($p<2.2 \times 10^{-16}$),

owing to an increase in acetylated α-tubulin signal in knockdown cells. These results

strongly support a ZNF423-dependent step in Shh signaling upstream of Smoothened

translocation in human DAOY cells.

Fig 5. *ZNF423* knockdown quantitatively alters *Smoothened* translocation. (*A*) RT-

PCR shows *Zfp423* expression in mouse P19 cells [32] and *ZNF423* RNA expression in

DAOY medulloblastoma cells. *GAPDH* was used as a positive control. (*B*) Structure of

pLKO-derived lentivirus constructs, expressing a shRNA from a U6 promoter (purple

arrow) and Smoothened-EGFP fusion (smoGFP, dark green) from a human *PGK*

promoter (light green). Vector backbone elements are indicated in blue. (*C*) Western

blots show reduced expression of ZNF423 protein in DAOY infected with lentivirus

expressing a *ZNF423* shRNA (hZ3) relative to control (Ctl). (*D*) Color-inverted images of

immunofluorescence for both GFP (red) and acetylated a-tubulin (blue) show

Smoothened translocation phenotypes of cells infected with lentivirus expressing both

smoGFP and an shRNA. Channel layers were shifted diagonally to allow independent

view of each. Three categorical phenotypes were observed, (1) strong acetylated α-

tubulin with little to no detected Smoothened, (2) intermediate levels, or (3) stronger

Smoothened signal relative to acetylated α-tubulin. (*E*) Stacked histogram of cilium

phenotypes summarizes five replicate experiments. Phenotypes are color-coded and

numbered as in (D). Number of cilia scored in each group (n), *ZNF423*-directed (ZNF) or

330   control (Ctl) shRNAs, and chi-square p-value for the difference between groups are

331   indicated.

ZNF423 deficiency reduces IFT88 translocation into cilia.

Because Smoothened is thought to use the intraflagellar transport (IFT) system

for translocation into cilia [35, 36], we tested whether IFT88 translocation was affected

by loss of *ZNF423* in the DAOY culture model or *Zfp423* mutation in vivo (Fig 6). IFT88

is an essential component of the IFT-B complex required for quantitative hedgehog

signaling in mice [11] and important for cerebellum development [37]. DAOY cells were

infected with viruses described in Fig 5B and selected for cilium expression of the EGFP

marker by direct fluorescence after fixation and staining. This necessarily restricted the

analysis to the subpopulation of cells with strong Smoothened-EGFP translocation, as

direct fluorescence is less sensitive than antibody staining and ZNF423-knockdown cells

had a higher proportion of cilia near the detection limit. Among these cells, those

transduced with *ZNF423*-directed shRNA showed both increased acetylated α-tubulin

intensity relative to control (as predicted by Figures 4 and 5) and dramatically decreased

ciliary IFT88 intensity (Fig 6A). The differences between *ZNF423* and control shRNAs

reached very high statistical significance for both mean intensities and ratios in cilia (Fig

6B-E, $p<2.2 \times 10^{-16}$, Kruskal-Wallis and Wilcoxon rank sum tests for full and pair-wise

comparisons, respectively, in each measure), despite similar overall IFT88 protein levels

in cellular extracts after *ZNF423* knockdown, as shown in Fig 5C. Strikingly, the IFT88

signal was also reduced relative to Smoothened-EGFP signal in *ZNF423*-knockdown

cells compared to control (Fig 6E), suggesting that loss of IFT88 staining was not a

consequence of selection for the high Smoothened-EGFP population or comparison to

acetylated α-tubulin levels.

To determine whether similar conditions affect granule precursor cells in situ, we

examined the relative distributions of IFT88 and acetylated α-tubulin in the EGL of

Zfp423^{nur12} and control littermate mice (Fig 6F-G). While most cilia from control

littermates showed robust IFT88 staining in the cilium, many cilia in mutant EGL had little

or no detectable IFT88 (Fig 6F). The proportion of cilia with little or no IFT88 staining

was approximately three times greater among mutant animals than among their control

littermates (Fig 6G). Similarly, the overall intensity of IFT88 within the cilium was

significantly reduced in GCPs within the EGL of *nur12* animals compared with littermate

controls (Fig 6H). Both measures showed highly significant *Zfp423*-dependence across

five littermate pairs (p=3.7 x 10^{-8} for proportion, chi-square test; p=2.2 x 10^{-15} for signal

intensities, Wilcoxon rank sum test with continuity correction). IFT88 translocation thus

appears impaired both in the human DAOY cell culture model and in mouse granule cell

precursors in tissue sections, providing a proximate cause for diminished Shh signaling

both ex vivo and in situ.

Fig 6. Reduced frequency and intensity of axonemal IFT88 in Zfp423-deificent

cells. (***A***) Confocal micrographs of DAOY cells transduced with the indicated shRNA

and Smoothened-EGFP. Immunofluorescence for acetylated α-tubulin and IFT88 and

direct fluorescence of EGFP in representative images are shown. Scale bar, 5 µm.

Distributions of mean axonemal measurements for (***B***) acetylated α-tubulin intensity, (***C***)

IFT88 intensity, (***D***) ratio of IFT88 to acetylated α-tubulin and (***E***) ratio of IFT88 to

Smoothened-EGFP fluorescence across three independent experiments with each

shRNA in DAOY cells are plotted. For each plot, differences are highly significant

among shRNA treatments (p<2.2x10^{-16}, Kruskal-Wallis test) and between the 6 *ZNF423*

and control populations (p<2.2x10^{-16}, Wilcoxon rank sum test with continuity correction).

(**F**) Confocal images of GCP cilia from EGL of control littermate and *Zfp423* null mutant

(*nur12*) mice. Scale bar, 2 μm.　(**G**) Localization of IFT88 relative to acetylated α-tubulin

in cilia from five littermate pairs shows a lower proportion of mutant cilia with IFT88

throughout the axoneme; p=3.7x10^{-8}, chi-square test.　(**H**) Mean axonemal intensities of

IFT88 are significantly reduced compared to littermate controls.　Differences between

each littermate pair were independently significant.　Difference between genotypes for

summed data of 5 pairs is highly significant (p=2.2x10^{-15}, Wilcoxon rank sum test).

TULP3 overexpression is a functional target of *ZNF423* depletion.

To link the functional requirement for human ZNF423 and mouse Zfp423 to

potential target genes required for Shh response via cilia, we performed high-throughput

cDNA sequencing (RNA-Seq) on the DAOY cell model, followed by confirmation of

specific genes in freshly isolated mouse GCPs (Fig 7). RNA-Seq data were processed

for normalized counts of annotated genes and then assessed for significant differences

between *ZNF423* and control knockdown samples in three biological replicates, using

tools in the HOMER software package [38, 39]. Gene set enrichment analysis [40, 41]

did not identify any highly enriched pathways, but showed a strong relationship to genes

up-regulated in DAOY cells after knockdown of *PCGF2*, a polycomb repressor complex

protein [42]. Of >1000 genes that passed this initial threshold, 12 stood out for their

known roles in primary cilia (Fig 7A). To determine whether these differences were

conserved between the human medulloblastoma cell line model and mouse GCPs, we

measured expression of mouse orthologs to five genes that had FDR < 0.01 and clean

assays by RT-qPCR using freshly isolated GCPs from *Zfp423*^{nur12} and littermate controls

(Fig 7B). *Tulp3*, *Nek9*, and *Arl4d* showed significantly elevated expression in mutant

GCPs, as predicted by the DAOY cell knockdown model, while the other two were not

significantly different between genotypes. In both DAOY and GCP cells, *TULP3*/*Tulp3*

showed the most significant difference, being overexpressed >2-fold in each context.

A

Gene	Fold change	P-value	FDR	Rank
TULP3	2.16	3.3E-16	3E-13	31
WDR34	0.55	2.0E-11	7E-09	61
CEP170B	0.68	1.5E-05	0.001	331
NEK9	1.33	3.8E-05	0.002	395
FKBP8	0.76	9.4E-05	0.004	475
IFT70	2.59	1.0E-04	0.005	487
ARL4D	2.11	1.2E-04	0.004	501
WNT5A	0.71	3.7E-04	0.011	651
IFT25	1.73	5.6E-04	0.016	720
CILD18	0.78	1.0E-03	0.026	832
JBTS13	1.53	1.1E-03	0.026	833
GLI1	2.01	1.9E-03	0.039	1016

B

Fig 7. Tulp3 is an expression target of Zfp423 deficiency. (*A*) RNA-Seq data from knockdown and control DAOY cells identified several canonical cilium genes at high confidence. *TULP3* was among the largest magnitude changes and highest statistical confidence. (*B*) RT-qPCR from acutely isolated GCPs confirmed significantly increased expression of *Tulp3*, *Nek9*, and *Arl4d* in mutant relative to non-mutant controls among 7 littermate pairs. Nonparametric p-values from the Wilcoxon signed rank test are shown.

Chromatin immunoprecipitation (ChIP) experiments with independent antibodies demonstrated physical interaction of Zfp423 with the *Tulp3* gene in GCPs. Previous ChIP experiments quantified by high-throughput sequencing (ChIP-Seq) from a P19 cell culture model that expressed high levels of Zfp423, provided only modest read depth relative to cell number [32]. However, these data provided suggestive support for binding ~7 kb 5' to *Arl4d* (Fig 8A) and for sites within and around *Tulp3* (Fig 8B). Because of the low yield in ChIP-Seq experiments and desire to test biological replicate samples, we quantified ChIP at these candidate sites relative to a previously described marker locus, 259C14S [43, 44], by ChIP-qPCR. Both a commercial antibody (goat) and a custom, affinity-purified antibody (rabbit) were previously characterized by Western blotting and immunofluorescence [32]. Both antibodies supported ChIP enrichment of a site in *Tulp3* intron 1 (Tulp3_3) that also had the strongest support in the previous ChIP-Seq data (Fig 8C). Although enrichment over background was modest with the commercial antibody, it was also uniquely significant at the Tulp3_3 site (p=0.037, Tukey Honest Significant Difference test after one-factor ANOVA). Enrichment with the custom antibody showed stronger support for the same site (p=0.012) and a non-significant trend at the Tulp3_2 site (p=0.28). Non-parametric analysis of pooled data from both antibodies supported the same result (p=0.0087, Wilcoxon rank sum test for Tulp3_3 vs. control site after Bonferroni correction, with no other pairwise combination p>0.2). Neither antibody

showed enrichment for the other two candidate sites. Taken together, six independent

ChIP experiments from primary GCPs support a direct and selective physical association

of Zfp423 with sequences in *Tulp3* intron 1.

Fig 8. Chromatin immunoprecipitation detects Zfp423 binding in *Tulp3* intron 1. (***A***)

UCSC browser window on *Arl4d* locus shows location of the ChIP-PCR assay (Arl4d_1)

and a ChIP-Seq peak call from P19 cells (chr11-190), 7 kb 5' to the annotated transcript

(Arl4d). The peak notation indicates the 190[th] best-supported peak on chromosome 11 in

that experiment [32]. (***B***) Similar window on *Tulp3* locus includes 5 potential peaks from

P19 ChIP-Seq data, three of which supported reliable quantitative PCR assays

(Tulp3_1, _2 and _3) within 150 bp of the peak. (**C**) Graphs show fold enrichment

relative to an IgG control for a control locus and the four candidate binding sites. Colored

dots indicate measured values from independent biological replicates, bars indicate

median values for each locus. A commercial anti-Zfp423 antibody developed in goat

(blue dots) showed modest but significant enrichment for Tulp3_3. A custom antibody

developed in rabbit and affinity purified against the immunogen showed strong

enrichment for Tulp3_3 and possible enrichment for Tulp3_2, 3 kb away. Non-parametric

analysis of the combined data provides strong and conservative statistical support for

binding at Tulp3_3 (p=0.0087, Wilcoxon rank sum test with Bonferroni correction).

To test the functional importance of TULP3 overexpression in effecting ZNF423

depletion phenotype, we compared knockdown of *ZNF423* to simultaneous knockdown

of both *ZNF423* and *TULP3.* Combinations of targeting or control shRNAs were

delivered to DAOY cells using lentiviral vectors. Knockdown of *ZNF423* with either of

two non-overlapping shRNA depleted ZNF423 protein to ~50% of control levels (by

Western blot) and produced ~2-fold increase in TULP3 protein in each culture (Fig 9A),

even though only 60-75% cells appeared to be productively infected based on the

fluorescent reporter protein. Of three shRNAs directed against *TULP3*, only T2 and T3

were effective in reducing measured TULP3 protein level (Fig 9B). T2 and T3 reduced

TULP3 levels in *ZNF423*-knockdown cells to 70-90% of levels in control cells with full

ZNF423 expression. We then compared Smoothened-GFP translocation in doubly-

infected cells, with ~100 cilia per group in two independent experiments (Fig 9C),

following the methods and scoring described for *ZNF423* knockdown in Fig 5.

Simultaneous knockdown of *TULP3* significantly reversed the reduction in Smoothened-

GFP translocation caused by knockdown *ZNF423* in each replicate experiment

(p=0.0023 - 6.2x10-5, Pearson's chi-squared test, 2 degrees of freedom for high vs. low

TULP3). Each of the effective *TULP3* shRNAs showed similar result on Smoothened-

GFP distribution. These results indicate a functional role for TULP3 overexpression in

mediating defective signaling in the primary cilium of ZNF423-deficient cells.

Fig 9. Tulp3 is a functional target of Zfp423. (***A***) Western blot shows loss of ZNF423

and increase of TULP3 levels in DAOY cells infected with lentiviral shRNA vectors

targeting *ZNF423*. Relative amounts are compared to control shRNA infections after

normalization to β-actin and GAPDH. (***B***) Western blot shows the effect of dual infection

by *ZNF423* and three independent *TULP3* shRNA vectors on TULP3 protein. Only

shRNAs T2 and T3 reduced TULP3 levels compared with control vector. (***C***) Co-infection

with *ZNF423* and *TULP3* shRNA reverses the decrease in Shh-stimulated Smoothened

GFP translocation into cilia seen with shRNA to *ZNF423* and either scrambled control or

an ineffective *TULP3* shRNA. Graphs indicate high (red), medium (purple) and low

(blue) translocation as in Fig 5D,E. Chi-square

DISCUSSION

Our work provides, in both mouse brain and human cell culture models, the first

mechanistic connection between ZNF423/Zfp423 ciliopathy-related phenotypes and

function of the primary cilium. While phenotype severity and range of organ involvement

vary, cerebellar vermis hypoplasia is a uniquely consistent feature among reported

ZNF423 families [30] and five *Zfp423* mouse mutations on a variety of strain

backgrounds [23-26]. Previous work in mice demonstrated loss of precursor proliferation

[23], though the site of action has remained unresolved between competing proposals

for cell intrinsic (potentially cell-autonomous) mechanisms in the granule cell lineage and

non-autonomous mechanisms through Shh expression in Purkinje cells. Analysis of cell

culture models previously uncovered defective DNA damage response as a novel

cellular mechanism in *ZNF423* patient-derived mutations [30], but did not assess effects

on primary cilia and left open the question of whether ZNF423 might also act upstream

of the cilium to regulate its function or downstream by mitigating its effects. Results

presented here demonstrate a role for human and mouse homologs in cilium function

and in intrinsic properties of granule cell precursors, acting – at least in part – through

elevated expression of TULP3. Our work further unites cilium function and DNA damage

response mechanisms by demonstrating that defects in both can occur through

mutations for the same molecular component.

Four lines of evidence implicate *Zfp423* as a bona fide ciliopathy gene. First,

both recessive loss of function and apparently dominant mutations in mice and humans

produce phenotypes within the ciliopathy spectrum, including cerebellar vermis

hypoplasia in both species, nephronophthisis in humans, and a variable degree of other

consequences that may reflect allele strength or genetic, environmental or stochastic

modifiers [23, 24, 26, 30]. Second, our imaging analyses show that while *Zfp423*-null

animals make primary cilia at a normal frequency, loss of Zfp423 results in an abnormal

distribution of morphologies. Cilia on mutant GCPs, as a group, appear wider near the

base (Fig 3 and Fig 4), where several Joubert Syndrome and nephronophthisis proteins

are localized [45-48], and tend to be longer. Structures at the base act to regulate traffic

of signaling cargoes into and out of the cilium [49, 50]. This striking difference in cilium

morphologies could be explained by an altered temporal profile of dynamic states, as the

ranges of measurements largely overlap between genotypes despite having different

distributions. Third, signaling through the primary cilium is functionally defective after

loss of Zfp423 activity. *Zfp423*-depleted primary GCPs are refractory to exogenous Shh

(Fig 2). *ZNF423*-depleted DAOY medulloblastoma cells are reproducibly deficient in

translocation into the cilium of both Smoothened (Fig 5) and IFT88 (Fig 6), which are

required for normal signal processing [12, 13, 51-53] and specifically for expansion of

GCPs [15]. Diminished IFT88 translocation was also seen in mouse GCPs in situ (Fig

6). Fourth, we find expression level changes in key cilium components that are

consistent between *ZNF423*-knockdown DAOY cells and *Zfp423*-mutant GCPs and

indicative of disrupted signaling, at least one of which is functionally important and likely

due to direct regulation. TULP3 was the best-supported target of Zfp423 in our analysis.

TULP3 is an inhibitor of Shh signaling and loss of *Tulp3* in mice results in increased Shh

signaling [54-56]. At a cellular level, TULP3 regulates ciliary trafficking at both retrograde

transport and cilium entry through its interaction with IFT-A [57, 58]. ChIP experiments

in primary GCPs with two different antibodies against Zfp423 both showed significant

enrichment of a site in *Tulp3* intron 1 that was predicted from an earlier analysis in a P19

cell culture model [32]. Functionally, reversing the overexpression of TULP3 caused by

ZNF423 depletion significantly reversed the Smoothened translocation defects of

ZNF423 knockdown cells (Fig 9).

Although Zfp423 is required for mitogenic response to Shh ex vivo, residual

proliferation both in knockdown cells and in the EGL of null animals away from the

midline shows that Zfp423 is not strictly required for all mitogenic response by GCPs.

Moreover, the DAOY model shows residual translocation of both IFT88 and

Smoothened-EGFP in knockdown cells. One possibility might be that mitogenic

response by GCPs is more sensitive to loss of *Zfp423* when cultured ex vivo than either

GCPs in situ or the transformed DAOY model–and it is not unreasonable to expect that

both environmental factors and intrinsic cell differences relevant to establishment of a

cell line might enhance proliferative response. Alternatively (or additionally), it may be

that IFT-independent diffusion of Smoothened into the cilium [50] provides a modest

level of signaling that is saturated at lower levels of Shh than the translocation-

dependent pathway, as might be suggested by the difference in sensitivity of

Smoothened-EGFP and IFT88 localization to DAOY cell cilia (Fig 6E).

Tulp3 overexpression may be a surprising candidate for limiting Smoothened

translocation. Although Tulp3 antagonizes Shh signaling, loss of *Tulp3* does not affect

Smoothened trafficking [57, 59]. Rather, Tulp3 is thought to inhibit Shh signaling through

localization of Gpr161, whose activation elevates cAMP level to promote cleavage of Gli

proteins to their repressor forms [58]. However, this well-documented mechanism is

based on Tulp3 loss-of-function may not be the only inhibitory effect when Tulp3 is

expressed substantially above its normal level, as in ZNF423/Zfp423 deficient cells (Fig

7 and 9). Excess Tulp3 could in principle recruit additional inhibitory molecules to the

cilium or titrate limiting components that promote Smoothened trafficking, such as

Gprasp2 or Pifo [60, 61]. It is also possible that excess Tulp3 affects cell state in a way

that indirectly affects IFT88 and Smoothened translocation to or concentration in the

primary cilium. More work is needed to test such hypotheses, which are not mutually

exclusive. In addition, since Zfp423 contributes to regulation of several ciliopathy-related

genes, the full extent of molecular consequences is likely to be more complex and

interplay among dysregulated components may be required to fully resolve molecular

mechanisms behind the anatomical phenotypes seen in both patients and mouse

models.

Our results have some similarities to those with ciliogenic transcription factors of

the RFX family and Foxj1 [62, 63], but also important differences. RFX proteins are

highly conserved regulators that activate core components for formation of motile cilia

and loss of RFX activity results in loss or truncation of cilia [64-68]. *Foxj1* is a Shh target

gene that also activates components for motile cilia and is sufficient to drive ciliogenesis,

though loss of *Foxj1* does not generally prevent cilium formation [69, 70]. Restoring

expression of target genes partially mitigates cell phenotypes [71]. By contrast, *ZNF423*

patients and *Zfp423* mouse models present phenotypes more typical of defects in non-

motile cilia, especially cerebellum vermis hypoplasia [23, 24, 30] and Zfp423 is not

required for appearance of cilia. Zfp423 is required for efficient signaling through cilia in

cerebellum and in the cell models we tested. Zfp423 appears to modulate components,

often as a negative regulator and we see improvement of cell phenotypes by mitigating

the overexpression of the first clear repression target, *TULP3*, in a DAOY cell model.

The results here demonstrate a signaling mechanism, complementary to the

DNA damage response (DDR) mechanism demonstrated by Chaki et al. [30], for

neurodevelopmental phenotypes in *ZNF423*-deficient patients and orthologous *Zfp423*

mutant mice. In addition to providing a structural and functional basis for diminished Shh

signaling in granule cell precursors, our results point to loss of Zfp423 repressive

functions on *Tulp3* and potentially other genes as a key component of the phenotype.

Establishment of both DDR and signaling mechanisms provides a more comprehensive

basis for understanding both the variety and variability of disease presentations in this

ciliopathy, and potentially in others.

MATERIALS AND METHODS

Mice. The *nur12* mutation and BALB/c–*Zfp423*^{nur12} and C57BL/6–*Zfp423*^{nur12} congenic

mice have been previously described [23, 26, 72]. For timed matings, midnight on the

night of conception was considered to be E0.0. All procedures were approved by the UC

San Diego Institutional Animal Care and Use Committee.

Antibodies. Anti-BrdU antibody (mouse monoclonal B44) was purchased from BD

Biosciences.   anti-GFP antibody (A11122) purchased from Molecular Probes.

Antibodies against acetylated α-tubulin (mouse monoclonal 6-11B-1, T6793) and γ-

tubulin (rabbit polyclonal, T5192) were purchased from Sigma-Aldrich. Rabbit anti-IFT88

(13967-1-AP), TULP3 (13637-1-AP), and Arl13b (17711-1-AP) were purchased from

Proteintech. Mouse anti-GAPDH antibody (GT239) was purchased from GeneTex.

Rabbit antiserum against Human ZNF423 (amino acid residues 247-407 relative to

reference sequence NP_055884.2), was prepared and affinity-purified against the

immunogen [32]. Commercial antibodies against ZNF423 (Santa Cruz Biotechnology

sc-48785, aa 1-105; Sigma-Aldrich SAB2104426, aa 1234-1283) showed similar pattern

(L. Flores-Garcia and B.A.H.). Western blots were developed with infrared-conjugated

secondary antibodies (Rockland), detected on a Li-Cor Odyssey Imaging Station, and

quantified in the ImageJ software package.

Primary cell culture and mitotic index. Primary GCPs were isolated from E18.5 cerebella

of BALB/c mice by Percoll gradient centrifugation essentially as described [73, 74], but

omitting the adhesion step. Transfection experiments were performed with a minimum of

three independent DNA preparations for each construct. 48 hrs after transfection, cells

were exposed to 4 µg/ml BrdU (Sigma-Aldrich) for 6 hrs prior to fixation with 4%

paraformaldehyde. Recombinant Shh was purchased from R&D Systems. TRC shRNAs

708 and 709 directed against mouse *Zfp423* were obtained from Sigma (Table 1).

Gene expression assays. Quantitative RT-PCR was carried out on a Bio-Rad CFX-96

using SYBR Green fluorescence as described [43]. TaqMan hydrolysis probe assays

(Life Technologies; *Shh* assay Mm00436528_m1, *Zfp423* assay Mm00473699_m1,

Gapdh assay 4352339E.) were performed on the same instrument, following

manufacturer's recommended conditions. Relative quantification used the $\Delta\Delta C_T$ method

and comparisons across isolated cell fractions were expressed relative to the same ratio

in whole cerebellum.

Immunofluorescence and cilium measurements. For tissues, 10 μm sections were

prepared from 4% paraformaldehyde fixed animals at age E18.5. Confocal (Olympus

FV1000) and structured illumination "super resolution" microscopy (Applied Precision

OMX) were performed in the UCSD School of Medicine Light Microscopy Facility.

Volume measurements on the acquired images were accessed by Volocity software

(PerkinElmer). For Smoothened translocation measurements, a Smoothened-EGFP

fusion gene [75] was obtained from Addgene and cloned into pLKO.1 vector (Sigma),

along with shRNA constructs targeting either *ZNF423*, scrambled control, or irrelevant

sequence controls directed against reporter genes not used in this study. Human

medulloblastoma-derived DAOY cells were obtained from Dr. Robert Wechsler-Reya.

TRC shRNA directed against human *ZNF423* were obtained from Sigma (Table 1).

Virus particles were packaged by co-transfection of 293FT cells with pCMVdR8.2 dvpr

and pCMV-VSV-G (Addgene plasmid 8455 and 8454) as described [76]. Replicate

cultures of DAOY cells were infected with each virus for 5 hr., cultured 7 days and serum

starved for 24 hr prior to treatment with 0.1 μg/ml Shh for 24 hrs. Cells were fixed in 4%

paraformaldehyde and imaged by confocal microscopy. Sixty-two to 100 consecutive

images were collected for each sample and images from paired samples randomly

ordered (knockdown vs. control) into a grid by the flip of a coin. Distribution of

Smoothened-GFP relative to acetylated α-tubulin was assessed as a nominal-scale

variable by 5 different investigators blinded to genotype and experimental condition.

Mean Intensity along the axoneme of each channel was measured by ImageJ (National

Institute of Health, USA).

RNA-seq

ZNF423 knockdown and control DAOY cells were generated with a modified pLKO

lentivirus that expressed shRNA to ZNF423 or a non targeting control and co-expressed

the fluorescent protein mCherry. Cells expressing high level of mCherry were enriched

by flow cytometry and cultured in three replicate plates per virus. Bar-coded RNA

libraries were prepared for sequencing using strand specific dUTP protocols [39] with

minor modifications. Briefly, total RNA was harvested using Trizol reagent. Poly(A) +

RNA was enriched with Dynabeads® mRNA Purification Kit (Life Technologies,

Cat#61006). For each library, 500 ng poly(A) + RNA was premixed with 0.2 µg of

Oligo(dT) and random hexamer primer, heated to 70 °C for 10 min., and chilled on ice.

Reverse transcription reactions were incubated at room temperature for 10 min., 42 °C

1h and 50 °C 20 min. First-strand cDNA was ethanol precipitated with 0.5 volumes 5 M

ammonium acetate. After second-strand cDNA synthesis, sequencing libraries were

constructed as described [38]. 250 to 500 bp size-selected, adaptor-ligated cDNA was

incubated with 1 U USER (NEB) at 37 °C for 15 min. followed by 5 min. at 95 °C before

PCR with Phusion High-Fidelity DNA polymerase. High-throughput sequencing was

performed by UCSD IGM Genomics Center. Reads were aligned to the human hg19

transcriptome using STAR version 2.3.1f [77]. Data analysis was performed using

HOMER [38] and the Bioconductor package edgeR [78, 79]. Gene set enrichment was

performed in GSEA [40, 41]

Chromatin Immunoprecipitation. ChIP was performed as described [32]. Briefly, 5-10 x

10^6 purified primary GCPs were crosslinked in 1% formaldehyde, sonicated, and

subjected to standard ChIP purification with the indicated antibodies at ~2 µg.

Quantitative PCR from ChIP samples used 0.5% input fraction and ChIP with pre-

immune IgG for relative quantification among samples.

Statistical tests. Standard statistical procedures were performed in the R base

environment (R 2.8.1 GUI 1.27 Tiger build 32-bit (5301)). Normality (Shapiro-Wilk test)

and equal variance (F test) of each data set was assessed prior to selecting parametric

(ANOVA, *t*) or non-parametric (Kruskal-Wallis, Wilcoxon) tests. Tukey box-and-whisker

plots indicate median (heavy line), quartile (box), and 1.5 times interquartile range

(whiskers). Notches where shown indicate ±1.58 times the interquartile range/square

root of *n*.

ACCESSION NUMBERS

RNA-seq data have been submitted and can be accessed by the Gene Expression

Omnibus (GEO) accession number GSE59598.

ACKNOWLEDGEMENTS

The authors gratefully acknowledge Robert Wechsler-Reya for DAOY cells, Young-Wook Cho for affinity-purified ZNF423 antibody, Wendy Alcaraz, Dorothy Concepcion, Lisbeth Flores-Garcia, Kevin Ross, Isabella Sanchez, Weica Xie, and Susan Zhang for manual scoring of de-identified images, Jennifer Santini of the UCSD Neuroscience Microscopy Shared Facility for advice and assistance with instrumentation, Sven Heinz and Christopher K. Glass for advice on RNA-Seq library preparation and HOMER software, the UCSD Institute for Genomic Medicine Genomics Core for high-throughput sequencing and Lawrence S. B. Goldstein and Xiang-Dong Fu for comments on draft manuscripts. The authors declare no competing financial interests.

REFERENCES

1.   Wechsler-Reya RJ, Scott MP. Control of neuronal precursor proliferation in the
cerebellum by Sonic Hedgehog. Neuron. 1999;22(1):103-14. PubMed PMID: 10027293.

2.   Wallace VA. Purkinje-cell-derived Sonic hedgehog regulates granule neuron
precursor cell proliferation in the developing mouse cerebellum. Curr Biol.
1999;9(8):445-8. PubMed PMID: 10226030.

3.   Dahmane N, Ruiz-i-Altaba A. Sonic hedgehog regulates the growth and
patterning of the cerebellum. Development. 1999;126(14):3089-100. PubMed PMID:
10375501.

4.   Corrales JD, Blaess S, Mahoney EM, Joyner AL. The level of sonic hedgehog
signaling regulates the complexity of cerebellar foliation. Development.
2006;133(9):1811-21. Epub 2006/03/31. doi: 10.1242/dev.02351. PubMed PMID:
16571625.

5.   Marino S. Medulloblastoma: developmental mechanisms out of control. Trends
Mol Med. 2005;11(1):17-22. Epub 2005/01/15. doi: 10.1016/j.molmed.2004.11.008.
PubMed PMID: 15649818.

6.   Roussel MF, Hatten ME. Cerebellum development and medulloblastoma. Curr
Top Dev Biol. 2011;94:235-82. Epub 2011/02/08. doi: 10.1016/B978-0-12-380916-
2.00008-5. PubMed PMID: 21295689.

7.   Wechsler-Reya R, Scott MP. The developmental biology of brain tumors. Annu
Rev Neurosci. 2001;24:385-428. PubMed PMID: 11283316.

8.   Stanton BZ, Peng LF. Small-molecule modulators of the Sonic Hedgehog
signaling pathway. Mol Biosyst. 2010;6(1):44-54. Epub 2009/12/22. doi:
10.1039/b910196a. PubMed PMID: 20024066.

9.   Hyman JM, Firestone AJ, Heine VM, Zhao Y, Ocasio CA, Han K, et al. Small-
molecule inhibitors reveal multiple strategies for Hedgehog pathway blockade. Proc Natl
Acad Sci U S A. 2009;106(33):14132-7. Epub 2009/08/12. doi:
10.1073/pnas.0907134106. PubMed PMID: 19666565; PubMed Central PMCID:
PMC2721821.

10.   Romer J, Curran T. Targeting medulloblastoma: small-molecule inhibitors of the
Sonic Hedgehog pathway as potential cancer therapeutics. Cancer Res.
2005;65(12):4975-8. Epub 2005/06/17. doi: 10.1158/0008-5472.CAN-05-0481. PubMed
PMID: 15958535.

11.   Huangfu D, Liu A, Rakeman AS, Murcia NS, Niswander L, Anderson KV.
Hedgehog signalling in the mouse requires intraflagellar transport proteins. Nature.
2003;426(6962):83-7. Epub 2003/11/07. doi: 10.1038/nature02061. PubMed PMID:
14603322.

12.   Corbit KC, Aanstad P, Singla V, Norman AR, Stainier DY, Reiter JF. Vertebrate
Smoothened functions at the primary cilium. Nature. 2005;437(7061):1018-21. Epub
2005/09/02. doi: 10.1038/nature04117. PubMed PMID: 16136078.

13.   Haycraft CJ, Banizs B, Aydin-Son Y, Zhang Q, Michaud EJ, Yoder BK. Gli2 and
Gli3 localize to cilia and require the intraflagellar transport protein polaris for processing
and function. PLoS Genet. 2005;1(4):e53. Epub 2005/10/29. doi:
10.1371/journal.pgen.0010053. PubMed PMID: 16254602; PubMed Central PMCID:
PMC1270009.

14.   Huangfu D, Anderson KV. Cilia and Hedgehog responsiveness in the mouse.
Proc Natl Acad Sci U S A. 2005;102(32):11325-30. Epub 2005/08/03. doi:
10.1073/pnas.0505328102. PubMed PMID: 16061793; PubMed Central PMCID:
PMC1183606.

15.   Spassky N, Han YG, Aguilar A, Strehl L, Besse L, Laclef C, et al. Primary cilia
are required for cerebellar development and Shh-dependent expansion of progenitor
pool. Dev Biol. 2008;317(1):246-59. Epub 2008/03/21. doi: 10.1016/j.ydbio.2008.02.026.
PubMed PMID: 18353302.

16.   Tsai RY, Reed RR. Cloning and functional characterization of Roaz, a zinc finger
protein that interacts with O/E-1 to regulate gene expression: implications for olfactory
neuronal development. J Neurosci. 1997;17(11):4159-69. PubMed PMID: 9151733.

17.   Hata A, Seoane J, Lagna G, Montalvo E, Hemmati-Brivanlou A, Massague J.
OAZ uses distinct DNA- and protein-binding zinc fingers in separate BMP-Smad and Olf
signaling pathways. Cell. 2000;100(2):229-40. PubMed PMID: 10660046.

18.   Ku MC, Stewart S, Hata A. Poly(ADP-ribose) polymerase 1 interacts with OAZ
and regulates BMP-target genes. Biochem Biophys Res Commun. 2003;311(3):702-7.
PubMed PMID: 14623329.

19.   Holzel M, Huang S, Koster J, Ora I, Lakeman A, Caron H, et al. NF1 is a tumor
suppressor in neuroblastoma that determines retinoic acid response and disease
outcome. Cell. 2010;142(2):218-29. Epub 2010/07/27. doi: 10.1016/j.cell.2010.06.004.
PubMed PMID: 20655465; PubMed Central PMCID: PMC2913027.

20.   Huang S, Laoukili J, Epping MT, Koster J, Holzel M, Westerman BA, et al.
ZNF423 is critically required for retinoic acid-induced differentiation and is a marker of
neuroblastoma outcome. Cancer Cell. 2009;15(4):328-40. PubMed PMID: 19345331.

21.   Masserdotti G, Badaloni A, Green YS, Croci L, Barili V, Bergamini G, et al.
ZFP423 coordinates Notch and bone morphogenetic protein signaling, selectively up-
regulating Hes5 gene expression. J Biol Chem. 2010;285(40):30814-24. PubMed PMID:
20547764.

22.   Signaroldi E, Laise P, Cristofanon S, Brancaccio A, Reisoli E, Atashpaz S, et al.
Polycomb dysregulation in gliomagenesis targets a Zfp423-dependent differentiation
network. Nat Commun. 2016;7:10753. Epub 2016/03/01. doi: 10.1038/ncomms10753.
PubMed PMID: 26923714; PubMed Central PMCID: PMC4773478.

23.   Alcaraz WA, Gold DA, Raponi E, Gent PM, Concepcion D, Hamilton BA. Zfp423
controls proliferation and differentiation of neural precursors in cerebellar vermis
formation. Proc Natl Acad Sci U S A. 2006;103(51):19424-9. PubMed PMID: 17151198.

24.   Cheng LE, Zhang J, Reed RR. The transcription factor Zfp423/OAZ is required
for cerebellar development and CNS midline patterning. Dev Biol. 2007;307(1):43-52.
PubMed PMID: 17524391.

25.   Warming S, Rachel RA, Jenkins NA, Copeland NG. Zfp423 is required for normal
cerebellar development. Mol Cell Biol. 2006;26(18):6913-22. PubMed PMID: 16943432.

26.   Alcaraz WA, Chen E, Valdes P, Kim E, Lo YH, Vo J, et al. Modifier genes and
non-genetic factors reshape anatomical deficits in Zfp423-deficient mice. Hum Mol
Genet. 2011;20(19):3822-30. Epub 2011/07/07. doi: 10.1093/hmg/ddr300. PubMed
PMID: 21729880; PubMed Central PMCID: PMC3168291.

27.   Cheng LE, Reed RR. Zfp423/OAZ participates in a developmental switch during
olfactory neurogenesis. Neuron. 2007;54(4):547-57. PubMed PMID: 17521568.

28.   Gupta RK, Arany Z, Seale P, Mepani RJ, Ye L, Conroe HM, et al. Transcriptional
control of preadipocyte determination by Zfp423. Nature. 2010;464(7288):619-23.
PubMed PMID: 20200519.

29.   Zhang LJ, Guerrero-Juarez CF, Hata T, Bapat SP, Ramos R, Plikus MV, et al.
Innate immunity. Dermal adipocytes protect against invasive Staphylococcus aureus
skin infection. Science. 2015;347(6217):67-71. Epub 2015/01/03. doi:
10.1126/science.1260972. PubMed PMID: 25554785; PubMed Central PMCID:
PMC4318537.

30.   Chaki M, Airik R, Ghosh AK, Giles RH, Chen R, Slaats GG, et al. Exome Capture
Reveals ZNF423 and CEP164 Mutations, Linking Renal Ciliopathies to DNA Damage
Response Signaling. Cell. 2012;150(3):533-48. Epub 2012/08/07. doi:
10.1016/j.cell.2012.06.028. PubMed PMID: 22863007.

31.   Gold DA, Baek SH, Schork NJ, Rose DW, Larsen DD, Sachs BD, et al. RORa
coordinates reciprocal signaling in cerebellar development through sonic hedgehog and
calcium-dependent pathways. Neuron. 2003;40(6):1119-31. PubMed PMID: 14687547.

32.   Cho YW, Hong CJ, Hou A, Gent PM, Zhang K, Won KJ, et al. Zfp423 binds
autoregulatory sites in p19 cell culture model. PloS One. 2013;8(6):e66514. Epub
2013/06/14. doi: 10.1371/journal.pone.0066514. PubMed PMID: 23762491; PubMed
Central PMCID: PMC3675209.

33.   Jacobsen PF, Jenkyn DJ, Papadimitriou JM. Establishment of a human
medulloblastoma cell line and its heterotransplantation into nude mice. J Neuropathol
Exp Neurol. 1985;44(5):472-85. Epub 1985/09/01. PubMed PMID: 2993532.

34.   Peyrl A, Krapfenbauer K, Slavc I, Yang JW, Strobel T, Lubec G. Protein profiles
of medulloblastoma cell lines DAOY and D283: identification of tumor-related proteins
and principles. Proteomics. 2003;3(9):1781-800. Epub 2003/09/16. doi:
10.1002/pmic.200300460. PubMed PMID: 12973738.

35.   Wang Y, Zhou Z, Walsh CT, McMahon AP. Selective translocation of intracellular
Smoothened to the primary cilium in response to Hedgehog pathway modulation. Proc
Natl Acad Sci U S A. 2009;106(8):2623-8. Epub 2009/02/07. doi:
10.1073/pnas.0812110106. PubMed PMID: 19196978; PubMed Central PMCID:
PMC2650314.

36.   Kuzhandaivel A, Schultz SW, Alkhori L, Alenius M. Cilia-mediated hedgehog
signaling in Drosophila. Cell Rep. 2014;7(3):672-80. Epub 2014/04/29. doi:
10.1016/j.celrep.2014.03.052. PubMed PMID: 24768000.

37.   Chizhikov VV, Davenport J, Zhang Q, Shih EK, Cabello OA, Fuchs JL, et al. Cilia
proteins control cerebellar morphogenesis by promoting expansion of the granule
progenitor pool. J Neurosci. 2007;27(36):9780-9. Epub 2007/09/07. doi:
10.1523/JNEUROSCI.5586-06.2007. PubMed PMID: 17804638.

38.   Heinz S, Benner C, Spann N, Bertolino E, Lin YC, Laslo P, et al. Simple
combinations of lineage-determining transcription factors prime cis-regulatory elements
required for macrophage and B cell identities. Mol Cell. 2010;38(4):576-89. Epub
2010/06/02. doi: 10.1016/j.molcel.2010.05.004. PubMed PMID: 20513432; PubMed
Central PMCID: PMC2898526.

39.   Levin JZ, Yassour M, Adiconis X, Nusbaum C, Thompson DA, Friedman N, et al.
Comprehensive comparative analysis of strand-specific RNA sequencing methods. Nat
Methods. 2010;7(9):709-15. Epub 2010/08/17. doi: 10.1038/nmeth.1491. PubMed PMID:
20711195; PubMed Central PMCID: PMC3005310.

40.   Mootha VK, Lindgren CM, Eriksson KF, Subramanian A, Sihag S, Lehar J, et al.
PGC-1alpha-responsive genes involved in oxidative phosphorylation are coordinately
downregulated in human diabetes. Nat Genet. 2003;34(3):267-73. Epub 2003/06/17. doi:
10.1038/ng1180. PubMed PMID: 12808457.

41.   Subramanian A, Tamayo P, Mootha VK, Mukherjee S, Ebert BL, Gillette MA, et
al. Gene set enrichment analysis: a knowledge-based approach for interpreting genome-
wide expression profiles. Proc Natl Acad Sci U S A. 2005;102(43):15545-50. Epub
2005/10/04. doi: 10.1073/pnas.0506580102. PubMed PMID: 16199517; PubMed Central
PMCID: PMC1239896.

42.   Wiederschain D, Chen L, Johnson B, Bettano K, Jackson D, Taraszka J, et al.
Contribution of polycomb homologues Bmi-1 and Mel-18 to medulloblastoma
pathogenesis. Mol Cell Biol. 2007;27(13):4968-79. Epub 2007/04/25. doi:
10.1128/MCB.02244-06. PubMed PMID: 17452456; PubMed Central PMCID:
PMC1951487.

43.   Concepcion D, Flores-Garcia L, Hamilton BA. Multipotent genetic suppression of
retrotransposon-induced mutations by Nxf1 through fine-tuning of alternative splicing.
PLoS Genet. 2009;5(5):e1000484. Epub 2009/05/14. doi:
10.1371/journal.pgen.1000484. PubMed PMID: 19436707; PubMed Central PMCID:
PMC2674570.

44.   Floyd JA, Gold DA, Concepcion D, Poon TH, Wang X, Keithley E, et al. A natural
allele of Nxf1 suppresses retrovirus insertional mutations. Nat Genet. 2003;35(3):221-8.
PubMed PMID: 14517553.

45.   Badano JL, Mitsuma N, Beales PL, Katsanis N. The ciliopathies: an emerging
class of human genetic disorders. Annu Rev Genomics Hum Genet. 2006;7:125-48.
Epub 2006/05/26. doi: 10.1146/annurev.genom.7.080505.115610. PubMed PMID:
16722803.

46.   Lee JE, Gleeson JG. Cilia in the nervous system: linking cilia function and
neurodevelopmental disorders. Curr Opin Neurol. 2011;24(2):98-105. Epub 2011/03/10.
doi: 10.1097/WCO.0b013e3283444d05. PubMed PMID: 21386674.

47.   Louie CM, Gleeson JG. Genetic basis of Joubert syndrome and related disorders
of cerebellar development. Hum Mol Genet. 2005;14 Spec No. 2:R235-42. Epub
2005/10/26. doi: 10.1093/hmg/ddi264. PubMed PMID: 16244321.

48.   Wolf MT, Hildebrandt F. Nephronophthisis. Pediatr Nephrol. 2011;26(2):181-94.
Epub 2010/07/24. doi: 10.1007/s00467-010-1585-z. PubMed PMID: 20652329.

49.   Hu Q, Milenkovic L, Jin H, Scott MP, Nachury MV, Spiliotis ET, et al. A septin
diffusion barrier at the base of the primary cilium maintains ciliary membrane protein
distribution. Science. 2010;329(5990):436-9. Epub 2010/06/19. doi:
10.1126/science.1191054. PubMed PMID: 20558667; PubMed Central PMCID:
PMC3092790.

50.   Milenkovic L, Scott MP, Rohatgi R. Lateral transport of Smoothened from the
plasma membrane to the membrane of the cilium. J Cell Biol. 2009;187(3):365-74. Epub
2009/12/02. doi: 10.1083/jcb.200907126. PubMed PMID: 19948480; PubMed Central
PMCID: PMC2779247.

51.   Liu A, Wang B, Niswander LA. Mouse intraflagellar transport proteins regulate
both the activator and repressor functions of Gli transcription factors. Development.
2005;132(13):3103-11. Epub 2005/06/03. doi: 10.1242/dev.01894. PubMed PMID:
15930098.

52.   May SR, Ashique AM, Karlen M, Wang B, Shen Y, Zarbalis K, et al. Loss of the
retrograde motor for IFT disrupts localization of Smo to cilia and prevents the expression
of both activator and repressor functions of Gli. Dev Biol. 2005;287(2):378-89. Epub
2005/10/19. doi: 10.1016/j.ydbio.2005.08.050. PubMed PMID: 16229832.

53.   Caspary T, Larkins CE, Anderson KV. The graded response to Sonic Hedgehog
depends on cilia architecture. Dev Cell. 2007;12(5):767-78. Epub 2007/05/10. doi:
10.1016/j.devcel.2007.03.004. PubMed PMID: 17488627.

54.   Cameron DA, Pennimpede T, Petkovich M. Tulp3 is a critical repressor of mouse
hedgehog signaling. Dev Dyn. 2009;238(5):1140-9. Epub 2009/04/01. doi:
10.1002/dvdy.21926. PubMed PMID: 19334287.

55.   Norman RX, Ko HW, Huang V, Eun CM, Abler LL, Zhang Z, et al. Tubby-like
protein 3 (TULP3) regulates patterning in the mouse embryo through inhibition of

Hedgehog signaling. Hum Mol Genet. 2009;18(10):1740-54. Epub 2009/03/17. doi:
10.1093/hmg/ddp113. PubMed PMID: 19286674; PubMed Central PMCID:
PMC2671991.

56.   Patterson VL, Damrau C, Paudyal A, Reeve B, Grimes DT, Stewart ME, et al.
Mouse hitchhiker mutants have spina bifida, dorso-ventral patterning defects and
polydactyly: identification of Tulp3 as a novel negative regulator of the Sonic hedgehog
pathway. Hum Mol Genet. 2009;18(10):1719-39. Epub 2009/02/19. doi:
10.1093/hmg/ddp075. PubMed PMID: 19223390; PubMed Central PMCID:
PMC2671985.

57.   Mukhopadhyay S, Wen X, Chih B, Nelson CD, Lane WS, Scales SJ, et al.
TULP3 bridges the IFT-A complex and membrane phosphoinositides to promote
trafficking of G protein-coupled receptors into primary cilia. Genes Dev.
2010;24(19):2180-93. Epub 2010/10/05. doi: 10.1101/gad.1966210. PubMed PMID:
20889716; PubMed Central PMCID: PMC2947770.

58.   Mukhopadhyay S, Wen X, Ratti N, Loktev A, Rangell L, Scales SJ, et al. The
ciliary G-protein-coupled receptor Gpr161 negatively regulates the Sonic hedgehog
pathway via cAMP signaling. Cell. 2013;152(1-2):210-23. Epub 2013/01/22. doi:
10.1016/j.cell.2012.12.026. PubMed PMID: 23332756.

59.   Qin J, Lin Y, Norman RX, Ko HW, Eggenschwiler JT. Intraflagellar transport
protein 122 antagonizes Sonic Hedgehog signaling and controls ciliary localization of
pathway components. Proc Natl Acad Sci U S A. 2011;108(4):1456-61. Epub
2011/01/07. doi: 10.1073/pnas.1011410108. PubMed PMID: 21209331; PubMed Central
PMCID: PMC3029728.

60.   Jung B, Messias AC, Schorpp K, Geerlof A, Schneider G, Saur D, et al. Novel
small molecules targeting ciliary transport of Smoothened and oncogenic Hedgehog
pathway activation. Sci Rep. 2016;6:22540. Epub 2016/03/05. doi: 10.1038/srep22540.
PubMed PMID: 26931153; PubMed Central PMCID: PMC4773810.

61.   Jung B, Padula D, Burtscher I, Landerer C, Lutter D, Theis F, et al. Pitchfork and
Gprasp2 Target Smoothened to the Primary Cilium for Hedgehog Pathway Activation.
PLoS One. 2016;11(2):e0149477. Epub 2016/02/24. doi: 10.1371/journal.pone.0149477.
PubMed PMID: 26901434; PubMed Central PMCID: PMC4763541.

62.   Thomas J, Morle L, Soulavie F, Laurencon A, Sagnol S, Durand B.
Transcriptional control of genes involved in ciliogenesis: a first step in making cilia. Biol
Cell. 2010;102(9):499-513. Epub 2010/08/10. doi: 10.1042/BC20100035. PubMed
PMID: 20690903.

63.   Choksi SP, Lauter G, Swoboda P, Roy S. Switching on cilia: transcriptional
networks regulating ciliogenesis. Development. 2014;141(7):1427-41. Epub 2014/03/20.
doi: 10.1242/dev.074666. PubMed PMID: 24644260.

64.   Swoboda P, Adler HT, Thomas JH. The RFX-type transcription factor DAF-19
regulates sensory neuron cilium formation in C. elegans. Mol Cell. 2000;5(3):411-21.
Epub 2000/07/06. PubMed PMID: 10882127.

65.   Dubruille R, Laurencon A, Vandaele C, Shishido E, Coulon-Bublex M, Swoboda
P, et al. Drosophila regulatory factor X is necessary for ciliated sensory neuron
differentiation. Development. 2002;129(23):5487-98. Epub 2002/10/31. PubMed PMID:
12403718.

66.   Bonnafe E, Touka M, AitLounis A, Baas D, Barras E, Ucla C, et al. The
transcription factor RFX3 directs nodal cilium development and left-right asymmetry
specification. Mol Cell Biol. 2004;24(10):4417-27. Epub 2004/05/04. PubMed PMID:
15121860; PubMed Central PMCID: PMC400456.

67.   Ashique AM, Choe Y, Karlen M, May SR, Phamluong K, Solloway MJ, et al. The
Rfx4 transcription factor modulates Shh signaling by regional control of ciliogenesis. Sci
Signal. 2009;2(95):ra70. Epub 2009/11/06. doi: 10.1126/scisignal.2000602. PubMed
PMID: 19887680.

68.   Piasecki BP, Burghoorn J, Swoboda P. Regulatory Factor X (RFX)-mediated
transcriptional rewiring of ciliary genes in animals. Proc Natl Acad Sci U S A.
2010;107(29):12969-74. Epub 2010/07/10. doi: 10.1073/pnas.0914241107. PubMed
PMID: 20615967; PubMed Central PMCID: PMC2919930.

69.   Yu X, Ng CP, Habacher H, Roy S. Foxj1 transcription factors are master
regulators of the motile ciliogenic program. Nat Genet. 2008;40(12):1445-53. Epub
2008/11/18. doi: 10.1038/ng.263. PubMed PMID: 19011630.

70.   Cruz C, Ribes V, Kutejova E, Cayuso J, Lawson V, Norris D, et al. Foxj1
regulates floor plate cilia architecture and modifies the response of cells to sonic
hedgehog signalling. Development. 2010;137(24):4271-82. Epub 2010/11/26. doi:
10.1242/dev.051714. PubMed PMID: 21098568; PubMed Central PMCID:
PMC2990214.

71.   Lu H, Toh MT, Narasimhan V, Thamilselvam SK, Choksi SP, Roy S. A function
for the Joubert syndrome protein Arl13b in ciliary membrane extension and ciliary length
regulation. Dev Biol. 2015;397(2):225-36. Epub 2014/12/03. doi:
10.1016/j.ydbio.2014.11.009. PubMed PMID: 25448689.

72.   Kile BT, Hentges KE, Clark AT, Nakamura H, Salinger AP, Liu B, et al.
Functional genetic analysis of mouse chromosome 11. Nature. 2003;425(6953):81-6.
PubMed PMID: 12955145.

73.   Hatten ME. Neuronal regulation of astroglial morphology and proliferation in vitro.
J Cell Biol. 1985;100(2):384-96. Epub 1985/02/01. PubMed PMID: 3881455; PubMed
Central PMCID: PMC2113456.

74.   Hatten ME, Gao W-Q, Morrison ME, Mason CA. The Cerebellum: Purification
and Coculture of Identified Cell Populations. In: Banker G, Goslin K, editors. Culturing
Nerve Cells. 2nd ed. ed. Cambridge, MA: MIT Press; 1998.

75.   Chen JK, Taipale J, Cooper MK, Beachy PA. Inhibition of Hedgehog signaling by
direct binding of cyclopamine to Smoothened. Genes Dev. 2002;16(21):2743-8. Epub
2002/11/05. doi: 10.1101/gad.1025302. PubMed PMID: 12414725; PubMed Central
PMCID: PMC187469.

76.   Stewart SA, Dykxhoorn DM, Palliser D, Mizuno H, Yu EY, An DS, et al.
Lentivirus-delivered stable gene silencing by RNAi in primary cells. RNA. 2003;9(4):493-
501. Epub 2003/03/22. PubMed PMID: 12649500; PubMed Central PMCID:
PMC1370415.

77.   Dobin A, Davis CA, Schlesinger F, Drenkow J, Zaleski C, Jha S, et al. STAR:
ultrafast universal RNA-seq aligner. Bioinformatics. 2013;29(1):15-21. Epub 2012/10/30.
doi: 10.1093/bioinformatics/bts635. PubMed PMID: 23104886; PubMed Central PMCID:
PMC3530905.

78.   Anders S, McCarthy DJ, Chen Y, Okoniewski M, Smyth GK, Huber W, et al.
Count-based differential expression analysis of RNA sequencing data using R and
Bioconductor. Nat Protoc. 2013;8(9):1765-86. Epub 2013/08/27. doi:
10.1038/nprot.2013.099. PubMed PMID: 23975260.

79.   Robinson MD, McCarthy DJ, Smyth GK. edgeR: a Bioconductor package for
differential expression analysis of digital gene expression data. Bioinformatics.
2010;26(1):139-40. Epub 2009/11/17. doi: 10.1093/bioinformatics/btp616. PubMed
PMID: 19910308; PubMed Central PMCID: PMC2796818.
996

998 **TABLES**

999 Table 1: shRNA sequences

shRNA	Target species	Source	Lab name	Full name	Sequence
mZ1	mouse	TRC/ Sigma	708	TRCN0000 084708	CCGGCCCTGAATGTAACGTGAAGTTCTCGAGAACTTCACGTTA CATTCAGGGTTTTT
mZ2	mouse	TRC/ Sigma	709	TRCN0000 084709	CCGGCGGTGCATTACATGACTACATCTCGAGATGTAGTCATGT AATGCACCGTTTTT
hZ3	human	TRC/ Sigma	8173	TRCN0000 018173	CCGGGCAACGTTTGTTCACGGACTTCTCGAGAAGTCCGTGAAC AAACGTTGCTTTTT
hZ4	human	TRC/ Sigma	8174	TRCN0000 018174	CCGGCCACATGATTGAGGAAGGCATCTCGAGATGCCTTCCTCA ATCATGTGGTTTTT
T1	human	TRC/ Sigma	45	TRCN0000 078045	CCGGCCTACTATATGTACTTGGAAACTCGAGTTTCCAAGTACA TATAGTAGGTTTTT
T2	human	TRC/ Sigma	46	TRCN0000 078046	CCGGCCCAGTCAGCATGTTTAGAAACTCGAGTTTCTAAACATG CTGACTGGGTTTTT
T3	human	TRC/ Sigma	47	TRCN0000 078047	CCGGCCAGTTGATTTATCTCGTGAACTCGAGTTCACGAGATAA ATCAACTGGTTTTT
luc	–	TRC/ Sigma	luc	SHC007	CCGGCGCTGAGTACTTCGAAATGTCCTCGAGGACATTTCGAAG TACTCAGCGTTTTT
scr	–	TRC/ Sigma	non-human or mouse	SHC002	CCGGCAACAAGATGAAGAGCACCAACTCGAGTTGGTGCTCTTC ATCTTGTTGTTTTT

1000