Discrimination between *Onchocerca volvulus* and *O. ochengi* filarial larvae in *Simulium damnosum s.l.* and their distribution throughout central Ghana using a versatile high-resolution speciation assay

Doyle SR [1,a,b,c], Armoo S [1,2,a], Renz A [3], Taylor MJ[4], Osei-Atweneboana MY [2], Grant WN [1]

[1.] Department of Animal, Plant and Soil Sciences, La Trobe University, Bundoora, 3086, Australia

[2.] Council for Scientific and industrial Research - Water Research Institute, Accra, Ghana

[3.] Institute of Evolution and Ecology, Department of Comparative Zoology, University of Tübingen, Auf der Morgenstelle 28, 74074, Tübingen, Germany

[4.] Department of Parasitology, Liverpool School of Tropical Medicine, Liverpool, United Kingdom

[a.] both authors contributed equally to this study

[b.] corresponding author

[c.] current address: Wellcome Trust Sanger Institute, Wellcome Genome Campus, Hinxton, Cambridgeshire, CB10 1SA, United Kingdom

Stephen R Doyle: stephenrdoyle1@gmail.com

Samuel Armoo: sarmoo@students.latrobe.edu.au

Alfons Renz: Alfons.Renz@uni-tuebingen.de

Mark J Taylor: mark.taylor@lstmed.ac.uk

Mike Yaw Osei-Atweneboana: oseiatweneboana@yahoo.co.uk

Warwick N Grant: w.grant@latrobe.edu.au

Corresponding Author:

Stephen R Doyle

E: stephenrdoyle1@gmail.com

Abstract

Background

Genetic surveillance of the human filarial parasite, *Onchocerca volvulus*, from

onchocerciasis endemic regions will ideally focus on genotyping individual infective

larval stages collected from their intermediate host, Simuliid blackflies. However,

blackflies also transmit other *Onchocerca* species, including the cattle parasite *O.*

ochengi, which are difficult to distinguish from the human parasite based on

morphological characteristics alone. This study describes a versatile approach to

discriminate between *O. volvulus* and *O. ochengi* that is demonstrated using parasite

infective larvae dissected from blackflies.

Results

A speciation assay was designed based on genetic differentiation between *O.*

volvulus and *O. ochengi* mitochondrial genome sequences that can be performed in

high-throughput high-resolution melt- or lower throughput conventional RFLP

analyses. This assay was validated on 185 *Onchocerca* larvae dissected from

blackflies captured from 14 communities in Ghana throughout 2011-13. The

frequency of *O. ochengi* was approximately 67% of all larvae analysed, which is

significantly higher than previously reported in this region. Furthermore, the species

distribution was not uniform throughout the study region, with 25%, 47%, and 93%

of *O. volvulus* being found in the western-most (Black Volta, Tain and Tombe), the

central (Pru) and eastern-most (Daka) river basins, respectively.

Conclusions

This tool provides a simple and cost-effective approach to determine the identity

and distribution of two *Onchocerca* species, and will be valuable for future genetic

studies that focus on parasites collected from blackflies. The results presented

highlight the need to discriminate *Onchocerca* species in transmission studies, as the

frequency of each species varied significantly between the communities studied.

Keywords:

Onchocerciasis, Onchocerca volvulus, Onchocerca ochengi, high-resolution melt

analysis (HRM), restriction fragment length polymorphism (RFLP), speciation, Ghana

Background

Onchocerca volvulus is a human filarial parasite responsible for the disease onchocerciasis, and is transmitted between hosts by blackflies of the genus Simulium. Collection of blackflies throughout disease endemic regions forms a significant part of onchocerciasis surveillance campaigns to determine the prevalence and transmission of the parasite. Two approaches are commonly used; the first, pool-screening, has been extensively used to determine the prevalence of a single parasite in a pool of low hundreds of blackfly heads [1], which can be detected using an Onchocerca-specific O-150 polymerase chain reaction (PCR) assay together with an O. volvulus-specific DNA probe. Pool-screening has been the predominant tool used for parasite surveillance by the African Programme for Onchocerciasis Control (APOC, Burkina Faso), and is suitable particularly when the prevalence of O. volvulus is low in areas where widespread use of ivermectin has taken place. Adaptation of the pool-screening approach to detect O. volvulus has recently been demonstrated using loop-mediated isothermal amplification (LAMP) targeting either the O. volvulus glutathione S-transferase 1a gene [2], or the O-150 repeat (Nankoberanyi et al., in prep), which promises to improve the throughput, cost and sensitivity of the conventional approach. Although pool-screening has the capacity to examine large numbers of flies, the primary objective of the assay is simply to determine the presence or absence of the parasite in a defined number of vectors. The second approach focuses on the isolation of parasites by the dissection of individual blackflies, and pre-dates the use of pool-screening (having been in use since the initial Onchocerciasis Control Programme (OCP) in West Africa). Dissection has the advantage that it allows the developmental stage and location of the parasite(s) within the blackfly to be determined [3]. Data on larval prevalence in the vector are used together with the biting density of the fly to calculate the transmission potential of the parasite [4, 5].

Genetic diagnostic tools are under development by us and others that aim to genetically define parasite populations, characterise transmission zones, and screen parasite populations for polymorphisms associated with variation in response to ivermectin [6]. Such analyses could be integrated with and complement existing entomological surveillance practices using the vector stages of the parasite as a non-

invasive source of parasite material. Genetic analysis of entomological derived

parasites will ultimately require dissection of infected blackflies to obtain individual

parasite larvae, and subsequently, an efficient and sensitive means to validate the

species of parasite collected.

Species characterisation of dissected larvae is challenging (although not

impossible) based on morphological characteristics [5, 7, 8]; it requires significant

time and expertise, and is therefore relatively low-throughput. Some studies report

using O-150-based diagnostic assays [9, 10], whilst others describe the prevalence of

O. volvulus without molecular confirmation [11-14]. Given that a number of

Onchocerca parasites, including the common cattle parasite *O. ochengi*, can be

transmitted by the same intermediate vector [5], it is likely that *O. volvulus* larval

stages may often be overestimated, and in turn, negatively influence the predictions

of disease transmission. In this study, we report the development of a versatile

molecular tool to discriminate the onchocerciasis parasite *O. volvulus* from other

Onchocerca species such as *O. ochengi* based on mitochondrial DNA (mtDNA)

sequence variation. This approach was validated using *Onchocerca* larvae dissected

from blackflies that were collected from three distinct regions in Ghana.

Methods

Blackflies were sampled from 14 communities in 5 river basins in central Ghana

during 2011 (number of blackflies (n) = 12031), 2012 (n = 9706) and 2013 (n = 9138)

using a human landing collection protocol as described previously [15]. Ethical

approval was obtained for the use of human vector collectors from the Institutional

Review Board of the Council for Scientific and Industrial Research, Ghana. All

blackflies caught were dissected to collect *Onchocerca* larvae (129 from the head, 51

from the thorax, and 5 from the abdomen used here), after which all larvae collected

were dried on microscope slides for preservation. In total, 160 infective (L3),

together with 21 pre-infective L2 and 4 L1 larvae were analysed (Supplementary

Table 1) using the molecular techniques described, which included 14 larvae

collected from 8 infected flies in the 2011 cohort, 24 larvae from 11 flies collected in

2012, and 147 larvae from 109 flies collected in the 2013 cohort. Individual larva

were recovered from slides by applying 2 µl of HPLC-grade water (Sigma) directly

onto the specimen and allowing it to rehydrate, so that the larva would detach from

the slide or could be dislodged by a gentle scrape with a pipette tip. Each larva was

individually transferred to 20 µl of sample lysis solution (DirectPCR Lysis reagent

[MouseTail; Viagen Biotech] supplemented with 0.3 mg/ml (w/v) proteinase K

[Roche]), and incubated for 18 h at 55°C followed by 85°C for 1 h. Larval lysates were

diluted with HPLC-grade water 1/10 prior to PCR. Adult *O. volvulus* samples were

obtained as previously described [16]. Adult *O. ochengi* were collected from infected

cattle udders obtained from an abattoir in Ngaoundéré, Cameroon. Nodules were

collected from the skin, from which worms were extracted using forceps via a small

incision before being fixed in 95% ethanol. DNA was extracted from individual adult

O. volvulus and *O. ochengi* samples using an Isolate II Genomic DNA extraction kit

(Bioline, Australia) following the manufacturers' instructions.

High resolution melt analysis was performed after real time PCR amplification

of a 79-bp product using the Bio-Rad CFX system (Bio-Rad laboratories). A 10 µl

reaction mix was prepared by combining 5 µl of 2X SsoAdvanced Universal SYBR

Green qPCR Supermix (Bio-Rad Laboratories), 0.5 µl of each of SP_OVOO_79bp (5'-

GTTTGGTTCTTGTTGATTTG-3') and ASP_OVOO_79bp (5'-ACATTAAACCCCCTTACC-3')

primers (0.5 µM working concentration), 2 µl of HPLC grade water, and 2 µl of the

diluted larva lysate. The amplification protocol was performed as follows: 98°C for 3

min, followed by 45 cycles of 98°C for 10 sec, 59.5°C for 10 sec and 72°C for 10 sec.

The melt protocol was performed immediately after the amplification, and was as

follows: 95°C for 10 sec, followed by a 65°C annealing step from which the

temperature was increased by 0.2°C/plate read until a final temperature of 95°C.

PCR products generated from adult *O. volvulus* and *O. ochengi* samples were cloned

into the plasmid pGem-T-Easy (Promega) and sequenced to confirm the nucleotide

variants. All lysates, together with positive (cloned products and adult *O. volvulus*

and *O. ochengi* genomic DNA samples) and no-template controls (NTC), were

analysed in duplicate. Raw melt data was analysed using Precision Melt software

(version 1.2; Bio-Rad Laboratories), from which the data was exported into Microsoft

Excel for further quantitative analysis. Spatial (within flies i.e., head, thorax or

abdomen, and between river basins), temporal (sampling year) and larval stage (L1,

L2 or L3) differences between *O. volvulus* and *O. ochengi* were each examined using

a 2 × 3 chi-square (χ^2) test with 2 degrees of freedom, and a p-value < 0.05 was

considered to represent significant deviation from the null hypothesis.

A subset of post-PCR and HRM products were analysed by ApaI restriction

digest to confirm the *O. volvulus* specific digestion. A 20 µl reaction consisting of 10

µl PCR product together with 0.5 µl ApaI (50,000 U/ml; New England Biolabs), 2 µl of

CloneSmart buffer and 7.5 µl of HPLC-grade water was incubated for 3 h at 25°C,

after which the products were run on a 2% agarose gel at 100 V for 50 mins and

visualised using GelRed DNA stain (Biotum).

Results

Whole mitochondrial genome alignments of the *O. volvulus* (NC_001861; [17]) and

O. ochengi (obtained from genomic sequence available at

http://www.nematodes.org/genomes/onchocerca_ochengi/ ; Blaxter Lab, University

of Edinburgh) sequences was performed to identify a PCR compatible region that

contained (i) a restriction site that was unique to one species, and (ii) additional

polymorphism(s) that would result in a difference in the melting temperature

between amplicons generated for each of the two species. A 79-bp region spanning

8654- to 8733-bp of the *O. volvulus* mitochondrial genome was chosen that

contained three nucleotide T > C transitions (*O. volvulus*: C,C,C; *O. ochengi*: T,T,T)

between the mtDNA alignment of the two species, one of which was found in an

ApaI restriction site that is present in *O. volvulus* but absent in *O. ochengi* (Figure

1A). Melt curve analysis demonstrated a shift in Tm between the two sequences,

with a mean Tm of 79.8°C and 78.6°C for the *O. volvulus*- and *O. ochengi*-derived

sequences, respectively (Figure 1B). This translated into a significant and consistent

deviation in melt curves as depicted in the fluorescence difference plots (Figure 1C).

To confirm that the difference in melt curves was consistent with the prediction that

the ApaI restriction site was present in the *O. volvulus* sequences but not in the *O.*

ochengi sequences, 42 larvae-derived, 4 adult-derived and 2 cloned PCR products

were analysed by restriction digest, of which a representative gel is shown in Figure

1D. The melt curve and restriction digest data were 100% concordant in the samples

analysed, demonstrating that both approaches were equally predictive of the

species in question. This result does not, however, exclude the possibility that

genetic variation may exist within either or both species of *Onchocerca* examined (or

other potential *Onchocerca* species endemic to onchocerciasis regions [18-20]) that

will result in melt curves that deviate from the *O. volvulus* and *O. ochengi* control

sequences described here. For example, an analysis of the mitochondrial sequence

of the *O. ochengi* Siisa variant [18] revealed a T,T,C haplotype that differed from the

T,T,T haplotype of the adult *O. ochengi* presented (Dr Adrian Streit, personal

communication). Although a single base change is unlikely to confound the

interpretation between the melt profiles of the *O. volvulus* and *O. ochengi* control

samples described here, this platform does offer further opportunity to explore

genetic variation among and within species. Therefore, amplicons giving rise to melt

curves that do not match controls should be sequenced.

Of the 141 flies from which *Onchocerca* larvae were recovered, 43 flies

contained more than a single larva (range: 2-6). Interestingly, 4 of the 43 flies (9.3%

of flies with multiple larva, 2.8% of total) contained both *O. volvulus* and *O. ochengi*

larvae, which must represent larval uptake by the blackfly from two blood meals

from different hosts at different times. This is not surprising, given the prevalence of

cattle and human cohabitation, and that blackflies will seek a blood meal from either

host [21, 22]. The overall proportion of each larval stage was similar between both

species (χ^2 = 3.397; df = 2; p = 0.091), however, the spatial distribution of each stage

within the fly was not equal; although a greater proportion of L3 were found in the

head relative to the thorax in both species, the difference between both head and

thorax was greatest in *O. volvulus* with a higher proportion of L3 collected from the

head (Supplementary Figure; χ^2 = 6.822; df = 2; p = 0.016), which suggests that while

the rate of development within the fly may be similar between the species, *O.*

volvulus may migrate towards the head of the fly earlier that *O. ochengi*. These

observations (co-infection and likely equivalent transmission potential (based on

prevalence of L3)), do suggest that the human and bovine hosts are constantly

exposed to both parasites which, in turn, raises an interesting question in regard to

the number of times host-switching may have occurred between human and cattle

(a speciation hypothesis whereby the most recent common ancestor of *O. onchengi*

and *O. volvulus* was a cattle parasite that established in the human host [23]).

The distribution of both species throughout the sampling region was not

uniform (Figure 2). Although no temporal differences between sampling years for

each species was seen (χ^2 = 1.075; df = 2, p = 0.292), a significant spatial trend was

observed that suggested that the western communities sampled had a lower

prevalence of *O. volvulus* (24.4%, n = 136; Black Volta, Tain and Tombe river basins),

a roughly equal prevalence of *O. volvulus* and *O. ochengi* in the central Pru river

basin (47.2% *O. volvulus*; n = 36), and a higher prevalence of *O. volvulus* (92.9%, n =

14) in the eastern-most Daka river basin (χ^2 = 29.888; df = 2, p = 1.618E-7). These

results are not necessarily in contrast to a recent investigation of persistent *O.*

volvulus with negligible *O. ochengi* transmission in Ghana [9]. Much of that study

was focused on two southern Ghanaian communities, and very few or no larvae

were found in the two communities that were shared with this study (Asubende &

Agborlekame, respectively); moreover, 3 of the 7 communities used by Lamberton *et*

al. were east of the Black Volta Lake, which would be consistent with high *O.*

volvulus prevalence observed in the eastern-most community (Wiae) sampled here.

We speculate that the high infection rate reported and difference in prevalence of *O.*

ochengi in a number of study regions presented here is correlated with the high

numbers of cattle in the north-western river basins, particularly during the dry

season (December to April). However, given that the sample size of larvae for many

communities was low (median = 6.5, range = 1-55 larvae/community; limited by the

number of larvae present in the blackfly populations and therefore by the number of

blackflies screened), further sampling is required to support these findings,

particularly in the Daka river basin where *Onchocerca* larvae were obtained from

only a single community out of the five communities in which flies were sampled.

Conclusions

A simple but versatile tool that discriminates between *O. volvulus* and *O. ochengi*

single larvae obtained by dissection of the blackfly intermediate host is described,

and offers a number of advantages over traditional O-150-based speciation assays.

By exploiting the multicopy nature of the mitochondrial genome, this assay is very

sensitive, requiring only 1/100th of a single larva, thus leaving sufficient material for

further genetic analysis. Although both *O. volvulus* and *O. ochengi* samples could be

distinguished easily via HRM or RFLP assays, no effort was made to identify other

described *Onchocerca* species that are potentially present [18-20]; further work is

required to characterise melt profiles and/or restriction sites from additional

Onchocerca spp. voucher specimens as they become available. In the context of

onchocerciasis entomological surveillance, which aims to detect transmission of the

human pathogen, this study demonstrates the need to accurately and efficiently

determine the identity of larvae in the vector in order to correctly estimate parasite

transmission. In Ghana, nationwide sampling of blackfly populations is necessary to

determine the distribution of *Onchocerca* spps. Implementation of genetic tools such

as those described here will inform the appropriate control strategy needed by

onchocerciais control programmes as they strive towards onchocerciasis elimination

in Ghana.

Competing Interests:

The authors have no competing interests to declare.

Author's Contributions:

SRD designed the study, performed the molecular biology and data analysis, and

drafted the manuscript. MYO-A, MT and SA designed the entomological study for

transmission assessment, SA was involved in the collection and dissection of

blackflies, and preparation of larvae samples. MYO-A and MT was involved in the

coordination of sample collection. AR provided samples. WG participated in the

analysis and helped draft the manuscript. All authors read and approved the final

version of the manuscript.

Acknowledgments:

We thank Mr. Francis Veriegh for contributing to the blackfly dissection, and PD Dr

Adrian Streit, Dr Annette Kuesel and Dr Aime Adjami for helpful suggestions toward

the manuscript. This investigation received financial support from TDR, the Special

Programme for Research and Training in Tropical Diseases, co-sponsored by UNICEF,

UNDP, The World Bank and WHO, the German Research Foundation (DFG Re

1536/5-1) and the European Foundation Initiative for Neglected Tropical Diseases.

We thank Dr Mark Blaxter and Dr Benjamin Makepeace for access to the

unpublished *O. ochengi* genome sequence, which was generated with support of the

EU-funded programme "Enhancing Protective Immunity Against Filariasis".

Supplementary Table 1. Distribution of communities and samples used in this study

Community	River Basin	Geographic coordinates		Larvae processed and species identified by HRM per sample collection year [1]						Total larvae per community
		Latitude	Longitude	2011 (12,031 flies dissected)		2012 (9,706 flies dissected)		2013 (9,138 flies dissected)		
				O. volvulus	*O. ochengi*	*O. volvulus*	*O. ochengi*	*O. volvulus*	*O. ochengi*	
Abua (ABA)	Pru	7.58587	-0.53485	1	-	-	-	-	-	*1*
Agbelekame I (AB)	Black Volta	8.14018	-2.12185	2	2	1	4	8	9 L2; 27 L3 ;	*53*
Akanyakrom (AKA)	Black Volta	8.289	-2.277	-	-	5	-	-	-	*5*
Asubende (ASU)	Pru	8.01085	-0.58528	-	-	2	-	5	2	*9*
Baaya (BAY)	Pru	8.01119	-0.59444	-	-	-	-	2	1 L2; 12 L3	*15*
Beposo (BEP)	Pru	8.0026	-0.57395	-	-	-	-	1	-	*1*
Fawoman (FAW)	Pru	8.0111	-1.01303	-	-	1	2	-	-	*3*
Fawoman-Banda (FAB)	Tombe	8.07108	-2.1441	-	-	-	6	2	-	*8*
Kyingakrom (KYN)	Black Volta	8.05578	-2.03266	-	3	-	-	-	-	*3*
Mantukwa (MAN)	Pru	8.01064	-1.0014	-	-	-	-	2	2	*4*
New Longoro (NLG)	Black Volta	8.08158	-2.02096	-	1	-	-	1	1 L2; 8 L3	*11*
Senyase (SEN)	Pru	8.01247	-0.59251	-	-	-	-	1 L1;1 L2;1 L3	-	*3*
Tainso (TAN)	Tain	8.0603	-2.06538	-	5	-	3	1 L1; 4 L2; 7 L3	1 L1; 5 L2; 29 L3	*55*
Wiae (WIA)	Daka	8.19219	-0.09429	-	-	-	-	1 L1; 12 L3	1	*14*
Total larvae processed per species per year				*3*	*11*	*3*	*15*	*49 (3 L1; 5 L2; 41 L3)*	*98 (1 L1; 16 L2; 81 L3)*	*185*

[1] L3 larvae unless otherwise stated

Figure 1. Genetic discrimination of *O. volvulus* and *O. ochengi*. (A) Sequence of the

PCR 79-bp amplicon used to discriminate between *O. volvulus* and *O. ochengi*.

Nucleotide differences between the two species are highlighted in bold, and the

ApaI restriction site present in the *O. volvulus* sequence is underlined. (B)

Normalised HRM data depicting melt curves from individual *O. volvulus* (green) and

O. ochengi (red) samples. Data presented includes cloned positive control 79-bp

products (pGem_Ov and pGem_Oo), adult DNA, and 4 larvae samples from both

species, each in duplicate. (C) The same samples are presented as in (B), showing the

difference curves normalised to the *O. ochengi* group. (D) Representative RFLP

analysis of amplicons generated in the HRM assay for each species, showing

digestion of the *O. volvulus* sequence, but not the *O. ochengi* sequence, with ApaI.

Figure 2. Distribution of *O. volvulus* and *O. ochengi* throughout Ghana. Pie charts

depict relative prevalence of *O. volvulus* (orange) and *O. ochengi* (purple) in each

community. Communities sampled in central Ghana (see insert) and the number of

samples analysed are as follows: Abua (ABA; n = 1), Agbelekame I (AB; n = 54),

Akanyakrom (AKA; n = 5), Asubende (ASU; n = 9), Baaya (BAY; n = 15), Beposo (BEP, n

= 1), Fawoman (FAW; n = 3), Fawoman-Banda (FAB, n = 8), Kyingakrom (KYN; n = 3),

Mantukwa (MAN; n = 4), New Longoro (NLG; n = 11), Senyase (SEN; n = 3), Tainso

(TAN; n = 55), Wiae (WIA; n = 14). The colour coding of communities reflect the river

basin in which they lie; Black Volta (black), Tain (dark blue), Tombe (light blue) Pru

(green) and Daka (red).

References

1. Katholi CR, Toe L, Merriweather A, Unnasch TR: **Determining the prevalence of Onchocerca volvulus infection in vector populations by polymerase chain reaction screening of pools of black flies**. *The Journal of infectious diseases* 1995, **172**(5):1414-1417.

2. Alhassan A, Makepeace BL, LaCourse EJ, Osei-Atweneboana MY, Carlow CK: **A Simple Isothermal DNA Amplification Method to Screen Black Flies for Onchocerca volvulus Infection**. *PloS one* 2014, **9**(10):e108927.

3. Cheke RA, Garms R: **Indices of onchocerciasis transmission by different members of the Simulium damnosum complex conflict with the paradigm of forest and savanna parasite strains**. *Acta tropica* 2013, **125**(1):43-52.

4. Duke BO: **Studies on factors influencing the transmisson of onchocerciasis. IV. The biting-cycles, infective biting density and transmission potential of "forest" Stimulium dannosum**. *Annals of tropical medicine and parasitology* 1968, **62**(1):95-106.

5. Eisenbarth A, Achukwi MD, Renz A: **Ongoing Transmission of Onchocerca volvulus after 25 Years of Annual Ivermectin Mass Treatments in the Vina du Nord River Valley, in North Cameroon**. *PLoS neglected tropical diseases* 2016, **10**(2):e0004392.

6. Osei-Atweneboana MY, Eng JK, Boakye DA, Gyapong JO, Prichard RK: **Prevalence and intensity of Onchocerca volvulus infection and efficacy of ivermectin in endemic communities in Ghana: a two-phase epidemiological study**. *Lancet* 2007, **369**(9578):2021-2029.

7. Wahl G, Schibel JM: **Onchocerca ochengi: morphological identification of the L3 in wild Simulium damnosum s.l., verified by DNA probes**. *Parasitology* 1998, **116 (Pt 4)**:337-348.

8. McCall PJ, Townson H, Trees AJ: **Morphometric differentiation of Onchocerca volvulus and O. ochengi infective larvae**. *Transactions of the Royal Society of Tropical Medicine and Hygiene* 1992, **86**(1):63-65.

9. Lamberton PH, Cheke RA, Winskill P, Tirados I, Walker M, Osei-Atweneboana MY, Biritwum NK, Tetteh-Kumah A, Boakye DA, Wilson MD *et al*: **Onchocerciasis transmission in Ghana: persistence under different control strategies and the role of the simuliid vectors**. *PLoS neglected tropical diseases* 2015, **9**(4):e0003688.

10. Adeleke MA, Mafiana CF, Sam-Wobo SO, Olatunde GO, Ekpo UF, Akinwale OP, Toe L: **Biting behaviour of Simulium damnosum complex and Onchocerca volvulus infection along the Osun River, Southwest Nigeria**. *Parasites & vectors* 2010, **3**:93.

11. Katabarwa MN, Eyamba A, Nwane P, Enyong P, Kamgno J, Kuete T, Yaya S, Aboutou R, Mukenge L, Kafando C *et al*: **Fifteen years of annual mass treatment of onchocerciasis with ivermectin have not interrupted transmission in the west region of Cameroon**. *Journal of parasitology research* 2013, **2013**:420928.

12. Katabarwa MN, Eyamba A, Nwane P, Enyong P, Yaya S, Baldiagai J, Madi TK, Yougouda A, Andze GO, Richards FO: **Seventeen years of annual distribution of ivermectin has not interrupted onchocerciasis transmission in North**

Region, Cameroon. *The American journal of tropical medicine and hygiene*
2011, **85**(6):1041-1049.

13.   Opara KN, Fagbemi OB, Ekwe A, Okenu DM: **Status of forest onchocerciasis**
in the Lower Cross River basin, Nigeria: entomologic profile after five years
of ivermectin intervention. *The American journal of tropical medicine and*
hygiene 2005, **73**(2):371-376.

14.   Wanji S, Kengne-Ouafo JA, Esum ME, Chounna PW, Tendongfor N, Adzemye
BF, Eyong JE, Jato I, Datchoua-Poutcheu FR, Kah E *et al*: **Situation analysis of**
parasitological and entomological indices of onchocerciasis transmission in
three drainage basins of the rain forest of South West Cameroon after a
decade of ivermectin treatment. *Parasites & vectors* 2015, **8**(1):202.

15.   Walsh JF, Davies JB, Le Berre R: **Entomological aspects of the first five years**
of the Onchocerciasis Control Programme in the Volta River Basin.
Tropenmedizin und Parasitologie 1979, **30**(3):328-344.

16.   Osei-Atweneboana MY, Awadzi K, Attah SK, Boakye DA, Gyapong JO, Prichard
RK: **Phenotypic evidence of emerging ivermectin resistance in Onchocerca**
volvulus. *PLoS neglected tropical diseases* 2011, **5**(3):e998.

17.   Keddie EM, Higazi T, Unnasch TR: **The mitochondrial genome of Onchocerca**
volvulus: sequence, structure and phylogenetic analysis. *Molecular and*
biochemical parasitology 1998, **95**(1):111-127.

18.   Eisenbarth A, Ekale D, Hildebrandt J, Achukwi MD, Streit A, Renz A: **Molecular**
evidence of 'Siisa form', a new genotype related to Onchocerca ochengi in
cattle from North Cameroon. *Acta tropica* 2013, **127**(3):261-265.

19.   Wahl G: **Identification of a common filarial larva in Simulium damnosum s.l.**
(type D, Duke, 1967) as Onchocerca ramachandrini from the wart hog. *The*
Journal of parasitology 1996, **82**(3):520-524.

20.   Wahl G, Renz A: **Transmission of Onchocerca dukei by Simulium bovis in**
North-Cameroon. *Trop Med Parasitol* 1991, **42**(4):368-370.

21.   Achukwi MD, Harnett W, Renz A: **Onchocerca ochengi transmission**
dynamics and the correlation of O. ochengi microfilaria density in cattle
with the transmission potential. *Veterinary research* 2000, **31**(6):611-621.

22.   Lamberton PH, Cheke RA, Walker M, Winskill P, Osei-Atweneboana MY,
Tirados I, Tetteh-Kumah A, Boakye DA, Wilson MD, Post RJ *et al*:
Onchocerciasis transmission in Ghana: biting and parous rates of host-
seeking sibling species of the Simulium damnosum complex. *Parasites &*
vectors 2014, **7**:511.

23.   Morales-Hojas R, Cheke RA, Post RJ: **Molecular systematics of five**
Onchocerca species (Nematoda: Filarioidea) including the human parasite,
O. volvulus, suggest sympatric speciation. *Journal of helminthology* 2006,
80(3):281-290.

A

O. ochengi   GTTTAGTTCTTGTTGATTTGCATCGGGC**T**CCTTTTGAT**T**TTTCTGAGTGTGAAAGTGAG**T**TGGTAAGGGGGTTTAATGT

O. volvulus GTTTGGTTCTTGTTGATTTGCATC<u>GGGC**C**C</u>CTTTTGAT**C**TTTCTGAGTGTGAAAGTGAG**C**TGGTAAGGGGGTTTAATGT

Supplementary Figure 1. Proportion of *O. volvulus* (orange) and *O. ochengi* (purple) larval life stages (L3, L2, & L1) found during dissection of the blackfly head, thorax and abdomen. Values reported are a percentage of the total larvae found within each species.