1

The promise of disease gene discovery in South Asia

Nathan Nakatsuka[1,2], Priya Moorjani[1,3], Niraj Rai[4], Biswanath Sarkar[5], Arti Tandon[1,6], Nick Patterson[6], Lalji Singh[4], David Reich[1,6,7,*], Kumarasamy Thangaraj[4,*]

[1]Department of Genetics, Harvard Medical School, New Research Building, 77 Ave. Louis Pasteur, Boston, MA 02115, USA

[2]Harvard-MIT Division of Health Sciences and Technology, Harvard Medical School, Boston, MA 02115, USA

[3]Department of Biological Sciences, Columbia University, 600 Fairchild Center, New York, NY 10027, USA

[4]CSIR-Centre for Cellular and Molecular Biology, Habsiguda, Hyderabad, Telangana 500007, India

[5]Anthrological Survey of India, Kolkata, West Bengal 700016, India

[6]Broad Institute of Harvard and Massachusetts Institute of Technology, Cambridge, MA 02141, USA

[7]Howard Hughes Medical Institute, Harvard Medical School, Boston, MA 02115, USA

It is tempting to think of the more than 1.5 billion people who live in South Asia as one large ethnic group, but in fact, South Asia is better viewed as comprised of very many small endogamous groups that usually marry within their own group (caste or tribe). To perform a high resolution assessment of South Asian demography, we assembled genome-wide data from over 2,000 individuals from over 250 distinct South Asian groups, more than tripling the number of diverse India groups for which such data are available, and including tribe and caste groups sampled from every state in India. We document shared ancestry across groups that correlates with geography, language, and caste affiliation, and characterize the strength of the founder events that gave rise to many of these groups. Over a third of the groups— including eighteen with census sizes of more than a million—descend from founder events stronger than those in Ashkenazi Jews and Finns, both of which have high rates of recessive disease due to their histories of strong founder events. These results highlight a major and unappreciated opportunity for reducing the disease burden among South Asians through the discovery of and genetic testing for recessive disease genes.

South Asia is a region of extraordinary cultural, linguistic, and genetic diversity, with a conservative estimate of over 4,600 anthropologically well-defined groups, many of which are endogamous communities with significant barriers to gene flow due to sociological and cultural factors that restrict intermarriage[1]. Of the small fraction of South Asian groups that have been characterized using genome-wide data, many exhibit large allele frequency differences from geographically proximal neighbors[2-4], indicating that they have experienced strong founder events, whereby a small number of ancestors gave rise to the many descendants that exist today[4]. The evidence that a substantial fraction of groups in South Asia might descend from founder events represents a major opportunity for improving health. Detailed studies of founder populations of European ancestry, including Ashkenazi Jews, Finns, Amish, Hutterites, Sardinians, and French Canadians, have resulted in the discovery of dozens of rare recessive diseases in each group, allowing genetic

counseling that has helped reduce disease burden in each of these communities[5].

Opportunities for improving health through founder event disease mapping in India

are even greater due to more widespread endogamy.

To characterize the medically relevant founder events in India, we carried out new

genotyping of 890 samples from 206 endogamous groups in India on the Affymetrix

Human Origins single nucleotide polymorphism (SNP) array[6]. Based on power

calculations to determine the number of samples needed to confidently detect a

founder event at least as strong as that in Ashkenazi Jews or Finns (Supplementary

Figure 1), we aimed in most cases to genotype up to five individuals per group.

Previous studies that sampled the genetic diversity of South Asia focused to a

disproportionate extent on tribal groups and castes with small census sizes in order

to capture the largest possible amount of anthropological diversity[3,4,7-9]. In this

study, our sampling included many groups with large census sizes to investigate the

prospects for future disease gene mapping. We combined the new data we collected

with previously reported data, leading to four datasets (Figure 1a). The Affymetrix

Human Origins SNP array data comprised 1,192 individuals from 231 groups in

South Asia, to which we added 7 Ashkenazi Jews. The Affymetrix 6.0 SNP array data

comprised 383 individuals from 52 groups in South Asia[4,8]. The Illumina SNP array

data comprised 188 individuals from 21 groups in South Asia[9] and 21 Ashkenazi

Jews[9,10]. The Illumina Omni SNP array data comprised 367 individuals from 20

groups in South Asia[7]. We merged 1000 Genomes Phase 3 data (2,504 individuals

from 26 different groups including 99 Finns) with each of these datasets. We

performed quality control to remove SNPs and individuals with a high proportion of

missing genotypes or those that were outliers in Principal Component Analysis

(PCA). To remove close relatives, we also removed one individual from each pair

that were outliers in their group for Identity-by-Descent (IBD) genomic segments,

and we removed all IBD segments that were larger than 20 centimorgans (cM).

We performed PCA on each of the three different datasets along with European

Americans (CEU), Han Chinese (CHB), and West Africans (YRI), and found that the

Siddi are strong outliers as previously reported (Supplementary Figure 2)[4,11,12]. We

next removed YRI, Siddi and indigenous Andamanese (another known outlier) from

the datasets and repeated PCA (Figure 1b, Supplementary Figure 3). The PCA

documents three broad clusters[4,8,7]. First, almost all Indian groups speaking Indo-

European and Dravidian languages lie along the "Indian Cline," reflecting the fact

that they are admixed, with different proportions of Ancestral Northern Indian

(ANI) ancestry related to Europeans, Central Asians, and Near Easterners; and

Ancestral Southern Indian (ASI) ancestry that is as different from ANI as Europeans

and East Asians are from each other[4]. The second major cluster includes groups that

speak Austroasiatic languages, as well as some non-Austroasiatic speaking groups

that have similar ancestry possibly due to gene flow with Austro-Asiatic speaking

neighbors or to a history of language shift. This set of groups cluster together near

the ASI end of the Indian cline, likely reflecting a large proportion of ASI-like

ancestry as well as a distinct ancestry that has some affinity to East Asians. The

Tibeto-Burmese speaking groups and other groups with high proportions of East

Asian ancestry such as the Austroasiatic speaking Khasi and Tharu form a gradient

of ancestry relating them to East Asian groups such as Han Chinese. These three

clusters are also evident in a neighbor-joining tree based on F_{ST} (Supplementary

Figure 4). We confirmed the East Asian related admixture in some groups using the

statistic f_3(Test; Mala, Chinese); significantly negative values of this statistic provide

unambiguous evidence that the *Test* population is admixed of populations related

(perhaps distantly) to Mala (an Indian Cline group with high ASI ancestry) and

Chinese[6] (Supplementary Table 1).

113
114
For each pair of individuals, we used GERMLINE to detect segments of the genome

where the individuals are likely to share a common ancestor within the last few

dozen generations, that is, where they are IBD[13]. We used HaploScore to filter out

segments that were likely to be false-positives[14]. After normalizing for sample size,

we estimated the distribution of IBD genome-wide within each group with at least

two samples (Figure 2, Online Data Table 1). We found systematic differences in the

inferred IBD on different platforms; however, by normalizing by the average IBD in

each group by that detected in European Americans (CEU) (present in all three

datasets), we were able to meaningfully compare groups across platforms

(Supplementary Figure 5). We confirmed the accuracy of this method for detecting

founder events by using two other methods that we found gave highly correlated

results (correlation r=0.86-0.98): first, we computed F_{ST} between each group and

other groups with similar ancestry sources, and second, we fit a formal model of

history using *qpGraph*[6] and measured the founder event as the population-specific

genetic drift post-admixture (Supplementary Figure 6)(Online Data Table 1). These

analyses suggest that over a third of the Indian groups we analyzed (111 in total)

have stronger founder effects than those that occurred in both Finns and Ashkenazi

Jews (Figure 3). These groups are geographically and anthropologically varied,

include diverse tribe, caste, and different religious groups, and also include eighteen

groups with census sizes of over a million (Figure 3; Table 1). However, the groups

with smaller census sizes are also medically important, as the per-individual rate of

recessive disease is expected to be higher in proportion to their IBD score. Study of

groups with small population size such as Amish, Hutterites, and the people of the

Saguenay Lac-St. Jean region have proven to be powerful, leading to the discovery of

dozens of novel disease variants that are specific to each group.

140
141
To better understand the history of the groups in which we detected founder events,

we computed IBD for all pairs of individuals across groups. After applying a cutoff

on the IBD score corresponding to ~1/3 of the founder event size of Ashkenazi Jews,

most of the groups either have no matches or only match other individuals in their

group. However, some groups also share IBD across groups, typically following a

religious affiliation (e.g. Catholic Brahmins) or a distinctive linguistic affiliation

(particularly Austroasiatic speakers) (Supplementary Table 3). These results point

to recent gene flows among some of these pairs of groups.

The strong founder events offer major opportunities for improving health in South Asia. The first opportunity lies in targeted discovery of new genetic risk factors for disease. It is already known that one group we identified as having a strong founder event, the Vysya, has over a 100-fold higher rate of butyrylcholinesterase deficiency than other Indian groups, so that in India, Vysya ancestry is a known counter-indication for the use of common muscle relaxants such as succinylcholine or mivacurium that are given prior to anasthesia[15]. The Agarwal community in North India, though not present in our study, is also known to have founder mutations causing higher rates of Hereditary Fructose Intolerance[16] and Megalencephalic Leukoencephalopathy[17]. Systematic studies in the Vysya and other founder event groups—involving collaboration with clinical geneticists, local pediatricians, obstetricians, midwives, and social workers to identify congenital syndromes that are common in these communities—would discover many more examples. Identification of pathogenic mutations responsible for such syndromes is straightforward with present technology. All that is required is collection of DNA samples from a small number of affected individuals and their families, usually followed by whole-exome sequencing to discover the causal changes. While rare recessive diseases would be a prime target for gene mapping, the founder groups we have identified may also be of substantial importance for disease gene mapping studies of common disease, as rare variant association analyses are known to have enhanced power in such groups[18,19].

Once group specific founder event disease mutations are discovered, they can be tested for prenatally, and indeed, much of the improvement in human health that has come from founder event disease gene mapping studies is due to prenatal testing. Another way that discovery of rare recessive disease genes is likely to be important, especially in India, is through pre-marriage counseling in traditional communities where arranged marriages are common. An example of the power of this approach is *Dor Yeshorim*, a community genetic testing program among orthodox Ashkenazi Jews in both the United States and Israel[20]. Matchmaking is the norm among the hundreds of thousands of traditional religious orthodox Ashkenazi

Jews in both the United States and Israel, and *Dor Yeshorim* has taken the approach

of visiting schools, genetically screening students for common recessive disease

causing mutations known to affect Ashkenazi Jews using an inexpensive test, and

entering the results into a confidential database. Match-makers query the *Dor*

Yeshorim database prior to making their suggestions to the families and receive

feedback about whether the potential couple is "incompatible" in the sense of both

being carriers for a recessive mutation at the same gene. The program is successful

in both the United States and Israel, such that ~95% of community members whose

marriages are arranged participate; as a result, recessive diseases like Tay Sachs

have virtually disappeared from these communities. A similar approach should

work as well in Indian communities where arranged marriages are common, and

where there is already recognition of the power of clinical screening to affect birth

outcomes. Given the potential for saving lives and ultimately financial and medical

resources, this or similar kinds of research could serve as an important investment

for future generations[21].

This study of more than 250 distinct groups represents the first systematic survey

for founder events in South Asia, and to our knowledge also presents the richest

dataset of genome-wide data from anthropologically well-documented groups

available for any region in the world. Despite the breadth of this data, the groups

surveyed here represent only about 5% of the anthropologically well-defined

groups in India. Extensions of the survey to all well-defined anthropological groups

would make it possible to identify large numbers of additional founder groups

susceptible to recessive diseases and to assess the extent to which the founder

events we have already detected are localized to the specific regions from which our

samples were drawn, or are shared across people of the same ethnic group across

different regions in India. An important priority for future work is also to carry out

pilot studies to find real disease genes in our groups, thereby proving by example

the power of this approach for directing future disease mapping studies.

Supplementary Data:

Supplementary Data include an excel spreadsheet detailing all groups and their scores on the IBD, Fst, and Population-specific drift analyses. Also included are 6 supplementary figures and 3 supplementary tables.

Acknowledgements:

We are grateful to the many Indian, Pakistani, Bangladeshi, and Nepalese communities and individuals who contributed the DNA samples analyzed here. We thank Raj Rajkumar (deceased) for his assistance in assembling the sample collection. We would also like to acknowledge Analabha Basu and Patha Majumdar for helping to share their data for analysis in this study. Funding for this project was provided by the NIGMS (T32GM007753) to NN. This work was supported by a Translational Seed Fund grant from the Dean's Office of Harvard Medical School to DR, who is also a member of the Howard Hughes Medical Institute. KT is supported by CSIR network project GENESIS (BSC0121). PM was supported by the National Institutes of Health (NIH) under a Ruth L. Kirschstein National Research Service Award F32 GM115006-01. Genotyping data for the samples collected for this study will be made available upon request from the corresponding authors.

References:

1. Mastana, S.S. Unity in diversity: an overview of the genomic anthropology of India. *Ann Hum Biol* **41**, 287-99 (2014).

2. Bamshad, M.J. *et al.* Female gene flow stratifies Hindu castes. *Nature* **395**, 651-2 (1998).

3. Basu, A. *et al.* Ethnic India: a genomic view, with special reference to peopling and structure. *Genome Res* **13**, 2277-90 (2003).

4. Reich, D., Thangaraj, K., Patterson, N., Price, A.L. & Singh, L. Reconstructing Indian population history. *Nature* **461**, 489-94 (2009).

5. Lim, E.T. *et al.* Distribution and medical impact of loss-of-function variants in the Finnish founder population. *PLoS Genet* **10**, e1004494 (2014).

6. Patterson, N. *et al.* Ancient admixture in human history. *Genetics* **192**, 1065-93 (2012).

7. Basu, A., Sarkar-Roy, N. & Majumder, P.P. Genomic reconstruction of the history of extant populations of India reveals five distinct ancestral components and a complex structure. *Proc Natl Acad Sci U S A* (2016).

8. Moorjani, P. *et al.* Genetic evidence for recent population mixture in India. *Am J Hum Genet* **93**, 422-38 (2013).

9. Metspalu, M. *et al.* Shared and unique components of human population structure and genome-wide signals of positive selection in South Asia. *Am J Hum Genet* **89**, 731-44 (2011).

10. Behar, D.M. *et al.* The genome-wide structure of the Jewish people. *Nature* **466**, 238-42 (2010).

11. Narang, A. *et al.* Recent admixture in an Indian population of African ancestry. *Am J Hum Genet* **89**, 111-20 (2011).

12. Shah, A.M. *et al.* Indian Siddis: African descendants with Indian admixture. *Am J Hum Genet* **89**, 154-61 (2011).

13. Gusev, A. *et al.* Whole population, genome-wide mapping of hidden relatedness. *Genome Res* **19**, 318-26 (2009).

262   14.   Durand, E.Y., Eriksson, N. & McLean, C.Y. Reducing pervasive false-positive
263   identical-by-descent segments detected by large-scale pedigree analysis. *Mol*
264   *Biol Evol* **31**, 2212-22 (2014).

265   15.   Manoharan, I., Wieseler, S., Layer, P.G., Lockridge, O. & Boopathy, R. Naturally
266   occurring mutation Leu307Pro of human butyrylcholinesterase in the Vysya
267   community of India. *Pharmacogenet Genomics* **16**, 461-8 (2006).

268   16.   Bijarnia-Mahay, S. *et al.* Molecular Diagnosis of Hereditary Fructose
269   Intolerance: Founder Mutation in a Community from India. *JIMD Rep* **19**, 85-
270   93 (2015).

271   17.   Shukla, P. *et al.* Molecular genetic studies in Indian patients with
272   megalencephalic leukoencephalopathy. *Pediatr Neurol* **44**, 450-8 (2011).

273   18.   Wang, S.R. *et al.* Simulation of Finnish population history, guided by empirical
274   genetic data, to assess power of rare-variant tests in Finland. *Am J Hum Genet*
275   **94**, 710-20 (2014).

276   19.   Sabatti, C. *et al.* Genome-wide association analysis of metabolic traits in a
277   birth cohort from a founder population. *Nat Genet* **41**, 35-46 (2009).

278   20.   Raz, A.E. Can population-based carrier screening be left to the community? *J*
279   *Genet Couns* **18**, 114-8 (2009).

280   21.   Rajasimha, H.K. *et al.* Organization for rare diseases India (ORDI) -
281   addressing the challenges and opportunities for the Indian rare diseases'
282   community. *Genet Res (Camb)* **96**, e009 (2014).

283   22.   Sudmant, P.H. *et al.* Global diversity, population stratification, and selection
284   of human copy-number variation. *Science* **349**, aab3761 (2015).

285   23.   Sudmant, P.H. *et al.* An integrated map of structural variation in 2,504 human
286   genomes. *Nature* **526**, 75-81 (2015).

287   24.   Patterson, N., Price, A.L. & Reich, D. Population structure and eigenanalysis.
288   *PLoS Genet* **2**, e190 (2006).

289   25.   Felsenstein, J. PHYLIP - Phylogeny Inference Package (Version 3.2). *Cladistics*
290   **5**, 164-166 (1989).

26.   Letunic, I. & Bork, P. Interactive Tree Of Life v2: online annotation and
display of phylogenetic trees made easy. *Nucleic Acids Res* **39**, W475-8
(2011).

27.   Browning, B.L. & Browning, S.R. Improving the accuracy and efficiency of
identity-by-descent detection in population data. *Genetics* **194**, 459-71
(2013).

28.   Browning, S.R. & Browning, B.L. Rapid and accurate haplotype phasing and
missing-data inference for whole-genome association studies by use of
localized haplotype clustering. *Am J Hum Genet* **81**, 1084-97 (2007).

Population	Sample Size	IBD Rank	F_{ST} Rank	Drift Rank	Census Size	Location
Pattapu_Kapu	4	15	17	12	13,697,000	Andhra Pradesh
Kumhar	5	20	28	18	3,144,000	Uttar Pradesh
Rajbanshi	4	29	NA	NA	3,801,677	West Bengal
Kallar	5	38	37	29	2,426,929	Tamil Nadu
Arunthathiyar	5	40	48	34	1,084,162	Tamil Nadu
Yadav	13	41	66	44	20,000,000	Puducherry
Naga	4	51	NA	NA	1,667,712	Nagaland
Scheduled_Caste_Haryana	4	53	66	47	5,113,615	Haryana
Reddy	8	56	70	53	22,500,000	Andhra Pradesh
Baniyas	4	60	81	56	4,848,000	Uttar Pradesh
Dawoodi	2	62	41	40	4,592,854	Gujarat
Muslim_Bihar	4	64	62	44	13,722,048	Bihar
Vysya	10	73	30	21	3,200,000	Andhra Pradesh
Rajput	4	75	139	136	51,137,113	Haryana
Garasia	5	77	70	43	4,215,603	Gujarat
Satnami	5	85	96	65	4,200,000	Chhattisgarh
Shani	3	87	139	80	4,029,411	Bihar
Newar	6	95	NA	NA	1,523,000	Nepal

310 **Table 1. Indian groups with strong IBD scores.** Eighteen Indian groups with IBD scores higher than Ashkenazi
311 Jews and census sizes over 1 million that are of particularly high interest for founder event disease gene mapping
312 studies.

314
315

317 **Figure 1. Dataset overview.** (a) Sampling locations for all analyzed groups. Each
318 point indicates a distinct group (random jitter added to help in visualization at
319 locations where there are many groups). (b) PCA of Human Origins dataset along
320 with European Americans (CEU) and Han Chinese (CHB).
321

322   A)   B)

323
324   C)   D)

325
326   **Figure 2.** Histogram of IBD in groups with founder events of different magnitudes:
327   (A) very large in Ulladan, (B) large in Birhor, (C) moderate in Ashkenazi Jews, and
328   (D) small in Mahadeo_Koli. In each plot, we showed for comparison the histogram of
329   IBD for European Americans (CEU) with a negligible founder event in blue, and that
330   in Finns (FIN) with a large founder event in black.
331

Figure 3. IBD scores normalized by that in European Americans (CEU).
Histogram ordered by IBD score, which is roughly proportional to the per-individual
risk for recessive disease due to the founder event. (These results are also given
quantitatively for each group in Online Table 1.) We restrict to groups with at least
two samples, combining data from all four genotyping platforms onto one plot. Data
from Ashkenazi Jews and Finns are highlighted in red, and from Indian groups with
stronger founder events and census sizes of more than a million in orange.

Online Methods:

Data Sets:

We used genotyping array data from multiple sources. We assembled a dataset of 1,182 individuals from 225 groups genotyped on the Affymetrix Human Origins array, of which data from 890 individuals from 206 groups is newly reported here (Figure 1a). We merged these data with a dataset published in Moorjani et al.[8], which consisted of 332 individuals from 52 groups genotyped on the Affymetrix 6.0 array. We also merged it with two additional datasets published in Metspalu et al.[9], consisting of 151 individuals from 21 groups genotyped on Illumina 650K arrays as well as a dataset published in Basu et al.[7], which consisted of 367 individuals from 20 groups generated on Illumina Omni 1-Quad arrays. These groups came from India, Pakistan, Nepal, Sri Lanka, and Bangladesh, and we refer to all of them here as the "South Asia" dataset.

We analyzed two different Jewish datasets, one consisting of 21 Ashkenazi Jewish individuals genotyped on Illumina 610K and 660K bead arrays[10] and one consisting of 7 Ashkenazi Jewish individuals genotyped on Affymetrix Human Origins arrays.

Our "Affymetrix 6.0" dataset consisted of 332 individuals genotyped on 329,261 SNPs, and our "Illumina_Omni" dataset consisted of 367 individuals genotyped on 750,919 SNPs. We merged the South Asia and Jewish data generated by the other Illumina arrays to create an "Illumina" dataset consisting of 172 individuals genotyped on 500,640 SNPs. Finally, we merged the data from the Affymetrix Human Origins arrays with the Ashkenazi Jewish data and data from the Simons Genome Diversity Project[22] to create a dataset with 1,225 individuals genotyped on 512,615 SNPs. We analyzed the four datasets separately due to the small intersection of SNPs between them and the possible systematic differences across genotyping platforms. We merged in the 1000 Genomes Phase 3 data[23] (2504 individuals from 26 different groups; notably, including 99 Finnish individuals) into

all of the datasets. We used genome reference sequence coordinates (hg19) for

analyses.

Quality Control:

We filtered the data on both the SNP and individual level. On the SNP level, we

required at least 95% genotyping completeness for each SNP (across all

individuals). On the individual level, we required at least 95% genotyping

completeness for each individual (across all SNPs).

To test for batch effects due to samples from the same group being genotyped on

different array plates, we studied instances where samples from the same group A

were genotyped on both plates 1 and 2 and computed an allele frequency difference

at each SNP, $Diff_A^i = \left(Freq_{PopA,Plate1}^i - Freq_{PopA,Plate2}^i\right)$. We then computed the

product of these allele frequencies averaged over all SNPs for two groups A and B

genotyped on the same plates, $\frac{1}{n}\sum_{i=1}^{n}\left(Diff_A^i\right)\left(Diff_B^i\right)$, as well as a standard error

from a Block Jackknife. This quantity should be consistent with zero within a few

standard errors if there are no batch effects that cause systematic differences across

the plates, as allele frequency differences between two samples of the same group

should just be random fluctuations that have nothing to do with the array plates on

which they are genotyped. This analysis found strong batch effects associated with

one array plate, and we removed this from analysis.

We used EIGENSOFT 5.0.1 smartpca[24] on each group. We also developed a

procedure to distinguish recent relatedness from founder effects so that we could

remove recently related individuals. We first identified all duplicates or obvious

close relatives by using Plink "genome" and removed all individuals who had both a

PI_HAT score greater than 0.45 and the presence of at least 1 IBD fragment greater

than 30cM long. We used an iterative procedure of identifying any pairs within each

group that had both total IBD and total long IBD (>20cM) that were greater than 2.5

SDs and 1 SD, respectively, from the group mean. After each round we repeated the

process if the new IBD score was at least 30% lower than the prior IBD score. Due to

their known very small census size (our sample consists of a substantial fraction of

the entire population) and exceptional anthropological interest, we excluded Onge, a

tribal population of the Andaman and Nicobar Islands, from this analysis. After data

quality control and merging with the 1000 Genomes Project data, the Affymetrix 6.0

dataset included 3,215 individuals genotyped on 326,181 SNPs, the Illumina dataset

included 2,789 individuals genotyped on 484,293 SNPs, the Illumina Omni dataset

included 2,834 individuals genotyped on 750,919 SNPs, and the Human Origins

dataset included 3,696 individuals genotyped at 500,648 SNPs.

Distance-Based Phylogenetic Tree:
413
414 We calculated genetic differentiation (F_{ST}) between all pairs of groups using

415 EIGENSOFT *smartpca* and created a neighbor-joining tree using PHYLIP[25] with

416 Yoruba as the outgroup. We used Itol[26] to display the tree.

418 **Power Calculations:**
419
We performed power calculations to determine the approximate number of samples

required to detect founder events of a pre-specified strength. We used Beagle 3.3.2

FastIBD[27] to identify all shared IBD segments between individuals within a group

with parameters *missing=0; lowmem=true; gprobs=false; verbose=true; fastIBD=true;*

ibdscale=scale (where scale = sqrt(#samples/100)). We used the output of this

program to calculate mean IBD sharing. We computed standard errors via jackknife

resampling over individuals for each group. These analyses demonstrate that only

3-5 individuals are needed to assess accurately the size of founder effects in groups

with strong founder events (Supplementary Figure 1). Weaker founder effects are

more difficult to detect, but these groups are of less interest from the perspective of

founder event disease mapping, so we aimed to sample ~5 individuals per group in

the new genotyping effort based on the Affymetrix Human Origins platform.

Phasing and IBD Detection:
434

We phased all datasets using Beagle 3.3.2 with the settings *missing=0; lowmem=true;*

gprobs=false; verbose=true[28]. We left all other settings at default. We determined IBD

segments using the GERMLINE algorithm[13] with the parameters *-bits 75 -err_hom 0 -*

err_het 0 -min_m 3. We used the genotype extension mode to minimize the effect of

any possible phasing heterogeneity amongst the different groups and used the

HaploScore algorithm to remove false positive IBD fragments with the

recommended genotype error and switch error parameters of 0.0075 and 0.003[14].

We chose a HaploScore threshold matrix based on calculations from Durand *et al.*

for a "mean overlap" of 0.8, which corresponds to a precision of approximately 0.9

for all genetic lengths from 2-10cM. In addition to the procedure we developed to

remove close relatives (Quality Control section), we also removed segments longer

than 20cM to ignore the effect of consanguinity and shorter than 3cM to minimize

false positives and better differentiate groups with stronger founder effects from

those with weaker effects. We treated all groups as subpopulations (e.g. Vysya and

Ashkenazi Jews) and only retained IBD segments within the subpopulation (e.g. only

IBD segments shared between Vysya individuals). We computed "founder effect

size" as the total length of IBD segments between 3-20cM divided by ((2*(# of

individuals in sample)) choose 2) to normalize for sample size. We also repeated

these analyses with FastIBD[27] for the Affy 6.0 and Illumina datasets and observed

that the results were highly correlated (r>0.96) (Supplementary Table 1). We chose

GERMLINE for our main analyses, however, because the FastIBD algorithm required

us to split the datasets into different groups, since it adapts to the relationships

between LD and genetic distance in the data, and these relationships differ across

groups. We used several different Jewish groups and all twenty-six 1000 Genomes

groups to improve phasing and calibration of the IBD scores, but of these groups we

only included results for two founder groups (Ashkenazi Jews and Finns for

comparison with Indian groups), and two outbred populations (CEU and YRI for

normalization) in the final IBD score ranking.

Between-Group IBD Calculations:
465

We determined IBD using GERMLINE as above. We collapsed individuals into

respective groups and normalized for between-group IBD by dividing all IBD from

each group by (2*# of individuals in the group). We normalized for within-group

IBD as described above. We defined groups with high shared IBD as those with an

IBD score greater than 3, corresponding to approximately three times the founder

effect size of CEU (and ~1/3 the effect size of Ashkenazi Jews).

f_3-statistics:
474
We used the f_3-statistic[6] $f_3(Test; Ref_1, Ref_2)$ to determine if there was evidence that

the *Test* group was derived from admixture of groups related to Ref_1 and Ref_2. A

significantly negative statistic provides unambiguous evidence of mixture in the

Test group. We assessed the significance of the f_3-statistic using a Block Jackknife

and a block size of 5 cM. We considered statistics over 3 standard errors below zero

to be significant.

Calculating Group Specific Drift:
483
We used ADMIXTUREGRAPH[6] to model each Indian population on the cline as a

mixture of ANI and ASI ancestry. Within the limits of our resolution, this model (YRI,

(Indian population, (Georgians, ANI)), [(ASI, Onge]]) proposed by Moorjani *et al*.[8] is

a good fit to the data, in the sense that none of the *f*-statistics relating the groups are

greater than three standard errors from expectation. This approach provides

estimates for post-admixture drift in each group (Supplementary Figure 6), which is

reflective of the strength of the founder event (high drift values imply stronger

founder events). We only included groups on the Indian cline in this analysis, and

we removed all groups with evidence of recent East Asian admixture.

PCA-Normalized F_{ST} Calculations:
495
To account for intermarriage across groups, we used clusters based on PCA to

estimate the minimum F_{ST} for each South Asian population (Supplementary Figure

6). Specifically, we calculated the F_{ST} between each group and the rest of the

individuals in their respective cluster based on EIGENSOFT *smartpca*. For these

analyses we only included groups with Austroasiatic-related genetic patterns (i.e.

those groups clustering near Austroasiatic speakers on the PCA) and those on the

Indian cline; we excluded all groups with recent East Asian admixture. For

Ashkenazi Jews and Finns, we used the minimum F_{ST} to their closest European

neighbors.

507