The promise of disease gene discovery in South Asia

Nathan Nakatsuka[1,2], Priya Moorjani[3,6], Niraj Rai[4], Biswanath Sarkar[5], Arti Tandon[1,6], Nick Patterson[6], Gandham SriLakshmi Bhavani[7], Katta Mohan Girisha[7], Mohammed S Mustak[8], Sudha Srinivasan[9], Amit Kaushik[10], Saadi Abdul Vahab[11], Sujatha M. Jagadeesh[12], Kapaettu Satyamoorthy[11], Lalji Singh[4,13], David Reich[1,5,14,*], Kumarasamy Thangaraj[4,*]

[1]Department of Genetics, Harvard Medical School, New Research Building, 77 Ave. Louis Pasteur, Boston, MA 02115, USA
[2]Harvard-MIT Division of Health Sciences and Technology, Harvard Medical School, Boston, MA 02115, USA
[3]Department of Biological Sciences, Columbia University, 600 Fairchild Center, New York, NY 10027, USA
[4]CSIR-Centre for Cellular and Molecular Biology, Habsiguda, Hyderabad, Telangana 500007, India
[5]Superintending Anthropologist (Physical) (Rtd.), Anthropological Survey of India, 27 Jawaharlal Nehru Road, Kolkata 700016, India
[6]Broad Institute of Harvard and Massachusetts Institute of Technology, Cambridge, MA 02141, USA
[7]Department of Medical Genetics, Kasturba Medical College, Manipal University, Manipal, India
[8]Department of Applied Zoology, Mangalore University, Mangalagangothri 574199, Mangalore, Karnataka, India
[9]Centre for Human Genetics, Biotech Park, Electronics City (Phase I), Bangalore 560100, India
[10]Amity Institute of Biotechnology, Amity University, Sector125, Noida 201303, India
[11]School of Life Sciences, Manipal University, Manipal 576104, India
[12]Fetal Care Research Foundation, 197 Dr. Natesan Road, Chennai 600004, India
[13]Present address: Genome Foundation, Hyderabad 500076, India
[14]Howard Hughes Medical Institute, Harvard Medical School, Boston, MA 02115, USA
*co-senior authors

The more than 1.5 billion people who live in South Asia are correctly viewed not as a single large population, but as many small endogamous groups. We assembled genome-wide data from over 2,800 individuals from over 260 distinct South Asian groups. We identify 81 unique groups, of which 14 have estimated census sizes of more than a million, that descend from founder events more extreme than those in Ashkenazi Jews and Finns, both of which have high rates of recessive disease due to founder events. We identify multiple examples of recessive diseases in South Asia that are the result of such founder events. This study highlights an under-appreciated opportunity for reducing disease burden among South Asians through the discovery of and testing for recessive disease genes.

South Asia is a region of extraordinary diversity, containing over 4,600 anthropologically well-defined groups many of which are endogamous communities with significant barriers to gene flow due to cultural practices that restrict marriage between groups[1]. Of the tiny fraction of South Asian groups that have been characterized using genome-wide data, many exhibit large allele frequency differences from close neighbors[2-4], consistent with strong founder events whereby a small number of ancestors gave rise to many descendants today[4]. The pervasive founder events in South Asia present a potential opportunity for reducing disease burden in South Asia. The promise is highlighted by studies of founder groups of European ancestry – including Ashkenazi Jews, Finns, Amish, Hutterites, Sardinians, and French Canadians – which have resulted in the discovery of dozens of recessive disease causing mutations in each group. Prenatal testing for these mutations has substantially reduced recessive disease burden in all of these communities[5,6].

We carried out new genotyping of 1,663 samples from 230 endogamous groups in South Asia on the Affymetrix Human Origins single nucleotide polymorphism (SNP) array[7]. We combined the data we newly collected with previously reported data, leading to four datasets (Figure 1a). The Affymetrix Human Origins SNP array data comprised 1,955 individuals from 249 groups in South Asia, to which we added 7 Ashkenazi Jews. The Affymetrix 6.0 SNP array data comprised 383 individuals from 52 groups in South Asia[4,8]. The Illumina SNP array data comprised 188 individuals from 21 groups in South Asia[9] and 21 Ashkenazi Jews[9,10]. The Illumina Omni SNP array data comprised 367 individuals from 20 groups in South Asia[11]. We merged 1000 Genomes Phase 3 data[12] (2,504 individuals from 26 different groups including 99 Finns) with each of these datasets. We removed SNPs and individuals with a high proportion of missing genotypes or that were outliers in Principal Components Analysis (PCA) (Figure 1b; Supplementary Text). The total number of unique groups analyzed in this study is 263 after accounting for groups represented in multiple datasets. To our knowledge, this represents the richest set of genome-wide data from anthropologically well-documented groups from any region in the world.

We devised an algorithm to quantify the strength of the founder events in each group based on Identity-by-Descent (IBD) segments, large stretches of DNA shared from a common founder in the last approximately one hundred generations (Figure

2). We computed an "IBD score" as a measure for the strength of the founder event
in each group's history: the average length of IBD segments between 3-20
centimorgans (cM) shared between two genomes normalized to sample size. Since
we are interested in characterizing the impact of recessive diseases that do not owe
their origin to consanguineous marriages of close relatives, we ignored self-matches
(internal homozygosity) in IBD calculations. We removed all individuals that had
evidence of recent relatedness (within a few generations) to others in the dataset by
computing IBD between all pairs of individuals in each group and removing one
individual from the pairs with outlying numbers of IBD segments (our focus on
founder events rather than recent relatedness also explains our choice to exclude
IBD segments of greater than 20 cM in size). We validated the effectiveness of this
procedure by simulation (Supplementary Table 1; Online Methods).
101
We expressed IBD scores for each group as a fraction of the IBD scores of the 1000
Genomes Project Finns merged into each respective dataset. Due to the fact that all
the SNP arrays we analyzed included more SNPs ascertained in Europeans than in
South Asians, the sensitivity of our methods to founder events is greater in
Europeans than in South Asians, our estimates of founder event strengths in South
Asian groups are conservative underestimates relative to that in Europeans
(Supplementary Figure 1 demonstrates this effect empirically and shows that it is
less of a bias for the strong founder events that are the focus of this study). We
computed standard errors for these ratios by a weighted Block Jackknife across
chromosomes and declared significance where the 95% confidence intervals did not
overlap 1. We carried out computer simulations to validate our procedure. The
simulations suggest that we are not substantially overestimating the magnitudes of
modest founder events, since for a simulated founder event that is half the
magnitude of that in Finns, we never infer the score to be significantly greater than
in Finns. The simulations also suggest that our procedure is highly sensitive to
detecting strong founder events, since for sample sizes of at least 5, the algorithm's
sensitivity is greater than 95% for determining that a group with two times the
bottleneck strength as Finns has an IBD score significantly greater than that of Finns
(Supplementary Figure 2 and Supplementary Table 2).
121
We infer that 81 out of 263 unique groups (96 out of 327 groups if not considering
the overlap of groups among datasets) have an IBD score greater than those of both
Finns and Ashkenazi Jews (Figure 3). These results did not change when we added
back the outlier samples that we removed in quality control. A total of 14 of these
groups have estimated census sizes of over a million (Figure 3; Table 1). However,
the groups with smaller census sizes are also very important – outside of South Asia,
small census size groups with extremely strong founder events such as Amish,
Hutterites, and people of the Saguenay Lac-St. Jean region have led to the discovery
of dozens of novel disease variants. We also searched for IBD across groups –
screening for cases in which the across-group IBD score is at least a third of the
within-group IBD score of Ashkenazi Jews – and found 77 cases of clear IBD-sharing,
which typically follow geography, religious affiliation, or linguistic grouping
(particularly Austroasiatic speakers) (Supplementary Table 3). Pairs of groups with

high shared IBD and descent from a common founder event will share risk for the
same recessive disease. However, these cross-group IBD sharing patterns are not
driving our observations, as we still identify 68 unique sets of groups without high
IBD to other groups that have significantly higher estimated IBD scores than both
Finns and Ashkenazi Jews.

Our documentation that very strong founder events affect a large fraction of South
Asian groups presents an opportunity for decreasing disease burden in South Asia.
This source of risk for recessive diseases is very different from that due to marriages
among close relatives, which is also a major cause of recessive disease in South Asia.
To determine the relative impact of these factors, we computed F_{ST}, a measurement
of allele frequency differentiation, between each group in the dataset and a pool of
other South Asian groups chosen to be closest in terms of ancestry proportions. We
find that inbreeding is not driving many of these signals, as 89 unique groups have
higher F_{ST} scores than those of Ashkenazi Jews and Finns even after reducing the F_{ST}
score by the proportion of allele frequency differentiation due to inbreeding. These
results show that while most recessive disease gene mapping studies in South Asia
have focused on families that are the products of marriages between close relatives,
recessive diseases are also likely to occur at an elevated rate even in non-
consanguineous cases because of shared ancestors more distantly in time.

As an example of the promise of founder event disease gene mapping in South Asia,
we highlight the case of the Vysya, who have a census size of more than 3 million
and who we estimate have an IBD score about 1.2-fold higher than Finns (Figure 3).
The Vysya have an approximately 100-fold higher rate of butyrylcholinesterase
deficiency than other groups, and Vysya ancestry is a known counter-indication for
the use of muscle relaxants such as succinylcholine or mivacurium that are given
prior to surgery[13]. This disease is likely to occur at a higher rate due to the founder
event in Vysya's history, and we expect that, like Finns, Vysya likely have a higher
rate of many other diseases compared to other groups. Other examples of recessive
disease genes with a likely origin in founder events are known anecdotally in South
Asia, highlighting the importance of systematic studies to find them[14].

To demonstrate how a new recessive disease in a founder event group could be
mapped, we carried out whole genome SNP genotyping in 12 patients from southern
India who had progressive pseudorheumatoid dysplasia (PPD), a disease known to
be caused by mutations in the gene *WISP3*[15,16]. Of the 6 individuals with the
Cys78Tyr mutation in *WISP3*,[15,16] 5 were from non-consanguineous marriages, and
we found a much higher fraction of IBD at the disease mutation site than in the rest
of the genome in these individuals (Supplementary Figure 3a; Supplementary Figure
4a), consistent with the Cys78Tyr mutation that causes PPD in these patients owes
its origin to a founder event. This pattern contrasts with the 6 other patients with
different disease variants and 6 patients with a mutation causing a different disease
(mucopolysaccharidosis (MPS) type IVA) who were from primarily consanguineous
marriages, and who lacked significant IBD across their disease mutation sites,
implying that in the case of these groups the driver for the recessive diseases was

181   marriage between close relatives (Supplementary Text). This example highlights
182   how not only marriages of close relatives, but also founder events, are substantial
183   causes of rare recessive disease in South Asia.
184
The evidence of widespread strong founder events presents a major opportunity for
disease gene discovery and public health intervention in South Asia that is not
widely appreciated. The current paradigm for recessive disease gene mapping in
South Asia is to collect cases in tertiary medical centers and map diseases in
individuals with the same phenotype, often blinded to information about group
affiliation as in the case of the PPD study where we do not have access to the ethnic
group information. However, our results suggest that collecting information on
group affiliation could be highly valuable for improving the success of these studies.
A fruitful way to approach gene mapping would be to proactively survey
communities known to have strong founder events, searching for diseases that
occur at high rates in these communities. This approach was pioneered in the 1950s
in studies of the Old Order Amish in the U.S., a founder population of approximately
100,000 individuals in whom many dozens of recessive diseases were mapped, a
research program that was crucial to founding modern medical genetics and that
was of extraordinary health benefit. Our study suggests that the potential for
disease gene mapping in South Asia would be orders of magnitude greater.
201
Mapping of recessive diseases may be particularly important in communities
practicing arranged marriages, which are common in South Asia. An example of the
power of this approach is given by *Dor Yeshorim*, a community genetic testing
program among religious Ashkenazi Jews[17], which visits schools, screens students
for common recessive disease causing mutations previously identified to be
segregating at a higher frequency in the target group, and enters the results into a
confidential database. Matchmakers query the database prior to making suggestions
to the families and receive feedback about whether the potential couple is
"incompatible" in the sense of both being carriers for a recessive mutation at the
same gene. Given that approximately 95% of community members whose marriages
are arranged participate in this program, recessive diseases like Tay-Sachs have
virtually disappeared in these communities. A similar approach should work as well
in South Asians communities. Given the potential for saving lives, this or similar
kinds of research could be a valuable investment for future generations[18].
216

Supplementary Data:

Supplementary Data include an Excel spreadsheet detailing all groups and their scores on the IBD, F_{ST}, and group-specific drift analyses. Also included are 7 supplementary figures and 5 supplementary tables.

Acknowledgements:

We are thankful to the many Indian, Pakistani, Bangladeshi, Sri Lankan, and Nepalese individuals who contributed the DNA samples analyzed here including the PPD and MPS patients. We are grateful to Analabha Basu and Partha P. Majumder for early sharing of data. Funding was provided by an NIGMS (GM007753) fellowship to NN, a Translational Seed Fund grant from the Dean's Office of Harvard Medical School to DR, Council of Scientific and Industrial Research, Government of India grant to KT, and an NIGMS grant 115006 to OPM. SS and SMJ acknowledge the funding from the Department of Biotechnology (BT/PR4224/MED/97/60/2011) and Department of Science and Technology (SR/WOS-A/LS-83/2011), Government of India. Funding for the mutation analysis of Indian patients with PPD was provided by Indian Council of Medical Research (BMS 54/2/2013) to KMG. DR is an Investigator of the Howard Hughes Medical Institute. Genotyping data for the samples collected for this study is available on request.

References:

1. Mastana, S.S. Unity in diversity: an overview of the genomic anthropology of India. *Ann Hum Biol* **41**, 287-99 (2014).

2. Bamshad, M.J. *et al.* Female gene flow stratifies Hindu castes. *Nature* **395**, 651-2 (1998).

3. Basu, A. *et al.* Ethnic India: a genomic view, with special reference to peopling and structure. *Genome Res* **13**, 2277-90 (2003).

4. Reich, D., Thangaraj, K., Patterson, N., Price, A.L. & Singh, L. Reconstructing Indian population history. *Nature* **461**, 489-94 (2009).

5. Lim, E.T. *et al.* Distribution and medical impact of loss-of-function variants in the Finnish founder population. *PLoS Genet* **10**, e1004494 (2014).

6. Arcos-Burgos, M. & Muenke, M. Genetics of population isolates. *Clin Genet* **61**, 233-47 (2002).

7. Patterson, N. *et al.* Ancient admixture in human history. *Genetics* **192**, 1065-93 (2012).

8. Moorjani, P. *et al.* Genetic evidence for recent population mixture in India. *Am J Hum Genet* **93**, 422-38 (2013).

9. Metspalu, M. *et al.* Shared and unique components of human population structure and genome-wide signals of positive selection in South Asia. *Am J Hum Genet* **89**, 731-44 (2011).

10. Behar, D.M. *et al.* The genome-wide structure of the Jewish people. *Nature* **466**, 238-42 (2010).

11. Basu, A., Sarkar-Roy, N. & Majumder, P.P. Genomic reconstruction of the history of extant populations of India reveals five distinct ancestral components and a complex structure. *Proc Natl Acad Sci U S A* (2016).

12. Genomes Project, C. *et al.* A global reference for human genetic variation. *Nature* **526**, 68-74 (2015).

13. Manoharan, I., Wieseler, S., Layer, P.G., Lockridge, O. & Boopathy, R. Naturally occurring mutation Leu307Pro of human butyrylcholinesterase in the Vysya community of India. *Pharmacogenet Genomics* **16**, 461-8 (2006).

14. Anju Shukla, M.H., Anshika Srivastava, Rajagopal Kadavigere, Priyanka Upadhyai, Anil Kanthi, Oliver Brandau, Stephanie Bielas, Katta Girisha. Homozygous c.259G>A variant in ISCA1 is associated with a new multiple mitochondrial dysfunctions syndrome. *bioRxiv* (2016).

15. Dalal, A. *et al.* Analysis of the WISP3 gene in Indian families with progressive pseudorheumatoid dysplasia. *Am J Med Genet A* **158A**, 2820-8 (2012).

16. Bhavani, G.S. *et al.* Novel and recurrent mutations in WISP3 and an atypical phenotype. *Am J Med Genet A* **167A**, 2481-4 (2015).

17. Raz, A.E. Can population-based carrier screening be left to the community? *J Genet Couns* **18**, 114-8 (2009).

18. Rajasimha, H.K. *et al.* Organization for rare diseases India (ORDI) - addressing the challenges and opportunities for the Indian rare diseases' community. *Genet Res (Camb)* **96**, e009 (2014).

19. Sudmant, P.H. *et al.* Global diversity, population stratification, and selection of human copy-number variation. *Science* **349**, aab3761 (2015).

284 20. Mallick, S. *et al.* The Simons Genome Diversity Project: 300 genomes from
285   142 diverse populations. *Nature* **538**, 201-206 (2016).
286 21. Sudmant, P.H. *et al.* An integrated map of structural variation in 2,504 human
287   genomes. *Nature* **526**, 75-81 (2015).
288 22. Patterson, N., Price, A.L. & Reich, D. Population structure and eigenanalysis.
289   *PLoS Genet* **2**, e190 (2006).
290 23. Chang, C.C. *et al.* Second-generation PLINK: rising to the challenge of larger
291   and richer datasets. *Gigascience* **4**, 7 (2015).
292 24. Gusev, A. *et al.* Whole population, genome-wide mapping of hidden
293   relatedness. *Genome Res* **19**, 318-26 (2009).
294 25. Hoaglin, B.I.a.D. *How to Detect and Handle Outliers*, (1993).
295 26. Palamara, P.F. ARGON: fast, whole-genome simulation of the discrete time
296   Wright-fisher process. *Bioinformatics* (2016).
297 27. Browning, S.R. & Browning, B.L. Rapid and accurate haplotype phasing and
298   missing-data inference for whole-genome association studies by use of
299   localized haplotype clustering. *Am J Hum Genet* **81**, 1084-97 (2007).
300 28. Durand, E.Y., Eriksson, N. & McLean, C.Y. Reducing pervasive false-positive
301   identical-by-descent segments detected by large-scale pedigree analysis. *Mol*
302   *Biol Evol* **31**, 2212-22 (2014).
303 29. Browning, B.L. & Browning, S.R. Improving the accuracy and efficiency of
304   identity-by-descent detection in population data. *Genetics* **194**, 459-71
305   (2013).
306 30. Bidchol, A.M. *et al.* GALNS mutations in Indian patients with
307   mucopolysaccharidosis IVA. *Am J Med Genet A* **164A**, 2793-801 (2014).
308

Group	Sample Size	IBD Score	IBD Rank	F_{ST} Rank	Drift Rank	Census Size	Location
Gujjar	5	11.6	19	33	46	1,078,719	Jammu and Kashmir
Baniyas	7	9.6	24	22	18	4,200,000	Uttar Pradesh
Pattapu_Kapu	4	9.5	25	24	21	13,697,000	Andhra Pradesh
Vadde	3	9.2	26	30	26	3,695,000	Andhra Pradesh
Yadav	12	4.4	48	87	67	1,124,864	Puducherry
Kshatriya_Aqnikula	4	2.4	75	109	NA	12,809,000	Andhra Pradesh
Naga	4	2.3	76	NA	NA	1,834,483	Nagaland
Kumhar	27	2.3	77	35	197	3,144,000	Uttar Pradesh
Reddy	7	2.0	84	129	106	22,500,000	Telangana
Brahmin_Nepal	4	1.9	86	63	141	4,206,235	Nepal
Kallar	27	1.7	94	87	73	2,426,929	Tamil Nadu
Brahmin_Manipuri	17	1.6	99	NA	NA	1,544,296	Manipur
Arunthathiyar	18	1.3	108	109	81	1,192,578	Tamil Nadu
Vysya	39	1.2	110	46	35	3,200,000	Telangana

Table 1. South Asian groups with estimated census sizes over 1 million and IBD scores significantly greater than
those of Ashkenazi Jews and Finns. Fourteen South Asian groups with IBD scores significantly higher than that of Finns,
census sizes over 1 million, and sample sizes of at least 3 that are of particularly high interest for founder event disease
gene mapping studies. For reference, Finns and Ashkenazi Jews (on the Human Originsarray) would have IBD scores of
1.0 and 0.9, IBD ranks of 121 and 135, and F_{ST} ranks of 109 and 129, respectively (the group-specific drift is difficult to
compare for groups with significantly different histories, so they were not calculated for Finns or Ashkenazi Jews).

317
318

Figure 1. Dataset overview. (a) Sampling locations for all analyzed groups. Each point indicates a distinct group (random jitter was added to help in visualization at locations where there are many groups). (b) PCA of Human Origins dataset along with European Americans (CEU) and Han Chinese (CHB). There is a large cluster (blue) of IndoEuropean and Dravidian speaking groups that stretch out along a line in the plot and that are well-modeled as a mixture of two highly divergent ancestral populations (the "Indian Cline"). There is another larger cluster of Austroasiatic speakers (light red) and groups that cluster with them genetically (dark red). Finally, there are groups with genetic affinity to East Asians that include Tibeto-Burman speakers (orange) and those that speak other languages (yellow).

331
332
333

Figure 2. Example histograms of IBD segments to illustrate the differences
between groups with founder events of different magnitudes: These histograms
provide visual illustrations of differences between groups with different IBD scores.
As a ratio relative to Finns (FIN; black), these groups (red) have IBD scores of: (A)
~26 in Ulladan, (B) ~3 in Birhor, (C) ~0.9 in Ashkenazi Jews, and (D) ~0.1 in
Mahadeo_Koli. In each plot, we also show European Americans (CEU) with a
negligible founder event in blue. Quantification of these founder events is shown in
Figure 3 and Online Table 1. The IBD histograms were normalized for sample size
by dividing their frequency by $\left\{\binom{2n}{2} - n\right\}$, where n is the number of individuals in
the sample. All data for the figure are based on the Human Origins dataset.

344
Figure 3. IBD scores relative to Finns (FIN). Histogram ordered by IBD score,
roughly proportional to the per-individual risk for recessive disease due to the
founder event. (These results are also given quantitatively for each group in Online
Table 1.) We restrict to groups with at least two samples, combining data from all
four genotyping platforms onto one plot. Data from Ashkenazi Jews and Finns are
highlighted in red, and from South Asian groups with significantly higher IBD scores
than that of Finns and census sizes of more than a million in brown. Error bars for
each IBD score are standard errors calculated by weighted block jackknife over each
chromosome. YRI=Yoruba (West African); CEU=European American.

Online Methods:

Data Sets:

We assembled a dataset of 1,955 individuals from 249 groups genotyped on the Affymetrix Human Origins array, of which data from 1,663 individuals from 230 groups are newly reported here (Figure 1a). We merged these data with the dataset published in Moorjani et al.[8], which consisted of 332 individuals from 52 groups genotyped on the Affymetrix 6.0 array. We also merged it with two additional datasets published in Metspalu et al.[9], consisting of 151 individuals from 21 groups genotyped on Illumina 650K arrays as well as a dataset published in [11], consisting of 367 individuals from 20 groups generated on Illumina Omni 1-Quad arrays. These groups come from India, Pakistan, Nepal, Sri Lanka, and Bangladesh. All samples were collected under the supervision of ethical review boards in India with informed consent obtained from all subjects.

We analyzed two different Ashkenazi Jewish datasets, one consisting of 21 individuals genotyped on Illumina 610K and 660K bead arrays[10] and one consisting of 7 individuals genotyped on Affymetrix Human Origins arrays.

Our "Affymetrix 6.0" dataset consists of 332 individuals genotyped on 329,261 SNPs, and our "Illumina_Omni" dataset consists of 367 individuals genotyped on 750,919 SNPs. We merged the South Asian and Ashkenazi Jewish data generated by the other Illumina arrays to create an "Illumina" dataset consisting of 172 individuals genotyped on 500,640 SNPs. We merged the data from the Affymetrix Human Origins arrays with the Ashkenazi Jewish data and data from the Simons Genome Diversity Project[19,20] to create a dataset with 4,402 individuals genotyped on 512,615 SNPs. We analyzed the four datasets separately due to the small intersection of SNPs between them. We merged in the 1000 Genomes Phase 3 data[21] (2,504 individuals from 26 different groups; notably, including 99 Finnish individuals) into all of the datasets. We used genome reference sequence coordinates (hg19) for analyses.

Quality Control:

We filtered the data at both the SNP and individual level. On the SNP level, we required at least 95% genotyping completeness for each SNP (across all individuals). On the individual level, we required at least 95% genotyping completeness for each individual (across all SNPs).

To test for batch effects due to samples from the same group being genotyped on different array plates, we studied instances where samples from the same group A were genotyped on both plates 1 and 2 and computed an allele frequency difference at each SNP, $Diff_A^i = \left(Freq_{PopA,Plate1}^i - Freq_{PopA,Plate2}^i\right)$. We then computed the product of these allele frequencies averaged over all SNPs for two groups A and B genotyped on the same plates, $\frac{1}{n}\sum_{i=1}^{n}\left(Diff_A^i\right)\left(Diff_B^i\right)$, as well as a standard error

from a weighted Block Jackknife across chromosomes. This quantity should be
consistent with zero within a few standard errors if there are no batch effects that
cause systematic differences across the plates, as allele frequency differences
between two samples of the same group should be random fluctuations that have
nothing to do with the array plates on which they are genotyped. This analysis
found strong batch effects associated with one array plate, and we removed these
samples from further analysis.
407
We used EIGENSOFT 5.0.1 smartpca[22] on each group to detect PCA outliers and
removed 51 samples. We also developed a procedure to distinguish recent
relatedness from founder events so that we could remove recently related
individuals. We first identified all duplicates or obvious close relatives by using
Plink[23] "genome" and GERMLINE[24] to compute IBD (described in more detail below)
and removed one individual from all pairs with a PI_HAT score greater than 0.45
and the presence of at least 1 IBD fragment greater than 30cM. We then used an
iterative procedure to identify additional recently related individuals. For sample
sizes above 5, we identified any pairs within each group that had both total IBD and
total long IBD (>20cM) that were greater than 2.5 SDs and 1 SD, respectively, from
the group mean. For sample sizes 5 or below, we used modified Z scores of
0.6745*(IBD_score - median(score))/MAD, where MAD is the median absolute
deviation, and identified all pairs with modified Z scores greater than 3.5 for both
total IBD and total long IBD as suggested by Iglewicz and Hoaglin[25]. After each
round, we repeated the process if the new IBD score was at least 30% lower than
the prior IBD score. Simulations showed that we were always able to remove a first
or second cousin in the dataset using this method (Supplementary Table 1).
Together these analyses removed 53 individuals from the Affymetrix 6.0 dataset, 21
individuals from the Illumina dataset, 43 individuals from the Illumina Omni
dataset, and 225 individuals from the Human Origins dataset.
428
429 After data quality control and merging with the 1000 Genomes Project data, the
430 Affymetrix 6.0 dataset included 2,842 individuals genotyped on 326,181 SNPs, the
431 Illumina dataset included 2,662 individuals genotyped on 484,293 SNPs, the
432 Illumina Omni dataset included 2,828 individuals genotyped on 750,919 SNPs, and
433 the Human Origins dataset included 4,177 individuals genotyped at 499,158 SNPs.
434

Simulations to Test Relatedness Filtering and IBD Analyses

We used ARGON[26] to simulate groups with different bottleneck strengths to test the
IBD analyses and relatedness filtering. We used ARGON's default settings, including
a mutation rate of $1.65*10^{-8}$ per base pair (bp) per generation and a recombination
rate of $1*10^{-8}$ per bp per generation and simulated 22 chromosomes of size 130 Mb
each. We pruned the output by randomly removing SNPs until there were 22,730
SNPs per chromosome to simulate the approximate number of positions in the
Affymetrix Human Origins array. For the IBD analyses, we simulated groups to have
descended from an ancestral group 1,800 years ago with N_e=50,000 and to have
formed two groups with N_e=25,000. These groups continued separately until 100

generations ago when they combined in equal proportions to form a group with
N_e=50,000. The group then split into 3 separate groups 72 generations ago that have
bottlenecks leading to N_e of either 400, 800, or 1600. The 3 groups then
exponentially expanded to a present size of N_e=50,000. We designed these
simulations to capture important features of demographic history typical of Indian
groups[4,8]. We chose the bottleneck sizes because they represent founder events with
approximately the strength of Finns (the bottleneck to 800), and twice as strong
(400) and half as strong (1600) as that group. We then performed the IBD analyses
described below with 99 individuals from the group with bottleneck strength
similar to that of Finns (198 haploid individuals were simulated and merged to
produce 99 diploid individuals) and different numbers of individuals from the other
groups. These analyses demonstrate that with only 4-5 individuals we can
accurately assess the strength of founder events in groups with strong founder
events (Supplementary Figure 2 and Supplementary Table 2). Weaker founder
events are more difficult to assess, but these groups are of less interest for founder
event disease mapping, so we aimed to sample ~5 individuals per group.
462
We wrote custom R scripts to simulate first and second cousin pairs. We took
individuals from the bottleneck of size 800 and performed "matings" by taking 2
individuals and recombining their haploid chromosomes assuming a rate of $1*10^{-8}$
per bp per generation across the chromosome and combining one chromosome
from each of these individuals to form a new diploid offspring. The matings were
performed to achieve first and second cousins. We then placed these back into the
group with group of size 800, and ran the relatedness filtering algorithms to
evaluate whether they would identify these individuals.
471
472   **Phasing, IBD Detection, and IBD Score Algorithm:**
473
We phased all datasets using Beagle 3.3.2 with the settings *missing=0; lowmem=true;*
gprobs=false; verbose=true[27]. We left all other settings at default. We determined IBD
segments using GERMLINE[24] with the parameters *-bits 75 -err_hom 0 -err_het 0 -*
min_m 3. We used the genotype extension mode to minimize the effect of any
possible phasing heterogeneity amongst the different groups and used the
HaploScore algorithm to remove false positive IBD fragments with the
recommended genotype error and switch error parameters of 0.0075 and 0.003[28].
We chose a HaploScore threshold matrix based on calculations from[28] for a "mean
overlap" of 0.8, which corresponds to a precision of approximately 0.9 for all genetic
lengths from 2-10cM. In addition to the procedure we developed to remove close
relatives (Quality Control section), we also removed segments longer than 20cM as
simulations showed that this increased sensitivity of the analyses (Supplementary
Table 2). We computed "IBD score" as the total length of IBD segments between 3-
20cM divided by $\left\{ \binom{2n}{2} - n \right\}$ where n is the number of individuals in each group to
normalize for sample size. We then expressed each group's score as a ratio of their
IBD score to that of Finns and calculated standard errors for this score using a
weighted Block Jackknife over each chromosome with 95% confidence intervals
defined as IBD score ±1.96*s.e.

492
We repeated these analyses with FastIBD[29] for the Affymetrix 6.0 and Illumina
datasets and observed that the results were highly correlated (r>0.96) (data not
shown). We chose GERMLINE for our main analyses, however, because the FastIBD
algorithm required us to split the datasets into different groups, since it adapts to
the relationships between LD and genetic distance in the data, and these
relationships differ across groups. We used data from several different Jewish
groups and all twenty-six 1000 Genomes groups to improve phasing, but of these
groups we only included results for Ashkenazi Jews and two outbred groups (CEU
and YRI) in the final IBD score ranking.
502
503
Disease patient analyses:
We use Affymetrix Human Origins arrays to successfully genotype 12 patients with
progressive pseudorheumatoid dysplasia (PPD) and 6 patients with
mucopolysaccharidosis (MPS) type IVA, all of whom had disease mutations
previously determined[15,16,30] (3 of the surveyed MPS patients are newly reported
here). A total of 6 of the PPD patients had Cys78Tyr mutations, 6 had Cys337Tyr
mutations (all 6 of the MPS patients had Cys78Arg mutations). We measured IBD as
described above and also detected homozygous segments within each individual by
using GERMLINE with the parameters *-bits 75 -err_hom 2 -err_het 0 -min_m 0.5 -*
homoz-only.
514
Haplotype sharing was assessed by analyzing phased genotypes for each mutation
group. At each SNP, we counted the number of identical genotypes for each allele
and computed the fraction by dividing by the total number of possible haplotypes (2
times the number of individuals). We took the larger value of the two possible
alleles (thus the fraction range was 0.5-1). We averaged these values over blocks of
10 or 25 SNPs and plotted the averages around the relevant mutation site.
521
522
Between-Group IBD Calculations:
We determined IBD using GERMLINE as above. We collapsed individuals into
respective groups and normalized for between-group IBD by dividing all IBD from
each group by $\left\{ \binom{2n}{2} \right\}$ where n is the number of individuals in each group. We
normalized for within-group IBD as described above. We defined groups with high
shared IBD as those with an IBD score greater than three times the founder event
strength of CEU (and ~1/3 the event strength of Ashkenazi Jews).
530
531
f_3-statistics:
We used the f_3-statistic[7] $f_3(Test; Ref_1, Ref_2)$ to determine if there was evidence that
the *Test* group was derived from admixture of groups related to *Ref$_1$* and *Ref$_2$*. A
significantly negative statistic provides unambiguous evidence of mixture in the
Test group. We determined the significance of the f_3-statistic using a Block Jackknife
and a block size of 5 cM. We considered statistics over 3 standard errors below zero
to be significant.
539

Computing Group Specific Drift:

We used qpGraph[7] to model each Indian group on the cline as a mixture of ANI and ASI ancestry, using the model (YRI, (Indian group, (Georgians, ANI)), [(ASI, Onge]]) proposed by Moorjani *et al.*[8] This approach provides estimates for post-admixture drift in each group (Supplementary Figure 5), which is reflective of the strength of the founder event (high drift values imply stronger founder events). We only included groups on the Indian cline in this analysis, and we removed all groups with evidence of East Asian related admixture (Figure 1b) because this admixture is not accommodated within the above model.

PCA-Normalized F_{ST} Calculations:

To account for intermarriage across groups, we used clusters based on PCA (Supplementary Text) to estimate the minimum F_{ST} for each South Asian group (Supplementary Figure 6). For the Affymetrix 6.0, Illumina, and Illumina_Omni datasets, we split the Indian cline into two different clusters and combined the Austroasiatic speakers and those with ancestry related to Austroasiatic speakers (according to the PCA of Figure 1b) into one cluster for a total of three clusters (all other groups were ignored for this analysis). For the Human Origins dataset we split the Indian cline into three different clusters and combined the groups with ancestry related to the main cluster of Austroasiatic speakers into one cluster for a total of four clusters (Khasi and Nicobarese were ignored in this analysis, because they do not cluster with the other Austroasiatic speaking groups). We then computed the F_{ST} between each group and the rest of the individuals in their respective cluster based on EIGENSOFT *smartpca* with inbreed set to YES to correct for inbreeding. For Ashkenazi Jews and Finns, we used the minimum F_{ST} to other European groups.

Code Availability:

Code for all calculations available upon request.