Transient structural variations alter gene expression and quantitative traits in *Schizosaccharomyces pombe*.

Authors

Daniel C. Jeffares[1,2], Clemency Jolly[1], Mimoza Hoti[1,2], Doug Speed[2], Charalampos Rallis[1,2,3], Christophe Dessimoz[1,4,5,6*], Jürg Bähler[1,2*], Fritz J. Sedlazeck[7*]

Affiliations:

1. Department of Genetics, Evolution and Environment, University College London, London WC1E 6BT, United Kingdom.
2. UCL Genetics Institute, University College London, London WC1E 6BT, United Kingdom.
3. Current address: School of Health, Sport and Biosciences, University of East London, London E15 4LZ, United Kingdom.
4. Department of Computer Science, University College London, London WC1E 6BT, United Kingdom.
5. Department of Ecology and Evolution & Center for Integrative Genomics, University of Lausanne, Biophore, 1015 Lausanne, Switzerland.
6. Swiss Institute of Bioinformatics, Biophore, 1015 Lausanne, Switzerland
7. Department of Computer Science, Johns Hopkins University, 21218 Baltimore, USA

* Correspondence should be addressed to C.D. (christophe.dessimoz@unil.ch), J.B. (j.bahler@ucl.ac.uk) or F.J.S. (fritz.sedlazeck@jhu.edu)

Abstract

The effects of structural variants on phenotypic diversity and evolution are poorly understood. We recently described the genetic and phenotypic variation among fission yeast strains and showed that genome-wide association studies are informative in this model. Here we extend this work by systematically identifying structural variations and investigating their consequences. We establish a curated catalog of copy number variants (CNVs) and rearrangements, including inversions and translocations. We find that SVs frequently vary within clonal populations and are weakly tagged by SNPs, consistent with rapid turnover. We show that CNVs produce measurable effects on gene expression both within and outside the duplicated regions. CNVs contribute to quantitative traits such as cell shape, cell growth under diverse conditions, sugar utilization in winemaking and antibiotic resistance, whereas rearrangements are strongly associated with reproductive isolation. Collectively, these findings have broad implications for evolution and for our understanding of quantitative traits and complex human diseases.

41 **Keywords**: structural variants, yeast, copy number variants, reproductive isolation,
42 quantitative genetics, next generation sequencing
43

INTRODUCTION
Structural variations (SVs), such as deletions, duplications, insertions, inversions or
translocations, affect organismal phenotypes and complex diseases (Klopocki and
Mundlos, 2011, Stefansson et al., 2014, Walters et al., 2013). SVs also influence
reproductive isolation within populations of yeast species (Avelar et al., 2013, Brown et
al., 2011, Rieseberg, 2001) as well as flies and mosquitoes (Ayala et al., 2013, McGaugh
and Noor, 2012). We, and others have recently begun to develop the fission yeast
Schizosaccharomyces pombe as a model for population genomics and quantitative trait
analysis (Avelar et al., 2013, Brown et al., 2011, Fawcett et al., 2014, Jeffares et al.,
2015, Zanders et al., 2014). This model organism combines the advantages of a small,
well-annotated haploid genome (Wood et al., 2002), abundant tools for genetic
manipulation and high-throughput phenotyping (Sabatinos and Forsburg, 2010), and
considerable resources of genome-scale and gene-centric data (Kim et al., 2010,
Marguerat et al., 2012, Ryan et al., 2012).
58

One facet of diversity studies in this species has been the discovery and analysis of a
small number of inversions and reciprocal translocations (Avelar et al., 2013, Zanders et
al., 2014, Brown et al., 2011). Given this evidence for SVs in this species, we recognized
that a systematic survey of SVs would progress our understanding of their biological
influence. Here, we build upon the recent availability of 161 fission yeast genomes
(Jeffares et al., 2015), representing the known population of this species worldwide, to
undertake such a survey. We show that rearrangements contribute towards reproductive
isolation, while copy number variants (CNVs) alter gene expression levels, contribute
significantly to quantitative traits, and are transient within clonal populations.
68

RESULTS
Genome- and population-wide detection of structural variations
To predict an initial set of SVs, we applied four inference software packages (Delly,
Lumpy, Pindel and cn.MOPs) (Klambauer et al., 2012, Layer et al., 2014, Rausch et al.,
2012, Ye et al., 2009) to existing short-read data (Jeffares et al., 2015), using parameters
optimized on simulated data (Methods). We then filtered these initial predictions,
accepting SVs detected by at least two callers, to obtain 315 variant calls (141 deletions,
112 duplications, 26 inversions, 36 translocations). We release this pipeline as an open
source tool called SURVIVOR (Methods). To ensure a high specificity, we further
filtered the 315 variants by removing SV calls whose breakpoints overlapped with low
complexity regions or previously annotated long terminal repeats (LTRs) (Jeffares et al.,
2015). Finally, we manually vetted all the remaining SVs by visual inspection of read

alignments in multiple strains for all 315 candidates. This meticulous approach aimed to
ensure a high quality call set, to mitigate against the high uncertainty associated with SV
calling (Layer et al., 2014).
This curation produced a set of 113 SVs, comprising 23 deletions, 64
duplications, 11 inversions and 15 translocations **(Figure 1a)**. Reassuringly, when our
variant calling methods were applied to an engineered knockout strain, we correctly
identified the known deletions and called no false positives. Attempts to validate all
rearrangements by PCR and BLAST searches of *de novo* assemblies positively verified
76% of the rearrangements, leaving only a few PCR-intractable variants unverified
(Methods).

91
92
Figure 1. Characteristics of SVs in *S. pombe*. (**a**) Relative proportions of SVs
identified. Duplications (DUP) were the most abundant SVs, followed by deletions
(DEL), inversions (INV), and translocations (TRA). (**b**) Population allele frequency
distribution of SVs, showing the frequencies of less abundant alleles in the population
(minor allele frequencies). (**c**) Length distributions of SVs, $\log_{10}$ scale. Deletions were
smallest (2.8–52 kb), duplications larger (2.6–510 kb), and inversions often very large,
spanning large portions of chromosomes (1.04 kb–5,374 kb, see (d)). Horizontal dotted
lines show the size of chromosome regions that contain an average of 1, 10 and 100 genes
in this yeast. (**d**) Locations of SVs on the three chromosomes compared to other genomic
features. From outside: density of essential genes, locations of *Tf*-type retrotransposons,
diversity (π, average pairwise diversity from SNPs), deletions (black), duplications (red),
and breakpoints of inversions and translocations as curved lines inside the concentric
circles (green and blue, respectively). Bar heights for retrotransposons, deletions and
duplications are proportional to minor allele frequencies. Diversity and retrotransposon

frequencies were calculated from 57 non-clonal strains as described by Jeffares et al.
(2015).
109
Most SVs were present at low frequencies, with 28% discovered in only one of
the strains analyzed (**Figure 1b**). The deletions were generally small (median length
595bp, **Figure 1c**), duplications showed a median length of 20 kb, with the largest
duplication extending to 510 kb and covering 200 genes (a singleton in strain
JB1207/NBRC10570). The majority of CNVs were present in copy numbers varying
between zero and sixteen (subsequently we refer to amplifications of two or more copies
as 'duplications').
Deletions and duplications and strongly biased towards the ends of chromosomes
(**Figure 1d, Supplementary Figure 1**), which are characterized by high genetic
diversity, frequent transposon insertions, and a paucity of essential genes (Jeffares et al.,
2015). All SVs preferentially occurred in positions of low gene density and showed a
strong tendency to not overlap with essential genes (**Supplementary Figure 2**). To
describe SVs further, we conducted gene enrichment analysis with the AnGeLi tool
(**Supplementary Table 1**), which interrogates gene lists for functional enrichments using
multiple qualitative and quantitative information sources (Bitton et al., 2015). The CNV-
overlapping genes were enriched for caffeine/rapamycin induced genes and genes
induced during meiosis (P = $4x10^{-7}$ and $1x10^{-5}$, respectively); they also showed lower
relative DNA polymerase II occupancy and were less likely to contain genes that are
known to produce abnormal cell phenotypes (P= $1.8x10^{-5}$ and $3x10^{-5}$, respectively). These
analyses are all broadly consistent with a paucity of CNVs in genes that encode essential
mitotic functions. Rearrangements disrupted only a few genes and showed no significant
enrichments.
132
Duplications are transient within clonal populations
Our previous work identified 25 clusters of near-clonal strains, which differed by <150
SNPs within each cluster (Jeffares et al., 2015). We expect that these clusters reflect
either repeat depositions of strains differing only at few sites (e.g. mating-type variants of
reference strains h^{90} and h^- differ by 14 SNPs) or natural populations of strains collected
from the same location. Such 'clonal populations' reflect products of mitotic propagation
from a very recent common ancestor, without any outbreeding. We therefore expected
that SVs should be largely shared within these clonal populations.
Surprisingly, our genotype predictions indicated that most SVs present in clonal
populations were segregating, i.e. were not fixed within the clonal population (68/95 SVs,
72%). Furthermore, we observed instances of the same SVs that were present in two or
more different clonal populations that were not fixed within any clonal population. These
SVs could be either incorrect allele calls in some strains, or alternatively, recent events
that have emerged during mitotic propagation. To distinguish between these two

scenarios, we re-examined the read coverage of all 49 CNVs present within at least one
clonal population. Since translocations and inversions were more challenging to
accurately genotype, we did not re-examine these variants. This analysis verified that 40
of these 49 CNVs (37 duplications, 3 deletions) were clearly segregating within at least
one clonal cluster (**Figure 2a, Supplementary Figure 3**). For example, one clonal
population of seven closely related strains, collected together in 1966 from grape must in
Sicily, have an average pairwise difference of only 19 SNPs (diversity $\pi = 1.5 \times 10^{-6}$).
Notably, this collection showed four non-overlapping segregating duplications. This
striking finding indicates that CNVs can arise or disappear frequently during evolution.
156
Transient duplications affect gene expression
Partial aneuploidies of 500-700 kb in the *S. pombe* reference strain are known to alter
gene expression levels within and, to some extent, outside of the duplicated region
(Chikashige et al., 2007). The naturally occurring duplications we described are typically
smaller (median length: 46 kb), including an average of 6.5 genes. To examine whether
naturally occurring CNVs have similar effects on gene expression, we examined eight
pairs of closely related strains (<150 SNPs among each pair) that contained at least one
unshared duplication (**Figure 2b, Supplementary Figure 3, Supplementary Table 2**).
Several of these strain pairs have been isolated from the same substrate at the same time,
and all pairs are estimated to have diverged approximately 50 to 65 years ago
(**Supplementary Table 2**). We assayed transcript expression from log phase cultures
using DNA microarrays, each time comparing a duplicated to a non-duplicated strain
from within the same clonal population. In seven of the eight strain pairs, the expression
levels of genes within duplications were significantly induced, although the degree of
expression changes between genes was variable (**Figure 2c**). The increased transcript
levels correlated with the increased genomic copy numbers, so that higher copy numbers
produced correspondingly more transcripts (**Supplementary Figure 4**). No changes in
gene expression were evident immediately adjacent to the duplications (**Supplementary
Figure 4**), suggesting that the local chromatin state was not strongly altered by the
CNVs.
Some genes outside the duplicated regions also showed altered expression levels
(**Figure 2d, Supplementary Table 3**). For example, two strain pairs differ by a single 12
kb duplication. Here, five of seven genes within the duplication showed induced
expression, while 45 genes outside the duplicated region also showed consistently altered
expression levels (38 protein-coding genes, 7 non-coding RNAs) (**Figure 2d**, arrays 7
and 8). As environmental growth conditions were tightly controlled, these changes in
gene expression probably reflect indirect and compensatory effects of the initial
perturbation caused by the duplication (**Supplementary Figure 5**). We conclude that
these evolutionary unstable duplications reproducibly affect the expression of distinct sets
of genes and thus have the potential to influence cellular function and phenotypes.

187
188
Figure 2. Transient duplications affect gene expression. (**a**) Duplications occur within
near-clonal strains. Plot showing average read coverage in 1 kb windows for two clonal
strains (JB760, JB886) with the duplication (red), five strains without duplication (green),
and two reference strains (h$^+$, and h-) (black). Genes (with exons as red rectangles) and
retrotransposon LTRs (blue rectangles) are shown on top. See **Supplementary Table 2**
for details. (**b**) Eight pairs of closely related strains, differing by one or more large
duplications, selected for expression analysis. The tree indicates the relatedness of these
strain pairs (dots colored as in d). The position of the reference strain (Leupold's 972,
JB22) is indicated with a black arrow. Black squares indicate the presence of a 32 kb
duplication that is associated with Brefeldin resistance (see below). (**c**) Gene expression
increases for most genes within duplicated regions. For each tested strain pair, we show
the relative gene expression (strains with duplication/strains without duplication) for all
genes outside the duplication (as boxplot) and for all genes within the duplication (red
strip chart). In all but one case (array 4), the genes within the duplication tend to be more
highly expressed than the genes outside of the duplication (all Wilcoxon rank sum test P-
values <1.5x10^{-3}). (**d**) Summary of expression arrays 1-8, with strains indicated as
colored dots (as in b), showing number of single-nucleotide polymorphism differences
between strains (SNP), sizes of duplications in kb (DUP, where '+X +Y' indicates two
duplications with length X and length Y, respectively). We show total numbers of

induced (up) and repressed (down) genes, both inside and outside the duplicated regions.
Arrays 2,3 and 7,8 (in yellow shading) are replicates within the same clonal population
that contain the same duplications, so we list the number of up- and down-regulated
genes that are consistent between both arrays. See **Supplementary Tables 3 and 4** for
details.
213
214
Copy number variants influence quantitative traits
To test whether SVs affect phenotypes, we examined the contributions of SNPs, CNVs
and rearrangements to 53 quantitative traits, including 11 cell shape parameters and
colony size on solid media assaying 42 stress and nutrient conditions (Jeffares et al.,
2015). For each phenotype, we used mixed model analysis to estimate the total proportion
of variance explained by the additive contribution of genomic variants (the narrow-sense
heritability).
When we determined heritability using only SNP data, estimates varied between 0% and
74%. After adding CNVs and rearrangements to SNPs in a composite model, the
estimated overall heritability increased for nearly all traits, in some cases by more than 2-
fold (e.g., resistance to Rapamycin) (**Fig 3a**). This finding indicates that the CNVs and
rearrangements can explain a substantial proportion of the trait variance. Using this
composite model, we quantified the individual contributions of heritability best explained
by SNPs, CNVs and rearrangements (**Fig 3b**). On average, SNPs explained 30% of trait
variance, CNVs 14%, and rearrangements 5% (**Supplementary Figure 6,**
Supplementary Table 5). Analysis of simulated data confirmed that the contribution of
CNVs could not be explained by linkage to causal SNPs alone (**Supplementary Figure**
6).
As some of the strains have recently been shown to have fermentation properties
that may be beneficial for winemaking (Benito et al., 2016), we examined three traits
related to wine fermentations (glucose/fructose utilization, malic acid degradation, acetic
acid content, all measured in grape must). Remarkably, the heritability of these wine-
fermentation traits was almost entirely due to SVs, with negligible contributions from
SNPs (**Supplementary Figure 7, Supplementary Table 6**). For glucose/fructose
utilization, the CNVs accounted for the entire heritability of 0.53 (**Supplementary**
Figure 8). Since many of these strains have been collected from fermentations
(**Supplementary Table 7**), the strong influence of CNVs may represent recent strong
selection and adaptation to fermentation conditions that has occurred via recent CNV
acquisition.
To locate specific SVs that affected these traits, we performed mixed model
genome-wide association studies, using all 68 SVs with minor allele counts >5 as well as
139,396 SNPs and 22,058 indels. Trait-specific significance thresholds for 5%
familywise error rates were computed via permutation analysis, and were approximately

10^{-4} (SVs), 10^{-6} (SNPs and indels). Five SVs were significantly associated with traits (3
duplications, 1 deletion, 1 translocation) (**Supplementary Table 8**). The median effect
size of these SVs was 9% (range 6-13%). The strongest signal was from a 32kb
duplication that covered 11 protein-coding genes and was significantly associated with 15
growth traits. This duplication is segregating (not fixed) within three clonal populations
(**Fig. 2b, Supplementary Figure 9**), often as part of a larger duplication. It was the most
significantly associated variant (from SNPs, indels and SVs) for growth in the antibiotic
Brefeldin, where it contributes 15% of the trait variance (**Fig 3c**). Three of the 11
duplicated genes encode transmembrane proteins, any of which could contribute to the
trait. Since some fungi produce Brefeldin to inhibit competitive growth, this duplication
is a striking example of a transient CNV that could provide a strong selective advantage
in a natural environment.
260   Our analysis of heritability showed that SNPs are generally able to capture most
of the genetic contribution of SVs (**Fig. 3**). To examine whether trait-influencing SVs
will be effectively detected by tagging SNPs in in this population, we examined the
linkage of all 113 SVs with SNPs. We found that only 63 of these SVs (55%) are in
strong linkage (r^2 >0.6), leaving 45% of the SVs weakly linked (**Supplementary Table
16**). This lack of linkage is consistent with SVs being transient, rather than persisting
within haplotypes. Such weakly-linked SVs will not be tagged well by SNPs (as shown
in **Fig. 3c**). Collectively, these analyses indicate that SVs, most notably CNVs, contribute
substantially to quantitative traits and suggests that GWAS analyses conducted without
genotyping SVs could fail to capture these genetic factors.
270

272
273
Figure 3. CNVs contribute to quantitative traits. (**a**) Heritability estimates obtained
using SNPs alone (grey bars) are generally lower than estimates obtained using SNPs,
CNVs and rearrangements (black lines extending above bars), consistent with CNVs and
rearrangements contributing to traits. The types of traits described are indicated by the
filled circles, including automated shape parameters (SHAPE/A), and growth rates on
solid media (SOL/M). (**b**) Contributions of CNVs (red), rearrangements (black) and
SNPs (open bars) to total heritability are shown with a stacked bar plot. CNVs in
particular make substantial contributions that will appear be associated with non-causal
SNPs if the causal CNVs have not been mapped in the genetic data. (**c**) Manhattan plot
for all variants vs. resistance to Brefeldin. Only variants with association P-values <0.1
are shown, including SNPs (black filled circles), indels (green crosses), and SVs (red
squares). Variants from the three chromosomes are shown, from left to right, followed by
variants from the mitochondrial genome. The 32 kb duplication on chromosome III is the
most significant variant. The horizontal dashed lines show the significance threshold

derived from 1000 permutations for SNPs (black), indels (green) and SVs (red). Each is
set to ensure a 5% family-wise error rate. The trait measured in this case was colony size
in the presence of 80 μM Brefeldin (as a ratio of colony size without Brefeldin). Similar
results, including a significant P-value for the 32kb duplication were found with 40 μM
and 120 μM Brefeldin (**Supplementary Table 8**).
293
294

Structural variations contribute to intrinsic reproductive isolation
Crosses between *S. pombe* strains produce between 90% and < 1% viable offspring
(Avelar et al., 2013, Zanders et al., 2014). We have previously shown that spore viability
correlates inversely with the number of SNPs between the parental strains (Jeffares et al.,
2015). This intrinsic reproductive isolation may be due to the accumulation of
Dobzhansky-Muller incompatibilities (variants that are neutral in one population, but
incompatible when combined) (Dobzhansky, 1933, Muller, 1939). However, genetically
distant strains also accumulate SVs, which are known to lower hybrid viability and drive
reproductive isolation (Rieseberg, 2001). In *S. pombe*, engineered inversions and
translocations reduce spore viability by ~40% (Avelar et al., 2013).
To analyse intrinsic reproductive isolation in more detail, we examined the
relationship between viability, SNPs and SVs. Both SV-distance (number of unshared
SVs between parents) and SNP-distance inversely correlated with hybrid viability
(Kendall correlation coefficients, SVs: $\tau = -0.26$, $P = 5.6 \times 10^{-3}$, SNPs: $\tau = -0.35$, $P =$
1.6×10^{-4}) (**Supplementary Figure 10**). While inversions and translocations are known to
lower hybrid viability as they affect chromosome pairing and segregation during meiosis
(Avelar et al., 2013, Delneri et al., 2003, Zanders et al., 2014), CNVs are not expected to
influence spore viability. Consistent with this view, we found that rearrangements
explained spore viability better than CNVs (rearrangements, $\tau = -0.36$, $P = 2.0 \times 10^{-4}$;
CNVs, $\tau = -0.10$, $P = 0.28$).
As the numbers of SNP and rearrangement differences between mating parents
are themselves correlated ($\tau = 0.53$, $P = 1.3 \times 10^{-8}$), we also estimated the influence of each
factor alone using partial correlations. When either SNPs or rearrangements were
controlled for, both remained significantly correlated with offspring viability ($P = 0.04$, P
$= 0.02$, respectively) (**Figure 4**). Taken together, these analyses indicate that both
rearrangements and SNPs contribute to reproductive isolation, but CNVs do not.
321

Figure 4. Both SNPs and rearrangements contribute to intrinsic reproductive isolation. Spore viability was measured from 58 different crosses from Jeffares et al. (2015) (black) or Avelar et al. (2013) (red), with each circle in the plots representing one cross. An additive linear model incorporating both SNP and rearrangement differences showed highly significant correlations with viability ($P = 1.2 \times 10^{-6}$, $r^2 = 0.39$). Both genetic distances measured using SNPs and rearrangements (inversions and translocations) significantly correlated with viability when controlling for the other factor (Kendall partial rank order correlations with viability SNPs|rearrangements $\tau = -0.20$, P = 0.03; rearrangements|SNPs $\tau = -0.22$, P = 0.02). Some strains produce low viability spores even when self-mated with their own genotype. The lowest self-mating viability of each strain pair is indicated by circle size (smaller circles indicate lower self-mating viability) to illustrate that low-viability outliers tend to include such cases (see **Supplementary Table 9** for details).

DISCUSSION

Here we present the first genome- and population-wide catalog of SVs among *S. pombe* strains. To account for the high discrepancy of available methods (Layer et al., 2014), we applied a consensus approach to identify SVs (SURVIVOR), followed by rigorous filtering and manual inspection of all calls. We focused on high specificity (the correctness of the inferred SV) rather than high sensitivity (attempting to detect all SVs).

Our previous analyses of these strains, conducted without SV data (Jeffares et al., 2015), attributed both trait variations and reproductive isolation to SNPs and/or small indels. Here we show that the small number of SVs we describe make substantial contributions to both of these factors. We demonstrate that CNVs (duplications and deletions) contribute significantly to our ability to describe quantitative traits, whereas variants that rearrange the order of the genome (inversions and translocations) produce much weaker effects on traits. In contrast, CNVs have no detectable influence on

reproductive isolation, while rearrangements contribute substantially to reproductive
isolation, similar to other species (Ayala et al., 2013, Noor et al., 2001).
A surprising aspect of our analysis is that duplications are generated and/or lost
frequently within clonal populations. For example, within seven strains that differ by as
few as 19 SNPs, we discovered four segregating duplications. A similar rapid occurrence
of duplications in *S. pombe* has been observed in laboratory conditions, where
spontaneous duplications suppress *cdc2* mutants at least 100 times more frequently than
SNPs, and these suppressor strains lose their duplications with equal frequency (Carr et
al., 1989). Similarly, duplications frequently occur during experimental evolution with
budding yeast (Dunham et al., 2002). Consistent with the transience of these variants,
they are frequently not well tagged by SNPs. These CNVs subtly alter the expression of
genes within and beyond the duplications, and contribute considerably to quantitative
traits. Within small populations, CNVs may produce larger effects on traits in the short
term than SNPs, as demonstrated by the 32 kb duplication that is associated with
resistance to the Brefeldin A (an antibiotic produced by entophytic fungi).
This analysis has relevance for human diseases, since *de novo* CNV formation in
the human genome occurs at a measurable rate (approximately one CNV/10 generations
(Itsara et al., 2010)), and CNVs are known to contribute to a wide variety of diseases
(Zhang et al., 2009). Indeed, both the population genetics and the effects of SVs within *S.*
pombe seem similar to human, in that CNVs are associated with stoichiometric changes
on gene expression, and SVs are in weak linkage with SNPs (Stranger et al., 2007,
Sudmant et al., 2015), and therefore may be badly tagged by SNPs in GWAS studies. We
show that CNVs in fission yeast contribute to quantitative traits and are also weakly
linked to SNPs. These findings highlight the need to identify SVs when describing traits
using GWAS, and indicate that a failure to call SVs can lead to an overestimation of the
impact of SNPs to traits or contribute to the problem that large proportions of the
heritable component of trait variation are not discovered in GWAS (the 'missing
heritability').
In summary, we show that different types of SVs have distinct influences at the
phenotype level and that a small number of SVs produce profound effects on the biology
of this species.

MATERIAL AND METHODS
Performance assessment of SV callers using simulated data
To identify filtering parameters for DELLY, LUMPY and Pindel for the *S. pombe*
genome, we simulated seven datasets (s1-s7) of 40x coverage with a range of different
SV types and sizes (**Supplemental Table 7**). The simulated read sets contained
sequencing errors (0.4%), SNPs and indels (0.1%) within the range of actual data from *S.*
pombe strains and between 30 and 170 SVs. These data sets were produced by modifying
the reference genome using our in-house software (SURVIVOR, described below), and
simulating reads from this genome with Mason software (Holtgrewe, 2010).
391   After mapping the reads and calling SVs, we evaluated the calls. We defined a SV
392   correctly predicted if: i) the simulated and reported SV were of the same type (e.g.
393   duplication), ii) were predicted to be on same chromosome, and iii) their start and stop
394   locations were with 1 kb. We then defined caller-specific thresholds to optimize the
395   sensitivity and false discovery rate (FDR) for each caller. FDRs on the simulated data
396   were low: DELLY (average 0.13), LUMPY (average 0.06) and Pindel (average 0.04).
397   Selecting calls that were present in at least two callers further reduced the FDR
(average of 0.01). DELLY had the highest sensitivity (average 0.75), followed by
SURVIVOR (average 0.70), LUMPY (average 0.62) and Pindel (0.55). We further used
simulated data to assess the sensitivity and FDR of our predictions. cn.mops was
evaluated with a 2 kb distance for start and stop coordinates. Our cn.mops parameters
were designed to identify large (above 12 kb) events and thus did not identify any SVs
simulated for s1-s6. Details of simulations and caller efficacy are provided in
Supplementary Table 10.
405
SURVIVOR (StructURal Variant majorIty VOte) Software Tool
We developed the SURVIVOR tool kit for assessing SVs for short read data that contains
several modules. The first module simulates SVs given a reference genome file (fasta)
and the number and size ranges for each SV (insertions, deletions, duplications,
inversions and translocations). After reading in the reference genome, SURVIVOR
randomly selects the locations and size of SV following the provided parameters.
Subsequently, SURVIVOR alters the reference genome accordingly and prints the so
altered genome. In addition, SURVIVOR provides an extended bed file to report the
locations of the simulated SVs.
415   The second module evaluates SV calls based on a variant call format (VCF) file
416   (Danecek et al., 2011) and any known list of SVs. A SV was identified as correct if i)
417   they were of same type (e.g. deletion); ii) they were reported on same chromosome, and
418   iii) the start and stop coordinates of the simulated and identified SV were within 1 kb
419   (user definable).
420   The third module of SURVIVOR was used to filter and combine the calls from
three VCF files. In our case, these files were the results of DELLY, LUMPY and Pindel.

This module includes methods to convert the method-specific output formats to a VCF
format. SVs were filtered out if they were unique to one of the three VCF files. Two SVs
were defined as overlapping if they occur on the same chromosome, their start and stop
coordinates were within 1 kb, and they were of the same type. In the end, SURVIVOR
produced one VCF file containing the so filtered calls. SURVIVOR is available at
github.com/fritzsedlazeck/SURVIVOR.
428
Read mapping and detection of structural variants
Illumina paired-end sequencing data for 161 *S. pombe* strains were collected as described
in Jeffares et al. (2015), with the addition of Leupold's reference 975 h^+ (JB32) and
excluding JB374 (known to be a gene-knockout version of the reference strain, see
below). Leupold's 968 h^{90} and Leupold's 972 h^- were included as JB50 and JB22,
respectively (**Supplementary Table 7**). For all strains, reads were mapped using
NextGenMap (version 0.4.12)(Sedlazeck et al., 2013) with the following parameter (-X
1000000) to the *S. pombe* reference genome (version ASM294v2.22). Reads with 20 base
pairs or more clipped were extracted using the script *split_unmapped_to_fasta.pl*
included in the LUMPY package (version 0.2.9)(Layer et al., 2014) and were then
mapped using YAHA (version 0.1.83)(Faust and Hall, 2012) to generate split-read
alignments. The two mapped files were merged using Picard-tools (version 1.105)
(http://broadinstitute.github.io/picard), and all strains were then down-sampled to 40x
coverage using Samtools (version 0.1.18)(Li et al., 2009).
443   Subsequently, DELLY (version 0.5.9, parameters: " –q 20 -r")(Rausch et al.,
444   2012), LUMPY (version 0.2.9, recommended parameter settings)(Layer et al., 2014) and
445   Pindel (version 0.2.5a8, default parameter)(Ye et al., 2009) were used to independently
446   identify SVs in the 161 strains using our SURVIVOR software. We then retained all
447   variants predicted by at least two methods. These SVs calls were genotyped using
448   DELLY.
449   To identify further CNVs, we ran cn.MOPS (Klambauer et al., 2012) with
parameters tuned to collect large duplications/deletions as follows: read counts were
collected from bam alignment files (as above) with *getReadCountsFromBAM* and
WL=2000, and CNVs predicted using *haplocn.mops* with minWidth= 6, all other
parameters as default. Hence, the minimum variant size detected was 12 kb. CNV were
predicted for each strain independently by comparing the alternative strain to the two
reference strains (JB22, JB32) and four reference-like strains that differed from the
reference by less than 200 SNPs (JB1179, JB1168, JB937, JB936).
457   After CNV calling, allele calling was achieved by comparing counts of coverage
in 100bp windows for the two reference strains (JB22, JB32) to each alternate strain
using custom R scripts. Alleles were called as non-reference duplications if the one-sided
Wilcoxon rank sum test p-values for both JB22 and JB32 vs alternate strain were less
than 1×10^{-10} (showing a difference in coverage) and the ratio of alternate/reference

coverage (for both JB22 and JB32) was >1.8 (duplications), or <0.2 (deletions). Manual
inspection of coverage plots showed that the vast majority of the allele calls were in
accordance with what we discerned by eye. These R scripts were also used to examine
CNVs predicted to be segregating within clusters (clonal populations). All such CNVs
were examined in all clusters that contained at least one non-reference allele call
(**Supplementary Table 11**).
Finally, we manually mapped two large duplications that did not satisfy these
criteria (DUP.I:2950001..3190000, 240kb and DUP.I:5050001..5560000, 510kb – both
singletons in JB1207), but were clearly visible in chromosome-scale read coverage plots
(Supplementary Figure 11).
472
Reduction of false discovery rate
This filtering produced 315 variant calls. However, because 31 of these 315 (~10%) were
called within the two reference strains (JB22, JB32), we expected that this set still
contained false positives. To further reduce the false positive rate, we looked for
parameters that would reduce calls made in reference strains (JB22 and JB32) but not
reduce calls in strains more distantly related to the reference (JB1177, JB916 and JB894
that have 68223, 60087 and 67860 SNP differences to reference(Jeffares et al., 2015)).
The reasoning was that we expected to locate few variants in the reference, and more
variants in the more distantly related strains. This analysis showed that paired end
support, repeats and mapping quality were of primary value.
We therefore discarded all SVs that had a paired end support of 10 or less. In
addition, we ignored SVs that appeared in low mapping quality regions (i.e. regions
where reads with MQ=0 map) or overlapped with previously identified retrotransposon
LTRs(Jeffares et al., 2015).
Finally, to ensure a high specificity call set, these filtered SVs were manually
curated using IGV (Thorvaldsdottir et al., 2013) (**Supplementary Tables 12,13**). We
assigned each SVs a score (0: not reliable, 1: unclear, 2: reliable based on inspection of
alignments through IGV). Only calls passing this manual curation as reliable (score 2)
were included in the final data set of 113 variants utilized for all further analyses.
These filtering and manual curation steps reduced our variant calls substantially, from
315 to 113. At this stage only 1/113 (~1%) of these variants was called within the two
reference strains (JB22, JB32).
495
PCR validation
PCR analysis was performed to confirm 10 of the 11 inversions and all 15 translocations
from the curated data set. One inversion was too small to examine by PCR
(INV.AB325691:6644..6784, 140 nt). Primers were designed using Primer3(Untergasser
et al., 2012) to amplify both the reference and alternate alleles. PCR was carried out with
each primer set using a selection of strains that our genotype calls predict to include at

least one alternate allele and at least one reference allele (usually 6 strains). Products
were scored according to product size and presence/absence (**Supplementary Tables 14,**
15).
505   Inversions: 9/10 variants were at least partially verified by either reference or
alternate allele PCR (3 variants were verified by both reference and alternate PCRs), and
7/10 inversions also received support from BLAST (see below). Translocations: 10/15
were at least partially verified by either reference or alternate allele PCR (5/15 variants
were verified by both reference and alternate PCRs). One additional translocation
received support from BLAST (see below), meaning that 11/15 translocations were
supported by PCR and/or BLAST. Three of the four translocations that could not be
verified were probably nuclear copies of mitochondrial genes (NUMTs) (Lenglez et al.,
2010), because one breakpoint was mapped to the mitochondrial genome.
514
Validation by BLAST of *de novo* assemblies
We further assessed the quality of the predicted breakpoints for the inversions and
translocations by comparing them to the previously created *de novo* assemblies for each
of the 161 strains (Jeffares et al., 2015). To this end, we created blast databases for the
scaffolds of each strain that were >1kb. We then created the predicted sequence for 1 kb
around each junction of the validated 10 inversions and 15 translocations. These
sequences were used to search the blast databases using BLAST+ with --gapopen 1 --
gapextend 1 parameters. We accepted any blast hsp with a length >800 bp as supporting
the junction (because these must contain at least 300 bp at each side of the break point).
Four inversions and three translocations gained support from these searches
(Supplementary File Tables2-PCR.xlsx).
526
Knockout strain control
Our sample of sequenced strains included one strain (JB374) that is known to contain
deletions of the *his3* and *ura4* genes. Our variant calling and validation methods
identified only two variants in this strain, both deletions that corresponded to the
positions of these genes, as below:
his3 gene location is chromosome II, 1489773-1488036, deletion detected at II:1488228-
1489646.
ura4 gene location is chromosome III, 115589-116726, deletion detected at III:115342-
117145.
This strain was not included in the further analyses of the SVs.
537
Microarray expression analysis
Cells were grown in YES (Formedium, UK) and harvested at OD_{600} =0,5. RNA was
isolated followed by cDNA labeling (Lyne et al., 2003). Agilent 8 x 15K custom-made *S.*
pombe expression microarrays were used. Hybridization, normalization and subsequent

washes were performed according to the manufacturer's protocols. The obtained data
were scanned and extracted using GenePix and processed for quality control and
normalization using in-house developed R scripts. Subsequent analysis of normalized
data was performed using R. Microarray data have been submitted to ArrayExpress
(accession number E-MTAB-4019). Genes were considered as induced if their expression
signal after normalization was >1.9, and repressed if <0.51.
548
Time to most recent common ancestor (TMRCA) estimates
Previously, based on the genetic distances between these strains and the 'dated tip' dating
method implemented in BEAST (Drummond et al., 2012), we have estimated the
divergence times between all 161 *S. pombe* strains sequenced (Jeffares et al., 2015). To
determine the TMRCA for pairs of strains, we re-examined the BEAST outputs using
FigTree to obtain the medium and 95% confidence intervals.
555
SNP and indel calling
SNPs were called as described (Jeffares et al., 2015). Insertions and deletions (indels)
were called in 160 strains using stampy-mapped, indel-realigned bams as described
previously (Jeffares et al., 2015). We accepted indels that were called by both the
Genome Analysis Toolkit HaplotypeCaller (DePristo et al., 2011) and
Freebayes(Garrison and Marth, 2012), and then genotyped all these calls with Freebayes.
Briefly, indels were called on each strains bam with HaplotypeCaller, and filtered
for call quality >30 and mapping quality >30 (bcftools filter --include 'QUAL>30 &&
MQ>30'). Separately, indels were called on each strains bam with Freebayes, and filtered
for call quality >30. All Freebayes vcf files were merged, accepting only positions called
by both Freebayes and HaplotypeCaller. These indels were then genotyped with
Freebayes using a merged bam (containing reads from all strains), using the --variant-
input flag for Freebayes to genotyped only the union calls. Finally indels were filtered for
by score, mean reference mapping quality and mean alternate mapping quality >30
(bcftools filter --include 'QUAL>30 && MQM>30 & MQMR>30'). These methods
identified 32,268 indels. Only 50 of these segregated between Leupold's h$^-$ reference
(JB22) and Leupold's h^{90} reference (JB50), whereas 12109 indels segregated between the
JB22 reference and the divergent strain JB916.
574
Heredity and GWAS
We selected 53 traits that contained at least values from 100 strains(Jeffares et al., 2015),
and so included multiple individuals from within clonal populations (growth rates on 42
different solid media and 11 cell shape characters measured with automated image
analysis). Trait values were normalized using a rank-based transformation in R, for each
trait vector y, normal.y =qnorm(rank(y)/(1+length(y))). Total heritability, and the
contribution of SNPs, CNVs and rearrangements were estimated using LDAK (version

5.94) (Speed et al., 2012), with kinship matrices derived from all SNPs, 87 CNVs, and 26
rearrangements. To assess whether the contribution of CNVs could be primarily due to
linkage with causal SNPs, we simulated trait data using the --make-phenos function of
LDAK with the relatedness matrix from all SNPs, assuming that all variants contributed
to the trait (--num-causals -1). We made one simulated trait data set per trait, for each of
the 53 traits, with total heritability defined as predicted from the real data. We then
estimated the heritability using LDAK, including the joint matrix of SNPs, CNVs and
rearrangements. To assess the extent to which the contribution of SNPs to heritability was
overestimated, we performed another simulation using the relatedness matrix from the 87
segregating CNVs alone, and then estimated the contribution of SNPs, CNVs and
rearrangements in this simulated data as above.
Genome-wide associations were performed with LDAK (version 5) using default
parameters, using a mixed model derived from kinship of all SNPs called previously
Jeffares et al. (2015). Association analysis was run separately for 68 SVs with a minor
allele count >5, for 139,396 SNPs and for 22,058 indels, both minor allele counts >5. We
examined the same 53 traits as for the heritability analysis (above). For each trait, we
carried out 1000 permutations of trait data, and define the 5^{th} percentile of these
permutations as the trait-specific P-value threshold.
600
Offspring viability and genetic distance
Cross spore viability data and self-mating viability were collected from previous analyses
(Avelar et al., 2013, Jeffares et al., 2015). The number of differences between each pair
was calculated using vcftools vcf-subset (Danecek et al., 2011), and correlations were
estimated using R, with the ppcor package. When calculating the number of CNVs
differences between strains, we altered our criteria for 'different' variants (to merge
variants whose starts and ends where within 1 kb), and merged CNVs if their overlap was
>50% and their allele calls were the same.
609

Acknowledgments

We thank Günter Klambauer for advice on cn.MOPS and Michael C. Schatz for helpful discussions and comments on the manuscript. F.S. was supported through National Science Foundation awards (DBI-1350041) and National Institutes of Health award (R01-HG006677). D.J., M.H., C.R. were supported by a Wellcome Trust Senior Investigator Award to J.B. (grant 095598/Z/11/Z). J.B. was supported by a Royal Society Wolfson Research Merit Award.

References

Avelar, A. T., Perfeito, L., Gordo, I. & Ferreira, M. G. 2013. Genome architecture is a selectable trait that can be maintained by antagonistic pleiotropy. *Nat Commun*, 4**,** 2235.

Ayala, D., Guerrero, R. F. & Kirkpatrick, M. 2013. Reproductive isolation and local adaptation quantified for a chromosome inversion in a malaria mosquito. *Evolution*, 67**,** 946-58.

Benito, A., Jeffares, D., Palomero, F., Calderon, F., Bai, F. Y., Bahler, J. & Benito, S. 2016. Selected Schizosaccharomyces pombe Strains Have Characteristics That Are Beneficial for Winemaking. *PLoS One*, 11**,** e0151102.

Bitton, D. A., Schubert, F., Dey, S., Okoniewski, M., Smith, G. C., Khadayate, S., Pancaldi, V., Wood, V. & Bahler, J. 2015. AnGeLi: A Tool for the Analysis of Gene Lists from Fission Yeast. *Front Genet*, 6**,** 330.

Brown, W. R., Liti, G., Rosa, C., James, S., Roberts, I., Robert, V., Jolly, N., Tang, W., Baumann, P., Green, C., Schlegel, K., Young, J., Hirchaud, F., Leek, S., Thomas, G., Blomberg, A. & Warringer, J. 2011. A Geographically Diverse Collection of Schizosaccharomyces pombe Isolates Shows Limited Phenotypic Variation but Extensive Karyotypic Diversity. *G3 (Bethesda)*, 1**,** 615-26.

Carr, A. M., Macneill, S. A., Hayles, J. & Nurse, P. 1989. Molecular cloning and sequence analysis of mutant alleles of the fission yeast cdc2 protein kinase gene: implications for cdc2+ protein structure and function. *Mol Gen Genet*, 218**,** 41-9.

Chikashige, Y., Tsutsumi, C., Okamasa, K., Yamane, M., Nakayama, J., Niwa, O., Haraguchi, T. & Hiraoka, Y. 2007. Gene expression and distribution of Swi6 in partial aneuploids of the fission yeast Schizosaccharomyces pombe. *Cell Struct Funct*, 32**,** 149-61.

Danecek, P., Auton, A., Abecasis, G., Albers, C. A., Banks, E., Depristo, M. A., Handsaker, R. E., Lunter, G., Marth, G. T., Sherry, S. T., Mcvean, G., Durbin, R. & Genomes Project Analysis, G. 2011. The variant call format and VCFtools. *Bioinformatics*, 27**,** 2156-8.

Delneri, D., Colson, I., Grammenoudi, S., Roberts, I. N., Louis, E. J. & Oliver, S. G. 2003. Engineering evolution to study speciation in yeasts. *Nature*, 422**,** 68-72.

Depristo, M. A., Banks, E., Poplin, R., Garimella, K. V., Maguire, J. R., Hartl, C.,
Philippakis, A. A., Del Angel, G., Rivas, M. A., Hanna, M., Mckenna, A.,
Fennell, T. J., Kernytsky, A. M., Sivachenko, A. Y., Cibulskis, K., Gabriel, S. B.,
Altshuler, D. & Daly, M. J. 2011. A framework for variation discovery and
genotyping using next-generation DNA sequencing data. *Nat Genet,* 43**,** 491-8.
Dobzhansky, T. 1933. On the Sterility of the Interracial Hybrids in Drosophila
Pseudoobscura. *Proc Natl Acad Sci U S A,* 19**,** 397-403.
Drummond, A. J., Suchard, M. A., Xie, D. & Rambaut, A. 2012. Bayesian phylogenetics
with BEAUti and the BEAST 1.7. *Mol Biol Evol,* 29**,** 1969-73.
Dunham, M. J., Badrane, H., Ferea, T., Adams, J., Brown, P. O., Rosenzweig, F. &
Botstein, D. 2002. Characteristic genome rearrangements in experimental
evolution of Saccharomyces cerevisiae. *Proc Natl Acad Sci U S A,* 99**,** 16144-9.
Faust, G. G. & Hall, I. M. 2012. YAHA: fast and flexible long-read alignment with
optimal breakpoint detection. *Bioinformatics,* 28**,** 2417-24.
Fawcett, J. A., Iida, T., Takuno, S., Sugino, R. P., Kado, T., Kugou, K., Mura, S.,
Kobayashi, T., Ohta, K., Nakayama, J. & Innan, H. 2014. Population genomics of
the fission yeast Schizosaccharomyces pombe. *PLoS One,* 9**,** e104241.
Garrison, E. & Marth, G. 2012. Haplotype-based variant detection from short-read
sequencing. *arXiv preprint arXiv,* 1207.3907
Holtgrewe, M. 2010. Mason-A Read Simulator for Second Generation Sequencing Data.
Institut für Mathematik und Informatik, Freie Universität Berlin.
Itsara, A., Wu, H., Smith, J. D., Nickerson, D. A., Romieu, I., London, S. J. & Eichler, E.
E. 2010. De novo rates and selection of large copy number variation. *Genome
Res,* 20**,** 1469-81.
Jeffares, D. C., Rallis, C., Rieux, A., Speed, D., Prevorovsky, M., Mourier, T.,
Marsellach, F. X., Iqbal, Z., Lau, W., Cheng, T. M., Pracana, R., Mulleder, M.,
Lawson, J. L., Chessel, A., Bala, S., Hellenthal, G., O'fallon, B., Keane, T.,
Simpson, J. T., Bischof, L., Tomiczek, B., Bitton, D. A., Sideri, T., Codlin, S.,
Hellberg, J. E., Van Trigt, L., Jeffery, L., Li, J. J., Atkinson, S., Thodberg, M.,
Febrer, M., Mclay, K., Drou, N., Brown, W., Hayles, J., Carazo Salas, R. E.,
Ralser, M., Maniatis, N., Balding, D. J., Balloux, F., Durbin, R. & Bahler, J. 2015.
The genomic and phenotypic diversity of Schizosaccharomyces pombe. *Nat
Genet,* 47**,** 235-41.
Kim, D. U., Hayles, J., Kim, D., Wood, V., Park, H. O., Won, M., Yoo, H. S., Duhig, T.,
Nam, M., Palmer, G., Han, S., Jeffery, L., Baek, S. T., Lee, H., Shim, Y. S., Lee,
M., Kim, L., Heo, K. S., Noh, E. J., Lee, A. R., Jang, Y. J., Chung, K. S., Choi, S.
J., Park, J. Y., Park, Y., Kim, H. M., Park, S. K., Park, H. J., Kang, E. J., Kim, H.
B., Kang, H. S., Park, H. M., Kim, K., Song, K., Song, K. B., Nurse, P. & Hoe, K.
L. 2010. Analysis of a genome-wide set of gene deletions in the fission yeast
Schizosaccharomyces pombe. *Nat Biotechnol,* 28**,** 617-23.

Klambauer, G., Schwarzbauer, K., Mayr, A., Clevert, D. A., Mitterecker, A., Bodenhofer,
U. & Hochreiter, S. 2012. cn.MOPS: mixture of Poissons for discovering copy
number variations in next-generation sequencing data with a low false discovery
rate. *Nucleic Acids Res,* 40**,** e69.
Klopocki, E. & Mundlos, S. 2011. Copy-number variations, noncoding sequences, and
human phenotypes. *Annu Rev Genomics Hum Genet,* 12**,** 53-72.
Layer, R. M., Chiang, C., Quinlan, A. R. & Hall, I. M. 2014. LUMPY: a probabilistic
framework for structural variant discovery. *Genome Biol,* 15**,** R84.
Lenglez, S., Hermand, D. & Decottignies, A. 2010. Genome-wide mapping of nuclear
mitochondrial DNA sequences links DNA replication origins to chromosomal
double-strand break formation in Schizosaccharomyces pombe. *Genome Res,* 20**,**
1250-61.
Li, H., Handsaker, B., Wysoker, A., Fennell, T., Ruan, J., Homer, N., Marth, G.,
Abecasis, G., Durbin, R. & Genome Project Data Processing, S. 2009. The
Sequence Alignment/Map format and SAMtools. *Bioinformatics,* 25**,** 2078-9.
Lyne, R., Burns, G., Mata, J., Penkett, C. J., Rustici, G., Chen, D., Langford, C., Vetrie,
D. & Bahler, J. 2003. Whole-genome microarrays of fission yeast: characteristics,
accuracy, reproducibility, and processing of array data. *BMC Genomics,* 4**,** 27.
Marguerat, S., Schmidt, A., Codlin, S., Chen, W., Aebersold, R. & Bahler, J. 2012.
Quantitative analysis of fission yeast transcriptomes and proteomes in
proliferating and quiescent cells. *Cell,* 151**,** 671-83.
Mcgaugh, S. E. & Noor, M. A. 2012. Genomic impacts of chromosomal inversions in
parapatric Drosophila species. *Philos Trans R Soc Lond B Biol Sci,* 367**,** 422-9.
Muller, H. J. 1939. Reversibility in Evolution Considered from the Standpoint of
Genetics. *Biological Reviews,* 14**,** 261–280.
Noor, M. A., Grams, K. L., Bertucci, L. A. & Reiland, J. 2001. Chromosomal inversions
and the reproductive isolation of species. *Proc Natl Acad Sci U S A,* 98**,** 12084-8.
Rausch, T., Zichner, T., Schlattl, A., Stutz, A. M., Benes, V. & Korbel, J. O. 2012.
DELLY: structural variant discovery by integrated paired-end and split-read
analysis. *Bioinformatics,* 28**,** i333-i339.
Rieseberg, L. H. 2001. Chromosomal rearrangements and speciation. *Trends Ecol Evol,*
16**,** 351-358.
Ryan, C. J., Roguev, A., Patrick, K., Xu, J., Jahari, H., Tong, Z., Beltrao, P., Shales, M.,
Qu, H., Collins, S. R., Kliegman, J. I., Jiang, L., Kuo, D., Tosti, E., Kim, H. S.,
Edelmann, W., Keogh, M. C., Greene, D., Tang, C., Cunningham, P., Shokat, K.
M., Cagney, G., Svensson, J. P., Guthrie, C., Espenshade, P. J., Ideker, T. &
Krogan, N. J. 2012. Hierarchical modularity and the evolution of genetic
interactomes across species. *Mol Cell,* 46**,** 691-704.
Sabatinos, S. A. & Forsburg, S. L. 2010. Molecular genetics of Schizosaccharomyces
pombe. *Methods Enzymol,* 470**,** 759-95.

Sedlazeck, F. J., Rescheneder, P. & Von Haeseler, A. 2013. NextGenMap: fast and
accurate read mapping in highly polymorphic genomes. *Bioinformatics,* 29**,** 2790-
1.
Speed, D., Hemani, G., Johnson, M. R. & Balding, D. J. 2012. Improved heritability
estimation from genome-wide SNPs. *Am J Hum Genet,* 91**,** 1011-21.
Stefansson, H., Meyer-Lindenberg, A., Steinberg, S., Magnusdottir, B., Morgen, K.,
Arnarsdottir, S., Bjornsdottir, G., Walters, G. B., Jonsdottir, G. A., Doyle, O. M.,
Tost, H., Grimm, O., Kristjansdottir, S., Snorrason, H., Davidsdottir, S. R.,
Gudmundsson, L. J., Jonsson, G. F., Stefansdottir, B., Helgadottir, I., Haraldsson,
M., Jonsdottir, B., Thygesen, J. H., Schwarz, A. J., Didriksen, M., Stensbol, T. B.,
Brammer, M., Kapur, S., Halldorsson, J. G., Hreidarsson, S., Saemundsen, E.,
Sigurdsson, E. & Stefansson, K. 2014. CNVs conferring risk of autism or
schizophrenia affect cognition in controls. *Nature,* 505**,** 361-6.
Stranger, B. E., Forrest, M. S., Dunning, M., Ingle, C. E., Beazley, C., Thorne, N., Redon,
R., Bird, C. P., De Grassi, A., Lee, C., Tyler-Smith, C., Carter, N., Scherer, S. W.,
Tavare, S., Deloukas, P., Hurles, M. E. & Dermitzakis, E. T. 2007. Relative
impact of nucleotide and copy number variation on gene expression phenotypes.
Science, 315**,** 848-53.
Sudmant, P. H., Rausch, T., Gardner, E. J., Handsaker, R. E., Abyzov, A., Huddleston, J.,
Zhang, Y., Ye, K., Jun, G., Hsi-Yang Fritz, M., Konkel, M. K., Malhotra, A.,
Stutz, A. M., Shi, X., Paolo Casale, F., Chen, J., Hormozdiari, F., Dayama, G.,
Chen, K., Malig, M., Chaisson, M. J., Walter, K., Meiers, S., Kashin, S., Garrison,
E., Auton, A., Lam, H. Y., Jasmine Mu, X., Alkan, C., Antaki, D., Bae, T.,
Cerveira, E., Chines, P., Chong, Z., Clarke, L., Dal, E., Ding, L., Emery, S., Fan,
X., Gujral, M., Kahveci, F., Kidd, J. M., Kong, Y., Lameijer, E. W., Mccarthy, S.,
Flicek, P., Gibbs, R. A., Marth, G., Mason, C. E., Menelaou, A., Muzny, D. M.,
Nelson, B. J., Noor, A., Parrish, N. F., Pendleton, M., Quitadamo, A., Raeder, B.,
Schadt, E. E., Romanovitch, M., Schlattl, A., Sebra, R., Shabalin, A. A.,
Untergasser, A., Walker, J. A., Wang, M., Yu, F., Zhang, C., Zhang, J., Zheng-
Bradley, X., Zhou, W., Zichner, T., Sebat, J., Batzer, M. A., Mccarroll, S. A.,
Genomes Project, C., Mills, R. E., Gerstein, M. B., Bashir, A., Stegle, O., Devine,
S. E., Lee, C., Eichler, E. E. & Korbel, J. O. 2015. An integrated map of structural
variation in 2,504 human genomes. *Nature,* 526**,** 75-81.
Thorvaldsdottir, H., Robinson, J. T. & Mesirov, J. P. 2013. Integrative Genomics Viewer
(IGV): high-performance genomics data visualization and exploration. *Brief
Bioinform,* 14**,** 178-92.
Untergasser, A., Cutcutache, I., Koressaar, T., Ye, J., Faircloth, B. C., Remm, M. &
Rozen, S. G. 2012. Primer3--new capabilities and interfaces. *Nucleic Acids Res,*
40**,** e115.

Walters, R. G., Coin, L. J., Ruokonen, A., De Smith, A. J., El-Sayed Moustafa, J. S.,
Jacquemont, S., Elliott, P., Esko, T., Hartikainen, A. L., Laitinen, J., Mannik, K.,
Martinet, D., Meyre, D., Nauck, M., Schurmann, C., Sladek, R., Thorleifsson, G.,
Thorsteinsdottir, U., Valsesia, A., Waeber, G., Zufferey, F., Balkau, B., Pattou, F.,
Metspalu, A., Volzke, H., Vollenweider, P., Stefansson, K., Jarvelin, M. R.,
Beckmann, J. S., Froguel, P. & Blakemore, A. I. 2013. Rare genomic structural
variants in complex disease: lessons from the replication of associations with
obesity. *PLoS One,* 8**,** e58048.
Wood, V., Gwilliam, R., Rajandream, M. A., Lyne, M., Lyne, R., Stewart, A., Sgouros,
J., Peat, N., Hayles, J., Baker, S., Basham, D., Bowman, S., Brooks, K., Brown,
D., Brown, S., Chillingworth, T., Churcher, C., Collins, M., Connor, R., Cronin,
A., Davis, P., Feltwell, T., Fraser, A., Gentles, S., Goble, A., Hamlin, N., Harris,
D., Hidalgo, J., Hodgson, G., Holroyd, S., Hornsby, T., Howarth, S., Huckle, E.
J., Hunt, S., Jagels, K., James, K., Jones, L., Jones, M., Leather, S., Mcdonald, S.,
Mclean, J., Mooney, P., Moule, S., Mungall, K., Murphy, L., Niblett, D., Odell,
C., Oliver, K., O'neil, S., Pearson, D., Quail, M. A., Rabbinowitsch, E.,
Rutherford, K., Rutter, S., Saunders, D., Seeger, K., Sharp, S., Skelton, J.,
Simmonds, M., Squares, R., Squares, S., Stevens, K., Taylor, K., Taylor, R. G.,
Tivey, A., Walsh, S., Warren, T., Whitehead, S., Woodward, J., Volckaert, G.,
Aert, R., Robben, J., Grymonprez, B., Weltjens, I., Vanstreels, E., Rieger, M.,
Schafer, M., Muller-Auer, S., Gabel, C., Fuchs, M., Dusterhoft, A., Fritzc, C.,
Holzer, E., Moestl, D., Hilbert, H., Borzym, K., Langer, I., Beck, A., Lehrach, H.,
Reinhardt, R., Pohl, T. M., Eger, P., Zimmermann, W., Wedler, H., Wambutt, R.,
Purnelle, B., Goffeau, A., Cadieu, E., Dreano, S., Gloux, S., et al. 2002. The
genome sequence of Schizosaccharomyces pombe. *Nature,* 415**,** 871-80.
Ye, K., Schulz, M. H., Long, Q., Apweiler, R. & Ning, Z. 2009. Pindel: a pattern growth
approach to detect break points of large deletions and medium sized insertions
from paired-end short reads. *Bioinformatics,* 25**,** 2865-71.
Zanders, S. E., Eickbush, M. T., Yu, J. S., Kang, J. W., Fowler, K. R., Smith, G. R. &
Malik, H. S. 2014. Genome rearrangements and pervasive meiotic drive cause
hybrid infertility in fission yeast. *Elife,* 3**,** e02630.
Zhang, F., Gu, W., Hurles, M. E. & Lupski, J. R. 2009. Copy number variation in human
health, disease, and evolution. *Annu Rev Genomics Hum Genet,* 10**,** 451-81.
801
802
803

Supplementary Figures

Supplementary Figure 1. Locations and minor allele frequencies of all structural variants in curated data set. Each of the three chromosomes is indicated by black bar, with scale (in megabases) at bottom. From top (same data as Fig 1): density of essential genes (blue), locations of *Tf*-type retrotransposons (green), and diversity (π, average pairwise diversity from SNPs, purple). Bar heights for deletions and duplications are proportional to minor allele frequency, the scale for retrotransposons is the frequency of the insertion in the 57 non-clonal strains. Diversity and retrotransposon were calculated from 57 non-clonal strains as described in Jeffares et al. (2015). Below, we show different types of SVs: deletions (black), duplications (red), inversions (green) and translocations (blue). The vertical lines terminating with open circles above dotted lines emit from the mid-point of each SV and indicate the minor allele frequencies in the population of 161 strains.

Supplementary Figure 2. Structural variations are biased towards chromosome ends and to low gene density regions. Top left panel, both CNVs and rearrangements (REA) are closer to the nearest chromosome end than the null distribution (rand), (Wilcoxon rank sum test, P-values 1.3×10^{-11} and 0.03, respectively). All other panels, calculated proportion of each duplication and deletion that contained all protein-coding or essential genes. Box plots show the distributions of these proportions for all genes (grey), and proportion of coverage by essential genes (red), compared to the null distribution (rand). All comparisons were significantly less than the null distributions (Wilcoxon rank sum test, P-values $<1.6 \times 10^{-4}$). The same analysis was performed with the junctions of inversions and translocations, by calculating the transcript coverage in the region 500 bp up- and down-stream of the predicted start and end junctions. These rearrangements are slightly biased away from genes ($P = 1.9 \times 10^{-3}$), but not significantly biased away from essential genes ($P > 0.05$). The null distributions were determined by selecting 10 regions for each actual variant/junction that were the same size, and were placed in random positions on the same chromosome and calculating the gene coverage of these regions. Essential genes were those with the Fission Yeast Phenotype Ontology term defined as FYPO:0002061 ("inviable") in PomBase.

Supplementary Figure 3. Duplications that segregate within closely related strains.
Plots show the average coverage in 1 kb non-overlapping windows for strains with a
duplication (red) and all closely related strains without duplication (green); all these
strains differ by <150 SNPs. The coverage of the two standard reference strains (h^+ and h^-
) is shown in black. Top row, from left: variant DUP.I:1216001..1300000 (cluster 12,
from Japan in 57), DUP.II:568001..698000 (cluster 12), DUP.II:1670001..1716000
(cluster 2, unknown origin), second row DUP.II:3240001..3260000 (cluster 2),
DUP.II:2116001..2134000 (cluster 1, includes reference strain from French grapes in
1947), DUP.III:1838001..1934000 (cluster 2, various locations 1921-22). Bottom row:
DUP.III:212001..258000 (cluster 6, Jamaica/USA), and DUP.III:274001..286000 (cluster
5, Sicily 1966). Genes are shown on top of plots with exons as red rectangles and
retrotransposon LTRs as blue rectangles.

Supplementary Figure 4. No significant increase in gene expression immediately

adjacent to duplications. For each duplication examined with DNA arrays, we show the

relative expression (strain 1 *vs* strain 2) near the duplication. P-values show the support

for the genes within the duplication (red vertical lines), or the 50 kb adjacent to the

duplication (green vertical lines) being more highly expressed than all other genes in the

chromosome (one-sided Wilcoxon rank sum tests). The grey horizontal lines show the 5[th],

50[th] and 95[th] percentiles for gene expression data on the chromosome. The bottom right

panel shows that the median increase in expression level within a duplication correlates

with the increase in genomic copy number. The solid back line shows the expected

increase for the 1:1 correspondence between genomic copy number and relative

expression (the line *y=x*), and the dashed line shows the linear model for the data.

868
Supplementary Figure 5. Chromosome-scale view of gene expression changes. The
relative gene expression levels (strain1/strain2) for arrays 2 and 3, and arrays 7 and 8 are
shown with their positions on the three chromosomes. Filled circles indicates genes that
we consider to be upregulated (red) or repressed (green). Those highlighted with open red
circles are consistently altered in both arrays (either 2+3, or 7+8). The blue lines show
where the segregating duplications are. Box plots at right show the spread of data.

real data

sim data: SNPs cause traits

sim data: CNVs cause traits

875
Supplementary Figure 6. Contributions of SNPs and CNVs to traits. Top panel: for
53 traits, we show the total heritability estimated by the combination of 243,289 SNPs
(green), 87 CNVs (red), and 26 rearrangements (grey). We then simulated data that was
entirely due to the effects of SNPs (middle panel), or entirely due to the effects of CNVs
(lower panel). In the middle panel, all estimates of the, usually minor, contribution of
CNVs is artefactual. In the lower panel all estimates of the contribution of SNPs is
artefactual. Again, these estimated are usually minor.
883
884

885
886
Supplementary Figure 7. Contributions to of CNVs, rearrangements and SNPs to
quantitative traits. For 223 traits we show the contribution of CNVs (red),
rearrangements (black) and SNPs (open bars) to the heritability. Traits are sorted by trait
type (indicated by coloured bars on *x* axis), and then by total heritability estimate.
Categories are wine traits (yellow), growth on liquid media (LIQ/M2; grey), growth on
solid media (SOL/M; magenta), automated shape parameters (SHAPE/A; cyan), manual
parameters (SHAPE/M; blue), growth on liquid media from this study (LIQ/M1; green)
and amino acid concentrations (AA; red). All traits are from(Jeffares et al., 2015), except
LIQ/M2(Brown et al., 2011) and wine(Benito et al., 2016).
896

898
Supplementary Figure 8. Details of the heritability of winemaking traits. Previously,
we examined the potential of the 57 non-clonal strains for winemaking(Benito et al.,
2016). Three parameters of fermentations in grape must, that are known to influence wine
quality, were measured for all strains; the acetic acid content (acetic), the residual glucose
+ fructose concentration (gluc+fruc), and the percentage of malic acid degradation
(malic). Bar plots show the estimated contribution of SNPs, CNVs and rearrangements to
these traits, using both normalized and raw (non-normalized) data. The final plot shows
the total heritability. Filled red circles show one standard deviation above the estimate,
open red circles show one standard deviation below. While estimates vary with
normalized/raw data, the SNP contributions are always low compared to SVs, and are not
significantly greater than zero for residual glucose + fructose concentration and malic
acid degradation.
911
912

Supplementary Figure 9. The 32kb duplication associated with resistance to

Brefeldin. Panels show relative coverage in 100bp windows (vs. reference genome) for

strains within clonal populations ('clonal clusters'). The coverage for strains that contain

the duplication are plotted in red, and those that do not are plotted in green. Reference

strain coverage (h^- and h^+) is shown in black. The red vertical dashed line indicates the

boundaries of the 32kb duplication – note that some clusters contain a larger duplicated

region. Clockwise from top left we show clusters 4, 11, 17 and 21 (see **Supplementary**

Table 7 for cluster members).

Supplementary Figure 10. Correlations between spore viability, parental SNP-genetic distance and parental SV-genetic distance. Spore viability was measured for 58 crosses in total, including data from both Jeffares et al. (2015) (black) and Avelar et al. (2013) (red), with each circle representing one cross. Unmerged CNV differences count any CNV as being different between parents when either start or end coordinates are more than 1 kb apart. Because this definition can cause us to count largely overlapping events as 'different', we also counted 'merged' differences where two CNVs were considered different only if their overlap was >50% of the total of both variants. This approach will exclude nested CNVs. CNV-genetic distance is not significantly correlated with viability in either case.

940
941
942 **Supplementary Figure 11. Chromosome-scale read coverage plots for three**
943 **chromosomes of strain JB1207.** Coverage is calculated relative to the reference strain
944 (JB22 in our collection). Two large duplications that did not satisfy the criteria used to
945 detect CNVs with cn.MOPs are indicated with blue arrows (DUP.I:2950001..3190000,
946 240kb and DUP.I:5050001..5560000, 510kb).