Physiological and biochemical changes associated with acute experimental

dehydration in the desert adapted mouse, *Peromyscus eremicus*

Lauren Kordonowy[1], Kaelina D. Lombardo[1], Hannah L. Green[2], Molly Dawson[2], Evice

A. Bolton[1], Sarah LaCourse[3], Matthew D. MacManes[1*]

[1] University of New Hampshire, Department of Molecular Cellular and Biomedical

Sciences

[2] University of New Hampshire, Department of Biological Sciences

[3] University of New Hampshire, Department of Psychology

[*] matthew.macmanes@unh.edu

Running title: Electrolyte change in acute dehydration

Keywords: Dehydration, desert, electrolyte, rodent

Summary statement: The establishment of baseline values for serum electrolytes

and water intake, as well as their response to acute dehydration is critical

for characterizing the physiology necessary for desert survival.

Conflict of Interest Statement: The authors declare no conflict of interest.

Abstract
Characterizing traits critical for adaptation to a given environment is an
important first step in understanding how phenotypes evolve. How animals adapt
to the extreme heat and aridity commonplace to deserts represents is an
exceptionally interesting example of these processes, and has been the focus
of study for decades. In contrast to those studies, where experiments are
conducted on either wild animals or captive animals held in non-desert
conditions, the study described here leverages a unique environmental chamber
that replicates desert conditions for captive *Peromyscus eremicus* (cactus
mouse). Here we establish baseline values for daily water intake and for serum
electrolytes, as well as the response of these variables to experimental
dehydration. In brief, *P. eremicus'* daily water intake is very low. It's serum
electrolytes are distinct from many previously studied animals, and its
response to acute dehydration is profound, though not suggestive of renal
impairment, which is atypical of mammals.

49
50

Introduction
Understanding the evolution of adaptive traits has long been one of the primary
goals in evolutionary biology. The study of the relationships between fitness
and phenotype, often powered by modern genomic techniques (59), has provided
researchers with insight into the mechanistic processes that underlie adaptive
phenotypes (15, 28). Systems in which the power of genomics can be combined
with an understanding of natural history and physiology are well suited for
the study of adaptation (9, 44) especially when researchers have the ability
to assay the link between genotype and phenotype in wild animals and then
conduct complementary experiments using representative animals in carefully
controlled laboratory environments. The study described here, characterizing
the physiology and serum biochemistry of *Peromyscus eremicus* is the first step
in a larger study aimed at understanding the genomics architecture of adaptation
to desert environments.
65

Desert adaptation has significant ecological, evolutionary, and biomedical significance. In contrast to humans and other mammals, desert rodents can survive in extreme environmental conditions and are resistant to the effects of dehydration. Physiological adaptions to deserts have been characterized in several rodents. Specifically, renal histology has been studied in multiple Heteromyid rodents (3), and the general conclusion is that these desert adapted animals have evolved elongate Loops of Henle (7, 10, 38) that are hypothesized to optimize water conservation. In addition to studies of renal histology, several studies have characterized pulmonary water loss (23, 51), water metabolism (26), and water consumption (12, 34, 35, 41, 46) in desert rodents. While desert animals possess specialized physiology that is efficient with regards to water metabolism and loss, whether or not specialized genomic adaptation exists is an active area of research (32, 36, 37).

Although the cactus mouse (*Peromyscus eremicus*) has not been a particular focus for the study of desert adaptation (but see (2, 32), this Cricetid rodent native to the arid regions of the Southwestern United States and Northern Mexico (57) offers a unique opportunity to understand physiological adaptations to deserts. *P. eremicus* is a member of a larger genus of animals known colloquially as the "*Drosophila* of mammals" (9), and *Peromyscus* species have been the focus of extensive study (25, 33, 52, 54). *P. eremicus* is a sister species to the non-desert adapted *P. californicus* (13), and it is closely related to *P. crinitus*, the canyon mouse, which is another desert adapted rodent native to Southwestern deserts.

Critical to desert survival is the ability to maintain water balance even when the acute loss of water exceeds dietary water intake (24). Indeed, the mammalian corpus consists of 60% water (30). Far from a static reservoir, proper physiologic function requires water for numerous processes, including nutrient transport (22), signal transduction, pH balance, thermal regulation (42) and the removal of metabolic waste. To accomplish these functions, a nearly constant supply of water is required to replace water loss (30), which occurs mainly via the gastrointestinal and genitourinary systems, and evaporative loss, which

is greatly accelerated in extreme heat and aridity (16). Because the body
possesses limited reserves, when loss exceeds intake during even a short period
of time, dehydration and death can occur. Mammals are exquisitely sensitive to
dehydration and possess limited compensatory mechanisms.

Characterizing desert adaptation requires careful and integrative physiological
studies, which should include a detailed characterization of water intake,
responses to dehydration, and the measurement of blood electrolytes. Indeed,
quantifying these metrics is one of the first steps in understanding how animals
survive in the extreme heat and aridity of deserts. In particular, the
electrolytes chloride and sodium are important markers of dehydration (18).
These molecules play essential roles in metabolic and physiological processes,
and they are integral to the functionally of a variety of transmembrane
transport pumps (11, 29), neurotransmission (62), and maintenance of tonicity
(19). Furthermore, hypernatremia causes restlessness, lethargy, muscle weakness,
or coma (1). Bicarbonate ion, in contrast, is primarily responsible for aiding
in the maintenance of the acid-base balance and is resorbed in the renal tubules
(39). Blood urea nitrogen (BUN) is a test that assays the abundance of urea –
the end-product for metabolism of nitrogen containing compounds. Urea is
resorbed in the glomerulus, and renal impairment is often inferred when BUN
becomes elevated (8). Importantly, the canonical model of urea resorption is
dependent on urine volume, which is markedly diminished in desert rodents, thus
limiting the utility of using BUN as an indicator of renal function. Lastly,
creatinine, a product of muscle breakdown, whose measured level does not depend
on urine volume is used as a measure of renal function (8).

Genes most frequently implicated in desert-adaptation include members of the
aquaporin family (27). However, previous work suggests that an alternative
gene family, the solute carriers, are more relevant for desert-adaptation in
the cactus mouse (32). As a first step towards fully elucidating the patterns
of adaptive evolution to deserts in *P. eremicus,* we characterized the normal
patterns of water intake and electrolyte levels as well as the physiologic
response to experimental dehydration. As such, this study provides critical

physiological and biochemical information about *P. eremicus* and its response
to dehydration and is generally useful as researchers begin to leverage large-
scale genome data against classic questions regarding the evolution of adaptive
phenotypes.
136
Materials and Methods
We used captive *P. eremicus* (n=44, 24 male, 20 female) that were descendant
from mice purchased from the University of South Carolina Peromyscus Genetic
Stock Center. The USC colony was founded using wild caught animals from a dry-
desert population in Arizona. For ongoing experimental purposes, animals are
housed in a large walk-in environmental chamber built to replicate the
environmental conditions in which this population has evolved. Specifically,
the animals experience a normal diurnal pattern of temperature fluctuation,
ranging from 90F during the daytime to 75F during the night. Relative humidity
(RH) ranges from 10% during the day to 25% during the night. Animals are housed
in standard lab mouse cages with bedding that has been dehydrated to match
desert conditions. They are fed a standard rodent chow, which has also been
dehydrated. Water is provided *ad lib* during certain phases of experimentation
and withheld completely during others. All animal care procedures follow the
guidelines established by the American Society of Mammalogy (53) and have been
approved by the University of New Hampshire Animal Care and Use Committee under
protocol number 103092.
154
All animals included in this study were sexually mature adults, as defined for
males as having scrotal testes and for females as having a perforate vaginal
meatus. A slight bias for the inclusion of males exists, as a concurrent study
of male reproductive genomics was occurring. Preliminary analyses conducted
suggest that no significant differences in any of the physiological measures,
and thus, males and females were analyzed as one group. For a subset of animals,
water intake was measured, which was accomplished via the use of customized
15ml conical tubes, wherein water intake was measured every 24 hours for a
minimum of 3 consecutive days (range 3-10 days). Animals selected for the
dehydration trial were weighed on a digital scale, housed without water for

three days, then re-weighed to determine the change in body mass due to
dehydration. At the conclusion of water measurement or after a three-day
dehydration animals were sacrificed via isoflurane overdose and decapitation.
Immediately after death, a 120uL sample of trunk blood was obtained for serum
electrolyte measurement. This was accomplished using an Abaxis Vetscan VS2
machine with a critical care cartridge, which measures the concentration of
several electrolytes (Sodium, Chloride, Bicarbonate ion, Creatinine, and Blood
Urea Nitrogen (BUN)) relevant to hydration status and renal function. Lastly,
the kidney, spleen, liver, lung, hypothalamus, testes, vas deferens and
epididymis were dissected out and stored in RNAlater (Ambion Inc.) for future
study. All statistical analyses were carried out in the statistical package,
R (50).
177
Results
We measured the daily water intake for 44 adult cactus mice (24 male, 20 female)
for between three and ten consecutive days. Mean water intake was 0.11 mL per
day per gram body weight (median=0.11, SD=0.05, min=0.033, max=0.23). We
measured levels of serum Sodium, Chloride, Bicarbonate ion, Creatinine, and
Blood Urea Nitrogen (BUN) for the same 44 adult mice, thereby establishing
normal (baseline) values for *P. eremicus* (Figure 1 and Table 1).
185
A comparison of mice provided with water *ad libitum* to mice that exposed to
experimental water deprivation for three days revealed that the dehydrated mice
lost an average of 23.2% of their body weight (median=23.9%, SD=5.3%, min=12.3%,
max=32.3%, n=13 dehydration treatment, 7 males, 6 females). Despite this
substantial weight loss, anecdotally, mice appeared healthy. They were active,
eating, and interacting with handlers and other mice, normally. The amount of
weight loss did not depend on daily water intake (p=0.63, R^2= 0.03), though the
trend suggests that animals that drink more water lost more weight). Furthermore,
body weight did not strongly influence the percent loss of body weight (Figure
2; p=0.68, R^2= 0.02).
196

In addition to a substantial loss in body weight, dehydration was associated
with differences in serum electrolytes (Figure 3; n=13 dehydrated, n=31
hydrated). These changes were subtle, but significant using a two-sample t-
test (p < 0.008 in all cases).

Lastly, the levels of serum electrolytes were tightly correlated with percent
body weight loss (Figure 4). Indeed, the relationship between the level of
serum sodium and weight loss was positive and significant, (ANOVA, F-statistic:
12.85, 11 DF, p= 0.004), as was the relationship between BUN and weight loss
(ANOVA, F-statistic: 9.089, 11 DF, p= 0.012). The relationships between weight
loss and chloride and bicarbonate levels respectively, were positive but not
significant. Of note, all data will be deposited in Dryad upon acceptance for
publication.

<u>Discussion</u>
Deserts are amongst the harshest environments on the planet. Indeed, animals
living in these areas must be highly adapted to the unique combination of
extreme heat and aridity. Given that our understanding of the physiology of
desert adapted animals is limited largely to studies in renal histology (38)
and on water intake and output (34, 56), an enhanced understanding of serum
electrolyte changes due to dehydration is informative. Because many of the
harmful effects of dehydration result from electrolyte abnormalities,
characterizing normal values and the electrolyte response to dehydration
represents a critical first step in garnering a deeper understanding of how
desert animals survive despite severe and prolonged dehydration.

In this study, normal (baseline) values for serum Sodium, Chloride, Bicarbonate
Ion, Creatinine, and Blood Urea Nitrogen were established in a captive colony
of lab animals housed in desert conditions. Although these measures may differ
in wild animals (see (14) for a brief review of such differences), establishing
normal values in captive animals is crucial, though future studies aim to
understand the patterns of electrolyte variation in wild animals. In *P. eremicus*,
we define the normal ranges for each electrolyte as those values falling between

the 1st and 3rd quartile. Serum Chloride and Sodium were significantly higher

than in published ranges for other mammals, including humans, a marsupial (58),

Cricetomys (48), and the porcupine (43). However, serum chloride and sodium

levels in our study were quite comparable to another wild rodent, *Neotoma*

fuscipes (61), a Mustelid (55), and the Hyrax (5). Values for BUN are generally

higher in this study; unfortunately, a direct comparison is not possible, as

measured values are dependent on the volume of urine produced. Serum Creatinine

is low, largely resulting from the general lack of muscle mass in *P. eremicus*

relative to other mammals. However, because the equipment used to analyze this

electrolyte does not effectively capture the lower end of the biological range,

direct comparisons are not made for this metric. Future measurements, using a

more sensitive HPLC method for quantitating serum Creatinine will improve our

ability to detect more subtle changed of change.

In addition to characterizing baseline electrolytes and their response to

experimental dehydration, the normative value for daily water intake was

estimated to be 0.11 mL per day per gram body weight. Though comparable measures

of water consumption are scarce, one study in two arid adapted *Limoys* (*L.*

pictus and *L. irroratus*) housed in non-desert captive settings were estimated

to be 0.18 and 0.17 mL per day per gram body weight respectively (17) – a value

much greater than in *P. eremicus*.

Animals that were exposed to experimental dehydration lost a substantial amount

of body weight. Dehydration in humans, resulting in loss of even a fraction of

this amount results in cardiovascular collapse and death (40). Indeed, even a

dehydration-related loss of a few percent of body weight may cause serious

renal impairment or renal failure. That the cactus mouse may lose so much

weight as a result of dehydration and remain active, and apparently healthy,

without renal impairment is a testament to their desert adaptation. The

magnitude of weight loss and the negative (though non-significant) relationship

between baseline weight and weight loss, coupled with the lack of behavioral

impairments suggests that metabolic water production via the oxidation of fat

may be an important and potentially adaptive mechanism preventing more serious

complications from acute dehydration. Indeed, water metabolism may produce a
substantial amount of water (reviewed in (31)), and had been demonstrated in a
diverse group of animals including marine mammals (49) and desert rodents (20).
Future studies of fat metabolism in *P. eremicus*, using computed tomography,
are planned. Because animals are essentially anuric, particularly when
dehydrated, means direct measurement of fax oxidation (*e.g.,* urine beta
hydroxybutryate) is not possible.

As described above, mice appear grossly behaviorally intact. Despite this, they
may be experiencing a degree of cognitive impairment, as is the case with human
dehydration, where even mild-dehydration is associated with cognitive
impairment (4). Future studies, using classical Y-maze and novel object
recognition tests aim to understand more fully the cognitive effects of
dehydration in cactus mouse.

In addition to weight loss, dehydrated animals demonstrated biochemical
evidence of physiological stress, in the form of increased Sodium, Chloride,
BUN, and Bicarb. There were no significant relationships between any
physiological value and Creatinine, suggesting that dehydration related stress
does not result in renal impairment or damage. Indeed, this is in contrast to
humans and other mammals where acute dehydration of the nature imposed on these
animals is universally related to renal failure and subsequent death. That *P.*
eremicus can withstand this level of dehydration is a testament to the processes
involved in adaptation.

In summary, we present here a set of physiological measurements that represent
the endpoints in the physiological management of acute dehydration in the
desert adapted cactus mouse. How these endpoint are achieved is an outstanding
question deserving future study, particularly in light of global climate change
(31). For instance Vasopressin, along with the Renin-Angiotensin-Aldosterone
system are thought to be a critically important to the regulation of water and
solute balance (6, 45, 47, 60, 63). Comparative genomic analysis, studies of

gene expression, and the measurement of protein levels will provide important insights into the actual mechanisms underlying these phenotypes.

In addition to understanding the mechanisms of salt and water balance, characterizing the ways in which desert animals prevent dehydration-linked renal failure is exceptionally important. Unlike humans, where repeated dehydration events leads to a progressive decline in renal function (21), it is hypothesized that repeated acute-dehydration is unlikely to be linked to renal failure in animals that have evolved in desert environments. Testing this hypothesis, along with understanding the mechanisms which limit renal damage (*e.g.,* modulating renal microcirculation, maintaining cell volume via organic osmolytes) could provide previously uncharacterized clinically-relevant insights into renal (dis)function.

1. **Adrogué HJ, Madias NE.** Hypernatremia. *New Engl J Med* 342: 1493–1499, 2000.

2. **al-Kahtani MA, Zuleta C, Caviedes-Vidal E, Garland T.** Kidney mass and relative medullary thickness of rodents in relation to habitat, body size, and phylogeny. *Physiol Biochem Zool* 77: 346–365, 2004.

3. **Altschuler EM, Nagle RB, Braun EJ, Lindstedt SL, Krutzsch PH.** Morphological study of the desert heteromyid kidney with emphasis on the genus *Perognathus*. *Anat Rec* 194: 461–468, 1979.

4. **Armstrong LE, Ganio MS, Casa DJ, Lee EC, McDermott BP, Klau JF, Jimenez L, Le Bellego L, Chevillotte E, Lieberman HR.** Mild dehydration affects mood in healthy young women. *Journal of Nutrition* 142: 382–388, 2012.

5. **Aroch I, King R, Baneth G.** Hematology and serum biochemistry values of trapped, healthy, free-ranging rock hyraxes (*Procavia capensis*) and their association with age, sex, and gestational status. *Vet Clin Pathol* 36: 40–48, 2007.

6. **Bankir L, Fernandes S, Bardoux P, Bouby N, Bichet DG.** Vasopressin-V2 receptor stimulation reduces sodium excretion in healthy humans. *J Am Soc Nephrol* 16: 1920–1928, 2005.

7. **Barrett JM, Kriz W, Kaissling B, De Rouffignac C.** The ultrastructure of the nephrons of the desert rodent (*Psammomys obesus*) kidney. II. Thin limbs of Henle of long-looped nephrons. *Am J Anat* 151: 499–514, 1978.

8.   **Baum N, Dichoso CC, Carlton CE.** Blood urea nitrogen and serum
creatinine. Physiology and interpretations. *Urology* 5: 583–588, 1975.

9.   **Bedford NL, Hoekstra HE.** Peromyscus mice as a model for studying natural
variation. *eLife* 4:e06813, 2015.

10.   **Beuchat CA.** Structure and concentrating ability of the mammalian kidney:
correlations with habitat. *Am J Physiol* 271: R157–79, 1996.

11.   **Blaustein MP, Lederer WJ.** Sodium/calcium exchange: its physiological
implications. *Physiol Rev* 79: 763–854, 1999.

12.   **Bradford DF.** Water stress of free-living *Peromyscus truei*. *Ecology* 55:
1407–1414, 1974.

13.   **Bradley RD, Durish ND, Rogers DS, Miller JR, Engstrom MD, Kilpatrick CW.**
Toward a molecular phylogeny for *Peromyscus*: Evidence from mitochondrial
cytochrome-b sequences. *J Mammal* 88: 1146–1159, 2007.

14.   **Calisi RM, Bentley GE.** Lab and field experiments: Are they the same
animal? *Horm Behav* 56: 1–10, 2009.

15.   **Castoe TA, de Koning APJ, Hall KT, Card DC, Schield DR, Fujita MK,**
Ruggiero RP, Degner JF, Daza JM, Gu W, Reyes-Velasco J, Shaney KJ,
Castoe JM, Fox SE, Poole AW, Polanco D, Dobry J, Vandewege MW, Li Q,
Schott RK, Kapusta A, Minx P, Feschotte C, Uetz P, Ray DA, Hoffmann FG,
Bogden R, Smith EN, Chang BSW, Vonk FJ, Casewell NR, Henkel CV,
Richardson MK, Mackessy SP, Bronikowsi AM, Yandell M, Warren WC, Secor
SM, Pollock DD. The Burmese python genome reveals the molecular basis
for extreme adaptation in snakes. *PNAS* 110: 20645–20650, 2013.

16.   **Cheuvront SN, Kenefick RW, Montain SJ, Sawka MN.** Mechanisms of aerobic
performance impairment with heat stress and dehydration. *Journal of*
Applied Physiology 109: 1989–1995, 2010.

17.   **Christian DP, Matson JO, Rosenberg SG.** Comparative water balance in two
species of *Liomys*. *Comp Biochem Physiol, Part A Mol Integr Physiol* 61A:
589–559, 1978.

18.   **Costill DL, Coté R, Fink W.** Muscle water and electrolytes following
varied levels of dehydration in man. *Journal of Applied Physiology* 40:
6–11, 1976.

19.   **Feig PU, McCurdy DK.** The hypertonic state. *N Engl J Med* 297: 1444–1454,
1977.

20.   **Frank CL.** Diet Selection by a Heteromyid Rodent: Role of Net Metabolic
Water Production. *Ecology* 69: 1943–1951, 1988.

21.   **Glaser J, Lemery J, Rajagopalan B, Diaz HF, García-Trabanino R, Taduri**
G, Madero M, Amarasinghe M, Abraham G, Anutrakulchai S, Jha V,

Stenvinkel P, Roncal-Jimenez C, Lanaspa MA, Correa-Rotter R, Sheikh-Hamad D, Burdmann EA, Andres-Hernando A, Milagres T, Weiss I, Kanbay M, Wesseling C, Sánchez-Lozada LG, Johnson RJ. Climate Change and the Emergent Epidemic of CKD from Heat Stress in Rural Communities: The Case for Heat Stress Nephropathy. *Clinical Journal of the American Society of Nephrology* 11: 1472–1483, 2016.

22. **Haussinger D.** The role of cellular hydration in the regulation of cell function. *Biochem J* 313 (Pt 3): 697–710, 1996.

23. **Hayes J, Bible C, Boone J.** Repeatability of mammalian physiology: Evaporative water loss and oxygen consumption of *Dipodomys merriami*. *J Mammal* 79: 475–485, 1998.

24. **Heimeier RA, Davis BJ, Donald JA.** The effect of water deprivation on the expression of atrial natriuretic peptide and its receptors in the spinifex hopping mouse, *Notomys alexis*. *Comparative Biochemistry and Physiology Part A: Physiology* 132: 893–903, 2002.

25. **Hoekstra H, Moghadam HK, Hoekstra J, Harrison PW, Berrigan D, Zachar G, Székely T, Vignieri S, Mank JE, Hoang A, Hill C, Beerli P, Kingsolver J.** Strength and tempo of directional selection in the wild. *P Natl Acad Sci Usa* 98: 9157–9160, 2001.

26. **Howell AB, Gersh I.** Conservation of water by the rodent *Dipodomys*. *J Mammal* 16: 1, 1935.

27. **Huang Y, Tracy R, Walsberg GE, Makkinje A, Fang P, Brown D, Van Hoek AN.** Absence of aquaporin-4 water channels from kidneys of the desert rodent *Dipodomys merriami merriami*. *Am J Physiol-Renal* 280: F794–F802, 2001.

28. **Huerta-Sánchez E, Jin X, Asan, Bianba Z, Peter BM, Vinckenbosch N, Liang Y, Yi X, He M, Somel M, Ni P, Wang B, Ou X, Huasang, Luosang J, Cuo ZXP, Li K, Gao G, Yin Y, Wang W, Zhang X, Xu X, Yang H, Li Y, Wang J, Wang J, Nielsen R.** Altitude adaptation in Tibetans caused by introgression of Denisovan-like DNA. *Nature* 512: 194–197, 2014.

29. **Jentsch TJ, Stein V, Weinreich F, Zdebik AA.** Molecular Structure and Physiological Function of Chloride Channels. *Physiol Rev* 82: 503–568, 2002.

30. **Jéquier E, Constant F.** Water as an essential nutrient: the physiological basis of hydration. *Eur J Clin Nutr* 64: 115–123, 2009.

31. **Johnson RJ, Johnson RJ, Stenvinkel P, Stenvinkel P, Jensen T, Jensen T, Lanaspa MA, Lanaspa MA, Roncal C, Roncal C, Song Z, Song Z, Bankir L, Bankir L, Sánchez-Lozada LG, Sanchez-Lozada LG.** Metabolic and Kidney Diseases in the Setting of Climate Change, Water Shortage, and Survival Factors. *Journal of the American Society of Nephrology* 27: 2247–2256, 2016.

32. **MacManes MD, Eisen MB.** Characterization of the transcriptome, nucleotide
sequence polymorphism, and natural selection in the desert adapted mouse
Peromyscus eremicus. *PeerJ* 2: e642, 2014.

33. **MacManes MD, Lacey EA.** Is promiscuity associated with enhanced selection
on MHC-DQα in mice (genus *Peromyscus*)? *PLOS ONE* 7: e37562, 2012.

34. **MacMillen RE, Lee AK.** Australian desert mice: independence of exogenous
water. *Science* 158: 383–385, 1967.

35. **Mares M.** Water Economy and Salt Balance in a South-American Desert
Rodent, *Eligmodontia typus*. *Comp Biochem Phys A* 56: 325–332, 1977.

36. **Marra NJ, Eo SH, Hale MC, Waser PM, DeWoody A.** A priori and a posteriori
approaches for finding genes of evolutionary interest in non-model
species: Osmoregulatory genes in the kidney transcriptome of the desert
rodent *Dipodomys spectabilis* (Banner-Tailed Kangaroo Rat). *Comparative
Biochemistry and Physiology - Part D: Genomics and Proteomics* (July 26,
2012). doi: 10.1016/j.cbd.2012.07.001.

37. **Marra NJ, Romero A, DeWoody JA.** Natural selection and the genetic basis
of osmoregulation in heteromyid rodents as revealed by RNA-seq. *Mol Ecol*
23: 2699–2711, 2014.

38. **Mbassa GK.** Mammalian renal modifications in dry environments. *Vet Res
Commun* 12: 1–18, 1988.

39. **McKinney TD, Burg MB.** Bicarbonate and fluid absorption by renal proximal
straight tubules. *Kidney Int* 12: 1–8, 1977.

40. **Mehta RL, Cerdá J, Burdmann EA, Tonelli M, García-García G, Jha V,
Susantitaphong P, Rocco M, Vanholder R, Sever MS, Cruz D, Jaber B,
Lameire NH, Lombardi R, Lewington A, Feehally J, Finkelstein F, Levin N,
Pannu N, Thomas B, Aronoff-Spencer E, Remuzzi G.** International Society
of Nephrology's 0by25 initiative for acute kidney injury (zero
preventable deaths by 2025): a human rights case for nephrology. *Lancet*
385: 2616–2643, 2015.

41. **Merkt JR, Taylor CR.** "Metabolic switch" for desert survival. *P Natl Acad
Sci Usa* 91: 12313–12316, 1994.

42. **Montain S, Latzka W, Sawka N.** Fluid replacement recommendations for
training in hot weather. *Mil Med* 164: 502–508, 1999.

43. **Moreau B, Vié JC, Cotellon P, Thoisy ID, Motard A, Raccurt CP.**
HEMATOLOGIC AND SERUM BIOCHEMISTRY VALUES IN TWO SPECIES OF FREE-RANGING
PORCUPINES (*COENDOU PREHENSILIS, COENDOU MELANURUS*) IN FRENCH GUIANA. *J
Zoo Wildl Med* 34: 159–162, 2003.

44. **Mullen LM, Vignieri SN, Gore JA, Hoekstra HE.** Adaptive basis of
geographic variation: genetic, phenotypic and environmental differences

among beach mouse populations. *Proc Biol Sci* 276: 3809–3818, 2009.

45. **Muñoz A, Riber C, Trigo P.** Dehydration, electrolyte imbalances and
renin-angiotensin-aldosterone-vasopressin axis in successful and
unsuccessful endurance horses Equine Veterinary Journal - Wiley Online
Library. *Equine Veterinary …* (2010). doi: 10.1111/j.2042-
3306.2010.00211.x/pdf.

46. **Nagy KA.** Seasonal patterns of water and energy balance in desert
vertebrates. *J Arid Environ* 14: 201–210, 1988.

47. **Nielsen S, Chou C, Marples D, Christensen E, Kishore B, Knepper M.**
Vasopressin increases water permeability of kidney collecting duct by
inducing translocation of aquaporin-CD water channels to plasma-
membrane. *P Natl Acad Sci Usa* 92: 1013–1017, 1995.

48. **Nssien M, Olayemi FO, Onwuka SK, Olusola A.** Comparison of some plasma
biochemcial parameters in two generations of african giant rat
(*Cricetomys gambianus*, waterhouse). *African Journal of Biomedical
Research* 5, 2002.

49. **Ortiz RM.** Osmoregulation in marine mammals. *Journal of Experimental
Biology* 204: 1831–1844, 2001.

50. **R Core Development Team F.** R: A Language and Environment for Statistical
Computing [Online]. R Development Core Team: 2011. http://www.R-
project.org.

51. **Schmidt-Nielsen B, Schmidt-Nielsen K.** Pulmonary Water Loss in Desert
Rodents. *Am J Physiol* 162: 31–36, 1950.

52. **Shorter KR, Crossland JP, Webb D, Szalai G, Felder MR, Vrana PB.**
Peromyscus as a Mammalian Epigenetic Model. *Genetics Research
International* 2012: 1–11, 2012.

53. **Sikes RS, Gannon WL, Animal Care and Use Committee of the American
Society of Mammalogists.** Guidelines of the American Society of
Mammalogists for the use of wild mammals in research. *J Mammal* 92: 235–
253, 2011.

54. **Steiner CC, Weber JN, Hoekstra HE.** Adaptive variation in beach mice
produced by two interacting pigmentation genes. *PLoS Biol* 5: e219, 2007.

55. **Thornton PC, Wright PA, Sacra PJ, Goodier TE.** The ferret, *Mustela
putorius furo,* as a new species in toxicology. *Lab Anim* 13: 119–124,
1979.

56. **Tracy R, Walsberg G.** Intraspecific variation in water loss in a desert
rodent, *Dipodomys merriami*. *Ecology* 82: 1130–1137, 2001.

57. **Veal R, Caire W.** *Peromyscus eremicus*. *Mammalian Species* 118: 1–6, 2001.

58. **Viggers KL, Lindenmayer DB**. Variation in Hematological and Serum
Biochemical Values of the Mountain Brushtail Possum, *Trichosurus caninus*
Ogilby (Marsupialia: Phalangeridae). *J Wildlife Dis* 32: 142–146, 1996.

59. **Vignieri SN, Larson JG, Hoekstra HE**. The selective advantage of crypsis
in mice. *Evolution* 64: 2153–2158, 2010.

60. **Weaver D, Walker L, Alcorn D, Skinner S**. The contributions of renin and
vasopressin to the adaptation of the Australian spinifex hopping mouse
(*Notomys alexis*) to free water deprivation. *Comp Biochem Physio* 108:
107–116, 1994.

61. **Weber DK, Danielson K, Wright S, Foley JE**. Hematology and serum
biochemistry values of dusky-footed wood rat (/). *J Wildlife Dis* 38:
576–582, 2002.

62. **Yu FH, Catterall WA**. Overview of the voltage-gated sodium channel
family. *Genome Biology* 4: 207, 2003.

63. **Zuber AM, Singer D, Penninger JM, Rossier BC, Firsov D**. Increased renal
responsiveness to vasopressin and enhanced V2 receptor signaling in
RGS2-/- mice. *J Am Soc Nephrol* 18: 1672–1678, 2007.

Table 1

	Normal	Min	Max	Mean
Sodium (mMol/L)	148-158	144	170	153
Chloride (mMol/L)	110-115	105	126	113
BUN (mg/dL)	29-46	22	64	37
Bicarb (mMol/L)	19-25	15	26	22
Creatinine (mg/dL)	>0.2-0.3	>0.2	0.4	0.22

506 **Table 1.** Normal values for serum electrolytes. Normal values (n=44, 24 male,

507 20 female) are defined as those values falling between the 1st and 3rd quartile.

508 Of note, the Abaxis VS2 electrolyte analyzer does not measure Creatinine below

509 0.2 mg/dL; therefore, the range for normal Creatinine is truncated at this
510 value.

524 **Figure 1**. Normal values (n=44, 24 male, 20 female) for serum electrolytes.
525 Human normal values (from Medline) are plotted for comparison in dotted red
526 lines. Of note, the Abaxis VS2 electrolyte analyzer does not measure Creatinine
527 below 0.2 mg/dL, and therefore the range for normal Creatinine is truncated at
528 this value.

538 **Figure 2.** Percent body weight loss as a function of initial body weight due to
539 experimental dehydration. No significant trend exists. (n=13, 7 males, 6
540 females)

Figure 3. Experimental dehydration resulted in increases in serum sodium, chloride, BUN and bicarbonate ion. Reported p-values are from a two-tailed t-test (n=13 dehydrated: DRY, n=31 hydrated: Wet)

567 **Figure 4.** The relationship between serum electrolytes is positive in all cases
568 and significant for Sodium (F-statistic: 12.85, 11 DF, p-value: 0.004283) and
569 BUN (F-statistic: 9.089, 11 DF, p-value: 0.01177). n=13, 7 males, 6 females

571