Physiological and biochemical changes associated with experimental dehydration in the

desert adapted cactus mouse, *Peromyscus eremicus*

Lauren Kordonowy[1], Kaelina Lombardo[1], Hannah Green[2], Evice Bolton[1], Sarah LaCourse[3], Matthew

MacManes[1]

[1] University of New Hampshire, Department of Molecular Cellular and Biomedical Sciences
[2] University of New Hampshire, Department of Biological Sciences
[3] University of New Hampshire, Department of Psychology

<u>**Abstract**</u>

Characterizing traits critical for adaptation to a given environment is an important first step in

understanding how phenotypes evolve. How animals adapt to the extreme heat and aridity

commonplace to deserts represents is an exceptionally interesting example of these processes, and has

been the focus of study for decades. In contrast to those studies, where experiments are conducted on

either wild animals or captive animals held in non-desert conditions, the study described here leverages

a unique environmental chamber that replicates desert conditions for captive *Peromyscus eremicus*

(cactus mouse). Here we establish baseline values for daily water intake and for serum electrolytes, as

well as the response of these variables to experimental dehydration. In brief, *P. eremicus'* daily water

intake is very low. It's serum electrolytes are distinct from many previously studied animals, and its

response to dehydration if profound, though not suggestive of renal impairment in the face of profound

dehydration, which is atypical of mammals.

Introduction

Understanding the evolution of adaptive traits has long been one of the primary goals in evolutionary

biology. The study of the relationships between fitness and phenotype, often powered by modern

genomic techniques (Vignieri et al. 2010), has provided researchers with insight into the mechanistic

processes that underlie adaptive phenotypes (Castoe et al. 2013; Huerta-Sánchez et al. 2014). Systems

in which the power of genomics can be combined with an understanding of natural history and

physiology are well suited for the study of adaptation (Mullen et al. 2009; Bedford and Hoekstra 2015)

especially when researchers have the ability to assay the link between genotype and phenotype in wild

animals and then conduct complementary experiments using representative animals in carefully

controlled laboratory environments. The study described here, characterizing the physiology and serum

biochemistry of *Peromyscus eremicus* is the first step in a larger study aimed at understanding the

genomics architecture of adaptation to desert environments.

Desert adaptation has significant ecological, evolutionary, and biomedical significance. In contrast to

humans and other mammals, desert rodents can survive in extreme environmental conditions and are

resistant to the effects of dehydration. Physiological adaptions to deserts have been characterized in

several rodents. Specifically, renal histology has been studied in multiple Heteromyid rodents (Altschuler

et al. 1979), and the general conclusion is that these desert adapted animals have evolved elongate

Loops of Henle (Barrett et al. 1978; Mbassa 1988; Beuchat 1996) that are hypothesized to optimize

water conservation. In addition to studies of renal histology, several studies have characterized

pulmonary water loss (Schmidt-Nielsen and Schmidt-Nielsen 1950; Hayes et al. 1998), water metabolism

(Howell and Gersh 1935), and water consumption (MacMillen and Lee 1967; Bradford 1974; Mares

1977; Nagy 1988; Merkt and Taylor 1994) in desert rodents. While desert animals possess specialized

physiology that is efficient with regards to water metabolism and loss, whether or not specialized

genomic adaptation exists is an active area of research (Marra et al. 2012; MacManes and Eisen 2014;

Marra et al. 2014) .

Although the cactus mouse (*Peromyscus eremicus*) has not been a particular focus for the study of

desert adaptation (but see (al-Kahtani et al. 2004; MacManes and Eisen 2014), this Cricetid rodent

native to the arid regions of the Southwestern United States and Northern Mexico (Veal and Caire 2001)

offers a unique opportunity to understand physiological adaptations to deserts. *P. eremicus* is a

member of a larger genus of animals known colloquially as the "*Drosophila* of mammals" (Bedford and

Hoekstra 2015), and *Peromyscus* species have been the focus of extensive study (Hoekstra et al. 2001;

Steiner et al. 2007; MacManes and Lacey 2012; Shorter et al. 2012). *P. eremicus* is a sister species to the

non-desert adapted *P. californicus* (Bradley et al. 2007), and it is closely related to *P. crinitus*, the canyon

mouse, which is another desert adapted rodent native to Southwestern deserts.

Critical to desert survival is the ability to maintain water balance even when the loss of water exceeds

dietary water intake (Heimeier et al. 2002). Indeed, the mammalian corpus consists of 60% water

(Jéquier and Constant 2009). Far from a static reservoir, proper physiologic function requires water for

numerous processes, including nutrient transport (Haussinger 1996), signal transduction, pH balance,

thermal regulation (Montain et al. 1999) and the removal of metabolic waste. To accomplish these

functions, a nearly constant supply of water is required to replace water loss (Jéquier and Constant

2009), which occurs mainly via the gastrointestinal and genitourinary systems, and evaporative loss,

which is greatly accelerated in extreme heat and aridity (Cheuvront et al. 2010). Because the body

possesses limited reserves, when loss exceeds intake during even a short period of time, dehydration

and death can occur. Mammals are exquisitely sensitive to dehydration and possess limited

compensatory mechanisms.

Characterizing desert adaptation requires careful and integrative physiological studies, which should

include a detailed characterization of water intake, responses to dehydration, and the measurement of

blood electrolytes. Indeed, quantifying these metrics is one of the first steps in understanding how

animals survive in the extreme heat and aridity of deserts. In particular, the electrolytes chloride and

sodium are important markers of dehydration (Costill et al. 1976). These molecules play essential roles

in metabolic and physiological processes, and they are integral to the functionally of a variety of

transmembrane transport pumps (Blaustein and Lederer 1999; Jentsch et al. 2002), neurotransmission

(Yu and Catterall 2003), and maintenance of tonicity (Feig and McCurdy 1977). Furthermore,

hypernatremia causes restlessness, lethargy, muscle weakness, or coma (Adrogué and Madias 2000).

Bicarbonate ion, in contrast, is primarily responsible for aiding in the maintenance of the acid-base

balance and is resorbed in the renal tubules (McKinney and Burg 1977). Blood urea nitrogen (BUN) is a

test that assays the abundance of urea – the end product for metabolism of nitrogen containing

compounds. Urea is resorbed in the glomerulus, and renal impairment is often inferred when BUN

becomes elevated (Baum et al. 1975). Importantly, the canonical model of urea resorption is dependent

on urine volume, which is markedly diminished in desert rodents, thus limiting the utility of using BUN

as an indicator of renal function. Lastly, creatinine, a product of muscle breakdown, whose measured

level does not depend on urine volume is used as a measure of renal function (Baum et al. 1975).

Genes most frequently implicated in desert-adaptation include members of the aquaporin family (Huang

et al. 2001). However, previous work suggests that an alternative gene family, the solute carriers, are

more relevant for desert-adaptation in the cactus mouse (MacManes and Eisen 2014). As a first step

towards fully elucidating the patterns of adaptive evolution to deserts in *P. eremicus*, we characterized

the normal patterns of water intake and electrolyte levels as well as the physiologic response to

experimental dehydration. As such, this study provides critical physiological and biochemical

information about *P. eremicus* and its response to dehydration and is generally useful as researchers

begin to leverage large-scale genome data against classic questions regarding the evolution of adaptive

phenotypes.

Materials and Methods

We used captive *P. eremicus* that were purchased from the University of South Carolina Peromyscus

Genetic Stock Center in 2013. These animals, which are descendant from wild caught animals from a

dry-desert population in Arizona, have been bred in captivity at the University of New Hampshire.

Animals are housed in a large walk-in environmental chamber built to replicate the environmental

conditions in which this population has evolved. Specifically, the animals experience a normal diurnal

pattern of temperature fluctuation, ranging from 90F during the daytime to 75F during the night.

Relative humidity (RH) ranges from 10% during the day to 25% during the night. Animals are housed in

standard lab mouse cages with bedding that has been dehydrated to match desert conditions. They are

fed a standard rodent chow, which has also been dehydrated. Water is provided *ad lib* during certain

phases of experimentation and withheld completely during others. All animal care procedures follow

the guidelines established by the American Society of Mammalogy (Sikes et al. 2011) and have been

approved by the University of New Hampshire Animal Care and Use Committee under protocol number

103092.

All animals included in this study were sexually mature adults. A slight bias for the inclusion of males

exists, as a concurrent study of male reproductive genomics was occurring. Preliminary analyses

conducted suggest that no significant differences in any of the physiological measures, and as a result,

males and females were analyzed as one group. For a subset of animals, water intake was measured,

which was accomplished via the use of customized 15ml conical tubes, wherein water intake was

measured every 24 hours for a minimum of 3 consecutive days (range 3-10 days). Animals selected for

the dehydration trial were weighed on a digital scale, housed without water for three days, then re-

weighed to determine the change in body mass due to dehydration. At the conclusion of water

measurement or after a three-day dehydration animals were sacrificed via isoflurane overdose and

decapitation. Immediately after death, a 120uL sample of trunk blood was obtained for serum

electrolyte measurement. This was accomplished using an Abaxis Vetscan VS2 machine with a critical

care cartridge, which measures the concentration of several electrolytes (Sodium, Chloride, Bicarbonate

ion, Creatinine, and Blood Urea Nitrogen (BUN)) relevant to hydration status and renal function. Lastly,

the kidney, spleen, liver, lung, hypothalamus, testes, vas deferens and epididymis were dissected out

and stored in RNAlater (Ambion Inc.) for future study. All statistical analyses were carried out in the

statistical package, R (R Core Development Team 2011).

Results

We measured the daily water intake for 22 adult cactus mice for between three and 10 consecutive

days. Mean water intake was 0.11 mL per day per gram body weight (median=0.11, SD= 0.05, min=

0.033, max=0.23). We measured levels of serum Sodium, Chloride, Bicarbonate ion, Creatinine, and

Blood Urea Nitrogen (BUN) for 44 adult mice, thereby establishing normal (baseline) values for *P.*

eremicus (Figure 1 and Table 1).

A comparison of mice provided with water *ad libitum* to mice that exposed to experimental water

deprivation for three days revealed that the dehydrated mice lost an average of 23.2% of their body

weight (median=23.9%, SD=5.3%, min=12.3%, max=32.3%). Despite this substantial weight loss,

anecdotally, mice appeared healthy. They were active, eating, and interacting normally. The amount of

weight loss did not depend on daily water intake (p=0.63, R^2= 0.03), though the trend suggests that

animals that drink more water lost more weight). Furthermore, body weight did not strongly influence

the percent loss of body weight (Figure 2; p=0.68, R^2= 0.02).

In addition to a substantial loss in body weight, dehydration was associated with differences in serum

electrolytes (Figure 3; n=19 dehydrated, n=24 hydrated). These changes were subtle, but significant

using a two-sample t-test (p < 0.008 in all cases).

Lastly, the levels of serum electrolytes were tightly correlated with percent body weight loss (Figure 4).

Indeed, the relationship between the level of serum sodium and weight loss was positive and significant,

(ANOVA, F-statistic: 12.85, 11 DF, p= 0.004), as was the relationship between BUN and weight loss

(ANOVA, F-statistic: 9.089, 11 DF, p= 0.012). The relationships between weight loss and chloride and

bicarbonate levels respectively, were positive but not significant.

Discussion

Deserts are amongst the harshest environments on the planet. Indeed, animals living in these areas

must be highly adapted to the unique combination of extreme heat and aridity. Given that our

understanding of the physiology of desert adapted animals is limited largely to studies in renal histology

(Mbassa 1988) and on water intake and output (MacMillen and Lee 1967; Tracy and Walsberg 2001), an

enhanced understanding of serum electrolyte changes due to dehydration is informative. Because many

of the harmful effects of dehydration result from electrolyte abnormalities, characterizing normal values

and the electrolyte response to dehydration represents a critical first step in garnering a deeper

understanding of how desert animals survive despite severe and prolonged dehydration.

In this study, normal (baseline) values for serum Sodium, Chloride, Bicarbonate Ion, Creatinine, and

Blood Urea Nitrogen were established in a captive colony of lab animals housed in desert conditions.

Although these measures may differ in wild animals (see (Calisi and Bentley 2009) for a brief review of

such differences), establishing normal values in captive animals is crucial, though future studies aim to

understand the patterns of electrolyte variation in wild animals. In *P. eremicus*, we define the normal

ranges for each electrolyte as those values falling between the 1^{st} and 3^{rd} quartile. Serum Chloride and

Sodium were significantly higher than in published ranges for other mammals, including humans, a

marsupial (Viggers and Lindenmayer 1996), *Cricetomys* (Nssien et al. 2002), and the porcupine (Moreau

et al. 2003). However, serum chloride and sodium levels in our study were quite comparable to another

wild rodent, *Neotima fuscipes* (Weber et al. 2002), a Mustelid (Thornton et al. 1979), and the Hyrax

(Aroch et al. 2007). Values for BUN are generally higher in this study; unfortunately, a direct comparison

is not possible, as measured values are dependent on the volume of urine produced. Serum Creatinine is

low, largely resulting from the general lack of muscle mass in *P. eremicus* relative to other mammals.

However, because the equipment used to analyze this electrolyte does not effectively capture the lower

end of the biological range, direct comparisons are not made for this metric.

In addition to characterizing baseline electrolytes and their response to experimental dehydration, the

normative value for daily water intake was estimated to be 0.11 mL per day per gram body weight.

Though comparable measures of water consumption are scarce, one study in two arid adapted *Limoys*

(*L. pictus* and *L. irroratus*) housed in non-desert captive settings were estimated to be 0.18 and 0.17 mL

per day per gram body weight respectively (Christian et al. 1978) – a value much greater than in *P.*

eremicus.

Animals that were exposed to experimental dehydration lost a substantial amount of body weight.

Dehydration in humans, resulting in loss of even a fraction of this amount results in cardiovascular

collapse and death. Indeed, even a dehydration-related loss of a few percent of body weight may cause

serious renal impairment or renal failure. That the cactus mouse may lose so much weight as a result of

dehydration and remain active, and apparently healthy, without renal impairment is a testament to

their desert adaptation. Yet, while anecdotally mice appear well, they may be experiencing substantial

cognitive impairment, as is the case with mild-human dehydration (Armstrong et al. 2012). Future

studies in the lab aim to understand the cognitive effects of dehydration in cactus mouse.

In addition to weight loss, dehydrated animals demonstrated biochemical evidence of physiological

stress, in the form of increased Sodium, Chloride, BUN, and Bicarb. There were no significant

relationships between any physiological value and Creatinine, suggesting that dehydration related stress

does not result in renal impairment or damage. Indeed, this is in contrast to humans and other

mammals where acute dehydration of the nature imposed on these animals is universally related to

renal failure and subsequent death. That *P. eremicus* can withstand this level of dehydration is a

testament to the processes involved in adaptation. Studies in progress aim to link patterns of

physiological change of the types described here to patterns of gene expression in both captive and wild

animals, further informing our understanding of renal failure due to dehydration in mammals.

ADROGUÉ, H. J. AND N. E. MADIAS. 2000. Hypernatremia. New England Journal Of Medicine 342:1493–1499.

AL-KAHTANI, M. A., C. ZULETA, E. CAVIEDES-VIDAL AND T. GARLAND. 2004. Kidney mass and relative medullary
thickness of rodents in relation to habitat, body size, and phylogeny. Physiological and Biochemical
Zoology 77:346–365.

ARMSTRONG, L. E. ET AL. 2012. Mild dehydration affects mood in healthy young women. Journal of
Nutrition 142:382–388.

AROCH, I., R. KING AND G. BANETH. 2007. Hematology and serum biochemistry values of trapped, healthy,
free-ranging rock hyraxes (Procavia capensis) and their association with age, sex, and gestational
status. Veterinary clinical pathology / American Society for Veterinary Clinical Pathology 36:40–48.

BARRETT, J. M., W. KRIZ, B. KAISSLING AND C. DE ROUFFIGNAC. 1978. The ultrastructure of the nephrons of the
desert rodent (Psammomys obesus) kidney. II. Thin limbs of Henle of long-looped nephrons. The
American journal of anatomy 151:499–514.

BAUM, N., C. C. DICHOSO AND C. E. CARLTON. 1975. Blood urea nitrogen and serum creatinine. Physiology
and interpretations. Urology 5:583–588.

BEDFORD, N. L. AND H. E. HOEKSTRA. 2015. Peromyscus mice as a model for studying natural variation. eLife
4:e06813.

BEUCHAT, C. A. 1996. Structure and concentrating ability of the mammalian kidney: correlations with
habitat. American Journal Of Physiology 271:R157–79.

BLAUSTEIN, M. P. AND W. J. LEDERER. 1999. Sodium/calcium exchange: its physiological implications.
Physiological reviews 79:763–854.

BRADFORD, D. F. 1974. Water stress of free-living Peromyscus truei. Ecology 55:1407–1414.

BRADLEY, R. D., N. D. DURISH, D. S. ROGERS, J. R. MILLER, M. D. ENGSTROM AND C. W. KILPATRICK. 2007. Toward a
molecular phylogeny for Peromyscus: Evidence from mitochondrial cytochrome-b sequences.
Journal Of Mammalogy 88:1146–1159.

CALISI, R. M. AND G. E. BENTLEY. 2009. Lab and field experiments: Are they the same animal? Hormones
And Behavior 56:1–10.

CASTOE, T. A. ET AL. 2013. The Burmese python genome reveals the molecular basis for extreme
adaptation in snakes. PNAS 110:20645–20650.

CHEUVRONT, S. N., R. W. KENEFICK, S. J. MONTAIN AND M. N. SAWKA. 2010. Mechanisms of aerobic
performance impairment with heat stress and dehydration. Journal of Applied Physiology
109:1989–1995.

CHRISTIAN, D. P., J. O. MATSON AND S. G. ROSENBERG. 1978. Comparative water balance in two species of
Liomys. Comparative Biochemistry and Physiology 61A:589–559.

COSTILL, D. L., R. COTÉ AND W. FINK. 1976. Muscle water and electrolytes following varied levels of

dehydration in man. Journal of Applied Physiology 40:6–11.

E. M. ALTSCHULER, R. B. NAGLE, E. J. BRAUN, S. L. LINDSTEDT AND P. H. KRUTZSCH. 1979. Morphological study of
the desert heteromyid kidney with emphasis on the genus *Perognathus*. The Anatomical Record
194:461–468.

FEIG, P. U. AND D. K. MCCURDY. 1977. The hypertonic state. The New England journal of medicine
297:1444–1454.

HAUSSINGER, D. 1996. The role of cellular hydration in the regulation of cell function. The Biochemical
Journal 313 (Pt 3):697–710.

HAYES, J., C. BIBLE AND J. BOONE. 1998. Repeatability of mammalian physiology: Evaporative water loss and
oxygen consumption of *Dipodomys merriami*. Journal Of Mammalogy 79:475–485.

HEIMEIER, R. A., B. J. DAVIS AND J. A. DONALD. 2002. The effect of water deprivation on the expression of
atrial natriuretic peptide and its receptors in the spinifex hopping mouse, *Notomys alexis*.
Comparative Biochemistry and Physiology Part A: Physiology 132:893–903.

HOEKSTRA, H. ET AL. 2001. Strength and tempo of directional selection in the wild. Proceedings Of The
National Academy Of Sciences Of The United States Of America 98:9157–9160.

HOWELL, A. B. AND I. GERSH. 1935. Conservation of water by the rodent *Dipodomys*. Journal Of
Mammalogy 16:1.

HUANG, Y. ET AL. 2001. Absence of aquaporin-4 water channels from kidneys of the desert rodent
Dipodomys merriami merriami. American Journal Of Physiology-Renal Physiology 280:F794–F802.

HUERTA-SÁNCHEZ, E. ET AL. 2014. Altitude adaptation in Tibetans caused by introgression of Denisovan-like
DNA. Nature 512:194–197.

J. R. MERKT AND C. R. TAYLOR. 1994. "Metabolic switch" for desert survival. Proceedings Of The National
Academy Of Sciences Of The United States Of America 91:12313–12316.

JENTSCH, T. J., V. STEIN, F. WEINREICH AND A. A. ZDEBIK. 2002. Molecular Structure and Physiological Function
of Chloride Channels. Physiological reviews 82:503–568.

JÉQUIER, E. AND F. CONSTANT. 2009. Water as an essential nutrient: the physiological basis of hydration.
European Journal of Clinical Nutrition 64:115–123.

MACMANES, M. D. AND E. A. LACEY. 2012. Is promiscuity associated with enhanced selection on MHC-DQα
in mice (genus *Peromyscus*)? PloS one 7:e37562.

MACMANES, M. D. AND M. B. EISEN. 2014. Characterization of the transcriptome, nucleotide sequence
polymorphism, and natural selection in the desert adapted mouse *Peromyscus eremicus*. PeerJ
2:e642.

MACMILLEN, R. E. AND A. K. LEE. 1967. Australian desert mice: independence of exogenous water. Science
158:383–385.

MARES, M. 1977. Water Economy and Salt Balance in a South-American Desert Rodent, Eligmodontia-
Typus. Comparative Biochemistry And Physiology A-Molecular & Integrative Physiology 56:325–332.

MARRA, N. J., A. ROMERO AND A. DEWOODY. 2014. Natural selection and the genetic basis of
osmoregulation in heteromyid rodents as revealed by RNA-seq. Molecular Ecology 23:2699–2711.

MARRA, N. J., S. H. EO, M. C. HALE, P. M. WASER AND A. DEWOODY. 2012. A priori and a posteriori approaches
for finding genes of evolutionary interest in non-model species: Osmoregulatory genes in the kidney
transcriptome of the desert rodent *Dipodomys spectabilis* (Banner-Tailed Kangaroo Rat).
Comparative Biochemistry and Physiology - Part D: Genomics and Proteomics:1–12.

MBASSA, G. K. 1988. Mammalian renal modifications in dry environments. Veterinary research
communications 12:1–18.

MCKINNEY, T. D. AND M. B. BURG. 1977. Bicarbonate and fluid absorption by renal proximal straight
tubules. Kidney International 12:1–8.

MONTAIN, S., W. LATZKA AND N. SAWKA. 1999. Fluid replacement recommendations for training in hot
weather. Military medicine 164:502–508.

MOREAU, B., J. C. VIÉ, P. COTELLON, I. D. THOISY, A. MOTARD AND C. P. RACCURT. 2003. Hematologic and serum
biochemistry values in two species of free-ranging porcupines (*Coendou prehensilis, Coendou*
melanurus) in french guiana. Journal of zoo and wildlife medicine : official publication of the
American Association of Zoo Veterinarians 34:159–162.

MULLEN, L. M., S. N. VIGNIERI, J. A. GORE AND H. E. HOEKSTRA. 2009. Adaptive basis of geographic variation:
genetic, phenotypic and environmental differences among beach mouse populations. Proceedings
of the Royal Society B 276:3809–3818.

NAGY, K. A. 1988. Seasonal patterns of water and energy balance in desert vertebrates. Journal Of Arid
Environments 14:201–210.

NSSIEN, M., F. O. OLAYEMI, S. K. ONWUKA AND A. OLUSOLA. 2002. Comparison of some plasma biochemcial
parameters in two generations of african giant rat (*Cricetomys gambianus*, waterhouse). African
Journal of Biomedical Research 5.

R CORE DEVELOPMENT TEAM, F. 2011. R: A Language and Environment for Statistical Computing. R
Development Core Team, Vienna, Austria. <http://www.R-project.org>.

R. S. SIKES, W. L. GANNONANIMAL CARE AND USE COMMITTEE OF THE AMERICAN SOCIETY OF MAMMALOGISTS. 2011.
Guidelines of the American Society of Mammalogists for the use of wild mammals in research.
Journal Of Mammalogy 92:235–253.

SCHMIDT-NIELSEN, B. AND K. SCHMIDT-NIELSEN. 1950. Pulmonary Water Loss in Desert Rodents. American
Journal Of Physiology 162:31–36.

SHORTER, K. R., J. P. CROSSLAND, D. WEBB, G. SZALAI, M. R. FELDER AND P. B. VRANA. 2012. *Peromyscus* as a
Mammalian Epigenetic Model. Genetics Research International 2012:1–11.

STEINER, C. C., J. N. WEBER AND H. E. HOEKSTRA. 2007. Adaptive variation in beach mice produced by two
interacting pigmentation genes. PLoS Biology 5:e219.

THORNTON, P. C., P. A. WRIGHT, P. J. SACRA AND T. E. GOODIER. 1979. The ferret, *Mustela putorius furo*, as a
new species in toxicology. Laboratory animals 13:119–124.

TRACY, R. AND G. WALSBERG. 2001. Intraspecific variation in water loss in a desert rodent, *Dipodomys*
merriami. Ecology 82:1130–1137.

VEAL, R. AND W. CAIRE. 2001. Peromyscus eremicus. Mammalian Species 118:1–6.

VIGGERS, K. L. AND D. B. LINDENMAYER. 1996. Variation in Hematological and Serum Biochemical Values of
the Mountain Brushtail Possum, *Trichosurus caninus* Ogilby (Marsupialia: Phalangeridae). Journal Of
Wildlife Diseases 32:142–146.

VIGNIERI, S. N., J. G. LARSON AND H. E. HOEKSTRA. 2010. The selective advantage of crypsis in mice. Evolution
64:2153–2158.

WEBER, D. K., K. DANIELSON, S. WRIGHT AND J. E. FOLEY. 2002. Hematology and serum biochemistry values of
dusky-footed wood rat (*Neotoma fuscipes*). Journal Of Wildlife Diseases 38:576–582.

YU, F. H. AND W. A. CATTERALL. 2003. Overview of the voltage-gated sodium channel family. Genome
Biology 4:207.

345   Table 1

	Normal	Min	Max	Mean
Sodium (mMol/L)	148-158	144	170	153
Chloride (mMol/L)	110-115	105	126	113
BUN (mg/dL)	29-46	22	64	37
Bicarb (mMol/L)	19-25	15	26	22
Creatinine (mg/dL)	>0.2-0.3	>0.2	0.4	0.22

348   **Table 1**. Normal values for serum electrolytes. Normal values are defined as those values falling

349   between the 1st and 3rd quartile. Of note, the Abaxis VS2 electrolyte analyzer does not measure

350   Creatinine below 0.2 mg/dL; therefore, the range for normal Creatinine is truncated at this value.

357   **Figure 1**. Normal values for serum electrolytes. Human normal values (from Medline) are plotted for

358   comparison in dotted red lines. Of note, the Abaxis VS2 electrolyte analyzer does not measure

359   Creatinine below 0.2 mg/dL, and therefore the range for normal Creatinine is truncated at this value.

369   **Figure 2**. Percent body weight loss as a function of initial body weight due to experimental dehydration.

370   No significant trend exists.

382 **Figure 3**. Experimental dehydration resulted in increases in serum sodium, chloride, BUN and

383 bicarbonate ion. Reported p-values are from a two-tailed t-test (n=19 dehydrated: DRY, n=24 hydrated):

384 WET.

393 **Figure 4**. The relationship between serum electrolytes is positive in all cases and significant for Sodium

394 (F-statistic: 12.85, 11 DF, p-value: 0.004283) and BUN (F-statistic: 9.089, 11 DF, p-value: 0.01177)

396